

HERB.

CAB INTERNATIONAL
MYCOLOGICAL INSTITUTE
LIBRARY

IMI / Books / STE ✓



UNITED STATES DEPARTMENT OF AGRICULTURE

OFFICE OF THE SECRETARY

CONTRIBUTION FROM THE FEDERAL HORTICULTURAL BOARD
C. L. MARLATT, CHAIRMAN, IN COLLABORATION WITH
THE BUREAU OF PLANT INDUSTRY
W. A. TAYLOR, CHIEF

[RAM, V, 508]

FOREIGN PLANT DISEASES

A MANUAL OF ECONOMIC PLANT DISEASES WHICH
ARE NEW TO OR NOT WIDELY DISTRIBUTED IN
THE UNITED STATES

PREPARED BY

JOHN A. STEVENSON

Chief Pathological Inspector, Federal Horticultural Board



WASHINGTON
GOVERNMENT PRINTING OFFICE

1926

UNITED STATES DEPARTMENT OF AGRICULTURE

OFFICE OF THE SECRETARY

CONGRESS HAS THE HONORABLE MEMBERS
OF THE SENATE AND HOUSE OF REPRESENTATIVES
AND THE SEVERAL DEPARTMENTS
OF THE GOVERNMENT

1904
FOREIGN PLANT DISEASES

A MANUAL OF ECONOMIC PLANT DISEASES WHICH
ARE NEW TO OR NOT WELL DISTRIBUTED IN
THE UNITED STATES

ADDITIONAL COPIES

OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.

AT

25 CENTS PER COPY



INDEX OF COMMON NAMES

Abaca.....	Musa.	Betal palm.....	Areca.
Aconite.....	Aconitum.	Betony.....	Stachys.
Aconite, winter.....	Eranthis.	Bindweed.....	Convolvulus.
Adder's-tongue.....	Erythronium.	Birch.....	Betula.
Adder's-tongue.....	Ophioglossum.	Bird-of-paradise flower.....	Strelitzia.
African cherry orange.....	Citropsis.	Bishop's cap.....	Mitella.
African lily.....	Agapanthus.	Bitter cress.....	Cardamine.
African millet.....	Eleusine.	Bittersweet.....	Celastrus.
African oil palm.....	Elaeis.	Blackberry.....	Rubus.
Agrimony.....	Agrimonia.	Blackberry lily.....	Belamacanda.
Aguacate.....	Persea.	Black bryony.....	Tamus.
Air potato.....	Dioscorea.	Black locust.....	Robinia.
Alder.....	Alnus.	Black salsify.....	Scorzonera.
Alder, white.....	Clethra.	Bladder fern.....	Cystopteris.
Alfalfa.....	Medicago.	Bladder nut.....	Staphylea.
Alfilaria.....	Erodium.	Bladder senna.....	Colutea.
Alkanet.....	Anchusa.	Bladderwort.....	Utricularia.
Alligator pear.....	Persea.	Blazing star.....	Mentzelia.
Almendro.....	Terminalia.	Bleeding heart.....	Dicentra.
Almond.....	Prunus.	Blister cress.....	Erysimum.
Almond, Spanish.....	Terminalia.	Blood leaf.....	Iresine.
Alumroot.....	Heuchera.	Blood lily.....	Haemanthus.
Alumroot, false.....	Tellima.	Blueberry.....	Vaccinium.
Amaranth.....	Amaranthus.	Blue-curls.....	Trichostema.
Amazon lily.....	Eucharis.	Blue-eyed grass.....	Sisyrinchium.
Amazon vine.....	Stigmaphyllon.	Blue grass.....	Poa.
Anatto.....	Bixa.	Bog asphodel.....	Narthecium.
Angel's-trumpet.....	Datura.	Boneset.....	Eupatorium.
Anise tree.....	Illicium.	Borage.....	Borago.
Apple.....	Malus.	Bottle brush.....	Callistemon.
Apricot.....	Prunus.	Bottle tree.....	Brachychiton.
Arborvitae.....	Thuja.	Bottle tree.....	Sterculia.
Arborvitae, false.....	Thujaopsis.	Box.....	Buxus.
Arrowhead.....	Sagittaria.	Bowstring hemp.....	Sansevieria.
Arrowroot.....	Maranta.	Boxwood.....	Maclura.
Artichoke, globe.....	Cynara.	Brake.....	Pteris.
Artichoke, Jerusalem.....	Helianthus.	Breadfruit.....	Artocarpus.
Ash.....	Fraxinus.	Bridal wreath.....	Spiraea.
Aspen.....	Populus.	Brome grass.....	Bromus.
Asphodel.....	Asphodelus.	Broom.....	Cytisus.
Aster, China.....	Callistephus.	Broom.....	Genista.
Autumn crocus.....	Colchicum.	Broomcorn millet.....	Panicum.
Avens.....	Geum.	Brussels sprouts.....	Brassica.
Avocado.....	Persea.	Bryony.....	Bryonia.
Azalea.....	Rhododendron.	Bryony, black.....	Tamus.
		Bugloss.....	Anchusa.
Baby's breath.....	Gypsophila.	Buckwheat.....	Fagopyrum.
Bachelor's-button.....	Centaurea.	Buckeye.....	Aesculus.
Bael fruit.....	Aegle.	Buckthorn.....	Rhamnus.
Balloon-flower.....	Platycodon.	Bugbane.....	Cimicifuga.
Balloon vine.....	Cardiospermum.	Bugle.....	Ajuga.
Balm.....	Melissa.	Bugle lily.....	Watsonia.
Balsam.....	Impatiens.	Burnet.....	Poterium.
Balsam apple.....	Momordica.	Burnet.....	Sanguisorba.
Bamboo.....	Bambuseae.	Bur, New Zealand.....	Acaena.
Bamboo.....	Phyllostachys.	Burning bush.....	Eutonymus.
Banana.....	Musa.	Bur reed.....	Sparganium.
Banberry.....	Actaea.	Burstwort.....	Herniaria.
Banyan.....	Ficus.	Bush clover.....	Lespedeza.
Barberry.....	Berberis.	Bush honeysuckle.....	Diervilla.
Barberson daisy.....	Gerbera.	Butcher's-broom.....	Ruscus.
Barley.....	Hordeum.	Butcher's-broom, climbing.....	Semele.
Barwood.....	Baphia.	Butterbur.....	Petasites.
Barnyard grass.....	Echinochloa.	Buttercup.....	Ranunculus.
Basil.....	Ocimum.	Butterfly bush.....	Buddleia.
Basswood.....	Tilia.	Butterfly flower.....	Schizanthus.
Bayberry.....	Myrica.	Butterfly pea.....	Clitoria.
Bay, sweet.....	Laurus.	Butterfly weed.....	Asclepias.
Beach grass.....	Ammophila.	Butternut.....	Juglans.
Bean.....	Phaseolus.	Butterwort.....	Pinguicula.
Beard grass.....	Andropogon.	Buttonbush.....	Cephalanthus.
Beardtongue.....	Pentstemon.		
Beard's-breech.....	Acanthus.	Cabbage.....	Brassica.
Beauty berry.....	Callicarpa.	Cacao.....	Theobroma.
Bedstraw.....	Galium.	Cainito.....	Chrysophyllum.
Beech.....	Fagus.	Calla.....	Zantedeschia.
Beefwood.....	Casuarina.	Camomile.....	Anthemis.
Beet.....	Beta.	Camphor tree.....	Cinnamomum.
Belladonna.....	Atropa.	Campion.....	Lychnis.
Bellflower.....	Campanula.	Campion.....	Silene.
Bent grass.....	Agrostis.	Canary grass, reed.....	Phalaris.
Bermuda grass.....	Cynodon.	Candlenut.....	Aleurites.

Candytuft.....	Iberis.	Cotton.....	Gossypium.
Canistel.....	Lucuma.	Cotton thistle.....	Onopordon.
Canterbury bells.....	Campanula.	Cottonweed.....	Diotis.
Cape jasmine.....	Gardenia.	Cottonwood.....	Populus.
Carambola.....	Averrhoa.	Cow parsnip.....	Heracleum.
Caraway.....	Carum.	Cowpea.....	Vigna.
Cardamom.....	Elektaria.	Crab.....	Malus.
Cardoon.....	Cynara.	Cranberry.....	Oxycoocus.
Carnation.....	Dianthus.	Crane's bill.....	Geranium.
Carob.....	Ceratonla.	Crape jasmine.....	Tabernaemontana.
Carrot.....	Daucus.	Crape myrtle.....	Lagerstroemia.
Cashew.....	Anacardium.	Cress, water.....	Nasturtium.
Cassava.....	Manihot.	Crested dogtail.....	Cynosurus.
Cassia bark tree.....	Cinnamomum.	Crocus, autumn.....	Colechicum.
Castor bean.....	Ricinus.	Crosswort.....	Crucianella.
Catbrier.....	Smilax.	Croton.....	Codiaeum.
Catchfly.....	Silene.	Crowberry.....	Empetrum.
Catnip.....	Nepeta.	Crowfoot.....	Ranunculus.
Cat-tail.....	Typha.	Crownbeard.....	Verbesina.
Cauliflower.....	Brassica.	Crown imperial.....	Fritillaria.
Cayenne pepper.....	Capsicum.	Crown vetch.....	Coronilla.
Ceara rubber.....	Manihot.	Cubeb.....	Piper.
Cedar.....	Cedrus.	Cuckoo-flower.....	Cardamine.
Cedar, white.....	Chamaecyparis.	Cucumber.....	Cucumis.
Cedar of Lebanon.....	Cedrus.	Cupang.....	Parkia.
Celandine poppy.....	Chelidonium.	Cupflower.....	Nierembergia.
Celery.....	Apium.	Curant.....	Ribes.
Century plant.....	Agave.	Custard apple.....	Annona.
Ceriman.....	Monstera.	Cypress.....	Cupressus.
Chamomile.....	Anthemis.		
Chaste tree.....	Vitex.	Daffodil.....	Narcissus.
Chayote.....	Sechium.	Daisy, Barberton.....	Gerbera.
Cherimoya.....	Annona.	Daisy, English.....	Bellis.
Cherry.....	Prunus.	Daisy, Michaelmas.....	Aster.
Cherry, ground.....	Physalis.	Daisy, Swan River.....	Brachycome.
Cherry orange.....	Citropsis.	Daisy tree.....	Olearia.
Chestnut.....	Castanea.	Dammur pine.....	Agathis.
Chestnut, water.....	Trapa.	Dame's rocket.....	Hesperia.
Chick-pea.....	Cicer.	Dandelion.....	Taraxacum.
Chickweed, mouse-ear.....	Cerastium.	Darnel.....	Lolium.
Chicory.....	Chicorium.	Dasheen.....	Colocasia.
Chilean crocus.....	Tecophilaea.	Date palm.....	Phoenix.
Chilean lily.....	Alstroemeria.	Dayflower.....	Commelina.
Chile bells.....	Lapageria.	Day lily.....	Hemerocallis.
China aster.....	Callistephus.	Dead nettle.....	Lamium.
Chinaberry.....	Melia.	Desert candle.....	Eremurus.
China fir.....	Cunninghamia.	Devil's-tongue.....	Amorphophallus.
Chinese laurel.....	Antidesma.	Dewberry.....	Rubus.
Chinese tallow tree.....	Sapium.	Dishcloth gourd.....	Luffa.
Chinese tallow tree.....	Stillingia.	Dock.....	Rumex.
Chinquapin.....	Castanopsis.	Dogwood.....	Cornus.
Chive.....	Allium.	Douglas fir.....	Pseudotsuga.
Chocolate tree.....	Theobroma.	Dovewood.....	Alchornea.
Christmas fern.....	Polystichum.	Dracena.....	Dracena.
Christmas rose.....	Helleborus.	Dragonhead.....	Dracoccephalum.
Cigar-box cedar.....	Cedrela.	Dragon-root.....	Arisaema.
Cinnamon.....	Cinnamomum.	Dragon tree.....	Dracena.
Cinquefoil.....	Potentilla.	Dropseed.....	Sporobolus.
Citron.....	Citrus.	Dune grass.....	Elymus.
Citronella grass.....	Cymbopogon.	Durian.....	Durio.
Cliff brake.....	Pellaea.	Dutchman's-pipe.....	Aristolochia.
Climbing hempweed.....	Mikania.		
Clock vine.....	Thunbergia.	East Indian trumpet flower.....	Oroxylum.
Clove tree.....	Eugenia.	Ebony.....	Diospyros.
Clover.....	Trifolium.	Eddoe.....	Colocasia.
Clover, bush.....	Lespedeza.	Edelweiss.....	Leontopodium.
Clover, holy.....	Onobrychis.	Eggplant.....	Solanum.
Clover, sweet.....	Mellilotus.	Elder.....	Sambucus.
Club moss.....	Selaginella.	Elephant's-ear.....	Colocasia.
Cocaine tree.....	Erythroxylon.	Elm.....	Ulmus.
Cockseggs.....	Salpichroa.	Endive.....	Cichorium.
Cocoa plum.....	Chrysobalanus.	English daisy.....	Bellis.
Cockscomb.....	Celosia.	Eryngio.....	Eryngium.
Coconut.....	Cocos.	Eugenia.....	Caryophyllus.
Coffee.....	Coffea.	Evening primrose.....	Oenothera.
Colocynth.....	Citrullus.	Everlasting.....	Helichrysium.
Coltsfoot.....	Tussilago.	Everlasting, pearl.....	Anaphalis.
Columbine.....	Aquilegia.		
Comfrey.....	Symphytum.	Fairy bells.....	Disporum.
Concehead.....	Strobilanthes.	False alumroot.....	Tellinia.
Congo pea.....	Cajanus.	False arborvitae.....	Thuopsis.
Copaiba.....	Copaifera.	False camomile.....	Matricaria.
Copper leaf.....	Acalypha.	False hellebore.....	Veratrum.
Coral blow.....	Russelia.	False indigo.....	Amorpha.
Coral tree.....	Erythrina.	False mallow.....	Malvastrum.
Coriander.....	Coriandrum.	False olive.....	Elaeodendron.
Cork tree.....	Phellodendron.	False Solomon's-seal.....	Smilacina.
Corn, Indian.....	Zea.	False spirea.....	Scorbaria.
Corn cockle.....	Lychnis.	False tamarix.....	Myricaria.
Corn flower.....	Centaurea.	Fan palm.....	Livistona.
		Featherfoil.....	Hottonia.

Feather grass.....	Stipa.	Harebell.....	Campanula.
Fennel.....	Foeniculum.	Hat palm.....	Carludovicia.
Fennel, giant.....	Ferula.	Hawaiian algarroba.....	Prosopis.
Fennugreek.....	Trigonella.	Hawkweed.....	Hieracium.
Fescue.....	Festuca.	Hawthorn.....	Crataegus.
Fig.....	Ficus.	Hazelnut.....	Corylus.
Fig marigold.....	Mesembryanthemum.	Heal-all.....	Brunella.
Figwort.....	Scrophularia.	Heartseed.....	Cardiospermum.
Filaree.....	Erodium.	Heath.....	Erica.
Filbert.....	Corylus.	Heather.....	Calluna.
Filmy fern.....	Hymenophyllum.	Hellebore, false.....	Veratrum.
Finger grass.....	Chloris.	Hellebore.....	Helleborus.
Fir.....	Abies.	Heliotrope.....	Heliotropium.
Fir, Douglas.....	Pseudotsuga.	Hemlock.....	Tsuga.
Fish poison tree.....	Piscidia.	Hempweed, climbing.....	Mikania.
Flamboyan.....	Poinciana.	Hemp.....	Cannabis.
Flat sedge.....	Cyperus.	Henbane.....	Hyoscyamus.
Flax.....	Linum.	Henequen.....	Agave.
Flax lily.....	Phormium.	Heronbill.....	Erodium.
Flax, New Zealand.....	Phormium.	Himalaya honeysuckle.....	Leycesteria.
Fleabane.....	Erigeron.	Hoarhound.....	Marrubium.
Flowering rush.....	Butomus.	Hoary pea.....	Tephrosia.
Foamflower.....	Tiarella.	Holly.....	Ilex.
Forget-me-not.....	Myosotis.	Hollyfern.....	Polystichum.
Four-o'clock.....	Mirabilis.	Hollygrape.....	Berberis.
Foxglove.....	Digitalis.	Hollyhock.....	Althaea.
Frangipani.....	Plumeria.	Holy clover.....	Onobrychis.
Fringe bell.....	Schizocodon.	Honesty.....	Lunaria.
Fringe tree.....	Chionanthus.	Honey locust.....	Gleditsia.
Fritillary.....	Fritillaria.	Honeysuckle.....	Lonicera.
Fumitory.....	Fumaria.	Honeysuckle, bush.....	Diervilla.
		Honeywort.....	Cerinth.
		Hop bush.....	Dodonaea.
		Hop.....	Humulus.
		Hop tree.....	Ptelea.
		Hop hornbeam.....	Ostrya.
		Horn poppy.....	Glaucium.
		Horned rampton.....	Phyteuma.
		Hornbeam.....	Carpinus.
		Horse-radish.....	Radicula.
		Horse-radish tree.....	Moringa.
		Horse-chestnut.....	Aesculus.
		Houseleek.....	Sempervivum.
		Huckleberry.....	Gaylussacia.
		Husk tomato.....	Physalis.
		Hyacinth.....	Hyacinthus.
		Hyacinth bean.....	Dolichos.
		Icaco.....	Chrysobalanus.
		India almond.....	Terminalia.
		India mulberry.....	Morinda.
		Indigo.....	Indigofera.
		Indigo, false.....	Amorpha.
		Iron tree.....	Metrosideros.
		Ironweed.....	Vernonia.
		Ivy.....	Hedera.
		Jaboticaba.....	Myrciaria.
		Jack bean.....	Canavalia.
		Jack fruit.....	Artocarpus.
		Jack-in-the-pulpit.....	Arisaema.
		Jacob's ladder.....	Polemonium.
		Jacob's rod.....	Asphodeline.
		Jasmine.....	Jasminum.
		Jasmine, cape.....	Gardenia.
		Jasmine, crane.....	Tabernaemontana.
		Jasmine, orange.....	Chalcas.
		Jerusalem artichoke.....	Helianthus.
		Jerusalem sage.....	Phlomis.
		Jessamine.....	Jasminum.
		Jet bead.....	Rhodotypos.
		Jicama.....	Pachyrhizus.
		Jimson weed.....	Datura.
		Jobo.....	Spondias.
		Job's-tears.....	Coix.
		Johnson grass.....	Holcus.
		Judas tree.....	Cercis.
		Joint fir.....	Ephedra.
		Jujube.....	Zizyphus.
		June berry.....	Amelanchier.
		Juniper.....	Juniperus.
		Jupiter's-beard.....	Centranthus.
		Jute.....	Corchorus.
		Kafir lily.....	Clivia.
		Kaki.....	Diospyros.
		Kale.....	Brassica.
		Kapok.....	Ceiba.
		Kauri pine.....	Agathis.
		Kidney vetch.....	Anthyllis.
		Knotweed.....	Polygonum.
		Kohlrabi.....	Brassica.
Hackberry.....	Celtis.		
Hair grass.....	Aira.		

Kudzu bean	Pueraria.	Meadow rue	Thalictrum.
Kumquat	Fortunella.	Meadowsweet	Filipendula.
Labrador tea	Ledum.	Medick	Medicago.
Lady bell	Adenophora.	Medlar	Mespilus.
Lady's-slipper	Cypripedium.	Mellic grass	Melica.
Ladies-tresses	Spiranthes.	Melon	Cucumis.
Lancepod	Lonchocarpus.	Mesquite	Prosopis.
Lansa	Lansium.	Mexican poppy	Argemone.
Larch	Larix.	Mexican rubber tree	Castilla.
Larkspur	Delphinium.	Michaelmas daisy	Aster.
Laurel	Laurus.	Mignonette	Reseda.
Lavender	Lavandula.	Milk vetch	Astragalus.
Lavender cotton	Santolina.	Milkweed	Asclepias.
Lead plant	Amorpha.	Millet, broomcorn	Panicum.
Leadwort	Plumbago.	Millet, pearl	Pennisetum.
Leek	Allium.	Millet	Setaria.
Lemon	Citrus.	Millet grass	Milium.
Lemon grass	Cymbopogon.	Mint	Mentha.
Lemon verbena	Lippia.	Mist flower	Eupatorium.
Lentil	Lens.	Mockasin flower	Cypripedium.
Leopard's-bane	Doronicum.	Mock cypress	Kochia.
Leopard flower	Belamacanda.	Mock orange	Philadelphus.
Lettuce	Lactuca.	Monkey comb	Pithecoctenium.
Lettuce, water	Pistia.	Monkey-face tree	Mallotus.
Licorice	Glycyrrhiza.	Monkey puzzle	Araucaria.
Lilac	Syringa.	Monkshood	Aconitum.
Lily	Lilium.	Moonflower	Calonyction.
Lily, African	Agapanthus.	Mooneed	Menispermum.
Lily, Amazon	Eucharis.	Morea	Moraea.
Lily, blackberry	Belamacanda.	Morning-glory	Ipomoea.
Lily, blood	Haemanthus.	Mosquito trap	Cynanchum.
Lily, Chilean	Alstroemeria.	Mountain ash	Sorbus.
Lily, day	Hemerocallis.	Mountain mahogany	Cercocarpus.
Lily, ginger	Hedychium.	Mourning bride	Scabiosa.
Lily of the valley	Convallaria.	Mouse-ear chickweed	Cerastium.
Lily, Peruvian	Alstroemeria.	Mulberry	Morus.
Lily, plaintain	Hosta.	Mullein	Verbascum.
Lily, spider	Hymenocallis.	Muskmelon	Cucumis.
Lily, St. Bernard	Anthericum.	Mustard	Brassica.
Lily, water	Nymphaea.	Myrrh	Myrrhis.
Lima bean	Phaseolus.	Myrtle	Myrtus.
Lime	Citrus.	Naibel	Hesperethusa.
Lime	Tilia.	Napier grass	Pennisetum.
Linden	Tilia.	Nasturtium	Tropaeolum.
Lion's-ear	Leonotis.	Necklace tree	Ormosia.
Litchi	Nephelium.	Nectarine	Prunus.
Lizard's-tail	Saururus.	New Zealand bur	Acaena.
Locust	Robinia.	New Zealand flax	Phormium.
Locust, honey	Gleditsia.	New Zealand spinach	Tetragonia.
Loosestrife	Lysimachia.	Night jasmine	Nyctanthes.
Loosestrife, purple	Lythrum.	Nightshade	Solanum.
Logan	Eriobotrya.	Nightshade, Malabar	Basella.
Lotus	Nelumbo.	Ninebark	Physocarpus.
Love grass	Eragrostis.	Niter bush	Nitraria.
Love-in-a-mist	Nigella.	Norfolk Island pine	Araucaria.
Loveage	Levisticum.	Nutmeg	Myristica.
Lousewort	Pedicularis.	Oak	Quercus.
Lucerne	Medicago.	Oat grass	Arrhenatherum.
Lungwort	Pulmonaria.	Oats	Avena.
Lupine	Lupinus.	Oil palm, African	Elaeis.
Lychee	Nephelium.	Oleander	Nerium.
Madar bush	Calotropis. *	Olive	Olea.
Madeira vine	Boussingaultia.	Onion	Allium.
Mahogany	Swietenia.	Orach	Atriplex.
Mahogany, mountain	Cercocarpus.	Orange	Citrus.
Maidenhair fern	Adiantum.	Orange jasmine	Chalcas.
Maidenhair tree	Ginkgo.	Orchard grass	Dactylis.
Maize	Zea.	Oregon grape	Berberis.
Malabar plum	Caryophyllus.	Osage orange	Maclura.
Malabar nightshade	Basella.	Otaheite gooseberry	Phyllanthus.
Malanga	Xanthosoma.	Oxeye	Bupththalmum.
Mallow	Malva.	Painted cup	Castilleja.
Mallow, false	Malvastrum.	Palm	Palmae.
Mamey colorado	Lucuma.	Palmyra palm	Borassus.
Mandarin orange	Citrus.	Pansy	Viola.
Mandioca	Manihot.	Papaya	Carica.
Mango	Mangifera.	Paper mulberry	Broussonetia.
Mangosteen	Garcinia.	Paper tree	Edgeworthia.
Maple	Acer.	Papyrus	Cyperus.
Marigold	Tagetes.	Pampas grass	Gynerium.
Marjorum	Origanum.	Para grass	Panicum.
Marram grass	Ammophila.	Para rubber	Hevea.
Marshmallow	Althaea.	Paraguay tea	Ilex.
Marsh marigold	Caltha.	Parrotbeak	Clianthus.
Masterwort	Astrantia.	Parsley	Petroselinum.
Maté	Ilex.	Parsnip	Pastinaca.
Matrimony vine	Lycium.	Passion flower	Passiflora.
May apple	Podophyllum.		

Patchouli	Pogostemon.	Rhubarb	Rheum.
Pea	Lathyrus.	Rice	Oryza.
Pea	Pisum.	Rice, wild	Zizania.
Pea, butterfly	Clitoria.	Rock brake	Cryptogramma.
Pea, chick	Cicer.	Rock cress	Arabis.
Pea, scurf	Psoralea.	Rock jasmine	Androsace.
Pea shrub	Caragana.	Rock purslane	Calandrinia.
Pea, sweet	Lathyrus.	Rockrose	Cistus.
Peach	Prunus.	Rosary bean	Rhynchosia.
Peanut	Arachis. *	Rosary pea	Abrus.
Pear	Pyrus.	Rose	Rosa.
Pear, alligator	Persea.	Rose apple	Jambosa.
Pearl everlasting	Anaphallis.	Rose mallow	Hibiscus.
Pearl millet	Pennisetum.	Rosemary	Rosmarinus.
Pearlwort	Sagina.	Rose myrtle	Rhodomyrtus.
Pelican flower	Aristolochia.	Rosewood	Dalbergia.
Pennycress	Thlaspi.	Rouge plant	Rivina.
Pentstemon	Pentstemon.	Royal fern	Osmunda.
Peony	Paeonia.	Royal poinciana	Poinciana.
Pepper	Piper.	Rubber, ceara	Manihot.
Pepper, cayenne	Capsicum.	Rubber, Mexican	Castilla.
Pepper, red	Capsicum.	Rubber, Para	Hevea.
Pepper tree	Schinus.	Rubber vine	Cryptostegia.
Periwinkle	Vinca.	Rue	Ruta.
Persimmon	Diospyros.	Russian olive	Flaeagnus.
Peruvian bark	Cinchona.	Rutabaga	Brassica.
Peruvian lily	Alstroemeria.	Rye	Secale.
Physic nut	Jatropha.	Rye grass	Lolium.
Pickeral weed	Pontederia.	St. Bernard's lily	Anthericum.
Pigeon pea	Cajanus.	St. John's-wort	Hypericum.
Pimpernel	Anagallis.	Safflower	Carthamus.
Pine	Pinus.	Sage	Salvia.
Pineapple	Ananas.	Sago palm	Cycas.
Pink	Dianthus.	Salal	Gaultheria.
Pinkroot	Spigelia.	Salsify, black	Scorzonera.
Pistache	Pistacia.	Saltbush	Atriplex.
Pitcher plant	Nepenthes.	Sandalwood	Santalum.
Plantain	Plantain.	Sand-box tree	Hura.
Plantain lily	Musa.	Sandwort	Arenaria.
Plum	Hosta.	Sapodilla	Achras.
Plume poppy	Prunus.	Sapote	Lucuma.
Plumeria	Bocconia.	Sarsaparilla	Aralla.
Poinsettia	Plumeria.	Saxifrage	Saxifraga.
Pokeberry	Euphorbia.	Saxifrage, golden	Chrysosplenium.
Pokeweed	Phytolacca.	Screw pine	Pandanus.
Polypody	Phytolacca.	Scurf pea	Psoralea.
Pomegranate	Polypodium.	Sea buckthorn	Hippophae.
Poplar	Punica.	Sea grape	Coccoloba.
Poplar, yellow	Populus.	Sea holly	Eryngium.
Poppy	Liriodendron.	Sea onion	Urginea.
Poppy, celandine	Papaver.	Sea pink	Armeria.
Poppy, horn	Chelidonium.	Sea pink	Statice.
Poppy, Mexican	Glaucium.	Self-heal	Brunella.
Potato	Argemone.	Sensitive fern	Onoclea.
Pot marigold	Solanum.	Sensitive plant	Mimosa.
Prayer bead	Calendula.	Serpent root	Scorzonera.
Prickle poppy	Abrus.	Service berry	Amelanchier.
Prickly ash	Argemone.	Sesame	Sesamum.
Prickly pear	Zanthoxylum.	Shad blow	Amelanchier.
Prickly thistle	Opuntia.	Shadbush	Amelanchier.
Primrose	Acanthollmon.	Shallot	Allium.
Primrose willow	Primula.	Shinleaf	Pyrola.
Privet	Jusslaea.	Siberian pea tree	Caragana.
Pumpkin	Ligustrum.	Silk cotton	Bombar.
Purple loosestrife	Ocucurbita.	Silk cotton	Ceiba.
Purslane	Lythrum.	Silk oak	Grevillea.
	Portulaca.	Silk-tassel bush	Garrya.
Quaking grass	Briza.	Silk vine	Periploca.
Quince	Cydonia.	Silver climber	Argyrea.
Quinine	Cinchona.	Silver tree	Leucadendron.
Quinoa	Chenopodium.	Sisal	Agave.
Rabbit-tail grass	Lagurus.	Skullcap	Scutellaria.
Radish	Raphanus.	Sky flower	Duranta.
Raisin tree	Hovenia.	Smartweed	Polygonum.
Ramie	Boehmeria.	Snail seed	Cocculus.
Rape	Brassica.	Snakebeard	Ophiopogon.
Raspberry	Rubus.	Snake gourd	Trichosanthes.
Rattlebox	Crotalaria.	Snake palm	Amorphophallus.
Ravenna grass	Erianthus.	Snapdragon	Antirrhinum.
Redbud	Cercis.	Snapweed	Impatiens.
Red cedar	Juniperus.	Sneezewort	Achillea.
Red pepper	Capsicum.	Snowbell	Styrax.
Redtop	Agrostis.	Snowberry	Symphoricarpos.
Red valerian	Centranthus.	Snowdrop	Galanthus.
Reed	Phragmites.	Snowflake	Leucojum.
Reed canary grass	Phalaris.	Snow-in-summer	Cerastium.
Reed, giant	Arundo.	Soapberry	Sapindus.
Rest-harrow	Ononis.	Soapwort	Saponaria.
Rhodes grass	Chloris.	Solomonseal	Polygonatum.

Solomonseal, false.....	Smilacina.	Toothwort.....	Dentaria.
Sorghum.....	Holcus.	Touch-me-not.....	Impatiens.
Sorrel.....	Rumex.	Towel gourd.....	Luffa.
Soursop.....	Annona.	Traveler's-tree.....	Ravenala.
Sow thistle.....	Sonchus.	Tree fern.....	Alsophila.
Soy bean.....	Soja.	Tree fern.....	Cyathea.
Spanish almond.....	Terminalia.	Treemallow.....	Lavatera.
Spanish moss.....	Tillandsia.	Tree-of-heaven.....	Ailanthus.
Spanish oyster plant.....	Scotymus.	Tree tomato.....	Cyphomandra.
Spanish plum.....	Spondias.	Trout lily.....	Erythronium.
Spatter-dock.....	Nymphaea.	Trumpet bush.....	Tecoma.
Speedwell.....	Veronica.	Trumpet creeper.....	Bignonia.
Spiderflower.....	Cleome.	Tuft root.....	Dieffenbachia.
Spider lily.....	Hymenocallis.	Tulip.....	Tulipa.
Spiderwort.....	Tradescantia.	Tulip tree.....	Liriodendron.
Spiked loosestrife.....	Lythrum.	Tunic flower.....	Tunica.
Spikenard.....	Aralia.	Turneric.....	Curcuma.
Spinach.....	Spinacia.	Turnip.....	Brassica.
Spinach, New Zealand.....	Tetragonia.	Twayblade.....	Liparis.
Spiraea.....	Spiraea.	Twinleaf.....	Jefersonia.
Spleenwort.....	Asplenium.	Twin spur.....	Melaspheerula.
Spring beauty.....	Claytonia.	Udo.....	Aralia.
Spruce.....	Picea.	Unicorn plant.....	Martynia.
Spurge.....	Euphorbia.	Valerian.....	Valeriana.
Spurry.....	Spergula.	Vegetable oyster.....	Tragopogon.
Squash.....	Cucurbita.	Venus-looking-glass.....	Specularia.
Squill.....	Scilla.	Vetch.....	Vicia.
Squirreltail grass.....	Hordeum.	Vine spinach.....	Basella.
Star apple.....	Chrysophyllum.	Violet.....	Viola.
Star flower.....	Trientalis.	Virginia creeper.....	Ampelopsis.
Star glory.....	Quamoclit.	Wahoo.....	Euonymus.
Star gooseberry.....	Phyllanthus.	Wallflower.....	Cheiranthus.
Star grass.....	Aletris.	Walnut.....	Juglans.
Star grass.....	Hypoxis.	Wand flower.....	Sparaxis.
Star-of-Bethlehem.....	Ornithogalum.	Water chestnut.....	Trapa.
Sting lily.....	Blumenbachia.	Water cress.....	Nasturtium.
Stock.....	Matthiola.	Water cress.....	Radiola.
Stone cress.....	Aethionema.	Water lettuce.....	Pistia.
Stonecrop.....	Sedum.	Water lily.....	Nymphaea.
Stone mint.....	Cunila.	Watermelon.....	Citrullus.
Strawberry.....	Fragaria.	Water soldier.....	Stratiotes.
Strawberry tree.....	Arbutus.	Water.....	Acacia.
Sudan grass.....	Holcus.	Wax plant.....	Hoya.
Sugar apple.....	Annona.	Weaversbroom.....	Spartium.
Sugar beet.....	Beta.	Wheat.....	Tritium.
Sugar cane.....	Saccharum.	Wheat grass.....	Agropyron.
Sugar palm.....	Arenga.	Weicelia.....	Diervilla.
Sunflower.....	Helianthus.	White slider.....	Clethra.
Sumac.....	Rhus.	White cedar.....	Chamaecyparis.
Summer cypress.....	Kochia.	White sapote.....	Castanroa.
Sunn hemp.....	Crotolaria.	Whitlow grass.....	Draba.
Sun rose.....	Helianthemum.	Wild calla.....	Calla.
Surinam cherry.....	Eugenia.	Wild ginger.....	Asarum.
Swan-River daisy.....	Brachycome.	Wild rice.....	Zizania.
Sweet alyssum.....	Alyssum.	Willow.....	Elymus.
Sweet bay.....	Laurus.	Willow weed.....	Salix.
Sweet clover.....	Melilotus.	Windflower.....	Epilobium.
Sweet flag.....	Acorus.	Wing nut.....	Anemone.
Sweetleaf.....	Symplocos.	Winter creeper.....	Pterocarya.
Sweet pea.....	Lathyrus.	Wisteria.....	Euonymus.
Sweet potato.....	Ipomoea.	Winter aconite.....	Eranthis.
Sweet shrub.....	Calycaanthus.	Winter cress.....	Barbarea.
Sword bean.....	Canavalia.	Witch-hazel.....	Hamamelis.
Sword fern.....	Nephrolepis.	Wintergr n.....	Gaultheria.
Sycamore.....	Platanus.	Woodbetony.....	Pedicularis.
Syringa.....	Philadelphus.	Wood fern.....	Aspidium.
Tail grape.....	Artabotrys.	Wood fern.....	Dryopteris.
Tallow tree, Chinese.....	Sapium.	Wood fern.....	Phegopteris.
Tamarack.....	Larix.	Wood-oil tree.....	Aleurites.
Tamarind.....	Tamarindus.	Woodruff.....	Asperula.
Tangerine.....	Citrus.	Wood rush.....	Luzula.
Tansy.....	Tanacetum.	Wormwood.....	Artemisia.
Tea.....	Thea.	Yam.....	Dioscorea.
Teak.....	Tectona.	Yam bean.....	Pachyrhizus.
Teasel.....	Dipsacus.	Yautia.....	Xanthosoma.
Ted.....	Eragrostis.	Yautia.....	Colocasia.
Teosinte.....	Euchisena.	Yarrow.....	Achillea.
Thatch palm.....	Thrinax.	Yellow mombin.....	Spondias.
Thistle, sow.....	Sonchus.	Yew.....	Taxus.
Thistle, globe.....	Echinops.	Yellow poplar.....	Liriodendron.
Thoroughwort.....	Eupatorium.	Yellow rocket.....	Barbarea.
Thrift.....	Statice.	Yellowwood.....	Cladrastis.
Thyme.....	Thymus.	Zephyr lily.....	Zephyranthes.
Tick clover.....	Meibomia.		
Timothy.....	Phleum.		
Toadflax.....	Linaria.		
Tobacco.....	Nicotiana.		
Tomato.....	Lycopersicum.		
Tomato, husk.....	Physalis.		

FOREIGN PLANT DISEASES

A MANUAL OF ECONOMIC PLANT DISEASES WHICH ARE NEW TO OR NOT WIDELY DISTRIBUTED IN THE UNITED STATES

This publication has been prepared for the purpose of supplying such available information as is required by the officers of the Federal Horticultural Board and others in the enforcement of plant quarantines and in the safeguarding of this country against foreign plant diseases. It is hoped that it will prove of value not only to inspectors in actually preventing entry of specific diseases, but that it will aid in a general way in pointing out the magnitude of the quarantine problem by presenting the very great number of plant diseases reported from foreign countries which are not known in the United States.

The manual is based on the data contained in the host index of foreign plant diseases maintained in the Office of Foreign Plant Quarantines of the Federal Horticultural Board. This file, which has now reached a total of several hundred thousand entries, is planned to cover the plant pathological literature of the world from the host standpoint, and, while by no means complete as yet, is of such extent as to give an excellent basis for the work in hand. This host index was started in 1914 by R. Kent Beattie, pathologist in charge, Office of Foreign Plant Quarantines, and has been continued by him and his assistants since that time.

As a matter of convenience, an alphabetical arrangement according to their scientific names has been followed both for the hosts and the causal agents of the diseases thereunder.

The selection of host genera has in large part been based on the Standard Cyclopedia of Horticulture, those genera being included which contain one or more species of economic importance. A number of genera are included, for instance *Chenopodium* and *Amaranthus*, which consist mostly of weeds, but since some members of such genera are of value, all the foreign diseases of the entire genus are given in each case, since these diseases are in large part capable of attacking both the economic and noneconomic species.

Common names have been taken in large part from Standardized Plant Names and include only those in common and general use or those most likely to be met with by inspectors. Common names of generic significance are given in caps and small caps immediately following the generic name. Other common names of specific value only then follow in ordinary type. Common names of only local significance are omitted. The characterization of each host genus is drawn in large part from the cyclopedia already referred to.

Throughout the course of the work an attempt has been made to select the included material on a uniform basis. Species of parasitic fungi or bacteria known to occur more or less widely in the United

States have been excluded. Species not so excluded and belonging to genera the members of which are parasitic for the most part, form the basis of the present work, together with species of other more or less saprophytic genera where specific mention of parasitism has been suggested. Some parasites are included which are reported from limited localities in the United States, it being advisable to guard against their further importation. A few fungi causing diseases of major importance and of comparatively recent introduction, such as *Endothia parasitica* which causes the chestnut blight, and *Cronartium ribicola*, which causes the white pine blister rust, are also listed. These still have interest or are of importance to inspectors. The records of the plant disease survey of the Bureau of Plant Industry have been helpful in determining established American plant diseases.

Under each fungus or bacterium listed there is given a brief summary of the symptoms of the disease produced by it, together with its specific hosts and the countries from which it has been reported as occurring. Diseases due to nematodes and to unknown causes supposed to be of an infectious nature are included. Limitations of space have made it impossible to cite the literature from which the information has been drawn. The citations are in the above-mentioned host index and are available for reference to anyone interested. For like reasons it has not been feasible to include illustrations.

It has not been possible to attempt a critical examination of the fungus species listed, although a few suggestions as to synonymy are made. It is realized that probably many fungi have been included which would prove harmless if introduced into America and that similarly many have been omitted which should properly find a place. The literature is so fragmentary as to make uniform treatment exceedingly difficult, if not impossible. The nomenclature used has been for the most part that of current usage. Certain monographic works, including Salmon on Erysiphaceae, Theissen and Sydow on Dothidiaceae, Sydow on Uredineae, Gauman on Peronospora, and others have been followed.

Because of the alphabetical arrangement of the material, indices other than the common-name index have been considered unnecessary.

The manuscript has been reviewed by the offices of Plant Disease Survey, Laboratory of Plant Pathology, Pathological Collections, Fruit Disease Investigations, Forest Pathology, Sugar Plant Investigations, Cotton, Truck, and Forage Crop Disease Investigations, and Cereal Investigations of the Bureau of Plant Industry. The nomenclature of the host genera has been checked by George B. Sudworth, dendrologist of the Forest Service and by F. V. Colville and S. F. Blake, of the Office of Economic and Systematic Botany. It has been necessary for the most part to use the host names given in the literature. Obviously it was not possible to handle the subject critically. Where the generic plant name thus used differs from that now recognized by the Department of Agriculture, the department name, together with the abbreviation Ag., follows it; for example, ABERIA. DOYYALIS. Ag.

- ABERIA. DOVYALIS** Ag. Small ornamental trees or shrubs of Africa and Ceylon.
Asterostromella aberiae Petch. On leaves of *A. gardneri* in Ceylon.
Phyllachora aberiae P. Henn. Tar spot on leaves of *A. affra* in the Union of South Africa.
- ABIES.** Fir. Ornamental coniferous evergreens and timber trees.
Bardleyella deformans (Bardl.) Diet. See *Picea*.
Camarsporium abietis Wilson. Attacks twigs and small branches of *A. lowiana* in Scotland.
Cucurbitaria pithyophila Fr. See *Pinus*.
Dasycephala abieticola Henn. and Shifr. Produces a canker on trunks and limbs of *A. homolepis* (*A. trachyphylla*) and *A. firma* in Japan.
Dasycephala willkommii Hart. See *Larix*.
Diplodia abiegna Maubl. On living needles of *A. concolor* in France.
Fusicoccum abietinum Hart. Prill and Delacr. *Dactylothea pitea* Sacc. This fungus produces cankers on the branches, which die as a result of girdling. Black stromata are produced on the cankered areas. On *A. firma*, *A. procurrens*, and *Abies* sp. in France and Germany. Reported from Ohio on *A. veitchii*, an introduced ornamental type.
Lophodermium abietis Rostr. See *Picea*.
Melampsora abietis-capreae Tub. See *Salix*.
Melampsorella symphyti Bub. See *Symphytum*.
Myosphaerella abietis (Rostr.) Lind. On leaves and shoots of *A. alba*, *A. arizonica*, *A. cephalonica*, *A. nordmanniana*, and *A. pinsapo* in Denmark.
Oospora abietum Oudem. A single row of minute greenish-gray tufts on each side of the median line and on both surfaces of needles of *A. nordmanniana*, *A. pinsapo*, *Picea abies*, and *Pseudotsuga taxifolia* in Holland. Defoliation may result.
Phomopsis pseudotsugae Wilson. See *Pseudotsuga*.
Physalospora abietina Prill and Delacr. On *A. alba* (*A. pectinata*) and *Picea abies* in France.
Phytophthora fagi R. Htg. See *Fagus*.
Behmiiopsis bohemica Bub. and Kab. (*Phoma bohemica* Bub. and Kab.) The leaves of the current year are attacked soon after expanding and rapidly shriveled. They turn red at first, then become darker, and remain hanging about a year. The fungus spreads into the shoots, which die back and with some host species twist and curve. The disease is so severe that planting of silver fir (*A. procurrens*) has been discontinued to a large extent in Scotland. Other hosts are *A. alba*, *A. cephalonica*, *A. nobilis*, *A. pinetorum*, and *A. pinsapo* in Scotland, Bohemia, and probably other European countries.
Rhizothrium abietis Naumoff. On needles of *A. sibirica* in Russia.
Sclerotopsis piceana (Karst.) Diet. See *Picea*.
- ABIES. ROSARY PEA.** Leguminous vine common throughout the Tropics, sometimes called "prayer tree."
Phyllachora perforans (Rehm.) Seec. and Syd. Tar spot on leaves of *A. procurrens* in Porto Rico.
Ravenelia ornata Syd. A rust producing yellow-brown to black pustules on lower surface of leaves of *A. procurrens* and *A. pauciflorus* in India, Ceylon, and the Philippines.
Uredo karimbachii P. Henn. This rust produces yellow-brown pustules on the lower (rarely on the upper) leaf surfaces of *A. procurrens* in New Guinea.
- ABUTILON.** See *Agapanthus*.
- ABUTILON.** Malvaceous shrubs or herbs, some species cultivated.
Ascochyta abutilonis Héb. Causing a leaf spot of *A. theophrasti* (*A. aricennae*) in Hungary.
Gloeosporium leptostromoides Bub. Small dark-brown pustules on stems of *Abutilon* sp. in Austria.
Meliola mollieriana Wint. Black sooty mold on leaves of *A. striatum* and *Abutilon* sp. in Brazil and Argentina.
Phylloticta abutilonis P. Henn. Circular, gray leaf spots with brown margins on *Abutilon* sp. in Brazil.
Puccinia abutili B. and Br. This rust appears as powdery brown pustules on both leaf surfaces of *A. procurrens*, *A. indicum*, *A. muticum*, and *A. sonnerati* in Ceylon, Abyssinia, Somali coast, central and south Africa.
- ACACIA.** Tropical and subtropical ornamental leguminous trees and shrubs. Several species of economic importance for gums, resins, tanning extracts, and gum arabic. A number of species are known as "wattle."
Aecidium esculente Bardl. This rust distorts and renders succulent the young shoots of *A. eburnea* in India.
Aecidium immersum P. Henn. The aecia of this rust occur on flesh-colored galls on young twigs of *A. abyssinica* in Eritrea.
Aecidium schweinfurthii P. Henn. A rust causing galls on twigs of *A. seyal* in Somali.
Aecidium torquens McAlp. Leaf rust on *A. abyssinica* in Abyssinia.
Ascochyta borjomi Bond. Circular, white leaf spots with dark brown margins on *Acacia* sp. in Russia.
Catacauma acaciae Theiss. and Syd. Forms numerous black, circular, shiny spots on upper surface of leaves of *A. leucophloea* in India.
Cercospora alemquerensis Speg. Leaf spot on *A. alemquerensis* in Brazil.
Cercospora theae Petch. See *Thea*.
Corticium salmonicolor B. and Br. See *Citrus*.
Haplophragmium acaciae Bacc. Leaf rust on *A. nubicola* in Abyssinia.
Haplophragmium ponderosum Syd. and Butl. The brown telial sori of this rust occur on woody parts 1 to 2 centimeters in diameter on twigs of *A. leucophloea* in India.
Marsonia acaciae Cke. and Mass. Produces irregular spots on the phyllodes, which are pale-brown to whitish, brown-margined, 0.5 to 1 centimeter in diameter and often confluent. On *Acacia* sp. in Australia.
Phyllachora acaciae P. Henn. Small, circular, shiny black spots on both sides of leaves of *A. amenzica*, *A. farnesiana*, and *A. tortuosa* in Ecuador, Mexico, and St. Thomas.
Phyllachora indica Theiss. and Syd. A similar tar-spot disease on leaves of *A. penninervis* in India, Ceylon, and adjoining islands.
Phyllachora parvicapsa Cke. Theiss. and Syd. This fungus produces black convex stromata on leaves of *Acacia* sp. in Australia.
Phylloticta acaciicola P. Henn. Amphigenous, dark brown spots, becoming lighter and drying out on leaves of *A. ramosissima* in Germany.
Phylloticta pedicellata Speg. Leaf spot on *A. pedicellata* in Brazil.
Phylloticta phyllodiorum Sacc. On phyllodes of *Acacia* sp. in Australia.
Physalospora phyllodii Cke. and Mass. On phyllodes of *A. ionifolia* and *A. suareolens* in Australia.
Ravenelia acaciae-melliferae Bacc. Rust on *A. mellifera* in Abyssinia.
Ravenelia acaciae-micranthae Diet. A rust producing dark-brown pustules on both surfaces of leaves of *A. micrantha* in Mexico.

- ACANTHOLIMON.** PRICKLY THRIFT. Hardy evergreen perennials.
Uromyces acantholimonis Syd. Brown to black rust pustules on leaves of *A. antheum* in Persia and Turkestan.
- ACANTHOPANAX.** Hardy shrubs or trees with ornamental foliage.
Aecidium acanthopanacis Diet. Rust on leaves of *A. spinosum* in Japan.
Mycosphaerella acanthopanacis Syd. Leaf spot of *A. spinosum* in Japan.
Phyllosticta acanthopanacis Syd. Leaf spot of *A. spinosum* in Japan.
Triphragmium thwaitesii B. and Br. Brown, powdery, rusty, ashy, or yellowish on yellow to brown spots, 1 to 2 centimeters in diameter on both leaf surfaces of *A. thwaitesii*, *A. macrophyllum*, *A. thwaitesii* sp., *Hedera stellata*, and *Heptapleurum emarginatum* in Japan, Java and Japan.
- ACANTHOPHOENIX.** See *Palmae*.
- ACANTHUS.** BEARS-BREECH. Hardy herbaceous perennials.
Aecidium acanthi Lott.-Sim. Rust on leaves of *Acanthus* sp. in east Africa.
Cercospora acanthi Pass. Brown spots on both leaf surfaces of *A. mollis*, *A. super*, *A. spinosus*, and *A. spinulosus* in France, Italy, and Algeria.
Cercospora compacta Trév. Brown leaf spots on *A. spinosus* in Italy.
Septoria acanthi Theism. Circular, white spots, occasionally concentric, on the upper sides of leaves of *A. mollis* in Algeria and Italy.
Septoria acanthina Sacc. and Magn. Subcircular, brown leaf spots on *A. mollis* in Italy.
- ACER.** MAPLE. Shade and timber trees.
Ascochyta acericola Massal. On *A. campestre* in Italy.
Ascochyta pallida Kst. and Bub. Circular to irregular brown spots on both sides of leaves of *A. platanoides* in Bohemia.
Ascochyta velata Kab. and Bub. On leaves of *A. platanoides* in Bohemia.
Cercospora acerina Hart. Attacks the cotyledons and stems of seedlings, often killing the plants. On *A. pseudoplatanus* in France and Germany.
Cercospora acerina Hart, Arn. Brown spots on cotyledons of *Acer* sp. in Europe, causing death of seedlings.
Cylindrosporium acerellum Sacc. Died. *Septoria acerella* Sacc. Minute, angular, white spots on upper surfaces of leaves of *A. campestre* in France and Denmark.
Cylindrosporium platanoidis (Kühn.) Died. *Septoria apfelii* Kühn. *Septoria acerina* Sacc. and Kueg. *Septoria acerina* Sacc. Small brown leaf spots on *A. campestre* and *A. platanoides* in Europe.
Cylindrosporium pseudoplatani R. and D. Died. *Septoria acerina* Theism. *S. pseudoplatani* R. and D. Subcircular to irregular reddish or brown spots without definite margins on leaves and cotyledons of *A. campestre*, *A. platanoides*, and *A. pseudoplatanus* in Italy, France, Denmark, and Germany.
Dermatea acerina Karst. On *Acer* sp. in Europe.
Diaporthelongiostis Sacc. *Gloeosporium acerinum* West. On leaves of *A. platanoides*, *A. pseudoplatanus*, and *Acer* sp. in Great Britain, Belgium, Denmark, and Austria.
Discomycopsis rhytismoides L. Mass. Said to produce black spots on the leaves of *A. pseudoplatanus* in Germany. Probably *Aspergillus*.
Discoa artoceas Todei Fr. On leaves of *A. campestre* in Italy.
Eoasus acericola Mass. and f. *pseudoplatani* Mass. Occurs on leaves of *A. campestre* and *A. pseudoplatanus* in Italy.
Eoasus acerinus Edsson. Forms roughened, discolored areas on leaves and branches of *A. platanoides* in Sweden.
Eoasus confusus Jacz. On leaves and branches of *A. campestre* in Russia.
Eoasus nikkoensis (Kus.) Sacc. and Trév. Forms grayish scurfs on paler or violet spots on under leaf surfaces of *A. purpurascens* in Japan.
Gnomonia inclinata (Desm.) Auerw. On *A. campestre* in Italy.
Guignardia acerifera (Cke.) Lind. *Laetia acerifera* (Cke.) Sacc. On *A. campestre* in Great Britain.
Guignardia rhytismophila Rehm. On upper surface of leaves of *A. pseudoplatanus* in Germany.
Laetia pseudoplatani Pass. On leaves of *A. pseudoplatanus* in France.
Leptothyrium acerinum Kie. Cke. Small, circular, black fruiting bodies on leaves of *A. confertum*, *A. campestre*, *A. opulifolium*, *A. palmatum*, *A. parvum*, and *A. platanoides* in middle and western Europe, the Balkan Peninsula, Great Britain, and Japan.
Marsonia acerina West. P. Magn. On *A. pseudoplatanus* in Europe.
Marsonia decolorans Kst. and Bub. On leaves, causing them to die back from the tips. On *A. negundo* in Bohemia and Denmark.
Marsonia truncatula Sacc. *Marsonia truncatula* (Sacc.) Magn. Obtrusive leaf spots on *A. campestre*, *A. monspessulanum*, *A. negundo*, and *A. tataricum* in Europe.
Mycosphaerella latebrosa Cke. Sacc. *Phyllosticta platanensis* Sacc. Leaf spot on *A. campestre*, *A. negundo*, *A. opulifolium*, and *A. platanoides* in Europe. Fossily occurs in a limited extent in the United States.
Mycosphaerella maculiformis Pers. Schroet. On leaves of *A. campestre*, *A. opulifolium*, *A. platanoides*, *A. pseudoplatanus*, and *Monia alba* in France, Switzerland, Denmark, Estonia, and Germany.
Myxosporium devastans Rostk. See *Betula*.
Parodiella aceris Kze. On leaves of *A. laetivum* in Java. The systematic position of this fungus is uncertain.
Phyllosticta aceris Sacc. Subcircular, pale-yellow spots on leaves of *A. campestre*, *A. monspessulanum*, *A. negundo*, and *A. platanoides* in France, Denmark, Italy, and Russia. Reported from Virginia and Minnesota.
Phyllosticta apatela Arn. and var. *perniciosa* Kst. and Bub. Large indefinite, pale-brown to gray spots on leaves of *A. platanoides* and *A. pseudoplatanus* in Bohemia and Germany.
Phyllosticta campestris Pass. Small, red to red-brown leaf spots on *A. campestre* and *A. pseudoplatanus* in France and Austria.
Phyllosticta negundicola Sacc. Small, pale-yellow to white spots on leaves of *A. negundo* in France.
Phyllosticta obtusula Sacc. and Br. On leaves of *A. campestre* in Germany.
Phyllosticta platanoidis Sacc. On leaves of *A. campestre* and *A. platanoides* in Europe.
Phyllosticta pseudoplatani Sacc. and var. *fallax* Sacc. and Rehm. On leaves of *A. campestre*, *A. pseudoplatanus* in Italy, France, Portugal, and Germany.
Phyllosticta tambowiensis Bub. and Sere. On leaves of *A. platanoides* in Russia.
Phytophthora fagi Hartig. See *Fagus*.
Phytophthora syringae Alet. See *Syringa*.
Schizothyrium annuliforme Syd. Large, irregular, discolored areas on leaves of *A. nitens* in India.

ACER.—Continued.

- Sclerotium scutellatum** A. S. Orbicular, dark-brown sclerotia, on leaves of *Acer*, *Frazinus*, and *Populus* in northern Europe.
- Septogloeum hartigianum** Sacc. The fungus causes the young twigs to dry up, preventing the buds from swelling. The mycelium spreads in the twigs during the summer, the effects not appearing until the following spring. This maple blight attacks *A. campestre* and *A. platanoides* in Italy, Russia, and Germany.
- Septoria schirajewskii** Bub. and Sere. On leaves of *A. platanoides* in Russia.
- Syngonium insigne** Penz. and Sacc. Forms black, cushion-shaped, carbonaceous fruiting bodies on leaves of *A. laurinum* in Java.
- Taphrina polyspora** (Sorok.) Johans. This fungus produces irregular, shiny, reddish-brown to black patches on leaves of *A. pseudoplatanus* and *A. tataricum* in Bulgaria, Serbia, Hungary, Scandinavia, Russia, and Germany.
- Uncinula aceris** (DC.) Sacc. and var. *tulasnei* Fekl. Powdery mildew on leaves of *A. campestre*, *A. monspessulanum*, *A. pictum*, *A. platanoides*, *A. pseudoplatanus*, *A. rubrum*, *A. spicatum*, and *A. tataricum* in Europe, Transcaucasia, and Japan.

ACHILLEA. YARROW. SNEEZEWORT. Herbaceous perennials.

- Ascochyta millefolii** Oud. On stems of *A. millefolium* in Holland.
- Cercospora achilleae** Jasp. Circular to oblong brown spots on leaves of *A. macrophylla* and *A. microphylla* in Switzerland.
- Entyloma achilleae** P. Magn. A smut producing spots on leaves of *A. millefolium* in Switzerland, Sweden, Denmark, and Germany.
- Peronospora radii** De By. See *Chrysanthemum*.
- Laestadia ptarmicae** Karst. and Starb. On *A. ptarmica* in Finland.
- Puccinia achilleae** Cke. Rust on stems and leaves of *A. albicaulis* in Kurdistan.
- Puccinia ptarmicae** Karst. Yellow to yellow-brown rust pustules on leaves of *A. cartilaginea* and *A. ptarmica* in Belgium, Denmark, Sweden, Finland, Russia, and Germany.
- Puccinia santolinae** P. Magn. Rust on stems and leaves of *A. santolina* in Egypt.
- Ramularia ptarmicae** Lind. Small, subcircular, dark-brown leaf spots on *A. ptarmica* in Germany.
- Schizothyrium ptarmicae** Desm. This fungus appears as small, flat, black fruiting bodies on the leaves of *A. ptarmica*, bringing about a stunting of infected plants in Great Britain, France, Belgium, Denmark, Finland, Siberia, and Germany.
- Septoria ptarmicae** Passer. Causes a leaf spot of *A. ptarmica* in Italy.
- Synchytrium globosum** Schroet. See *Potentilla*.
- Tuberculina microstigma** Sacc. On leaves of *A. millefolium* in Germany.
- Tylenchus millefolii** Loew. This nematode produces numerous small galls and hypertrophied areas on stems of *A. millefolium* and *A. tanacetifolium* in Germany.

ACHRAS. Sapodilla. Fruit and ornamental tree. Furnishes the chicle of commerce.

- Phyllosticta sapotae** Sacc. Deep-brown, circular to irregular spots appearing on both leaf surfaces with ashen-brown margins. On *A. zapota* in Ceylon and the Bahamas.
- Uredo sapotae** Arth. and Johnston. Rust on leaves of *A. zapota* in Cuba and the Bahamas.

ACIDANTHERA. See *Tritonia*.**ACNISTUS**. Ornamental tropical shrubs.

- Puccinia acnisti** Arth. Rust on leaves of *A. aggregatus* in Trinidad and Costa Rica.

ACONITUM. MONKSHOOD. *Aconite*. Hardy perennial herbs.

- Coleosporium aconiti** Thuem. Rust on leaves of *A. barbatum* in Siberia.
- Fuscladium aconiti** Bres. On leaves of *A. clusium* in Hungary.
- Phyllosticta aconitifolia** Petr. Large, usually irregular, dark olive-green or brownish-green spots on leaves and sometimes on stems of *A. moldavicum* in Bohemia.
- Puccinia lycoctoni** Fekl. Yellow to yellow-brown rust pustules on leaves of *A. lycoctonum* in France, Italy, Switzerland, Norway, Austria, Hungary, India, and Germany.
- Ramularia monticola** Speg. On leaves of *A. lycoctonum*, *A. moldavicum*, and *A. napellus* in Italy and Austria.
- Septoria aconiti** Bacc. Circular, dark-brown leaf spots with definite margins on *Aconitum* sp. in China.
- Septoria fuscomaculans** Kab. and Bub. Oblong to irregular, often confluent, dark-brown spots on lower leaf surfaces of *A. paniculatum* in Tyrol.
- Septoria lycoctoni** Speg. Including var. *sibirica* Sacc. and var. *macrospora* C. Mass. Ashen, irregular spots, with dark-brown margins, on leaves of *A. lycoctonum* and *Aconitum* sp. in Italy, Russia, and Siberia.
- Septoria napelli** Speg. On leaves of *A. napellus* in Italy and Yugoslavia.

ACORUS. SWEET FLAG. Aquatic plants.

- Ascochyta acori** Oud. On leaves of *A. calamus* in Holland.
- Eriospora leucostoma** B. and Br. On leaves of *A. gramineus* in Europe.
- Leptosphaeria microspica-calami** Karst. On leaves of *A. calamus* in Italy.
- Phyllosticta acorella** Sacc. and Penz. Linear spots on both surfaces of leaves of *A. calamus* in France.
- Phyllosticta acori** Oud. Elongate black spots on leaves of *A. calamus* in Holland.
- Physoderma calami** Krieg. Subcircular to oblong brown spots on leaves of *A. calamus* in Germany.
- Physoderma tenue** Nowak. See *Nymphaea*.
- Uredo acori** Racib. Brown-rust pustules on leaves of *A. calamus* and *A. terrestris* in Java, India, and the Philippines.

ACRISTA. See *Palmae*.**ACROCOMIA**. See *Palmae*.**ACTAEA**. BANEERRY. Herbaceous perennials.

- Actinonema actaeae** Allesch. Irregular, brown to black spots on leaves of *A. spicata* in Germany.
- Coleosporium actaeae** Karst. Yellow rust pustules on leaves of *A. erythrocarpa* and *A. rubra* in Siberia.
- Didymaseus kitmanoffi** Sacc. On leaves of *A. spicata* in Siberia.
- Marsonia actaeae** Bres. Leaf spot of *A. spicata* in Denmark and Germany.

ACTINIDIA. Ornamental woody climbers.

- Aecidium actinidiae** Syd. Rust on leaves of *A. arguta* in Japan.
- Gloeosporium** sp. Anthracnose on leaves of *A. arguta* in Japan.

ACTINOSTEMMA. Ornamental vines.

- Marsonia actinostemmae** F. Tassi. Circular, dirty-brown, zonate spots on both surfaces of leaves of *A. paniculatum* in Italy.

ADENOCARPUS. Ornamental woody shrubs.

- Aecidium bubakii** Gz. Frag. On leaves of *A. intermedius* in Spain.

- ADENOPHORA.** LADYBELL. Hardy herbaceous perennials.
Aecidium adenophorae Jacz. Yellow rust pustules on yellow or brown leaf spots on *A. latifolia* and *A. verticillata* in Japan and Russia.
Aecidium adenophorae-verticillatae Syd. Leaf rust on *A. verticillata* in Japan.
Septoria adenophorae Thuem. Gray spots on leaves of *A. tricuspidata* in Siberia.
- ADENOSTYLES.** Herbaceous perennials.
Ascochyta adenostylis Kab. and Bub. Large, irregular, dirty-gray to dark-brown spots on leaves of *A. albifrons* in Tyrol.
Cercospora septorioides Sacc. Angular gray spots on leaves of *A. albifrons* and *A. alliariae* in Switzerland and Austria.
Colosporium cacaliae Oth. See *Cacalia*.
Phyllosticta adenostylis Allesch. Subcircular, often confluent, dark-brown spots on leaves of *A. alpina* in Germany.
Ramularia filaris Fres. Dull yellow, irregular leaf spots on *A. albifrons*, *A. alliariae*, *A. alpina*, and *Doronicum austriacum* in Italy, Bohemia, and Austria.
Uromyces cacaliae (DC.) Ung. Yellow-brown rust pustules on *A. albifrons*, *A. alliariae*, *A. alpini*, *A. auriculata*, *A. glabra*, *A. kerneri*, and *Cacalia* spp. in Japan, Rumania, Italy, Switzerland, Austria, Hungary, and Germany.
Uromyces veratri (DC.) Schroet. (*Aecidium adenostylis* Syd.) See *Veratrum*.
- ADESMIA.** Herbs or shrubs.
Protomyces vagabundus Speg. On stems of *A. punctata* and *Medicago denticulata* in Argentina.
Puccinia adesmiae P. Henn. Rust-producing globular galls on branches of *A. trijuga* in Argentina.
Puccinia bergii Speg. Cinnamon-colored rust pustules on leaves of *A. punctata* in Uruguay.
Uredo solitaria Diet. and Neg. Dark-brown rust pustules on leaves of *A. radiculata* in Chile.
- ADIANTUM.** MAIDENHAIR FERN.
Aphelenchus oleosistis Ritz. Bos. See *Begonia*.
Catharinia paszchkeana Rehm. Greenish-gray to whitish irregular spots on leaves of *A. curvatum* in Brazil.
Hyalospora adiantum-capilli-veneris Syd. Yellow rust pustules on leaves of *A. capillus-veneris* in Spain, France, Italy, Austria, and Dalmatia.
Mycosphaerella rehmania Jaap. On leaves of *A. capillus-veneris* in Dalmatia.
Phyllosticta adianticola Young. Brown leaf spots on *A. tenerum* in Porto Rico.
Septoria filicum (Desm.) Auersw. On leaves of *A. capillus-veneris* in Italy.
Uredinopsis adiantum Kom. Rust pustules on leaves of *Adiantum* sp. in Manchuria.
Uredo gymnogammes P. Henn. See *Dryopteris*.
- ADONIS.** Hardy annual and perennial herbs.
Aecidium solms-laubachii Boy. and Jacz. Rust on leaves of *A. aestivale* in France.
Urocystis leimbachii Oertel. A smut forming galls at the base of the stems of *A. aestivale* in Germany.
- AEGLE.** Bael fruit. Tropical fruit tree.
Bacterium citri Hassé. See *Citrus*.
Cercospora undulata (Ch. Bern.) Sacc. Leaf spots, blackish above, yellow beneath, on *A. odorata* and *Citrus* sp. in Java.
- AEGOPIDIUM.** GOUTWEED. Herbaceous perennials.
Ascochyta podagrariae Bres. On leaves of *A. podagraria* in Germany.
Mycosphaerella aegopodii A. Pot. Whitish or pale spots on leaves of *A. podagraria*, widespread in Europe.
Peronospora podagrariae Oth. On leaves of *A. podagraria* in Switzerland.
Phyllosticta podagrariae Oud. Irregular, indefinite dark-brown spots on leaves of *A. podagraria* in Holland.
Protomyces macrosporus Unger. See *Coriandrum*.
Puccinia aegopodii (Schum.) Mart. Dark-brown rust pustules on leaf blades and petioles of *A. podagraria* in Europe.
Puccinia leioderma Lindr. Brown powdery rust pustules on under side of leaves of *A. alpestre* and *A. tenera* in Turkestan, Siberia, and Japan.
Septoria aegopodii (Pr.) Sacc. Causes spotting of leaves of *A. podagraria* in Russia.
Septoria aegopodina Sacc. Angular, brown, finally white, leaf spots on *A. podagraria* in Italy and Serbia.
- AERIDES.** See *Orchidaceae*.
- AESCULUS.** HORSECHESTNUT. Buckeye. Shade trees.
Ascochyta aesculi Kab. and Bub. Large, irregular, yellow to brown spots, which coalesce, destroying the leaves of *A. hippocastanum* in Bohemia.
Ascochyta grandinaeulans Kab. and Bub. Large, dark-yellow, irregular spots with reddish-purple margins on leaves of *A. hippocastanum* in Bohemia.
Phleospora parvissima v. Hoehn. Numerous subcircular brown spots on leaves of *A. hippocastanum* in Austria and Germany.
Phyllosticta aesculana Oud. Leaf spots on *A. hippocastanum* in Holland.
Phyllosticta aesculina Sacc. Ochraceous leaf spots on *A. hippocastanum* in France.
Phyllosticta hippocastani Oud. Indefinite brown areas on *A. hippocastanum* in Holland.
Phyllosticta socialis Bub. and Kab. Small, angular, purple-brown spots on leaves of *A. hippocastanum* in Bohemia.
Phytophthora syringae Kleb. See *Syringa*.
Septoria aesculi (Lib.) West. Leaf spots on *A. hippocastanum* in Belgium, France, Italy, Russia, and Austria. Of limited distribution in the United States.
Septoria aesculina Thuem. Leaf spots on *A. hippocastanum* in France and Austria. Said to be serious in California on *A. californica*.
Stereum purpureum Pers. See *Fraxinus*.
- AETHIONEMA.** STONE CRESS. Herbaceous perennials.
Aecidium aethionematis P. Magn. Rust pustules on leaves of *A. buzbaumi* in Asia Minor.
- AEXTOXICON.** South American trees.
Hendersonia aetoxiei Speg. Circular, grayish spots with purple areas surrounding, on the upper surface of leaves of *A. punctatum* in Chile.
Phyllosticta aetoxiei Speg. Whitish spots, with narrow dark-purple borders, on leaves of *A. punctatum* in Chile.
- AFZELIA.** INTSIA. Ag. Tropical timber trees.
Pomes lignosus Klotzsch. See *Hevea*.
Hyaloderma afzeliae V. Keissl. Small black globular fruiting bodies in groups on leaves of *Afzelia* sp. in Samoa and the Solomon Islands.
Phyllachora afzeliae Syd. Shining black convex areas on leaves of *A. bijuga* in the Philippines.
Trabutia stephaniae Banc. On leaves of *A. retusa* in the Straits Settlements.

- AGAPANTHUS. ABUMON** Ag. Herbaceous perennials with tuberous root stalks. The name African lily is sometimes applied to *A. umbellatus*.
- Ascochyta hyacinthi** Tassi. See *Hyacinthus*.
- Mycosphaerella agapanthi** (K. and Cke.) Lind. Large, irregular, dark leaf spots on *A. umbellatus* in south Africa.
- AGATHIS. DAMMARA** Ag. DAMMAR PINE. Karui pine. Coniferous trees.
- Aecidium balansae** Cor. On *A. alba* in Straits Settlements and New Caledonia.
- Dielsiella pitzelii** P. Henn. Circular, black tar spots on leaves of *A. palmerstoni* in Queensland.
- AGAVE.** Century plant. Sisal. Henequen. Ornamental and fiber plants.
- Botryosphaeria agaves** (P. Henn.) Butl. Dark-brown spots, becoming pale at the center, on leaves of *A. sisalana*, in India and Tanganyika.
- Gloeosporium agaves** Syd. Irregular, black, diseased areas which finally coalesce on leaves of *Agave* sp. in Brazil.
- Hendersonia agaves** Maubl. Dirty-white spots with brown margins on upper surface of leaves of *Agave* sp. in Algeria.
- Leptosphaeria agaves** Syd. and Butl. On leaves of *A. sisalana* in India.
- Marsonia agaves** Earle. Sunken yellow spots on leaves of *Agave* sp. in Colombia.
- Phyllosticta agaves** Maubl. Yellowish spots, with brown margins, on upper surfaces of leaves of *Agave* sp. in Algeria.
- Septoria megaspora** Speg. On leaves of *A. americana* in Argentina.
- Tubercularia agaves** Pat. On leaves of *Agave* sp. in Costa Rica.
- Tubercularia concentrica** Mont. and Fr. Large brown spots on leaves of *Agave* sp. in France.
- AGERATUM.** Garden annuals.
- Aecidium agerati** P. Henn. Rust pustules on brown leaf spots on *A. conyzoides* in central Africa.
- Albugo brasiliensis** Speg. On leaves of *A. conyzoides* in Colombia and Brazil.
- Puccinia rosea** (Diet. and Holw.) Arth. Brown rust pustules on leaves of *A. conyzoides*, *A. corymbosum*, *A. strictum*, *Eupatorium deltoideum*, *E. glabratum*, *E. gonzalessii*, *E. polyodon*, *E. trinervium*, *E. tubiflorum*, and *Stevia rhombifolia* in Mexico, Porto Rico, and Guatemala.
- Uredo agerati** Mayor. Rust on leaves of *A. conyzoides* in Colombia.
- AGLAONEMA.** Variegated-leaved aroid.
- Gloeosporium graffii** Syd. Anthracnose on leaves of *A. dorsinervium* in the Philippines.
- AGONIS.** See *Leptospermum*.
- AGRIMONIA. AGRIMONY.** Perennial herbs.
- Discosia artocreas** (Tode) Fr. On leaves of *A. eupatoria* in Austria.
- Phyllosticta agrimoniae** Lasch. On leaves of *A. eupatoria* in Denmark.
- Ramularia agrimoniae** Sacc. White, moldlike growth on under sides of leaves of *A. eupatoria* in Siberia.
- AGROPYRON. WHEAT GRASS.**
- Ascochyta graminum** Lasch. Pale-brown spots on leaves of *Agropyron* sp. in Germany.
- Hypochnus fuciformis** (Berk.) McAlp. See *Lolium*.
- Ophiobolus cariceti** (B. and Br.) Sacc. See *Triticum*.
- Uromyces agropyri** Barcl. Yellow-brown to brown rust pustules on leaves of *Agropyron* sp. in India.
- Uromyces fragilipes** Trenchsch. Chestnut-brown rust pustules on leaves of *A. squarrosus* in Transcaspiia (Russia).
- AGROSTIS. BENT GRASS. REDTOP.** Forage grasses.
- Ascochyta graminicola** Sacc. See *Bromus*.
- Diophia graminis** (Fekl.) Sacc. See *Triticum*.
- Endodothella helvetica** (Fekl.) Theiss. and Syd. Black, slightly raised areas on leaves of *Agrostis* sp. in Switzerland and Germany.
- Endodothella toensis** (P. Henn.) Theiss. and Syd. Black fruiting bodies on leaves of *A. perennans* and *A. tenuiflora* in Japan.
- Hypochnus fuciformis** (Berk.) McAlp. See *Lolium*.
- Leptosphaeria sparsa** (Fekl.) Sacc. See *Poa*.
- Ophiobolus cariceti** (B. and Br.) Sacc. See *Triticum*.
- Phyllachora fuscescens** Speg. Black, linear fruiting bodies on leaves of *Agrostis* sp. in Argentina.
- Physoderma agrostidis** Lagerh. Oblong spots on leaves of *A. alba* in Sweden.
- Puccinia agrostidis** Plowr. (*Aecidium aquilegiae* Pers.) Yellow to brown rust pustules on leaves of *Aquilegia alpina*, *A. atrata*, *A. atropurpurea*, *A. castellana*, *A. glandulosa*, *A. haenkeana*, *A. nigricans*, *A. pyrenaica*, *A. sternbergii*, *Agrostis alba*, *A. perennans*, *A. tenuiflora*, and *A. vulgaris* in Europe, India, and Madeira. Reported from New York.
- Puccinia borealis** Juel. Rust with its aecial stage on *Thalictrum alpinum*, and the uredo and telia stages on *A. borealis*, *Anthoxanthum odoratum*, *Calamagrostis stricta*, and *Hierochloë odorata* in Sweden, Norway, and Iceland.
- Puccinia moyanoi** Speg. Brown rust pustules on leaves of *A. moyanoi* in Patagonia.
- Septoria triseti** Speg. Small, linear ashen-colored spots on leaves of *A. magellanica* and *Trisetum phleoides* in Chile.
- Tilletia decipiens** (Pers.) Koern. A smut forming black spore masses in the ovaries and stunting the plants of *A. alba*, *A. stolonifera*, and *A. vulgaris* (*A. pumila*) in Denmark, Italy, Sweden, and Germany.
- Tilletia separata** Kunze. Forms black spore masses in ovaries of *A. spica-venti* and *A. vulgaris* in France, Switzerland, Great Britain, Russia, and Germany.
- Uromyces poae** Rabh. See *Poa*.
- AILANTHUS.** Ornamental trees, *A. altissima*, sometimes called "tree-of-heaven."
- Ascochyta allanthi** Boud. and Fautr. Ashen-colored pustules on leaves of *A. altissima* (*A. glandulosa*) in France.
- Cercospora allanthi** Syd. Circular black spots, changing to brown, on both leaf surfaces of *A. altissima* in Germany.
- Phyllosticta allanthi** Sacc. Irregular, pale-yellow spots with reddish margins, on leaves of *A. altissima* in Italy, Japan, and China.
- Uncinula delavayi** Pat. Powdery mildew on leaves of *Ailanthus* sp. in China.
- AIRA. HAIRGRASS.** Ornamental annual grasses.
- Entioma catenulatum** Rostk. Oblong gray spots on leaves of *A. caespitosa* in Denmark.
- Leptosphaeria insignis** Karst. On leaves of *A. alpina* and *Alopecurus oratus* in Spitzbergen.
- Leptosphaeria nigrans** (Desm.) Ces. and De N. On leaf sheaths of *A. caespitosa*, *Alopecurus pratensis*, *Avena pratensis*, *Dactylis glomerata*, and *Milium effusum* in Finland, France, Italy, and Great Britain.
- Mycosphaerella recutita** (Fr.) Johans. Black fruiting bodies on leaves of *A. caespitosa*, *Bromus benekei*, *B. ramosus*, *Dactylis glomerata*, and *Glyceria fluitans* in Sweden, Denmark, Great Britain, and Germany.
- Puccinia borealis** Juel. See *Agrostis*.

ATRA—Continued.

Tilletia airae Blytt. Black spore masses in ovaries of *A. caespitosa* in Sweden.

Tilletia airae-caespitosae Lindr. Black, powdery spore masses produced in discolored areas in leaves of *A. caespitosa* in Finland.

Tilletia flectens Lagh. Dark-brown fruiting areas in leaves of *A. flexuosa* in Sweden.

Uredo airae Lagh. Rust pustules on leaves of *A. caespitosa* and *A. flexuosa* in Denmark.

Uredo airae-flexuosae Liro. Brown rust pustules on leaves of *A. flexuosa* in Finland.

Ustilago graminica B. and Br. Black spore masses occurring in short parallel transverse areas on leaves of *A. caespitosa* and *Glyceria aquatica* in Great Britain.

AJUGA. BUGLE. Herbaceous perennials.

Acidium ajugae Syd. Yellow rust pustules in circular brown spots on leaves of *Ajuga* sp. in India.

Phyllosticta ajugae Sacc. and Speg. Dull-yellow spots, with brown margins, on leaves of *Ajuga* sp. and *A. reptans* in France and Italy.

Ramularia ajugae (Niessl) Sacc. and var. *ajugae-pyramidalis* Sacc. Subcircular brown spots on leaves of *A. generensis*, *A. latmanni*, *A. mucronata*, *A. pyramidalis*, and *A. reptans* in France, Switzerland, Russia, Denmark, Italy, Yugoslavia, Bulgaria, Bohemia, Austria, and Germany.

AKEBIA. Twining woody plants.

Acidium akebiae P. Henn. Yellow rust pustules in yellow or brown leaf spots on *A. lobata* and *A. quinata* in Japan.

Ascochyta akebiae Bres. Circular to angular brown spots, fading out to white with black borders, on leaves of *A. quinata* in Germany.

Triphragmium thwaitesii B. and Br. See *Acanthopanax*.

ALANGIUM. Tropical trees or shrubs.

Chaetosphaeria indica Niessl. On leaves of *A. decapetalum* in India.

ALBIZZIA. Leguminous ornamental woody plants.

Ascochyta julibrissin F. Tassi. On pods of *A. julibrissin* in Italy.

Diplodia cacaocicola P. Henn. See *Theobroma*.

Endodothelia albizziae Syd. Black stromata on leaves of *A. marginata* in the Philippines

Fomes lamaricensis Murr. See *Hevea*.

Fomes pseudofereus Wakef. See *Hevea*.

Heminthosporium albizziae Petch. Small spots on leaves of *A. lebbek* in Ceylon.

Microstroma albizziae Syd. On living leaves of *A. fastigiata* in the Union of South Africa.

Phragmocaulis vivitensis (Cke.) Theiss. and Syd. Black concave areas on upper surfaces of leaflets of *Albizzia* sp. and *A. fastigiata* in the Union of South Africa.

Phyllosticta divergens Sacc. Subcircular pale-yellow spots, with darker margins, on leaves of *A. anthelmintica* and *A. lebbek* in Abyssinia and Porto Rico.

Ravenelia albizziae Diet. Brown rust pustules on leaves of *A. anthelmintica* in Abyssinia.

Ravenelia albizziae-amarae Bacc. Brown leaf rust on *A. amara* in Abyssinia.

Ravenelia japonica Diet. and Syd. Cinnamon-brown to dark-brown rust pustules on leaves and branches of *A. julibrissin* in Japan.

Ravenelia minima Cke. Brown rust pustules on leaves of *A. fastigiata* in Uganda and the Union of South Africa.

Ravenelia papillosa Speg. Rust on leaves of *A. julibrissin* and *Acacia platensis* in Argentina.

Ravenelia sessilis Berk. Yellow-brown to black rust pustules on leaves and pods of *A. lebbek* in Ceylon and India.

Sphaerophragmium acaciae (Cke.) P. Magn. Brown rust pustules on leaves of *A. lebbek* in China and India.

Sphaerophragmium luzonicum Yates. Rust on *A. procera* in the Philippines.

Uredo ngamboensis P. Henn. Rust on leaves of *A. lebbek* in east Africa and the Philippines.

Uromyadadium tepperianum (Sacc.) McAlp. See *Acacia*.

ALBUCA. Bulbous plants.

Uromyces albucae Kalchbr. and Cke. Cinnamon-brown rust pustules on leaves of *A. altissima*, *A. aurea*, *A. juncea*, and *A. minor* in Angola and the Union of South Africa.

Ustilago albucae Syd. Dark-brown, powdery masses of smut spores replacing the ovaries and anthers of *Albuca* sp. in Central Africa.

ALCHEMILLA. Hardy herbaceous perennials.

Gymnoconia alchemillae Bacc. Rust on leaves of *A. pedata* in Abyssinia.

Laetstidia alchemillae Rostr. On leaves of *A. vulgaris* in Greenland.

Ovularia alpina Mass. On leaves of *A. hoppeana* in Switzerland.

Ovularia haplospora (Speg.) Magn. Subcircular, often confluent, dull-brown spots, with brownish-purple margins, on leaves of *A. alpestris*, *A. fissca*, *A. montana*, *A. pratensis*, *A. pubescens*, *A. subcrenata*, and *A. vulgaris* in Russia, Finland, Sweden, Serbia, Bulgaria, Switzerland, Italy, Austria, Denmark, and Germany.

Ovularia schroeteri (Kuehn.) Sacc. Ashen leaf spots on *A. vulgaris* in France, Switzerland, Austria, and Germany.

Pernospora alchemillae Oth. Downy mildew on leaves of *A. vulgaris* in Europe and Siberia.

Phyllosticta alchemillae (Vgr.) Allesch. On leaves of *A. vulgaris* in Denmark.

Puccinia aliena Syd. Dark-brown powdery rust pustules on leaves of *A. pedata* in tropical Africa.

Uromyces alchemillae (Pers.) Lev. Yellow to brown powdery rust pustules on under sides of leaves of *A. acutangula*, *A. acutiloba*, *A. alpestris*, *A. fissca*, *A. montana*, *A. pastoralis*, *A. pratensis*, *A. pubescens*, *A. splendens*, and *A. vulgaris* in all of Europe, Greenland, and Asia Minor.

Uromyces melosporus (Therry.) Syd. Dark-brown rust pustules on leaves of *A. alpina*, *A. hoppeana*, *A. pedata*, *A. pentaphylla*, and *A. saxatilis* in Uganda, Switzerland, Austria, and Germany.

Venturia alchemillae (Grev.) B. and Br. Pale spots on leaves of *A. vulgaris* in Italy, Belgium, France, Scotland, Finland, and Germany.

ALCHORNEA. DOVEWOOD. Tropical shrubs or trees.

Acidium alchorneae Sacc. Leaf rust on *Alchornea* sp. and *A. rugosa* in the Philippines.

Gloeosporium alchorneae Syd. Anthracnose on leaves of *A. javanica* and *A. rugosa* in the Philippines and Amboina.

Oliva capituliformis (P. Henn.) Arth. Rust on leaves of *A. iricurana* and *A. latifolia* in Porto Rico, Tortola, and Brazil.

ALETRIS. STAR GRASS. Small hardy herbs.

Colletotrichum aletridis P. Henn. Anthracnose on leaves of *A. japonica* in Japan.

Gloeosporium thuemenii Sacc. See *Anthurium*.

ALEURITES. CANDLENUT. WOOD-OIL TREE. Oil-producing trees.

Cercospora aleuritidis Miy. Leaf spot of *A. cordata* in China.

Gloeosporium aleuriticum Sacc. Anthracnose of *A. moluccana* in the Philippines.

ALLAMANDA. Tropical shrubs and climbers.

Septoria allamanda Wint. On leaves of *A. hendersoni* in Portugal.

ALLIUM. ONION, GARLIC, LEEK, CHIVE, SHALLOT. Bulbous herbaceous perennials, cultivated for the most part as food plants, but with some ornamental species.

Bacillus cepivorax Delw. Said to cause a soft rot of bulbs of *A. vpa* in France.

Bacillus croci Mizz. See Crocus.

Cercospora duddiae Walsb. Yellow, circular to irregular spots, on leaves of *A. cepa* and *A. sativum* in the Philippines. The lesions are numerous and rapidly coalesce, killing leaves, with resulting losses to the crop often as high as 50 per cent.

Cercospora victorialis Thum. Large, sunken, circular, greenish-black leaf spots on *A. victorialis* in Siberia.

Chlorospora vastatrix Spag. Produces a "very serious" decay of bulbs of *A. cepa* and *A. sativum* in Argentina.

Fusarium atro-virens (Berk.) On bulbs of *A. sativum* in Great Britain.

Fusarium cepae Bacc. Found in connection with root rot and bulb decay of *Allium* in Japan.

Heterosporium allii-cepae Rensch. Produces gray, elongated spots, with greenish margins over the surface of leaves of *A. cepa* in Yugoslavia.

Hypochnus cucumeris Frank. See Cucumis.

Metamspora allii-fragilis Kleb. See Salix.

Metamspora allii-populina Kleb. See Populus.

Metamspora salicis-albae Kleb. See Salix.

Myosphaerella allioria Auerw. Found on leaves of *A. asperum* in Yugoslavia.

Myosphaerella schoenoprasii (Rosenk.) Sacc. On leaves of *A. fistulosum*, *A. monanthum*, *A. porrum*, and *A. schoenoprasum* in China, Japan, and throughout Europe.

Mycosporium alliorum Berk. Forms dark patches on bulbs of *A. sativum* in Great Britain.

Phyodermis allii Hesse. On leaves of *A. schoenoprasum* and *Allium* sp. in Germany.

Phytophthora allii S. Saw. This fungus forms watery, white, mycelial masses on leaves and flower stalks of *A. fistulosum* in Japan. A soft rot ensues.

Puccinia allii-japonici Diet.

Dark-brown, powdery, rust pustules on leaves of *A. japonicum* in Japan.

Puccinia permixta Syd. Rust on leaves of *A. scirpense*, *A. monanthum*, *A. sativum*, and *A. aphaeocarpum* in Russia.

Puccinia porri (S. Wint.) Wint. Yellow to dark-brown rust pustules on leaves of *Allium* spp. Over 40 species are reported as hosts in New Zealand, Formosa, Japan, Syria, and Europe. Sparingly introduced into the United States.

Puccinia winteriana P. Magn. See Phalaris.

Sclerotinia bulborum (Wak.) Balm. On *A. sativum* and other species in Great Britain. Probably not distinct from the following species.

Sclerotium cepivorum Berk. This fungus causes a serious rot of onion bulbs. Small black fruiting bodies (sclerotia) appear on the surface of diseased bulbs. The disease attacks *A. cepivorum*, *A. cepa*, *A. porrum*, and *A. sativum* in New Zealand, Argentina, Great Britain, Spain, France, Italy, Holland, and Germany. Confused in American literature with *Botrytis allii* Munn. Definitely reported from Oregon and Virginia.

Septoria allieae Gies. On leaves of *Allium* sp. in India.

Septoria allieola Bacc. On leaves of *A. sativum* and *A. aphaeocarpum* in Spain.

Septoria allii-striatellii Spag. Pale, indefinite spots on leaves of *A. striatellum* in Argentina.

Septoria ranajevicii Bub. On leaves of *A. monanthum* in Yugoslavia.

Stemphylium allii Oud. On leaves of *Allium* sp. in Holland.

Tylenchus dipsaci Kuehn. See Narcissus.

Urocystis coelestis (Schlecht.) Rab. See Colchicum.

Urocystis magica Pass. Black masses of smut spores formed in leaves of *A. magicum* in Italy.

Uromyces ambiguus (DC.) Lev. Yellow to black powdery rust pustules on leaves and stems of *A. schoenoprasum*, *A. schoenoprasum*, *A. schoenoprasum*, and *A. aphaeocarpum* in Turkey, Yugoslavia, Russia, Finland, Sweden, Denmark, Holland, France, Austria, and Germany.

Uromyces durus Diet. Rust-forming hard, black, crustlike masses of telial spores on leaves of *A. cepivorum* in Japan.

Uromyces primaverilis Spag. Linear brown rust pustules on leaves of *A. striatellum* in Argentina.

Uromyces reticulatus (Thum.) Bub. Yellow to dark-brown rust pustules on *A. victorialis* in Japan, Siberia, Spain, Portugal, France, Switzerland, Austria, and Hungary.

Ustilago allii Molin. Black spore masses in pustules on bulbs of *A. cepa* in Australia.

Ustilago ceparum Glaw. Black spore masses in bulbs of *A. cepa* in Germany.

ALNUS. ALDER. Ornamental woody plants and timber trees.

Eoasus alni-glutinosa Tub. On *A. glutinosa* in Italy, Denmark, and Sweden.

Eoasus epiphyllus Syd. Causes a wither-brown and forms a gray fruiting layer on leaves, which are distorted, of *A. glutinosa* and *A. incana* in Russia, Switzerland, Portugal, Sweden, Bulgaria, Denmark, Italy, and Austria.

Gloeosporium alneum West. Anthracnose on leaves of *A. glutinosa* and *A. incana* in Albania, Yugoslavia, and Austria.

Gnomonia setacea (Pers.) Ces. and De N. On leaves of *A. glutinosa* in Italy.

Gnomoniella albomaculans Neger. Chalky-white areas on leaves of *A. incana* in Norway.

Gnomoniella tubiformis Thum. Sacc. On leaves of *A. glutinosa* in Italy, Russia, and Germany.

Hypoderma roseum Pers. Fr. See Populus.

Leptothyrium alneum (Lév.) Sacc. On leaves of *A. glutinosa*, *A. incana*, and *A. viridis* in Europe.

Metamsporioidium alni (Thum.) Kleb. Yellow to brown rust pustules on lower leaf surfaces of *A. glutinosa*, *A. firma*, *A. glutinosa*, *A. incana*, *A. montana*, and *A. viridis* in Ecuador, Guatemala, Great Britain, Japan, and Siberia. Also reported from California.

Myosphaerella alnobetulae Jaap. On leaves of *A. viridis* (*A. alnobetula*) in Switzerland.

Passalora alnobetulae Jaap. On leaves of *A. viridis* in Austria.

Passalora bacillifera (Mont.) Fr. Subcircular spots, yellow-brown above, gray-green below, on leaves of *A. viridis* and *A. glutinosa* in Austria.

Phyllosticta alnea Oud. Pale rust-colored spots on leaves of *A. glutinosa* in Holland.

Phyllosticta alniola C. Mass. Small, crowded, angular, often confluent, rust-brown spots on leaves of *A. glutinosa* in Russia and Italy.

Phyllosticta alnigena Thum. Large, yellow-brown, circular, concentrically zoned spots on leaves of *A. glutinosa* and *A. glutinosa* in Italy and Portugal.

Phyllosticta alni-glutinosa Syd. Irregular, yellow-brown spots, 0.5 to 1 centimeter in diameter, on leaves of *A. glutinosa* in Italy, Russia, and Germany.

Phyllosticta alniparda Oud. Circular to angular brown spots, with black margins, on leaves of *A. glutinosa* in Holland.

Phytophthora syringae Kleb. See Syringa.

ALNUS—Continued.

- Ramularia alnicola** Cke. Circular gray leaf spots on *A. glutinosa* in Russia and Great Britain.
- Rhytisma nervale** (Alb. and Schw.) Rehm. Linear rough, black fruiting areas on leaves of *Alnus* sp., *Betula* sp., and *B. alba* in Portugal.
- Sclerotinia alni** Maubl. Forms sclerotia in place of seed in catkins of *A. glutinosa* and *A. incana* in Denmark, Germany, and Russia.
- Septoria alni** Sacc. Subcircular ochraceous spots on upper leaf surfaces of *A. glutinosa* in France and Italy.
- Septoria alnicola** Cke. Brown circular leaf spots on *A. glutinosa* in Great Britain and Italy.
- Septoria alnigena** Sacc. On leaves of *A. cordifolia* and *A. glutinosa* in Italy.
- Septoria carisoleensis** Kab. and Bub. Circular to angular, often confluent, purplish-brown spots with red-brown margins on leaves of *A. viridis* in Italy.
- Taphrina japonica** Kus. Produces a witches'-broom effect on *A. japonica* in Japan. The diseased branches are triangular in cross section. Leaves on the broomed branches appear early but are soon checked in their growth and remain small and pale in color until they wither and fall. The mycelium of the fungus overwinters in the buds.
- Taphrina sadebeckii** Johans. Circular yellowish or white leaf spots on *A. glutinosa*, *A. incana*, and *A. tinctoria* in Europe.
- Taphrina viridis** Sadeb. Forms leaf spots on *A. viridis* in Switzerland.
- Uncinula miyabei** (Salm.) Sacc. and Syd. White powdery mildew on leaves of *A. incana*, *A. japonica*, *A. maritima*, and *A. pendula* in Japan.
- ALOCASIA.** Tropical aroids used as decorative foliage plants.
- Cacoma anthurii** Har. var. *alocasiae* Racib. Rust on leaves of *A. metallica* in Java.
- Gloeosporium thumenii** (Von Th.) Sacc. See Anthurium.
- Mycesphaerella alocasiae** Syd. Leaf spot on *A. indica* and *A. macrorrhiza* in Java and the Philippines.
- Phyllosticta portiana** Sacc. On leaves of *A. portei* in the Philippines.
- Uredo alocasiae** P. Henn. Yellow rust pustules on leaves of *Alocasia* sp. in New Guinea.
- ALOE.** Acaulescent perennial succulents.
- Dothidea aloicola** P. Henn. On leaves of *Aloe* sp. in Abyssinia.
- Gloeosporium affine** Sacc. See Hoya.
- Macrophoma aloës** Scalia. On leaves of *A. latifolia* in Italy.
- Montagnella maxima** Mess. Raised (6 to 8 mm.) circular black areas, 4 to 5 centimeters broad, on leaves of *A. ferox* and *A. platylepis* in the Union of South Africa.
- Montagnella uberata** Sacc. Gall-like fruiting areas on leaves of *A. abyssinica* and *A. flavovirens* in Abyssinia and Eritrea.
- Phyllosticta aloës** Kalchbr. On leaves of *A. latifolia*, *A. purpurascens*, and *A. volkensi* in Spain and the Union of South Africa.
- Septoria helynickii** West. Leaf spot of *A. variegata* in Belgium.
- Uromyces aloës** (Cke.) P. Magn. Brown rust pustules on large sunken spots on leaves of *A. abyssinica*, *A. arborescens*, *A. baumi*, *A. eru*, *A. maculata*, *A. saponaria*, *A. schimperi*, *A. spicata*, *A. transvaalensis*, and *A. vera* in India and Africa.
- ALPINIA.** Tropical decorative plants of the ginger family.
- Catacauma alpiniae** (Sacc. and Berl.) Theiss. and Syd. Black, slightly raised stromata on long elliptical whitish spots on leaves of *A. coerula* in Australia.
- Catacauma renalmiae** (Rehm.) Theiss. and Syd. Tar spot on leaves of *A. antillarum* and *Renealmia* sp. in Porto Rico and Brazil.
- Cercospora alpiniae** Syd. Leaf spot on *Alpinia* sp. in the Philippines.
- Leptosphaeria alpiniae** Maubl. Leaf spot on *A. nutans* in Brazil.
- ALSINE.** Herbaceous annuals, mostly weeds, but a few species cultivated in rock gardens. See, also, *Stellaria* and *Arenaria*.
- Septoria alsines** Rostr. On leaves and stems of *A. verna* in Denmark.
- Septoria vandasia** Bub. Blackened areas on branches, leaves, bracts, and calyces of *A. glomerata* in Bulgaria.
- Uromyces alsines** Tranzsch. Rust on leaves of *A. setacea* in Russia.
- ALSPHILA.** TREE FERN.
- Monorhizina filicina** (B. and Br.) Theiss. and Syd. Large black circular to irregular crustlike fruiting areas on leaves of *Alsphila* sp. in Ceylon and Queensland.
- Pachypatella alsophilae** (Rac.) Theiss. and Syd. See Cyathea.
- Phyllosticta alsophilae** Syd. Subcircular brown spots, becoming pale at the center, on leaves of *Alsphila* sp. in Brazil.
- Polystomella puichella** (Speg.) Theiss. On leaves of *Alsphila* and *Blechnum* in Brazil and Australia.
- ALSTONIA.** PALA Ag. Ornamental trees.
- Ascochyta alstoniae** P. Henn. Circular to oblong pale spots, with brown margins, on leaves of *A. scholaris* in Brazil.
- ALSTROEMERIA.** Tuberous-rooted herbs, sometimes called "Peruvian lily" and "Chilean lily."
- Puccinia alstroemeriae** Syd. Produces light-brown to dark-brown rust pustules on dead irregular areas on both surfaces of leaves of *A. revoluta* in Chile.
- Scolecotrichum alstroemeriae** Allesch. On the lower surfaces of leaves of *Alstroemeria* sp. in Brazil.
- Uromyces alstroemeriae** (Diet.) P. Henn. Yellow sunken, circular to elliptic spots on both leaf surfaces, in which appear yellow-brown to deep-brown rust pustules. The hosts are *A. aurantiaca*, *A. isabellana*, *A. revoluta*, and *A. subrosulacea* in Brazil and Chile.
- ALTERNANTHERA.** Herbaceous bedding plants.
- Laestadia cephalariae** (Awd.) Sacc. and var. *alternantherae* Sacc. Small, pale, sunken spots on leaves of *A. sessilis* and *Cephalaria alpina* in Germany.
- ALTHAEA.** MARSHMALLOW HOLLYHOCK. Annual, biennial, or perennial herbs.
- Cercospora nebulosa** Sacc. On leaves and stems of *A. rosea* in Italy.
- Erysiphe taurica** Léw. Powdery mildew on leaves of *Acanthophyllum glandulosum*, *Althai camelorum*, *A. maurorum*, *Althaea ficifolia*, *A. kurdica*, *Artemisia dracunculus*, *Astragalus* sp., *Capparis spinosa*, *Cercidiphyllum crispum*, *Carlinia corymbosa*, *C. lanata*, *Carthamus lanatus*, *Carlinthe major*, *Chondrilla juncea*, *Cicer*, *Clematis sibirica*, *Cnicus arvensis*, *C. cardunculus*, *C. lanceolatus*, *Coccinea dubia*, *Cynara cardunculus*, *Daucus maximus*, *Eryngium campestre*, *E. noëanum*, *Euphorbia lanata*, *Ezochorda alberti*, *Foeniculum vulgare*, *Hedysarum falconeri*, *Helianthemum oelandicum*, *Inula nervosa*, *Nepeta podostachys*, *Odontospermum aquaticum*, *Phlomis herba-venti*, *P. tuberosa*, *Pieris hieracoides*, *Psoralea drupacea*, *Saussurea*, *Scutellaria multicaulis*, *Taraxacum montanum*, *Teucrium chamaedrys*, *Verbascum blattaria*, *V. phlomisoides*, *V. speciosum*, *Vicia tenuifolia*, and *Zygophyllum fabago* in Europe, Algeria, Turkey, Syria, Persia, Turkestan, and India.
- Phyllosticta althaeicola** Passer. Definite subcircular to irregular, often confluent, whitish spots on leaves of *A. officinalis* in France.

ALTHAEA—Continued.

Phyllosticta althaeae Sacc. Irregular brown spots on leaves of *A. rosea* in France and Italy. Reported from Ohio.

Phyllosticta destructiva Desm. Subcircular pale-yellow leaf spots, with darker margins, on *A. rosea*, *Eunymus europaeus*, *Eunymus* sp., *Lycium barbarum*, *Malva neglecta*, *M. nigrescens*, *M. sylvestris*, *Malva* sp., and *Menyanthes* sp. in Great Britain, France, Denmark, Belgium, Italy, Russia, Austria, and Germany. The diseased areas fall out, causing a "shot-hole" effect.

Phyllosticta puccinospila Mass. Small whitish spots on leaves of *A. rosea* in Italy.

Puccinia heterogena Lagh. See *Malva*.

Septoria althaeae Thum. Pale brown spots on leaves of *A. rosea* in Bohemia.

Septoria heterochroa Desm. See *Malva*.

Septoria parasitica Fautr. White spots on upper surfaces of leaves of *A. rosea* and *Althaea* sp. in France.

ALYSSUM. SWEET ALYSSUM. Lowherbs, cultivated in gardens.

Peronospora alyssi calycini Gäum. Downy mildew on leaves of *A. calycinum* in Switzerland and Germany.

Peronospora alyssi incani Gäum. As above in Bohemia.

Puccinia alyssi Lindr. Dark-brown rust pustules on leaves and stems of *A. halimifolium* and *A. spinosum* in Italy and Spain.

Septoria alyssi Bres. Small, subcircular to oblong pale spots on leaves of *A. saratilis* in Germany.

AMARANTHUS. AMARANTH. Coarse annuals; some species grown for colored foliage and showy flower clusters.

Ascochyta amarantii Allesch. Subcircular to irregular pale or white spots with brown margins on leaves of *Amaranthus* sp. in Germany.

Coniothyrium amaranti (Fl. Tassi.) Sacc. and D. Sacc. Circular spots on leaves of *A. caudatus* in Italy.

Peronospora amarantii Gäum. Downy mildew on leaves of *A. bliti* in Switzerland and Holland.

Phyllosticta moleriana Thum. Large dull-yellow leaf spots, with dark-red margins, on *A. melancholicus* in Portugal.

Phyllostictiella amaranti Fl. Tassi. On leaves of *A. caudatus* in Italy.

Rhizoctonia napi West. See *Brassica*.

AMARYLLIS. Bulbous plants.

Aecidium amaryllidis Syd. and Butl. Yellow rust pustules appear on long sunken spots on the lower surfaces of leaves, more rarely on the upper, of *Amaryllis* sp. (cultivated forms) in India.

Phyllosticta amaryllidis Bres. Small black spots which become elongated or confluent are produced on the leaves of *Amaryllis* sp. in Brazil.

AMBROSINIA. Dwarf perennial bulbous herbs.

Entyoloma dietelaum Bub. Black spore masses in red-brown areas on leaves of *A. bassii* in Sardinia.

AMELANCHIER. SHAD BLOW. JUNE BERRY. SHADBUSH. SERVICE BERRY. Trees and shrubs grown for ornament and fruit.

Coleopuccinia sinensis Pat. See *Cotoneaster*.

Gymnosporangium amelancheris (DC.) Ed. Fisch. See *Juniperus*.

Ochropsora ariae (Eckl.) Syd. See *Sorbus*.

Phyllosticta mespili Sacc. var. **amelanchieri** Jacz. On leaves of *A. rotundifolia* (*A. vulgaris*) in Russia.

AMMOPHILA. BEACH GRASS. MARRAM GRASS. Sand binding grasses.

Ascochyta perforans (Rob.) Sacc. On leaves of *A. arenaria* in Belgium.

Heterosporium graminis McAlp. On leaves and culms of *A. arenaria* in Australia.

Leptosphaeria littoralis Sacc. On leaves and stems of *A. arenaria* and *Calamagrostis* sp. in Belgium and Germany.

Mycosphaerella lineolata (Desm.) Schroet. See *Calamagrostis*.

Septoria ammophila Syd. Long, narrow spots, limited by the veins, on leaves of *A. arenaria* in Germany.

Thecaphora ammophilae Oud. Black powdery spore masses on leaves of *A. arenaria* in Holland.

Uredo ammophilae Syd. Golden-yellow rust pustules over entire leaf surfaces of *A. arenaria* in Germany.

Uredo ammophilina Kleb. Brown rust pustules on leaves of *A. arenaria* in Europe.

AMORPHA. FALSE INDIGO. LEAD PLANT. Ornamental shrubby plants, grown for flowers and foliage.

Systemma amorphae (Rabh.) Theiss. and Syd. On *A. fruticosa* in Germany.

AMORPHOPHALLUS. DEVIL'S-TONGUE. Sometimes known as "snake palm." Giant aroids.

Cercospora amorphophalli P. Henn. Dull-brown circular spots on leaves of *A. campanulatus* and *Amorphophallus* sp. in Java and the Philippines.

Cercospora chevalieri Sacc. Circular to angular, scattered or confluent, white spots on leaves of *Amorphophallus* sp. in Uganda.

Hypochnus cucumeris Frank. See *Cucumis*.

Puccinia paulula Syd. Rust pustules on leaves of *A. campanulatus* in the Philippines.

Rhizoctonia destruens Tass. See *Brassica*.

AMPELOPSIS. VIRGINIA CREEPER. Ornamental woody vines.

Cercospora vitis-heterophyllae P. Henn. On leaves of *A. heterophylla* in Japan.

Phakospora ampelopsidis Diet. and Syd. Golden-yellow to brown rust pustules on under surfaces of leaves of *A. heterophylla*, *A. japonica*, *A. leoides*, *Vitis flexuosa*, and *V. inconstans* in Japan and China.

Phyllosticta allescheri Syd. Circular, fuscous-brown spots on leaves of *A. quinquefolia* and *A. tricuspidata* (*A. viticuli*) in Germany.

Phyllosticta quinquefoliae Allesch. Irregular gray spots, becoming confluent on both surfaces of leaves of *A. heteracea* in Germany.

Phyllosticta speschniewiana Sacc. and Syd. Small circular to irregular spots on both surfaces of leaves of *A. heteracea* in Transcaucasia.

AMSONIA. Perennial herbs.

Phyllosticta amsoniae Fl. Tass. Small subcircular to irregular spots on leaves of *A. satifolia* in Italy.

AMYGDALUS. See *Prunus*.**ANACARDIUM**. CASHEW. Tropical fruit and shade trees.

Atichia millardeti Racib. See *Cinnamomum*.

Dendrodochium paraense Vinc. On leaves of *A. occidentale* in Brazil.

Fusarium udum Butl. See *Dianthus*.

Parodiella melioloides (B. & C.) Wint. On leaves of *Anacardium* sp. in Brazil.

- ANAGALLIS.** PIMPERNEL. Annual or perennial herbs.
Peronospora anagallidis Schroet. Downy mildew on lower sides of leaves of *A. arvensis* and *A. coerulea* in Germany.
Peronospora candida Fekl. White effused fungus patches (downy mildew) on lower surfaces of leaves of *A. arvensis*, *A. coerulea*, and *A. phoenica* in Tunis, Great Britain, France, Switzerland, Denmark, Belgium, Italy, and Germany.
- ANANAS.** PINEAPPLE.
Asterinella stuhlmanni (P. Henn.) Theiss. Brown spots on leaves extending rapidly to cover entire area with black superficial fruiting bodies. On *A. sativus* in Tanganyika, Philippines and India.
Bacterial rot. A rot of pineapple (*A. sativus*) slips and suckers, thought to be due to bacteria, has caused considerable loss in plantings in Costa Rica.
Diplodia cacaotcola P. Henn. See Theobroma.
"New disease." A disease of *A. sativus* in the Philippines and Queensland, due to an unknown cause, is characterized by a hypertrophy of the tissues, giving a rough corrugated appearance. Many diseased suckers develop a heart rot. A similar disease is known in Hawaii.
Thielaviopsis paradoxa (De Seyn.) v. Hoeh. See Saccharum.
- ANAPHALIS.** PEARL EVERLASTING. Hardy, herbaceous border plants.
Miyagia anaphalidis Miy. Leaf rust on *A. margaritacea* and *A. yedoensis* in Japan.
- ANARRHINUM.** Biennial and perennial herbs.
Septoria anarrhini Syd. Small circular or elongated spots on leaves of *A. bellidifolium* in Portugal.
- ANCHUSA.** BUGLOSS. Sometimes, but erroneously, called "alkanet." Hardy biennial or perennial herbs.
Ramularia anchusae Massal. Circular fuscous spots on leaves of *A. barrelieri*, *A. italica*, *A. officinalis*, and *Echium vulgare* in Serbia, Bulgaria, Italy, Denmark, Russia, and Germany.
Ramularia anchusae-officinalis Elias. Irregular dark-brown spots, with indefinite sunken margins, on leaves of *A. officinalis* in Sweden.
- ANDIRA.** VOUCAPOUA Ag. ANGELIN. Tropical shade trees.
Cercospora stevensii Young. Small leaf spots, reddish-brown above, black beneath, on leaves of *Andira* sp. in Porto Rico.
Phyalospora andirae Stevens. Conspicuous, irregular, tan-colored spots on both surfaces of leaves of *A. inermis*, (*A. jamaicensis*) in Porto Rico.
Polystigma pusillum Syd. Small circular to irregular yellowish spots on leaves of *A. creelsa* in Guatemala.
Ravenelia goyazensis P. Henn. Dark-brown rust pustules on leaves of *A. pisonis* in Brazil.
- ANDROMEDA.** See Pieris and Zenobia.
- ANDROPOGON.** BEARD GRASS. Ornamental grasses. See also Holcus and Cymbopogon.
Balanisia andropogonis Syd. Aborts inflorescences of *A. aciculatus* in India.
Balanisia sessilis P. Henn. Forming black sessile subglobose or puvinate stromata on culms of *Andropogon* sp. in Africa.
Cerebellaria andropogonis Ces. Black spore masses in spikelets of *A. pertusus* in Ceylon.
Cintractia columellifera (Tul.) McAlp. Dark-brown spore masses in ovaries, enclosed at first by glumes and ovary walls, finally exposed. On *A. australis*, *A. hirtus*, and *Pennisetum cenchroides* in Australia, Mauritius, and Madeira.
Claviceps pusilla Ces. Ergot on *Andropogon* sp. in Italy.
Endodothella andropogonis (P. Henn.) Theiss. and Syd. Black elliptical stromata on both leaf surfaces of *Andropogon* sp. in the Congo.
Endodothella dispar Syd. Black elliptical stromata on both surfaces of leaves of *A. contortus* in India.
Entyloma obesum Syd. Elongate smut pustules on lower leaf surfaces of *A. annulatum* in India.
Epheis pallida Pat. Gray or black stromata, deforming spikelets of *A. aciculatus* in Tonkin and the Philippines.
Leptosphæria herpetchoides De Not. See Triticum.
Phaeodothis fallax (Sacc.) Theiss. and Syd. Long black stromata on both leaf surfaces of *A. gryllus* and *A. ischaemum* in Italy, Yugoslavia and Austria.,
Phyllachora andropogonicola Speg. Small black erumpent fruiting bodies on leaves of *A. saccharatus* in Argentina.
Phyllachora andropogonis Karst. and Har. Tar spot on leaves of *A. candolleana*, *Andropogon* sp., and *Cymbopogon confertiflorus* in Timor, Ceylon, and the Philippines.
Phyllachora assimilis Theiss. and Syd. Linear black stromata on both leaf surfaces of *A. assimilis* in India.
Phyalospora festucae (Lib.) Sacc. On leaves of *Andropogon* sp., *Arundo* sp., *Festuca* sp., *Holcuschaenus* sp., and *Phragmites communis* in France, Italy, and Germany.
Puccinia andropogonicola Har. and Pat. Linear brown to black rust pustules on both leaf surfaces of *Andropogon* sp. in the Congo.
Puccinia andropogonis-macranthi Diet. Rust on leaves of *A. macranthus* in Japan.
Puccinia andropogonis-micranthi Diet. Brown rust pustules on leaves of *A. micranthus* in Japan.
Puccinia cesatii Schroet. Cinnamon-brown to dark-brown rust pustules on leaves of *A. angustifolius*, *A. annulatus*, *A. gryllus*, *A. hirtus*, *A. ischaemum* and *A. pubescens* in Albania, Yugoslavia, Dalmatia, Rumania, Bulgaria, Russia, Portugal, France, Switzerland, Italy, Austria, Hungary, and Germany.
Puccinia chrysopogi Barel. Yellow rust pustules in sunken yellow spots on leaves of *Jasminum humilis* and brown pustules on leaves of *A. (Chrysopogon) gryllus* in India.
Puccinia citrata Syd. Brown leaf rust on *A. citratus* in the Philippines.
Puccinia duthiae Ell. and Tr. Rust pustules on leaves of *A. intermedius* and *A. pertusus* in India.
Puccinia erythraeensis Pazschke. Rust on leaves of *A. commutatum* and *Andropogon* sp. in Abyssinia.
Puccinia hookeri Syd. Brown rust pustules on both sides of leaves of *A. echinulatus* in north India.
Puccinia kojukensis Diet. Brown to black rust pustules on leaves of *Andropogon* sp. in Japan and India.
Puccinia nakanishikii Diet. Brown, linear, powdery rust pustules on leaves of *A. intermedius*, *Cymbopogon confertiflorus*, *C. schoenanthus*, and *C. nardus* in Ceylon, India, Japan, and the Philippines.
Puccinia posadensis Sacc. and Trott. Linear brown rust pustules on leaves, culms and sheaths of *A. condensatus* in Argentina.
Puccinia propinqua Syd. and Butl. Rust on leaves of *Andropogon* sp. in India.
Puccinia prunicolor Syd. and Butl. Rust pustules on purple sunken spots on leaves and sheaths of *A. seratus* in India.
Puccinia pusilla Syd. Black rust pustules on yellow or brown indefinite spots on leaves of *A. assimilis* in India.

ANDROPOGON—Continued.

- Puccinia venestula* Arth. Brown leaf rust on *A. brevifolius* in Porto Rico and Costa Rica.
- Puccinia versicolor* Diet. and Holw. Brown to black rust pustules on leaves of *A. contortus* and *A. melanocarpus* in Argentina, Mexico, and India.
- Septoria grylli* Sacc. On leaves of *A. gryllus* in Italy.
- Sorosporium andropogonis-aciculati* Petch. Smut forming black spore masses in inflorescence of *A. aciculatus* in Ceylon and the Philippines.
- Sorosporium dembiansense* Bacc. Smut on *A. rufus*, *A. papillipes*, and *A. arrhenobasis* in Abyssinia.
- Sorosporium geminellum* Syd. and Butl. Elongate dark-brown smut sori replacing ovaries of *Andropogon* sp. in India.
- Sorosporium icosiense* Maire. Smut on *A. hirtus* in French north Africa.
- Sorosporium maranguensis* P. Henn. Black powdery spore masses in inflorescences of *A. lepidus* in tropical Africa.
- Sorosporium tembuti* P. Henn. and Evans. Black spore masses in ovaries of *A. cymbosum* in the Union of South Africa.
- Sorosporium wildemanianum* P. Henn. Smut sori, with ashen coverings, formed in ovaries and stamens of *Andropogon* sp. in the Congo.
- Thecaphora berkeleyana* Fisch. Spherical black smut sori in spikelets of *A. perforatus* in Ceylon.
- Tilletia calospora* Pass. Smut on *A. agrestis* in Italy.
- Tolyposporium philippinense* Syd. Black powdery smut sori replacing ovaries of *A. contortus* in the Philippines.
- Uredo andropogonis-hirti* R. Maire. Brown rust pustules on small sunken brown spots on leaves of *A. hirtus* and *A. pubescens* in France, Spain, and Greece.
- Uredo andropogonis-lepidi* P. Henn. Elongate yellow rust pustules on leaves of *A. lepidus* in tropical Africa.
- Uromyces andropogonis-annulati* Syd. and Butl. Brown rust pustules on leaves of *A. annulatus* in India.
- Uromyces dignyi* Pat. and Har. Brown rust pustules on leaves of *A. hirtiflorus*, *A. proximus*, and *Andropogon* sp. in Niger Territory, Abyssinia, and Guatemala.
- Ustilago amadelpa* Syd. and Burl. Black, powdery, smut sori in panicles of *Andropogon* sp. in India.
- Ustilago andropogonis-aciculati* Petch. Smut on *A. aciculatus* in the Philippines and Ceylon.
- Ustilago andropogonis-annulati* Bref. Powdery, dark brown, spore masses in ovaries of *A. annulatus* in India.
- Ustilago andropogonis-finitimi* Maubl. Black spore masses in inflorescences of *A. finitimus* in east Africa.
- Ustilago andropogonis-hirtifolii* P. Henn. Smut sori in the spikelets, protected at first by long cylindrical, cinnamon-colored membranes which rupture, exposing the black spore masses. On *A. hirtifolius* in Mexico.
- Ustilago andropogonis-tuberculati* Bref. Compact, hard, black sori in ovaries of *A. tuberculatus* in India.
- Ustilago bicornis* P. Henn. Sori black, in panicles of *A. bicornis* in Brazil.
- Ustilago congensis* Syd. Powdery, black smut sori, 1 to 2 centimeters long, in ovaries of *Andropogon* sp. in the Congo.
- Ustilago culmiperda* Schroet. Smut sori in culms of *A. bicornis* in Brazil.
- Ustilago dinteri* Syd. Sori developing in inflorescences, 1 to 5 centimeters long, inclosed by gray or brown membranes, which rupture, exposing dark fuscous or black spore masses, on *A. papillosum* in southwest Africa.
- Ustilago duthiei* Ricker. Smut sori, olive-brown, deforming the spikelets of *A. bladii* in India.
- Ustilago effusa* Syd. Black, smut sori in sheaths, culms, and unexpanded leaves of *A. muricatus* in India.
- Ustilago filiformis* P. Henn. Long smut sori (up to 4 centimeters) destroying ovaries of *A. contortus* in east Africa.
- Ustilago furcata* Pat. and Har. Powdery, dark chestnut-colored sori, protected by white membranes, in ovaries of *Andropogon* sp. in west Africa.
- Ustilago globulifera* Sacc. and Trott. Hard, black, smut sori in inflorescences of *Andropogon* sp. in the Congo.
- Ustilago guaranitica* Speg. Smut on *A. condensatus* in Brazil and Argentina.
- Ustilago ischaemonoides* P. Henn. Black smut sori, covered at first with ashen membranes, destroying the inflorescences of *Andropogon* sp. in the Congo.
- Ustilago lanigeri* P. Magn. Black smut sori destroying inflorescences of *A. lanigerus* in Persia.
- Ustilago leucostachys* P. Henn. Sori in panicles of *A. leucostachys* in Brazil, yellowish at first, then black, due to the rupturing of the protecting membranes.
- Ustilago nyassal* Syd. Dark-brown smut sori in inflorescences of *Andropogon* sp. in Nyassa.
- Ustilago occulta* P. Henn. Black smut masses in culms of *Andropogon* sp. in Brazil.
- Ustilago stuhlmanni* P. Henn. Powdery, black smut masses on *Andropogon* sp. in central Africa.
- Ustilago superflua* Syd. Smut sori destroying ovaries of *A. foveolatus* in India.
- Ustilago tenuis* Syd. Long, dark brown sori in inflorescences of *A. pertusus* in India.
- Ustilago tonkinensis* P. Henn. Sori in red-brown spots on both surfaces of leaves of *Andropogon* sp. in Tonkin.
- Ustilago tumefaciens* P. Henn. Smut sori gall-like, causing abortion of inflorescences of *A. rufus* in tropical Africa.
- Ustilago vanderysti* P. Henn. Black, powdery sori destroying spikelets of *Andropogon* sp. in the Congo.
- Ustilago warneckeana* P. Henn. Dark-brown smut sori, deforming and destroying inflorescences of *A. contortus* in Camerun and India.

ANDROSACE—ROCK JASMINE. Rock plants.

- Peronospora agrorum* Gaum. Downy mildew on leaves of *A. septentrionalis* in Sweden.
- Puccinia volkartiana* Ed. Fisch. Dark-brown rust pustules on leaves and stems of *A. chamaejasme* in Switzerland.
- Puccinia dubyi* Muell. Rust on leaves of *A. alpina*, *A. glacialis*, and *A. laggeri* in Switzerland.
- Septoria androsaces* Pat. Sunken black spots on leaves of *A. rotundifolia* in China.
- Ustilago androsaces* Karst. Powdery, dark-brown smut sori in ovaries of *Androsace* sp. in Siberia.

ANEMONE (including *Pulsatilla*). Hardy perennials, sometimes called "windflower."

- Aphelenchus olestinus* Ritz. Bos. See Begonia.
- Ascochyta anemones* Kab. and Bub. Circular to irregular, yellow to dark-brown leaf spots on *A. nemorosa* in Bohemia.
- Coleosporium pulsatillae* Lev. Golden rust pustules in brown sunken spots on leaves of *A. cernua*, *A. halleri*, *A. montana*, *A. patens*, *A. pratensis*, *A. vulgaris*, and on the needles of *Pinus silvestris* in Japan, Siberia, Russia, Switzerland, Denmark, France, Italy, Austria, Hungary, and Germany.

ANEMONE—Continued.

- Micropeltis wettsteinii** v. Hoehn. Black fruiting bodies on upper surfaces of leaves of *A. wettsteinii* in Brazil.
- Ochropsora ariae** (Fekl.) Syd. See Sorbus.
- Phyllosticta anemones** P. Brun. Circular to irregular, gray or gray-brown, spots on leaves of *A. japonica* in France.
- Phyllosticta nigro-maculans** Sacc. Black spots, often involving entire leaf area, on *A. nemorosa* in France, Spain, and Italy.
- Physoderma deformans** Rostk. On *A. nemorosa* and *A. ranunculoides* in Denmark.
- Puccinia japonica** Diet. Chestnut-colored rust pustules on leaves and petioles of *A. flaccida* in Japan.
- Puccinia schelliana** Thuem. Powdery, dark-brown rust pustules on yellow leaf spots of *A. narcissiflora* in Russia.
- Puccinia singularis** P. Magn. Powdery, cinnamon-brown rust pustules, on pale circular sunken spots of *A. altaica* and *A. ranunculoides* in Japan, Yugoslavia, Rumania, Spain, France, Denmark, Austria, and Hungary.
- Puccinia vesiculosa** Schlecht. Powdery, brown rust pustules, in circular spots, on leaves of *A. narcissiflora* in Alaska.
- Ramularia pulsatillae** Höll. Dull yellow, indefinite spots, generally near the tips of leaves of *A. (Pulsatilla) nigricans* in Hungary.
- Sclerotinia tuberosa** (Hedw.) Fekl. Attacks the roots of *A. nemorosa* in Great Britain, Bohemia, Russia, and Austria, transforming them into hard, black sclerotia resembling the roots.
- Septocylindrium anemones** Delacr. Large, dirty, white spots on leaves of *A. sylvestris* and *A. sylboatica* in France.
- Septoria altaica** Thuem. On leaves of *A. altaica* in Siberia.
- Septoria brunaudii** Sacc. and Syd. Large, subcircular to irregular, gray or ashen spots, with narrow brown margins, on leaves of *A. japonica* in France.
- Septoria ficariae** Desm. Ashen spots with brown margins on leaves of *A. nemorosa* and *A. ranunculoides* in Denmark, France, Belgium, and Great Britain.
- Septoria silvicola** Desm. On leaves of *A. nemorosa* in Russia, France, and Belgium.
- Synchytrium anemones** Wor. Small galls on under sides of leaves and stems of *A. nemorosa*, *A. ranunculoides*, *A. silvestris*, and *A. virginiana* in Europe and North Africa. Also collected once in Iowa on *A. cylindrica*.

ANGELICA. Perennial herbs.

- Mycosphaerella angelicae** (Fr.) A. Pot. On *A. montana* in Switzerland and Italy.
- Phyllachora angelicae** (Fr.) Fekl. On leaves of *A. decursiva* and *A. sylvestris* in Japan, Siberia, Switzerland, France, and Finland. A doubtful species reported by Theissen as probably only an immature *Mycosphaerella*.
- Puccinia angelicae-edulis** Miy. Powdery brown to black rust pustules, on blades and petioles of leaves of *A. matsumurae*, *A. miqueliana*, *A. polyclada*, *A. polymorpha*, *A. shikokiana*, *A. ursina*, and *A. vulgata* in Japan.
- Puccinia angelicae-mamilata** Kleb. Brown rust pustules on leaves and petioles of *A. sylvestris*, *Polygonum bistorta* and *P. viviparum* in Helvetia and Italy.
- Puccinia angelicicola** P. Henn. Black rust pustules on yellow spots on leaves of *A. miqueliana* in Japan.
- Puccinia karstenii** Lindr. Brown rust pustules on leaves and petioles of *A. sylvestris* in Sweden, Denmark, and Finland.
- Puccinia miyabeana** Miy. Dark-brown rust pustules on leaves of *A. hakonensis*, *A. kiusiana*, and *A. utilis* in Japan.
- Puccinia psoroderma** Lindr. Brown rust pustules on *A. purpurascens* in the Caucasus.
- Ramularia angelicae** v. Hoeh. On leaves of *A. sylvestris* in Spain, Austria, and Denmark.
- Ramularia archangelica** Lindr. White spots with brown borders on leaves of *A. archangelica* in Lapland.
- Septoria bondarzewii** P. Henn. Yellow leaf spots on *A. sylvestris* in Russia.

ANGOPHORA. GUM MYRTLE. Trees or shrubs.

- Gloeosporium angophorae** F. Tassi. Anthracnose on branches and petioles of *A. costata* (*A. lanceolata*) in Australia.

ANGRAECUM. See Orchidaceae.**ANISACANTHUS**. Ornamental shrubs.

- Puccinia anisacanthi** Diet. and Holw. Brown rust pustules on leaves of *Anisacanthus* sp. in Mexico.

ANNONA. SOURSOP. CUSTARD APPLE. CHERIMOYA. SUGAR APPLE. Tropical fruit trees.

- Aecidium anonae** P. Henn. Yellow rust pustules on leaves of *Annona* sp. in Brazil.
- Ascochyta cherimolae** Thuem. Irregular white or yellow leaf spots, on *A. cherimola* in Ecuador and Portugal.
- Colletotrichum ananicola** Speg. Subcircular, gray spots, with fuscous margins, on leaves of *A. cherimola* in Argentina.
- Corticium salmonicolor** B. and Br. See Citrus.
- Fomes lamaeensis** Murr. See Hevea.
- Phyllachora atro-maculans** Syd. Dark-brown, circular to irregular, sunken-leaf spots on *Annona* sp. in Costa Rica. Black opaque stromata are produced on the spots.
- Phyllosticta anonae** P. Henn. Circular to oblong, pale spots, with dark-brown margins, on leaves of *Annona* sp. in Brazil.
- Phyllosticta cherimolae** Alm. and Cam. Elongate gray spots, with brown margins, on leaves of *A. cherimola* in Portugal.
- Phyllosticta insularum** Sacc. Causes leaves to die, beginning at margins and extending irregularly inward. On *A. muricata* in the Philippines.
- Pocosphaeria anonae** Rangel. On leaves of *A. reticulata* in Brazil.
- Uredo cherimolae** Lagh. Light-brown, effused spore masses on lower surfaces of leaves of *A. cherimola*, *A. reticulata*, *A. squamosa*, and *Rollinia multiflora* in Ecuador, Cuba, Trinidad, and sparingly in Florida.

ANOMATHECA. See Lapeyrousia.**ANTHEMIS**. CAMOMILE, sometimes spelled "chamomile." Annual, biennial and perennial herbs.

- Peronospora anthemidis** Gäum. Downy mildew on leaves of *A. austriaca* and *A. cotula* in Norway, Holland, Denmark, Bohemia, Austria, and Germany.
- Peronospora radii** De By. See Chrysanthemum.
- Puccinia anthemidis** Syd. Dark-brown, rust pustules on stems and leaves of *A. altissima* in France.
- Puccinia baumleriana** Bub. Dark-brown, rust pustules on leaves of *A. tinctoria* in Hungary.
- Ramularia anthemidis** Höll. Indefinite spots on leaves of *A. ruthenica* in Hungary.

ANTHERICUM. ST. BERNARD'S LILY.

Aecidium antherici P. Henn. and Evans. Yellow pustules on effuse oblong yellow spots on leaves of *Anthericum* sp. in the Union of South Africa.

Puccinia asphodeli Duby. See *Asphodelus*.

ANTHOLYZA. Cormous plants related to *Gладиолус*.

Aecidium antholyzae Bub. Yellow rust pustules on leaves of *A. aethiopica* in the Union of South Africa.

Uromyces antholyzae Syd. This rust produces dark-brown pustules on both leaf surfaces of *A. abyssinica* in Abyssinia.

ANTHOXANTHUM. Aromatic grasses.

Dilophia graminis (Eckl.) Sacc. See *Triticum*.

Helmintosporium dematioideum Bub. and Wrobl. On leaves of *A. odoratum* in Bohemia. Reported from New York.

Puccinia anthoxanthi Fekl. Brown to dark-brown rust pustules on leaves, leaf-sheaths and stems of *A. odoratum* in Great Britain, Belgium and Germany.

Puccinia borealis Juel. See *Agrostis*.

Puccinia fujiensis S. Ito. Brown rust pustules on leaves of *A. japonicum* in Japan.

Septoria oxyspora Penz. and Sacc. On leaves of *A. odoratum*, *Arundo donax*, and *Hierochloa borealis* in Italy and Denmark.

Uredo anthoxanthina Bub. Powdery, yellow rust pustules on leaves of *A. odoratum* in Great Britain, Switzerland, and Germany.

ANTHURIUM. Ornamental tropical aroids, much cultivated in greenhouses.

Colletotrichum anthurii Delacr. Dark-brown spots on leaves of *Anthurium* sp. in France.

Gloeosporium anthurii Allesch. On leaves and petioles of *A. martianum* in Germany. Probably the same as the following species.

Gloeosporium thumenii (von Th.) Sacc. Black, oblong acervuli on leaves of *Aletris fragrans*, *Anthurium leuconeurum*, *Alocasia cucullata*, *A. rotunda*, *Caridodora atrovirens*, *Dracaena draco*, *Dieffenbachia concolor* and *Smilax atrovirens* in France, Denmark, and Austria.

Mycosphaerella anthurii Miles. Large, irregular, white areas on leaves of *A. acule* in Porto Rico. The diseased areas become dry and papery, often dropping out.

Phyllachora engleri Speg. Black, circular, scattered, slightly raised, shining stromata on leaves of *A. dominicense*, *A. scandens*, *Philodendron* sp. and *Spathycarpa* sp. in the West Indies, Central and South America.

Phyllosticta cavarae Trinch. Irregular, yellow to brown spots, with ashen brown or dirty white distinct margins, on leaves of *A. crassinervium* in Italy.

Physalospora inanis (Schw.) Sacc. On leaves of *Anthurium* sp. and other aroids in Ecuador and Surinam.

Uredo anthurii Har. Rust on leaves of *A. scandens* in Porto Rico and France.

ANTHYLLIS. KIDNEY VETCH. Perennial herbs, including ornamental and forage plants.

Ascochyta vulnerariae Fekl. Purple spots on leaves of *A. vulneraria* in Portugal and Germany.

Cercospora radiata Fekl. Dull-brown spots on leaves of *A. dillenii* var. *dalmatica* and *A. vulneraria* in Spain, Italy, Switzerland, Denmark, Russia, and Germany.

Ramularia schulzeri Bauml. Dull-yellow to reddish spots, on leaves of *A. vulneraria*, *Lotus corniculatus*, and *L. uliginosus* in France, Austria, Hungary, and Germany.

Septoria anthyllidis Sacc. Brown spots, becoming whitish, on leaves of *A. dillenii* and *A. vulneraria* in Dalmatia, Sardinia, Russia, Austria, and Hungary.

Septoria henryana Trav. Circular to irregular, grayish-white spots with yellow-gray margins, on leaves of *A. vulneraria* in Italy.

Typhula trifolii Rostr. See *Trifolium*.

Uromyces anthyllidis Grex. Cinnamon-brown to dark-brown rust pustules on leaves of *A. dillenii*, *A. maritima*, *A. polyphylla*, *A. tetraphylla*, *A. vulneraria*, *Hippocrepis biconcolor*, *H. ciliata*, *H. multisiliquosa*, *H. unisiliquosa*, *Lotus aegyptiacus*, *L. cytoides*, *L. ornithopodioides*, *Lupinus albus*, *L. luteus*, *Medicago arabica*, *M. hispida*, *M. orbicularis*, *Trigonella foenum-graecum*, *T. monspeliaca*, and *T. occulta* in Madeira, Tripoli, Tunis, Morocco, Egypt, India, and Europe.

ANTIDESMA. CHINESE LAUREL. Tropical trees.

Cronartium antidesmae-dioicae Syd. Leaf rust on *A. ghaesemilla* in the Philippines.

Guignardia fusco-coriacea Rehm. On leaves of *A. buniis* and *Antidesma* sp. in the Philippines.

Phyllosticta ghaesemillae Koord. Subcircular, solitary or confluent, brown spots, becoming whitish, on leaves of *A. ghaesemilla* in Java.

Puccinia aequalis P. Henn. Yellow-brown to dark-brown rust pustules on leaves of *A. venosum* and *Antidesma* sp. in tropical Africa.

Uredo antidesmae Rae. Golden-yellow rust pustules on leaves of *A. buniis* in Java.

Uredo antidesmae-dioicae Rae. Brown rust pustules on circular sunken spots on leaves of *A. dioica* and *A. ghaesemilla* in Java and the Philippines.

ANTIRRHINUM. SNAPDRAGON. Flower-garden and greenhouse herbs.

Ascochyta decipiens Pass. Dark-brown spots on stems and branches of *A. majus* in Italy.

Cercospora sp. Shot-hole disease. Minute, buff-colored spots appear on the lower leaves, increasing in size rapidly and spreading to the upper leaves, calyces, seed pods and stems of younger shoots, the latter being girdled causing them to die back rapidly. The spots are more or less circular in shape, with a well defined margin, and are surrounded by a ring of discolored tissue. The central areas, which are slightly sunken, dry out and fall, giving a shot hole effect. On cultivated snapdragon (*A. majus*) in Great Britain.

Cercospora antirrhini Wakef. Small, sunken patches on leaves and stems of *A. majus* in Great Britain. These patches have a dark-green, water-soaked appearance at first, later becoming dull yellow.

Fusarium udum Butl. See *Dianthus*.

Peronospora antirrhini Schroet. Downy mildew on leaves of *A. majus* and *A. orontium* in Denmark, Switzerland, and Germany.

Phyllosticta linariae Sacc. Subcircular brown spots, becoming whitish, with narrow reddish margins on leaves of *A. hispanicum* and *Linaria catiline* in France.

Pseudomonas antirrhini Takimoto. Bacterial leaf spot of *A. majus* in Japan.

Septoria antirrhini Desm. Yellow spots on stems and leaves of *A. majus* in Great Britain, France, Italy, Portugal, Dalmatia and Australia.

AOTUS. Leguminous shrubs.

Cronartium jacksoniae P. Henn. See *Jacksonia*.

APHELANDRA. Tropical ornamental shrubs.

Aecidium aphelandrae P. Henn. Circular dull-brown spots, with dark margins, on leaves of *Aphelandra* sp. in Brazil.

APIUM. CELERI Ag. Celery.

Bacillus apii Mig. Bacterial rot of *A. graveolens* in Italy.

Bacillus apiovorus Worm. Reported as the cause of a heart rot of celery (*A. graveolens*) in Great Britain. Probably not distinct from *B. caratovorus* Jones.

Cercospora pastinacae Karst. On leaves of *A. graveolens* in Finland.

Phytaeota magnusiana (Allesch.) Bres. On leaves of *A. graveolens* in Russia, Italy and Germany. Reported from New York.

Puccinia apii Desm. Powdery, cinnamon-brown to dark-brown rust pustules on leaves and petioles of *A. graveolens*, *A. prostratum*, and *Selinum japonicum* in Tasmania, India, Japan, Spain, Italy, Finland, Sweden, Denmark, Switzerland, France, Holland, Belgium, Great Britain, Austria, Hungary, and Germany.

Puccinia thumeni (Thuem.) McAlp. Cinnamon-brown to dark-brown rust pustules on both surfaces of leaves of *A. graveolens* and *A. prostratum* in Victoria and Tasmania.

Septoria apicalia Speg. Leaf spot on *A. australe* in Chile and Argentina.

Uredo cundinamarcensis Mayor. Brown rust pustules on leaves of *A. ternatum ranunculifolium* in Colombia.

APOSERIS. See Hyoseris.**AQUILEGIA.** COLUMBINE. Hardy herbaceous perennials.

Actinonema pallens Sacc. and Cav. On leaves of *A. vulgaris* in Italy.

Haplobasidium pavonium v. Hoeh. Large, solitary or marginal yellow spots, with purple-brown margins on leaves of *A. vulgaris* in Spain and Austria.

Marssonina aquilegiae (Rbh.) Rostr. Subcircular, ashen colored spots, with dark margins, on leaves of *A. vulgaris* in Great Britain, France, and Denmark.

Phyllosticta aquilegiola Brun. Dull-brown spots on leaves of *A. vulgaris* in Italy.

Puccinia agrostidis Plowr. See Agrostis.

Puccinia melasmoides Franzsch. Brown rust pustules on indefinite spots on leaves and petioles of *A. aktiensis* and *A. vulgaris* in Japan and Turkestan.

Septoria penzigii Cav. and Mor. Whitish leaf spots, with broad brown margins, on *A. vulgaris* in Italy and Russia.

ARABIS. ROCK CRESS. Perennial or annual herbs, mostly rock plants.

Peronospora arabis alpinae Gäum. Downy mildew on leaves of *A. alba* and *A. alpina* in Switzerland.

Peronospora arabis hirsutae Gäum. As above on *A. arenosa* and *A. hirsuta* in France, Switzerland, Norway, Denmark, Austria, and Germany.

Peronospora arabis turritae Gäum. As above on *A. turrita* in Switzerland.

Phyllosticta alpina Allesch. and var. *helvetica* Jaap. Large, circular or elongate definite spots on leaves of *A. alpina* in Bavaria and Switzerland.

Puccinia thlaspeos Schubert. Brown rust pustules on leaves of *A. halleri*, *A. hirsuta*, *A. thaliana*, *Thlaspi alpestre*, *T. arvense*, *T. calamitarius*, *T. montanum*, *T. perfoliatum*, *T. praecox*, and *T. rotundifolium*, in Alaska, Russia, Sweden, Switzerland, Holland, Belgium, France, Austria, Hungary, and Germany.

Septoria arabis Sacc. On leaves of *A. ciliata* in Italy.

Ustilago thlaspeos (Beck.) Lagh. A smut forming black spore masses in the ovules in pods of *A. hirsuta*, *A. petraea* and *Thlaspi alpestre* in Switzerland and Austria.

ARACHIS. PEANUT.

Cercospora sp. A leaf spot, due to a *Cercospora* sp., said to be distinct from *C. personata*, has caused considerable losses to the peanut (*A. hypogaea*) in India.

Hypochnus cucumeris Frank. See Cucumis.

Mosaic or Curl. An obscure disease of *A. hypogaea*, probably of the mosaic group of diseases, occurs in the Union of South Africa, Java, and other parts of the Orient. It is characterized by a dwarfing of infected plants and the yellowing and crinkling of the foliage. The disease is called "crinkle leaf" or rosette in south Africa.

Rhizoctonia destruens Tassi. See Solanum.

Sclerotinia arachidis Hanzawa. Stem rot of *A. hypogaea* in Japan.

Sclerotinia miyabeana Hanzawa. Stem rot of *A. hypogaea* in Japan.

Uredo arachidis Lagh. Numerous dull-brown rust pustules on the under sides of leaves of *A. hypogaea* throughout the West Indies, Central America, and South America. This disease has proven itself the limiting factor of production in some of the West Indies. It has been found in one locality in Florida, but is thought to have been eradicated.

ARALIA. Spikenard. Sarsaparilla. Udo. Herbs, shrubs, and trees, often spiny.

Cercospora araliae P. Henn. Effuse, dull-brown spots on leaves of *A. sinensis* and *A. spinosa* in Japan.

Phyllosticta araliae Sacc. and Berl. Large yellow to ashen spots, with dull yellow margins, on leaves of *Aralia* sp. in Algeria.

Physalospora araliae Pat. On leaves of *Aralia* sp. in Ecuador.

Puccinia araliae-cordatae P. Henn. Brown to black rust pustules on brown spots on leaves of *A. cordata* in Japan.

ARAUCARIA. MONKEY PUZZLE. NORFOLK ISLAND PINE. Ornamental coniferous trees.

Coniothyrium pallido-fuscum Sacc. See Chamaecyparis.

Gloeosporium araucariae Karst. and Har. Anthracnose on leaves and cones of *A. rulei* in New Caledonia.

Haplodothis araucariae (Rehm.) v. Hoeh. Wart-like stromata on needles of *A. brasiliana* in Brazil.

Phyllosticta aaraucariae Wor. On needles of *A. imbricata* in Russia.

ARBUTUS. STRAWBERRY TREE. Broad-leaved evergreen trees and shrubs.

Ascochyta unedonis Sacc. (*Ascochyta unedonis* [Sacc.] Fl. Tass.) Ashen colored spots with reddish margins on leaves of *A. unedo* in France and Spain.

Cercospora molleriana Wint. Circular to irregular, often confluent, gray leaf spots on *A. andrachne*, *A. longifolia*, and *A. unedo* in Russia, Portugal, and Italy.

Corynebium microstictum B. and Br. On *A. unedo* in Italy.

Exobasidium unedonis Maire. The fungus deforms the branches and destroys the leaves of *A. unedo* in Algeria. It fruits on the under surface of the leaves and branches.

Mycosphaerella unedonis Jaap. On leaves of *A. unedo* in Dalmatia.

Phyllosticta arbuti (Desm.) Sacc. Small fuliginous-colored spots on leaves of *A. unedo* in France.

Phyllosticta arbuti-unedonis Pass. On leaves of *Arbutus* sp. in Portugal.

Septoria arbuti Pass. On leaves of *A. unedo* in Italy.

Septoria hirsens Sacc. Small dirty-white to dark-purple spots, with slightly raised margins, on leaves of *Arbutus* sp. in Mexico.

Septoria unedonis Rob. and Desm. Small, irregular, numerous whitish spots with purple margins on leaves of *A. canariensis* and *A. unedo* in Algeria, Albania, Yugoslavia, Portugal, France, Italy, Great Britain, and Austria.

ARDISIA. ICACOREA Ag. Ornamental shrubs.**Helminthosporium flagellatum** Yates. On *A. disticha* in the Philippines.**Phyllachora ardisiae** P. Henn. Black tar spot on leaves of *A. candolleana* in the Philippines.**Phyllosticta ardisiae** Trinch. On leaves of *A. humilis* in Italy.**Physalospora circinana** Pat. Small fuscous circular spots on leaves of *Ardisia* sp. in Brazil.**Uromyces myrsines** Diet. Brown rust pustules on leaves of *A. compressa* and *Myrsine* sp. in Brazil and Costa Rica.**ARECA.** Betel palm. See *Palmae*.**AREGELIA.** Epiphytic hothouse plants of the *Bromeliaceae*.**Lembosia bromeliacearum** Rehm. Black fruiting bodies on dark-brown circular spots on leaves of *A. compacta* and *A. cruenta* in Brazil.**ARENARIA.** SANDWORT. See, also, *Alsine*. Low herbs, suitable for borders and rockwork.**Helminthosporium nubigenum** Speg. On leaves of *A. tetraquetra* in Italy.**Laestadia arctica** Rostr. On leaves of *A. (Honckenia) peploides* in Greenland.**Peronospora campestris** Gaum. Downy mildew on lower surfaces of leaves of *A. cibota*, *A. serylifolia*, and *A. trinervia* in France, Switzerland, Belgium, Denmark, Russia, Austria, and Germany.**Peronospora honckenyaee** Syd. Downy mildew on leaves of *A. (Honckenia) peploides* in Scotland.**Puccinia modica** Holw. Brown to black powdery rust pustules on leaves of *A. alsinoides*, *A. lanuginosa*, *A. peyritschii*, and *A. reptans* in Mexico and Guatemala.**Uredo arenaricola** P. Henn. Yellow-brown to brown rust pustules on leaves of *A. diffusa* in Argentina.**Uromyces arenariae** Tranzsch. Linear dark-brown rust pustules on leaves of *A. capillaris* and *A. longifolia* in Russia.**Uromyces arenariae-grandiflora** Mayor. Leaf rust on *A. grandiflora* in Switzerland.**ARENGA.** Sugar palm. See *Palmae*.**ARGEMONE.** PRICKLE POPPY. Mexican poppy. Prickly herbs.**Peronospora indica** Güm. Downy mildew on leaves of *A. mexicana* in India. The name is occupied by *P. indica* Syd. on *Calceolaria*.**Rhizoctonia napi** West. See *Brassica*.**Septoria chelidonii** Desm. See *Chelidonium*.**ARGYREA.** SILVER CLIMBER. Ornamental climbers.**Aecidium argyreae** B. and Br. Yellow rust pustules on leaves of *A. argentea*, *A. elliptica*, *A. pomacea*, *A. populiifolia*, and *A. speciosa* in India and Ceylon.**Cercospora cordobensis** Speg. Yellow circular spots on leaves of *A. megapotamica* in Ceylon and Argentina.**Uredo argyreae** Petch. Dark-brown rust pustules on leaves of *A. tiliaefolia* in Ceylon.**ARISAEMA.** DRAGON-ROOT. JACK-IN-THE-PULPIT. Aroids with tuberous roots.**Septoria arisaemae** Petch. Leaf spots on *A. leschenaultii* in Ceylon.**ARISARUM.** Ornamental aroids.**Phyllosticta arisari** Bres. Subcircular dull-brown spots on leaves of *A. vulgare* in Portugal.**Phyllosticta arisaricola** Sacc. and Syd. Circular brown spots, with light-brown margins, on leaves of *A. vulgare* in Tunis.**Septoria arisari** (Dur. and Mont.) Sacc. Dark irregular spots on leaves of *A. simorhinum* and *A. vulgare* in France and Algeria.**Septoria arisaricola** Pat. Circular scattered or confluent whitish spots on leaves of *A. vulgare* in Tunis.**ARISTEA.** Greenhouse herbs.**Puccinia dehiscens** Syd. Brown rust pustules on leaves of *Aristea* sp. in the Union of South Africa.**ARISTOLOCHIA.** DUTCHMAN'S-PIPE. PELICAN FLOWER. Ornamental perennial vines.**Ascochyta aristolochiae** Sacc. Subcircular dark-brown to black leaf spots on *A. clematidis* in Yugoslavia and Italy.**Ascochyta aristolochicola** Höll. On capsules of *A. clematidis* in Hungary.**Ascochyta siphonis** Allesch. Subcircular to irregular black spots, becoming ashen, with black margins, on both sides of leaves of *A. siphon* in Germany.**Ascochyta versicolor** Bub. Circular to angular dark-brown spots, with raised black margins, on leaves of *A. clematidis* in Tyrol.**Cercospora olivascens** Sacc. Olive-gray spots on leaves of *A. clematidis* and *A. longa* in France and Italy.**Meliola atricapilla** Starb. Black mold on leaves of *Aristolochia* sp. in Brazil.**Mycosphaerella aristolochiae** Syd. On leaves of *A. tagala* in the Philippines.**Phyllosticta aristolochiae** F. Tassi. Round or angular to irregular dirty-white spots, with brown margins, on leaves of *A. clematidis*, *A. sempervirens*, and *A. siphon* in Italy and Germany. Reported once from the United States.**Phyllosticta siphonis** Kab. and Bub. Small angular, finally confluent, spots on upper surfaces of leaves of *A. siphon* in Bohemia.**Puccinia aristolochiae** (D. C.) Wint. Yellow to dark-brown rust pustules on leaves of *A. botta*, *A. clematidis*, *A. longa*, *A. maurorum*, *A. pallida*, *A. poicilantha*, and *A. rotunda* in Russia, Kurdistan, Anatolia, Syria, Spain, Italy, France, Austria, and Germany.**Sclerotium zeylanicum** (B. and Br.) Petch. Root and collar rot of *A. leuconeura* in Ceylon. See, also, *Thea*.**Septoria aristolochiae** Sacc. On leaves of *A. clematidis* in Italy.**ARISTOTELIA.** Evergreen trees or shrubs.**Cercospora aristoteliae** Cke. Subcircular brown spots, with red margins, on leaves of *A. racemosa* in New Zealand.**Phyllosticta aristoteliae** Speg. Indefinite angular dull-brown spots on leaves of *A. macqui* in Chile.**Septoria maqui** P. Henn. Dark-brown spots, with red margins, on leaves of *A. maqui* in Chile and Germany.**ARMERIA.** STATICE Ag. THRIFT. Sometimes called "sea pink." Perennial herbs, used as border and rock garden plants.**Gloeosporium armeriae** Allesch. Anthracnose on scapes of *A. vulgaris* (*Statice armeria*) in Greenland.**Phyllosticta armeriae** Allesch. On leaves of *A. vulgaris* in Greenland.**Septoria armeriae** Allesch. and P. Henn. On leaves of *A. (Statice) plantaginea* in Greenland, Italy, and France.**Uromyces armeriae** (Schlecht.) Lev. Yellow to dark-brown rust pustules on leaves of *A. alpina*, *A. elongata*, *A. longibracteata*, *A. martinia*, *A. plantaginea*, *A. splendens*, *A. stenophylla*, and *A. vulgaris* in Europe. Reported from California.**ARMORACIA.** See *Radicula*.

AENICA. Perennial herbs.

Entyloma calendulae (Oud.) De By. See *Calendula*.

Phyllosticta arnicae Fekl. Indefinite yellow spots on upper surfaces of leaves of *A. montana* in Switzerland. Reported from Colorado.

Septoria arnicae Fekl. Dark-brown spots on leaves of *A. montana* in Switzerland.

ARRACACIA. Umbelliferous herbs. Roots of one species eaten.

Puccinia arracacharum Lindr. Yellow to brown powdery pustules on leaves of *A. bracteata* in Ecuador and Guatemala.

Puccinia imperspicua Syd. Black rust pustules on leaves of *A. multifida* in Mexico.

ARRHENATHERUM. Oatgrass. Cultivated as a meadow grass.

Dilophia graminis (Fekl.) Sacc. See *Triticum*.

Exobasidium graminicolum Bres. A doubtful species reported as causing indefinite yellow spots on leaves of *A. elatius* and *Bromus* sp. in Germany.

Phyllosticta stomaticola Bäuml. Circular to irregular brown spots, becoming whitish, with red margins, on leaves of *A. elatius* in Hungary.

Puccinia arrhenatheri (Kleb.) Erikss. Yellow aecial stage producing witches'-broom of *Berberis heteropoda*, *B. hispanica*, and *B. vulgaris*; brown uredo and telial stages on *Arrhenatherum elatius* in Turkestan, Algeria, Spain, France, Sweden, Austria, Hungary, and Germany.

Septoria culmifida Lind. See *Pileum*.

Uromyces seseli-graminis Ed. Fisch. Brown rust pustules on leaves of *Arrhenatherum elatius*, *Melica ciliata*, *Poa* sp., and *Seseli glaucus* in Europe.

Ustilago dura App. and Gassn. Smut in inflorescences of *A. elatius* in Europe.

ARTABOTRYS. TAIL GRAPE. Woody tropical climbers.

Uredo artabotrydis Syd. Brown rust pustules on leaves of *A. odoratissima* in Formosa.

ARTEMISIA. WORMWOOD. Aromatic and bitter herbs and shrubs.

Cercospora ferruginea Fekl. On leaves of *A. vulgaris* in Japan, Poland, Bohemia, Italy, Denmark, Austria, and Germany.

Cercospora fuscescens Niessl. On stems of *A. vulgaris* in Europe.

Cercospora olivacea Oth. On leaves of *A. absinthium* in Switzerland.

Erysiphe taurica Lev. See *Althaea*.

Nematostoma artemisiae Syd. On living leaves of *A. vulgaris* in Japan.

Puccinia artemisiae-keiskeana Miura. Rust on leaves of *A. keiskeana* in Japan.

Puccinia artemisiella Syd. Black rust pustules on leaves of *A. vulgaris* in Japan, India, Russia, and Germany.

Puccinia artemisiicola Syd. Brown rust pustules on leaves and stems of *A. austriaca* and *A. campestris* in Denmark, Russia, Austria, Hungary, and Germany.

Puccinia ferruginosa Syd. Powdery, dull-brown rust pustules on small, sunken, brown leaf spots on *A. vulgaris* var. *japonica* and *Artemisia* sp. in India and Japan.

Septoria artemisiae Pass. Dull-brown spots on leaves of *A. vulgaris* in Italy.

Septoria kriegeriae Bres. On leaves of *A. vulgaris* in Saxony.

Uredo artemisiae-japonica Diet. Brown rust pustules on leaves of *A. japonica* in Japan.

ARTOCARPUS. BREADFRUIT. JACKFRUIT. Tropical fruit trees.

Catacauma microcentrum Theiss. and Syd. See *Ficus*.

Cercospora artocarpi Syd. On leaves of *A. communis* (*A. incisa*) in the Philippines.

Corticium salmonicolor B. and Br. See *Citrus*.

Diplodia artocarpi Sacc. Causes twig die-back and fruit rot of *Artocarpus* sp. in Java and the Philippines. Probably not distinct from *D. cacaocicola*.

Fomes lamaensis Murr. See *Hevea*.

Fomes lignosus Klotzsch. See *Hevea*.

Marchalia constellata (B. and Br.) Sacc. Circular, flat, black stromata on leaves of *Artocarpus* sp. in the Philippines and *A. integra* (*A. integrifolia*) in Ceylon.

Phyllosticta sp. On leaves of *A. communis* in India.

Phyllosticta (?) *artocarpi* Speg. Circular, determinate white spots, with broad dark margins, on leaves of *A. integra* (*A. integrifolia*) in Argentina.

Phytophthora sp. Causes fruit rot of *A. incisa* in Ceylon.

Phytophthora faberi Maubl. See *Theobroma*.

Rhizopus artocarpi Rac. Causes a soft rot of fruit and male inflorescences of *A. integra* in the Philippines.

Rosellinia pepo. Pat. See *Citrus*.

Septoria artocarpi Cke. Circular, brown sunken spots on leaves of *A. integrifolia* in India.

Septoria eburnea v. Hoch. Circular to angular or irregular leaf spots on *A. communis* in Samoa.

Sphaerostilbe repens B. and Br. See *Hevea*.

Uredo artocarpi B. and Br. (*Physopella artocarpi* [B. and Br.] Arth.) Numerous dull-brown rust pustules covering lower surfaces of leaves of *A. camansi*, *A. communis*, *A. laccucha*, and *Castilla-elastica* in Ceylon, Cuba, and Porto Rico.

ARUM. Variegated leafed, herbaceous aroids.

Ascochyta arophila Bub. Circular to elliptical brown spots, becoming white at the centers, on leaves of *A. italicum* in Montenegro.

Colletotrichum montemartirii Togn. Anthracnose on *A. italicum* in Italy.

Phyllosticta aricola Bub. Brown, circular, or elongate leaf spots, becoming white at the centers, on *A. italicum* in central Europe.

Phyllosticta scrophulariae-bosniacae Bub. On leaves of *A. italicum* and *A. bosnia* in Central Europe.

Phyllosticta tuzsonii Bub. Elongate to irregular yellow spots with white centers, on leaves of *A. italicum* in Hungary.

Ramularia ari Fautr. Gray leaf spots on *A. italicum* and *A. maculatum* in Spain, France, and Delmatia.

Septoria ari Desm. Subcircular brown spots on leaves of *A. italicum* and *A. maculatum* in Algeria, Yugoslavia, Bulgaria, France, and Italy.

Ustilago plumbea Rostr. A smut producing black spore masses on leaves and petioles of *A. maculatum* in Denmark.

ARUNCUS. GOATSBEARD. Ornamental, tall perennial herbs.

Ascochyta arunci Sacc. On stems of *A. sylvester* in Italy.

Colletotrichum exiguum Penz. and Sacc. Numerous, small, brown leaf spots on *A. sylvester* in Italy and Austria.

Ochropsora ariae (Fekl.) Syd. See *Sorbus*.

Phyllosticta aruncina Sacc. Small, grayish-white leaf spots on *A. sylvester* in Italy.

Phyllosticta griseo-fusca Bub. Small, angular to confluent leaf spots dull green, then brown above, gray beneath on *A. sylvester* in Bohemia.

Ramularia spiraeae-arunci (Sacc.) Allesch. On leaves of *A. sylvester* in Switzerland and Austria.

Septoria arunci Pass. Gray-white leaf spots with dark-red margins on *A. sylvester* in Italy.

ARUNDINA. See Orchidaceae.

ARUNDINARIA. See Bambuseae.

ARUNDO. GIANT REED. Tall perennial grasses.

Apospora montagni Sacc. Linear black stromata on leaves of *A. donax* and *A. mauritanica* in Algiers, Portugal, and France.

Ascochyta donacina Sacc. On culms of *A. donax* in Italy and France.

Phyllachora atrofurans Rehm. Black stromata on leaves of *A. cannaeformis* in the Philippines.

Phyllachora cyperi Rehm. var. **donacis** Berl. and F. Sacc. "Tar spot" disease on culms of *A. donax* in India and Portugal.

Phyllachora donacina Rehm. Black, convex, orbicular or ellipsoid stromata in parallel series on leaves of *A. cannaeformis* in the Philippines.

Physalospora festucae (Lib.) Sacc. See Andropogon.

Puccinia isiacae (Thum.) Wint. Rust-brown to dark-brown rust pustules on leaves of *A. isiacae* and *Phragmites communis* in Egypt, Asia Minor, and Russia. Aecia may possibly occur on *Raphanus sativus*.

Puccinia torosa Thum. Brown rust pustules on leaves of *A. donax* in the Union of South Africa.

Septoria donacina Sacc. Indefinite spots on culms of *A. donax* in Italy.

Septoria donacis Pass. Small dry spots on leaves of *A. donax* in Dalmatia, Italy, France, and Portugal.

Septoria oxyspora Penz. and Sacc. See Anthoxanthum.

ASARUM. WILDGINGER. Low, nearly stemless herbs.

Septoria asari Sacc. Circular to angular whitish spots, with black borders, on leaves of *A. europaeum* in Italy.

Septoria asaricola Allesch. Subcircular to angular, brown then gray spots, often confluent, on leaves of *A. europaeum* in Russia and Bavaria.

ASCLEPIAS. MILKWEED. BUTTERFLY WEED. Perennial, milky-juiced herbs.

Cercospora asclepiadis P. Henn. Numerous circular leaf spots on *Asclepias* sp. in Brazil.

Cronartium flaccidum (Alb. and Schw.) Wint. See Paonia.

Mycosphaerella asclepiadis Siem. On leaves of *A. incarnata* in Caucasia.

Puccinia concrescens E. and E. Leaf rust on *A. curassavica* and *A. nitida* in Porto Rico, Cuba, Bahamas, and South America.

Septoria schlechteriana P. Henn. Pale leaf spots, with dark brown margins, on *A. fruticosa* in the Union of South Africa.

Septoria syriacae Kab. and Bub. Numerous circular to angular often confluent spots on leaves of *A. syriaca* in Bohemia.

ASPARAGUS. Herbaceous perennials and tender woody shrubs and vines. One species used for food.

Aecidium schlechterianum P. Henn. Round yellow spots in which yellow rust pustules appear, are formed on the leaves of *A. medeoloides* in south Africa.

Botryosphaeria dispersa De Not. On *A. officinalis* in Europe.

Cercospora asparagi Sacc. Dirty-white to gray spots on false leaves and branches of *A. officinalis* in Brazil, Cuba, Argentina, and Italy.

Cercosporina asparagicola Speg. On *A. officinalis* in Argentina.

Colletotrichum sp. Anthracnose on stems of *Asparagus* sp. in Brazil.

Coniothyrium sp. Produces stem cankers on stems of *Asparagus* sp. in Brazil.

Hysterostomella tenella Syd. On leaves and stems of *A. stenotus* and *Asparagus* sp. in the Union of South Africa.

Leptosphaeria circinans (Fekl.) Sacc. On roots and stems of *A. officinalis*. *Beta vulgaris*, *Medicago sativa*, *Phaseolus* sp., *Solanum tuberosum*, *Trifolium pratensis*, and *T. repens* in Italy, Russia, and Germany.

Leptosphaeria indica Syd. and Butl. On *Asparagus* sp. in India.

Melanospora asparagi Arnaud. Attacks the base of the stalks of *A. officinalis* in France, preventing further development.

Ophiobolus compar Karst. Purple sunken spots, in which small, black, fruiting bodies appear, are formed on stems of *A. officinalis* in Finland.

Puccinia asparagi-lucidi Diet. Chestnut-colored rust pustules, becoming dark brown to black on both leaf surfaces and on stems of *A. lucidus* and *A. schoberioides* in Japan.

Puccinia phyllostachiae Cke. This rust produces powdery brown pustules on the lower leaf surfaces and on stems of *A. falcatus* in Natal and Ceylon.

Zopfia variopora (Boud.) Arnaud (*Z. rhizophila* Rabenh.) On *A. acutifolius*, *A. horridus*, and *A. officinalis* in France, Italy, and Algeria.

ASPERULA. WOODRUFF.

Mycosphaerella asperula Roum. and Fautr. On leaves of *A. tinctoria* in Esthonia.

Peronospora calotheca De By. Downy mildew on leaves of *A. odorata* and *A. tinctoria* in central and north Europe.

Phyllosticta asperulae Sacc. Indefinite, yellow spots on leaves of *A. odorata* in France and Hungary.

Pseudopeziza autumnalis (Fr.) Karst. See Galium.

Puccinia asperulae-cynanchicae Wurtb. Rust on leaves of *A. cynanchica* and *A. odorata* in Spain, Yugoslavia, Switzerland, and Germany.

Puccinia asperulae-odoratae Wurtb. On leaves of *A. odorata* in Denmark, Sweden, Switzerland, Yugoslavia, and Japan.

Puccinia asperulina (Juel) Lagh. Golden-yellow rust pustules, becoming brown, causing distortion of leaves of *A. ciliata* and *A. tinctoria* in Sweden, Russia, and Hungary.

Puccinia helvetica Schroet. Brown rust pustules on leaves of *A. taurina* in Switzerland, Italy, and Hungary.

Puccinia oliganthae McAlp. Brown to dark-brown rust pustules on leaves and stems of *A. oligantha* in Victoria (Australia).

Puccinia spillogena Lindr. Yellow-brown to dark-brown rust pustules on leaves of *A. mollugo* in Persia.

Septoria asperulae Bauml. and var. **italica** Mass. Irregular gray spots with fuscous margins, soon involving the entire leaf, on *A. odorata* and *A. taurina* in Denmark, Italy, and Hungary.

Septoria asperulae-arensis Maire. On leaves of *A. creensis* in French north Africa.

Septoria asperulae-taurinae Bub. Dirty-white spots on upper surfaces of leaves of *A. taurina* in Hungary.

Thekopsora guttata (Schroet.) Syd. Yellow to dark brown rust pustules on leaves of *A. galioides*, *A. odorata*, *Galium aristatum*, *G. hercynicum*, *G. mollugo*, *G. rotundifolium*, *G. saxatile*, *G. schultesium*, *G. silvestre*, *G. sylvestris*, *G. spurium*, *G. uliginosum*, *G. terum*, and *Sherardia creensis* in Europe. Reported from Oregon.

Uromyces asperulae McAlp. Cinnamon-brown to dark-brown rust pustules on leaves and stems of *A. oligantha* in Australia.

- ASPHODELINE.** **WJACOB'S ROD.** Hardy, herbaceous plants.
Septoria asphodelines Syd. On leaves of *Asphodeline taurica* in Bulgaria.
ASPHODELUS. **ASPHODEL.** Hardy, herbaceous, stemless plants.
Cladochytrium asphodeli De By. (*Physoderma asphodeli* [De By.] Vesterg.) On leaves of *Asphodelus* sp. causing dark brown or black irregular spots, in Tunis and Algeria.
Diplodia uncinjensis Bub. On stems of *A. microcarpus* in Montenegro.
Phoma asphodeli Sacc. (*P. asphodelina* Thuem.) On stems and leaves of *Asphodelus* sp. in Spain.
Phomopsis asphodelina (Thuem.) Trav. and Sp. On scapes and capsules of *A. microcarpus* in Spain.
Phyllosticta caulicola Pat. Broad, brown spots with black margins are produced on the stems of *A. microcarpus* in Algeria.
Puccinia asphodeli Moug. Yellow to yellow-brown rust pustules, becoming darker on both surfaces of the leaves of *Anthericum ramosum*, *Asphodelus albus*, *A. cerasiferus*, *A. microcarpus*, *A. occidentalis*, *A. ramosus*, *A. subulpinus*, and *A. tenuifolius* in France, Italy, Portugal, Spain, Greece, Yugoslavia, Syria, Algeria, Canary Islands, Corsica, and Balearic Islands.
Puccinia barbeyi (Roum.) P. Magn. Sunken elongate spots occur on both leaf surfaces, and on the stems, in which yellow and brown rust pustules appear. On leaves of *A. fistulosus*, *A. microcarpus*, and *A. tenuifolius* in France, Greece, Syria, Algeria, Spain, and India.
Puccinia unanum Gz. Frag. Yellow to brown rust sori on leaves and scapes of *A. albus* in Spain.
Septoria asphodelina Sacc. On leaves of *A. albus*, *A. fistulosus* and *A. microcarpus* in Spain and Belgium.
Septoria asphodeli-ramosi Pat. Ashen elongate spots on stems of *A. ramosus* in Tunis.
ASPIDISTR. Florist's plant grown for its stiff, shining foliage which is often variegated.
Ascochyta aspidistriae Mass. Large, irregular, bleached areas on leaves of *A. lurida* in Great Britain. Reported from Minnesota.
Phyllosticta aspidistriae Oud. Large, rust-colored spots on leaves of *A. elatior* in Holland.
Septoria transversalis Sacc. On leaves of *A. elatior* in Italy.
ASPIDIUM. DRYOPTERIS Ag. WOOD FERN.
Exoascus cornu-cervi Sadeb. On leaves of *A. aristatum* in New Caledonia.
Exoascus fuscus (Gieslen.) Sacc. and Syd. Produces fleshy, dull-brown galls on leaves of *A. pallidum* in Sicily and Albania.
Exoascus verstergrenii (Gieslen.) Sacc. and Syd. Brown galls on both surfaces of leaves of *A. filix-mas* in the Swedish Baltic Islands, and Switzerland.
Exobasidium brevieri Boud. On leaves of *A. filix-mas* in Estonia.
Gleosporium filicinum Rostk. Anthracnose of leaves of *A. filix-mas* in Sweden.
Hyalopora filicum Diet. See *Asplenium*.
Milesina carpatica A. Wrób. Yellow rust pustules on leaves of *A. filix-mas* in Austria.
Milesina kriegeriana P. Magn. Yellow rust pustules on leaves of *A. aculeatum*, *A. dilatatum*, *A. filix-mas* and *A. spinulosum* in Dalmatia, Finland, Denmark, Austria, Hungary and Germany.
Milesina vogesiaca Syd. Yellow rust pustules on leaves of *A. lobatum* in Austria and Germany.
Mycosphaerella aquilina (Fr.) Schroet. and f. *aspidiorum* Sacc. On leaves of *A. filix-mas* and *Asplenium adnigrum* in Switzerland and Estonia.
Mycosphaerella thelypteris Syd. On *A. thelypteris* in Germany.
Trabutella filicina (Sacc. and Sc.) Theiss. and Syd. Black stromata on leaves of *A. lonchitis* in Alaska.
ASPENIUM. SPLEENWORT. Ferns.
Aphelenchus olesistus Ritz. Bos. See *Begonia*.
Cercospora asplenii Jaap. On leaves of *A. trichomanes* in Dalmatia.
Exobasidium brevieri Boud. Small, irregular white spots, often confluent, on leaves of *A. filix-femina* in France.
Hyalopora asplenii-wichuriae Diet. Brown rust pustules on indefinite dirty-brown spots on leaves of *A. wichuriae* in Japan.
Hyalopora filicum Diet. Yellow rust pustules on leaves of *A. decursivae-pinnatum*, *A. japonicum*, *A. patens*, *Athyrium nipponicum*, *Dryopteris laza*, and *Pteris semipinnata* in Japan.
Milesina feurichii P. Magn. Yellow rust pustules on leaf blades and petioles of *A. septentrionalis* in Finland, Switzerland, and Germany.
Milesina magnusiana Jaap. Rust pustules on yellow-brown spots on leaves of *A. nigrum* in Dalmatia, Italy, and Corsica.
Milesina murariae (P. Magn.) Grove. Yellow rust pustules on leaves of *A. ruta-muraria* in Scotland, Switzerland, Austria, and Germany.
Mycosphaerella asplenii (Auersw.) Lind. On leaves of *A. septentrionale* in Europe.
Mycosphaerella filicum (Desm.) Starb. See *Aspidium*.
Phyllosticta asplenii Jaap. On leaves of *A. ruta* in Switzerland.
Ramularia asplenii Jaap. On leaves of *A. ruta* in Switzerland and Germany.
Sclerodothis confuens (Starb.) Theiss. and Syd. Black stromata on gray leaf spots on *Asplenium* sp. in Sweden.
ASSONIA. See *Dombeya*.
ASTER. MICHAELMAS DAISY. Perennial herbs.
Aecidium microsporium Diet. Rust on leaves of *A. divaricatus* in Brazil.
Bacillus asteracearum Pava. Small, scattered, ochre-colored spots, first appear on the lower leaves, increasing in size and spreading up the stalk to the flowers which, together with the leaves, turn brown and wither on *Aster* spp. in Italy.
Cephalosporium asteris Dow. A serious wilt disease on cultivated *Aster* in Great Britain is attributed to this fungus. Infected plants show bright-lemon or orange-colored mottling of the lower leaves which finally become brown and shrivelled. An excessive suckering is also in evidence.
Cercospora asterina Speg. Ashen-brown spots, with darker margins, on leaves of *A. unifolius* in Argentina.
Coleosporium asterum (Diet.) Syd. Yellow rust pustules on leaves of *A. glenhi*, *A. hispidus*, *A. pinnatifidus*, *A. scaber*, *A. tataricus*, *A. trinervius*, *Boltonia cantoniensis*, *B. incisa*, *B. indica*, *Callistephus chinensis*, *A. sinensis*, and *Pinus densiflora* in China, Formosa, and Japan. Reported from California.
Didymaria asteris Oud. Broad, irregular white spots on leaves of *Aster* sp. in Holland.
Entyloma calendulae (Oud.) De By. See *Calendula*.
Puccinia asteris-alpini Syd. Black rust pustules on small brown sunken spots on leaves of *A. alpinus* in Switzerland.
Ramularia asteris-tripolii Jaap. Subcircular to oblong, often indefinite, yellow, finally gray or brown spots on leaves of *A. tripolium* in Denmark.
Septoria tatarica Syd. Greenish-gray, circular then confluent, leaf spots on *A. tataricum* in Japan.

ASTILBE. Ornamental perennial herbs.

Puccinostele darkiana (Bard.) Diet. Yellow rust pustules on circular or elongate spots, on leaves and petioles of *A. congesta*, *A. philippinensis*, *A. rivularis*, *A. sinensis*, and *A. thunbergii* in India, Japan, and the Philippines.

Puccinostele mandshurica Diet. Powdery, yellow rust pustules on lower surfaces of leaves and on petioles of *A. chinensis* in Japan, Korea, and Manchuria.

Septoria hotteiae F. Tassi. On leaves of *A. japonica* in Italy.

Triphragmium nishidanum Diet. Chestnut-colored rust pustules on leaves of *A. chinensis* var. *albiflora* and *A. sinensis* in Japan.

ASTRAGALUS. MILKWEED. Hardy herbs or subshrubs.

Cylindrosporium astragalii E. Rostr. On leaves of *A. oroboides* in Norway.

Erysiphe taurica Lev. See *Althaea*.

Eryachora tragacanthae (Lev.) Theiss. and Syd. Black, irregular stromata on thorax and leaves of *A. tragacantha* in Persia.

Microsphaera astragalii (DC.) Trev. Powdery mildew on leaves of *A. cicer*, *A. glycyphyllos*, and *A. onobrychis* in Europe.

Mycosphaerella magnusiana Jaap. On leaves of *A. alpinus* in Austria.

Oularia sphaeroidea Sacc. On leaves of *A. sinicus* in Japan.

Oularia tuberculiformis v. Hoeh. Small, often confluent, dirty-yellow leaf spots on *A. cicer* in Yugoslavia and Austria.

Peronospora astragalina Syd. Downy mildew on leaves of *A. alpinus* and *A. oroboides* in Norway, Lapland and Austria.

Phylachora melaena (Rabh.) Sacc. Black stromata on brown definite leaf spots on *A. glutaceus* in Kurdistan.

Phyllosticta astragalicola Mass. Olivaceous spots, with brown margins, on leaves of *A. glycyphylloides* in Italy.

Phyllosticta desertorum Sacc. Subcircular small brown spots on both surfaces of leaves of *A. alopecurus* in Siberia.

Phyllosticta exscapi Höll. On leaves of *A. exscapus* in Hungary.

Physalosporina astragalina (Rehm) Wor. On leaves of *A. cicer* in Russia and Germany.

Physalosporina obscura (Juel.) Wer. On leaves of *A. alpinus* and *A. arenarius* in Denmark and Russia.

Placosphaerella tragacanthae Pat. Black, rough, fruiting areas on leaves of *A. fontanesius* and *A. tragacantha* in Persia and Arabia.

Polystigma obscurum Juel. Stromata occupying most of leaf surface, white beneath, yellowish-red, then gray-brown above. On *A. alpinus* and *A. oroboides* in Sweden and Norway.

Septoria astragalii Desm. On leaves of *A. alpinus* and *A. glycyphyllos* in Spain, Denmark, Russia, Bohemia, and Austria.

Septoria henningsiana Wint. On leaves of *A. onobrychis* in Asia Minor and Yugoslavia.

Septoria serebriankowii Sacc. Causes yellowing of leaves of *A. onobrychis* in Russia.

Uromyces carneus (Nees.) Har. Brown, powdery, rust pustules covering entire leaf surfaces of *A. alpinus* and *A. australis* in Norway, Sweden, Switzerland, and Austria.

Uromyces gordianus Bub. Brown rust pustules on leaves of *A. exscapus* in Switzerland, Austria, Hungary and Germany.

Uromyces patagonicus Speg. Cinnamon-brown rust pustules on leaves of *Astragalus* sp. in Patagonia.

Uromyces persicus Syd. Dark-brown rust pustules on leaves of *A. remotifolius* in Persia.

Uromyces splendens Wott. Yellow to brown rust pustules on leaves of *A. oroboides* in Norway.

ASTELANTIA. MASTERWORT. Perennial garden plants.

Ascochyta astrantiae Roum. On leaves of *A. major* in France.

Fabraea astrantiae (Ces.) Rehm. On *Astrantia* sp. in Switzerland.

Phyllosticta astrantiaeola Gz. Frag. On leaves of *A. major* in Spain.

Pseudopeziza sanculae-astrantiae Niessl. On leaves of *A. major* in Germany.

Puccinia astrantiae Kalchb. Dark-brown rust pustules on leaves and petioles of *A. major* and *A. minor* in Portugal, France, Belgium, Austria, Hungary and Germany.

Puccinia astrantiae-rivipari Semad. Rust pustules on leaves of *A. minor* and *Polygonum viviparum* in Switzerland.

Ramularia oreophila Sacc. Angular, dull yellow-brown spots on leaves of *A. major* and *A. minor* in Italy, Spain, Switzerland, Russia and Austria.

ASTROCARYUM. See *Palmae*.**ASTYSTASIA.** Hothouse evergreen herbs and shrubs.

Puccinia phaeosticta Pat. and Har. Brown rust pustules on circular brown spots on leaves of *Asystasia* sp. in Indo-China.

Uredo asystasiae P. Henn. Brown rust pustules on yellow spots on leaves of *A. gangetica* in east Africa.

ATALANTIA. Small, usually spiny trees and shrubs related to *Citrus*.

Bacterium citri Hass. See *Citrus*.

Phyllosticta disciformis Penz. See *Citrus*.

ATAMOSCO. See *Zephyranthes*.**ATRIPLEX.** SALTBUSH. Orach. Ornamental shrubs, garden vegetables and many weedy species.

Ascochyta atriplicis Diet. Subcircular, often confluent, dull-yellow spots, with pale centers and dark margins, on leaves of *A. hastata* in Germany.

Ascochyta chenopodii Rostk. Circular, pale-brown spots, with yellow margins, on leaves of *A. bonariensis*, *A. littoralis*, *Chenopodium album*, *C. glaucum*, and *Spinacia oleracea* in Denmark, China, Russia, and Germany.

Cercospora dubia (Riess.) Wt. On leaves of *A. hastata*, *A. hortensis*, *A. laciniata*, *A. littoralis*, *A. nitens*, and *A. patula* in France, Bohemia, Russia, Denmark, and Germany. Reported from Montana.

Peronospora minor (Casp.) Güm. Downy mildew on leaves of *A. nitens*, *A. patula* and *A. roseum* in England and north Europe.

Phyllosticta atriplicicola Speg. Circular, ashen-colored spots on leaves of *A. hastata* in Argentina.

Phyllosticta halophila Speg. Circular, often confluent, white spots, with purplish margins, on leaves of *A. hastata* in Argentina.

Ramularia dubia Wint. Spots on leaves of *A. hastata*, *A. nitens* and *A. patula* in Russia and Germany.

Septoria atriplicis (West.) Fekl. Greenish or dirty yellow to whitish leaf spots on *A. patula*, *Chenopodium murale* and *C. cultaria* in Siberia, Tunis, Bulgaria, France, Spain, Italy, Belgium, Germany, and Central America. Reported from a few localities in the United States.

Uromyces atriplicis McAlp. Pale-brown to dark-brown rust pustules on leaves of *A. semibaccata* in Victoria.

ATRIPLEX—Continued.

Urophylletis pulposa (Wallr.) Schroet. Swellings on stems and leaves of *A. hastata*, *A. patula*, *Beta vulgaris*, *Chenopodium album*, *C. cubicum*, *C. glaucum*, *C. rubrum* and *C. urticum* in Argentina, Denmark, Siberia, and Germany.

ATROPA. **BELLADONNA**. Medicinal herbs.

Ascochyta atropae Bres. Pale-brown spots with darker margins on leaves of *A. belladonna* in Germany.

Phyllosticta atropa F. Tassi. Circular, white spots, with yellow borders, on leaves of *A. belladonna* in Italy.

Phytophthora erythroseptica Pethyb. See Solanum.

Puccinia atropae Mont. Yellow to black rust pustules on leaves of *A. aristata* in the Canary Islands.

Ramularia atropae Allesch. Irregular, dull-brown spots on both leaf surfaces of *A. belladonna* in Switzerland and Germany.

AUCUBA. Ornamental evergreen shrubs, often with variegated leaves, and with bright scarlet fruit.

Ascochyta aucubae Sacc. and Speg. On leaves of *A. japonica* in Italy and France. Var. *brunaudiana* Sacc. differs in the more definite spots with black margins.

Ascochyta aucubicola Wint. Irregular, whitish spots with fuscous margins on leaves of *A. japonica* in Portugal.

Colletotrichum pollacci Magn. Anthracnose on *A. japonica* in Japan.

Phyllosticta ampla P. Brun. Large, irregular brownish-green or gray-green spots on leaves of *A. japonica* in France.

Phyllosticta aucubae Sacc. and Speg. On leaves of *A. japonica* in Italy.

Phyllosticta aucubicola Sacc. Pale, indefinite spots on leaves of *A. japonica* in Denmark and Italy.

Phyllosticta indica Roum. and Karst. Large, whitish spots, with dull-brown margins, on leaves of *A. japonica* and *A. lanceolata* in Indo-China.

Pleospora infectoria Fekl. var. *aucubicola* Mont. Black patches appear on tips and along edges of the leaves. These areas increase in size, soon involving the entire blade which withers and dies. New leaves are attacked as they unfold. On *A. japonica* in Italy.

Septoria aucubae West. Large, irregular, pale brown spots with darker margins on leaves of *A. japonica* in Italy and Belgium.

Sphaerulina aucubae Shir. and Har. On leaves of *A. japonica* in Japan.

AURICULA. See Primula.**AVENA**. OATS. Annual grasses cultivated for grain and forage.

Acremonella verrucosa Togn. Producing black spots on culms and sheaths of *A. sativa* and *Triticum* sp. in Italy.

Ascochyta graminicola Sacc. var. *ciliolata* Sacc. On *A. filifolia* and *A. pratensis* in Spain.

Cladochytrium graminis Buesgen. See Festuca.

Heterosporium cerealinum Oud. On leaves of *Avena* sp. in Holland and Belgium.

Leptosphaeria nigrans (Desm.) Ces. and De N. See Aira.

Metasphaeria avenae (Auersw.) Sacc. On leaves of *Avena* sp. and *Calamagrostis* sp. in France and Germany.

Ophiobolus cariceti (B. and Br.) Sacc. See Triticum.

Puccinia pratensis Blytt. Powdery, brown pustules on leaves of *A. pratensis* in Sweden and Norway.

Rhizoctonia napi West. See Brassica.

Sclerospora macrospora Sacc. See Triticum.

Septoria planiuscula Diet. Leaf spots on *A. planiuscula* in Germany.

Tylenchus hordei Schoeyen. This nematode is said to cause small galls on the roots of *A. sativa*, *Elymus arenarius*, *Hordium vulgare* and *Poa pratensis* in northern Europe. Possible should be referred to *Heterodera radiculola*.

Tylenchus tritici Bast. See Triticum.

AVERRHOA. Carambola. Tropical fruit trees.

Cercospora averrhiae Petch. Circular, white leaf spots with purple-red margins, on *A. carambola* in Ceylon.

Cercospora averrhoi Welles. Irregular yellow lesions on leaves 3 to 5 millimeters in diameter, which may coalesce to form large irregular spots. Center of diseased areas die, becoming grayish-brown, surrounded by band of yellow tissue. Black lesions form on the fruit. On *A. carambola* in the Philippines. Probably not distinct from the preceding species.

Phyllosticta sp. On leaves of *A. bilimbi* in Cuba.

AZALEA. See Rhododendron. The diseases of *Azalea* and *Rhododendron* are interchangeable to such an extent that it does not appear feasible to separate them, just as the host genera themselves are botanically inseparable.**AZARA**. Small, ornamental evergreen trees or shrubs.

Uredo azarae Neg. Golden-yellow rust pustules on leaves of *A. integrifolia* in Chile.

BACKHOUSIA. Evergreen shrubs or trees.

Ascochyta apiospora Cke. and Mass. Brown spots with purple borders on leaves of *Backhousia* sp. and *Myrtus* sp. in Australia.

BACTRIS. See Palmae.**BAHIA**. Herbaceous perennials.

Phragmodothis minutissima (Starb.) Theiss. and Syd. On stems of *Bahia* sp. in Brazil.

BAMBUSA. See Bambusae.**BAMBUSEAE**. BAMBOO. Large, ornamental, and economic perennial grasses. It has seemed desirable to consider under one heading the diseases of the various genera comprising the tribe *Bambuseae* of the *Gramineae*, not only since the diseases are, to a considerable extent, interchangeable, but because the host plants themselves are so difficult of determination that in many cases disease reports give only the common name bamboo, making exact host determination impossible, and a separation of the diseases, therefore, out of the question.

Adiclosporium take Miy. Produces witches'-brooms of *Phyllostachys* sp. in China and Japan.

Ascochyta arundinariae Gz. Frag. On leaves of *Arundinaria nitida* in Spain.

Ascopolyporus polychrous A. Moell. On living twigs of *Bambusa* sp. in Brazil.

Balanisia chusqueola P. Henn. Black, carbonaceous crusts, about one centimeter long, surrounding stems of *Chusquea* sp. in Costa Rica.

Balanisia claviceps Speg. On *Bambusa* sp. in Costa Rica and Brazil.

Balanisia regularis Moell. Producing witches'-broom effects on *Bambusa* sp. in Brazil.

Cenangella bambusicola Rick. White, then greenish-yellow, sessile fruiting bodies on branches of *Bambusa* sp. in Brazil.

Chaetophoma pellicula Sacc. and Syd. Black effused areas on leaves of *Chusquea* sp. in Chile.

Coccidiella arundinariae Hara. Fleshy, disk-shaped, scattered fruiting bodies on leaves of *Arundinaria simoni* and *Sasa borealis* in Japan.

BAMBUSEAE—Continued.

- Colletotrichum septorioides** Sacc. On culms of *Bambusa* sp. in China.
- Cylindrosporium bambusae** Miy. and Hara. On leaves of *Bambusa* sp. and *Phyllostachys bambusoides* in Japan.
- Dothidella gigantochloae** (Rehm.) Theiss. and Syd. On stems and leaves of *Gigantochloa* sp., *G. levis*, and *G. scribneriana* in the Philippines. The exact position of this fungus is uncertain.
- Endodothella bambusae** (Rabh.) Theiss. and Syd. Linear black stromata on rust-brown leaf spots on *Bambusa spinosa* and *Bambusa* sp. in India.
- Epichloe bambusae** Pat. Fleshy, black, effused (8 to 10 centimeters) stromata on leaves of *Bambusa* sp. and *Sasa spiculosa* in Japan, Java, and India.
- Epichloe sasae** Hara. Causes witches' brooms on *Sasa spiculosa* in Japan.
- Fomes lignosus** Klotzsch. See Hevea.
- Gilletiella chusqueae** (Pat.) Sacc. and Syd. Small, black, circular or elliptical stromata on brown leaf spots on *Chusquea* sp. in Ecuador.
- Gloeosporium sphacelatum** Hara. Anthracnose on *Phyllostachys reticulata* in Japan.
- Gloniella chusqueae** P. Henn. Leaf spot on *Chusquea* sp. in Chile.
- Gloniella chusqueicola** P. Henn. Black, erumpent fungus pustules on young twigs of *Chusquea* sp. in Chile.
- Guignardia bambusae** Miy. and Hara. On leaves of *Bambusa* sp. and *Phyllostachys* sp. in Japan.
- Guignardia blumeanae** (Sacc.) Yates. On leaves of *Bambusa blumeana* and *B. spinosa* in the Philippines.
- Helminthosporium bdellomorphum** Speg. On culms of *Chusquea valdiviensis* in Chile.
- Helminthosporium blumeanum** Sacc. On leaves of *Bambusa blumeana* and *B. spinosa* in the Philippines.
- Homostema fusispora** Syd. Small, yellow, indefinite spots on leaves of *Bambusa* sp. in the Philippines.
- Hypocrepis phyllostachydis** (Syd.) M. and H. In the ovaries of *Phyllostachys puberula* in Japan.
- Kusanobotrys bambusae** P. Henn. Round, black, often confluent, spots on upper surfaces of leaves of *Bambusa veticii* in Japan.
- Leptosphaeria schneideriana** Rick. Small black spots on leaves of *Bambusa* sp. in Brazil.
- Loculistroma bambusae** Pat. and Charles. Sessile, dark-green or black fruiting bodies, resembling sclerotia of *Claviceps purpurea*, produced at the nodes. Diseased branches show a witches'-broom effect due to shortening of internodes. The fungus attacks *Phyllostachys* sp. in China.
- Malmomyces pulchellus** Starb. On young branches of *Bambusa blumeana* and *B. spinosa* in Brazil and the Philippines.
- Melasmia phyllostachydis** Hara. On leaves of *Phyllostachys edulis* and *Phyllostachys* sp. in Japan.
- Miyoshia fusispora** Kawamura. Elliptical brown spots on stems of *Arundinaria narihira* in Japan.
- Mosaic.** A typical Mosaic disease has been found on *Phyllostachys* plants imported from China. The disease is related to, and may be identical with, the sugar-cane mosaic.
- Mycosphaerella bambusae** Pat. On culms of *Phyllostachys reticulata* and *Bambusa* sp. in Japan and Venezuela.
- Mycosphaerella bambusicola** Miy. and Hara. Circular, elliptical, or irregular dull-brown spots, with black borders, often coalescing and causing death of surrounding areas, on leaves of *Phyllostachys bambusoides* and *P. puberula* in Japan and India.
- Mycosphaerella bambusifolia** Miy. and Hara. Circular to irregular, fuscous, often coalescent, leaf spots on *Phyllostachys bambusoides*, *P. puberula*, and *Phyllostachys* sp. in Japan.
- Ophiodothis linearis** Rehm. Black stromata formed on lower leaf surfaces which die and split open. On *Chusquea* sp. in Brazil.
- Phallosphaeria bambusae** Miy. and Hara. Brown, later grayish, spots along veins of leaves, which are ultimately killed. On *Arundinaria simonii*, *Bambusa* sp., and *Sasa paniculata* in Japan.
- Phyllisella graminicola** v. Hoehn. Black stromata on leaves of *Chusquea wetsteinii* in Brazil.
- Phyllachora bambusae** Syd. and Butl. Circular, black, slightly raised stromata on linear brown leaf spots on *Bambusa spinosa* and *B. blumeana* in the Philippines and Malabar.
- Phyllachora bambusina** Speg. Black, linear, stromata on leaves of *Bambusa* sp. in Brazil.
- Phyllachora caespiticia** Theiss. and Syd. Black stromata on upper surfaces of leaves of *Bambusa* sp. in Indo-China.
- Phyllachora chusqueae** P. Henn. and Lind. Black, linear, cushion-shaped stromata on leaves of *Chusquea quila* in Chile.
- Phyllachora eximia** Syd. Black convex stromata on sunken yellow or strawberry colored spots on leaves of *Arundinaria alpina* in Tanganyika.
- Phyllachora gracilis** Speg. Black stromata on lower surfaces of leaves of *Bambusa* sp. in Brazil.
- Phyllachora maculans** (Karst.) Theiss. and Syd. On leaves of *Bambusa* sp. in Indo-China.
- Phyllachora malabarensis** Syd. and Butl. Black stromata on yellowish-white spots on upper surfaces of leaves of *Bambusa* sp. in India.
- Phyllachora megastroma** Pat. On leaves of *Bambusa* sp. in Congo.
- Phyllachora orbicula** Rehm. Black stromata on small, round, yellowish spots on leaves of *Bambusa blumeana* and *B. spinosa* in China and the Philippines.
- Phyllachora phyllostachydis** Hara. Black stromata on leaves of *Phyllostachys reticulata* and *P. shiraiana* in Japan.
- Phyllachora shiraiana** Syd. Oblong to irregular black stromata on upper surfaces of leaves of *Arundinaria japonica*, *A. simonii*, *Bambusa* sp., *Phyllostachys bambusoides*, *Sasa paniculata*, *S. spiculosa*, *Schizostachyum acutiflorum*, and *S. diffusum* in India, the Philippines, and Japan.
- Phyllachora sinensis** Sacc. On *Bambusa* sp. in China.
- Phyllachora tjankorreh** Rac. Black circular, slightly convex, stromata on leaves of *Bambusa* sp., *Dinochloa scandens*, *D. tjankorreh*, and *Schizostachyum rotundifolium* in Java and the Philippines.
- Phyllosticta phyllostachydis** Siem. Leaf spot on *Phyllostachys kumazae* in southern Russia.
- Phyllosticta take** Miy. and Hara. On leaves of *Bambusa* sp. and *Phyllostachys bambusoides* in Japan.
- Puccinia bambusoides** S. and Z. Brown rust pustules on leaves of *Phyllostachys bambusoides* in Japan.
- Puccinia corticioides** B. and Br. A rust producing yellowish-brown leathery crusts, up to 10 centimeters long, on stems of *Arundinaria hindsii*, *A. japonica*, *A. narihira*, *A. simonii*, and *Phyllostachys bambusoides* in Japan.
- Puccinia gracilenta** Syd. and Butl. Brown rust pustules on large, dull-brown sunken leaf spots on *Bambusa* sp. in India.
- Puccinia ignava** Arth. Brown powdery rust pustules on leaves of *Bambusa vulgaris*, *Bambusa* sp., and *Dendrocalamus giganteus* in Trinidad, Porto Rico, and Cuba.
- Puccinia kusanol** Diet. Brown to black rust pustules on *Arundinaria fortunei*, *A. japonica*, *A. narihira*, *A. simonii*, *A. variabilis*, *Bambusa veticii*, *Phyllostachys bambusoides*, *P. nigra*, *P. nigro-punctata*, *Sasa nipponica*, and *S. ramosa* in Japan.

BAMBUSEAE—Continued.

- Puccinia longicornis** Pat. and Har. Yellow-brown to dark-brown rust pustules on leaves of *Arundinaria japonica*, *A. simoni*, *Bambusa spinosa*, *B. tessellata*, *B. vulgaris*, *Sasa albo-marginata*, and *S. paniculata* in Japan, Ceylon, China, Indo-China, and the Philippines.
- Puccinia melanocephala** Syd. Brown to dark-brown rust pustules, covering under surfaces of leaves of *Arundinaria* sp., *Bambusa* sp., and *Phyllostachys* spp. in Japan and India. This rust has been introduced into Florida and Georgia.
- Puccinia mitriformis** S. Ito. Dark-brown to black rust pustules on leaves of *Sasa paniculata* in Japan.
- Puccinia phyllostachydis** Kusano. Powdery rust-brown to dark-brown or black rust pustules on leaves of *Phyllostachys aurea*, *P. bambusoides*, *P. henonis*, and *P. maritima* in Japan.
- Puccinia sasa** Kusano. Powdery rust-brown pustules on leaves of *Sasa borealis* in Japan.
- Puccinia xanthosperma** Syd. Dark-brown rust pustules on yellow, then gray, sunken leaf spots on *Bambusa* sp. in India and Japan.
- Puccinospora chusqueae** Speg. Elliptical dark brown spots, with yellow margins, on leaves of *Chusquea tenuiglumis* in Brazil.
- Pycnotherma bambusinum** Syd. More or less thickened circular spots on upper surfaces of leaves of *Bambusa vulgaris* and *Schizostachyum* sp. in the Philippines.
- Rosellinia rachidis** Rehm. On rachides of *Gigantochloa scribneriana* in the Philippines.
- Scirrhia bambusae** Malusio. (*Melanconium bambusae* Mal.) Brown spots and streaks appear on the branches and twigs and extend rapidly, soon involving the entire area. These spots turn whitish and dry out, finally being covered by the characteristic small black fruiting bodies. On *Bambusa gracilis* and *B. mitis* in Italy.
- Septoria bambusae** Brun. Pale-brown irregular spots, with broad black margins, on leaves of *Bambusa arundinacea* and *Phyllostachys puberula* in Japan.
- Septoria bambusella** Speg. Elliptical, indefinite, dark-brown spots on leaves of *Chusquea tenuiglumis* in Brazil.
- Shiraia bambusicola** P. Henn. Fleshly stromata on culms of *Bambusa* sp., *Phyllostachys bambusoides*, *P. puberula*, and *Sasa speculosa* in Japan.
- Trichonectria bambusicola** Rehm. On leaves of *Bambusa blumeana* and *B. spinosa* in the Philippines.
- Uredo arundinarieae** Syd. Brown rust pustules on leaves of *Arundinaria fortunei* and *A. japonica* in Japan and Ceylon.
- Uredo bambusarum** P. Henn. Powdery brown rust pustules, on round to linear spots, covering both leaf surfaces, of *Bambusa* sp. in Brazil.
- Uredo dendrocalami** Petch. Leaf rust on *Dendrocalamus strictus* in Ceylon.
- Uredo inflexa** Ito. Small powdery yellow-brown rust pustules on leaves of *Bambusa* sp. and *Sasa* sp. in Japan.
- Uredo ochlandrae** Petch. Brown rust pustules on leaves of *Ochlandra stridula* in Ceylon.
- Ustilago phyllostachydis** Syd. Black sclerotia formed in place of ovaries of *Phyllostachys bambusoides* in Japan.
- Ustilago shiraiana** P. Henn. This smut attacks the growing points of young branches and causes a witches'-broom effect. Infected branches are somewhat swollen and their growth stopped, the sori developing beneath the leaf sheaths and bracts, the powdery deep-brown spore masses finally breaking through. The hosts known for this very serious bamboo disease are *Arundinaria simonii*, *Arundinaria* sp., *Bambusa paniculata*, *B. veitchii*, *Phyllostachys bambusoides*, *P. henonis*, *P. puberula*, *P. quilioides*, *Sasa paniculata*, and *S. ramosa* in Japan and China. The disease has appeared at several points in California and Florida on imported plants but is now thought to have been eradicated.
- BANKSIA. SIRMUELLERA** Ag. Ornamental Australian evergreen shrubs or trees.
- Parodiella banksiae** Sacc. and Bizz. Numerous globose black aggregated perithecia on leaves of *B. marginata* in New South Wales.
- Phyllosticta banksiae** P. Henn. Brown spots, becoming pale, with yellow-brown margins, on leaves of *B. verticillata* in Germany.
- BAPHIA**. Barwood. Shrubs or small trees.
- Uredo baphiae** P. Henn. Black powdery rust pustules on leaves of *B. nitida* in central Africa.
- Uredo kampuluvensis** P. Henn. Powdery brown rust pustules on leaves of *B. cornifolia* and *Combretum baumium* in the Union of South Africa.
- BARBAREA. BARBAREA** Ag. WINTER CRESS. Yellow-flowered hardy biennials, sometimes called yellow rocket.
- Cercospora nasturtii** Pass. subsp. *barbareae* Sacc. On leaves of (*Campe barbarea*) *B. vulgaris* in France, Italy, and Russia.
- Peronospora barbareae** Güm. Downy mildew on leaves of *Campe stricta* (*B. stricta*) and *C. barbarea* (*B. vulgaris*) in central and north Europe.
- Ramularia coehleariae** Cke. Circular white spots on leaves of *C. stricta* (*B. stricta*) and *Armoracea rusticana* (*Radicula officinalis*) in Great Britain and Germany.
- BARRINGTONIA**. Broad-leaved evergreen trees.
- Cercospora barringtoniae** Syd. Circular yellow to yellow-brown spots on leaves of *B. luzonensis* in the Philippines.
- Phyllachora barringtoniae** (B. and Br.) Sacc. On *B. speciosa* in Ceylon. A doubtful species.
- Phyllosticta microstegia** Syd. On leaves of *B. racemosa* in the Philippines.
- Phyalospora barringtoniae** Syd. On *Barringtonia* sp. in the Philippines.
- BASELLA**. VINE-SPINACH. Annual or biennial tropical pot herbs, sometimes called Malabar nightshade.
- Ascochyta basellae** P. Henn. Circular dull-brown spots, drying out lighter in color, on leaves of *B. rubra* in Brazil.
- Phyllosticta basellae** Rangel. Associated with *Stagonospora basellae* Rangel.
- Stagonospora basellae** Rangel. Circular to elliptical white leaf spots, with dark-purple margins, on *B. rubra* in Brazil.
- Uromyces basellae** Syd. Yellow to brown rust pustules on leaves of *B. rubra* in the Philippines.
- BAUHINIA**. Ornamental tropical trees and shrubs.
- Cercospora bauhiniarum** Syd. On leaves of *B. malabarica* in the Philippines.
- Cercospora latimaculans** Wakef. Circular to irregular, often confluent, pale-brown leaf spots on *B. reticulata* and *Bauhinia* sp. in Gold Coast.
- Corticium salmonicolor** B. and Br. See Citrus.
- Phyllachora bauhiniarum** (Wint.) Theiss. and Syd. "Tar-spot" disease on leaves of *B. vahlii* in India.
- Phyllachora bahnerae** Racib. Circular black stromata on small yellow-green spots on leaves of *B. glauca* in Java.
- Phyllachora tenuis** (Berk.) Sacc. Black stromata on upper surfaces of leaves of *B. vahlii* and *Bauhinia* sp. in Mexico, Nicaragua, and Java.

BELLIDIASTRUM. Herbaceous perennials.

Entyloma bellidiastri Maire. Circular, whitish spots, turning brown, on leaves of *B. michelii* in Tyrol.

Entyloma calendulae (Oud.) De By. See *Calendula*.

Puccinia firma Diet. Brown rust pustules on leaves of *B. michelii* and *Carex firma* in Tyrol and Germany.

Septoria bellidiastri Allesch. On leaves of *B. michelii* in Bavaria.

BELLIS. *ENGLISH DAISY.* Low perennial herbs.

Entyloma bellidis Kræger. Circular white or yellowish spots, which are often confluent so as to cover the entire area of leaf blades of *B. perennis* in Dalmatia, Denmark, and Germany.

Puccinia distincta Motay. Orange-yellow and brown rust pustules on leaves of *B. perennis* in Australia and Tasmania.

Ramularia bellidis Sacc. Subcircular dull yellow spots on leaves of *B. perennis* in Italy.

Septoria bellidicola Rob. & Desm. White spots with red margins on leaves of *B. perennis* in Australia and Italy.

BERBERIS. Including *Mohavia*. Oregon grape, holly grape. *BARBERT.* Deciduous or evergreen shrubs cultivated for their ornamental leaves, flowers, and fruits.

Aecidium aridum Diet. and Neg. Rust on leaves of *B. burifolia* and *B. heterophylla* in Chile.

Aecidium berberis-ruscifolia P. Henn. Yellow rust pustules covering the under sides of leaves of *B. ruscifolia* in Argentina.

Aecidium berberidis-thunbergii P. Henn. Yellow rust pustules on round red spots on under surfaces of leaves of *B. thunbergii* in Japan.

Aecidium haussknechtianum P. Henn. Golden-yellow rust pustules on leaves, petioles, and fruits of *B. crataegina* in Persia.

Aecidium leucanum P. Magn. Rust on leaves of *B. burifolia* in Chile.

Aecidium montanum Burt. Rust pustules on red leaf spots on *B. aristata*, *B. coccinea*, and *B. ibericum* in India.

Aecidium tubiforme Diet. and Neg. Rust on leaves of *B. burifolia* in Chile.

Ascochyta australis Speg. Circular to irregular gray-brown spots, with yellow margins on leaves of *B. glauca* in Argentina.

Ascochyta berberidis Sacc. On branches of *B. vulgaris* in France and Russia.

Didymotheca berberidis Rehm, Thüss, and Syd. On branches of *B. laevigata* in Chile.

Didymosphaeria epidermidis (Fr.) Feck. var. *macrospora* Elias. On branches of *B. vulgaris* in Sweden.

Dothidea carneofusca v. Hoch. On branches of *B. vulgaris* in Austria.

Gambelia cornuta Mass. On leaves of *B. nepalensis* in India.

Gloeosporium japonicum Hennen. Anthracnose on *Mohavia japonica* in Japan.

Heterosporium berberidis Ranof. On branches of *B. vulgaris* in Serbia.

Laestadia berberidis Delacr. On branches of *B. vulgaris* in France.

Melasma berberidis Thüss and Wint. Small brown spots on leaves of *B. vulgaris* in Austria and Germany.

Microsphaera berberidis DC. Lev. Powdery mildew on leaves of *B. aquifolium* and *B. vulgaris* in Europe, Cyprus, Transcaucasia, Turkey, and Japan.

Mycosphaerella australis Speg. Lind. On leaves of *B. laevigata* in Chile.

Mycosphaerella berberidis Auerw. Lind. On leaves of *B. vulgaris* in Belgium and Switzerland.

Phoma berberidicola Vesterg. On branches of *B. vulgaris* in Sweden and Switzerland.

Phragmodothia berberidis Alm. and Cam. Thüss, and Syd. On branches of *B. vulgaris* in Sweden.

Phyllosticta aquifolii Allesch. On leaves of *Mohavia japonica* in Denmark.

Phyllosticta asiatica (Fr.) Grev. Irregular dull-brown spots, with dark, purplish-brown margins, on leaves of *B. asiatica* in Great Britain.

Phyllosticta berberidicola Speg. Spots on leaves of *B. laevigata* in Uruguay.

Phyllosticta berberidis Feck. Irregular gray spots, sometimes whitish, on leaves of *B. laevigata* and *B. coccinea* in France, Italy, Denmark, Russia, Austria, and Germany.

Phyllosticta mahoniae Thüss, Kell. On leaves of *Mohavia aquifolium* in Italy, France, Russia, and Denmark.

Phyllosticta mahoniaecola Pass. Large brown indefinite spots on leaves of *Mohavia aquifolium*, *M. laevigata*, and *M. japonica* in Italy, France, and Great Britain. Var. *aquifolia* P. Henn. is the same as the species, except that the leaf spots have definite brown margins.

Phyllosticta spegazziniana Kell. On leaves of *Mohavia japonica* in Argentina.

Phyllosticta wetendorpii Thüss. Angular white spots, with red margins, on leaves of *B. citreola* and *B. vulgaris* in Yugoslavia, Siberia, Belgium, and Germany.

Puccinia antarctica Speg. Dark-brown rust pustules on leaves of *B. burifolia* (*B. glauca*) in Patagonia.

Puccinia arhenatheri Kell., Erik. See *Arhenatherum*.

Puccinia barri-aranae Diet. and Neg. Chestnut-colored rust pustules on leaves of *B. burifolia* in Chile.

Puccinia berberidis Mot. Yellow to yellow-brown rust pustules on angular or irregular leaf spots on *B. aristata*, *B. droopingii*, *B. emarginata*, *B. japonica*, *B. heterophylla*, and *B. apiculata* in Chile.

Puccinia berberidis-trifoliae Diet. and Neg. Brown powdery rust pustules on small brown sunken spots on leaves and petioles of *B. trifolia* in Mexico.

Puccinia droogensis Burt. Rust on leaves of *B. coccinea* in Ceylon and India.

Puccinia meyeri-alberti P. Magn. Brown rust pustules on angular leaf spots on *B. burifolia*, *B. coccinea*, *B. laevigata*, *B. linearefolia*, and *B. perennis* in Chile.

Puccinia naumanniana P. Magn. Diet. Black powdery rust pustules on leaves of *B. burifolia* in Chile.

Puccinia stolpiana (P. Magn.) Diet. and Neg. Yellow to black powdery pustules on leaves of *B. burifolia* and *B. ilicifolia* in Chile.

Septoria berberidis Noss. Subcircular dull-brown spots, with dark purple margins on leaves of *B. vulgaris* in Yugoslavia, Russia, Italy, Switzerland, and Austria.

Septoria mahoniae Kasser. Small brown spots on leaves of *Mohavia aquifolium* in Russia, Italy, and Germany.

Uromyces quitenis Berk. Rust on leaves of *Berberis* sp. in Ecuador.

BETA. *BEET.* *BEET ROOT.* Cultivated for their edible leaves, and for sugar production.

Aetionomyces spp. Several species of *Aetionomyces* including *A. albus* (R. D. Guss.) *A. intermedius* Kunt. (W.) and *A. melleocoloratus* W., said to cause rot of beet roots (*B. vulgaris*) have been reported from Europe. Their relation to *A. albus* is uncertain.

Aphanomyces laevis de By. Causes a soft rot of sugar beet (*B. vulgaris*) roots in Europe.

Ascochyta betae Prill and Delacr. On petioles of *B. vulgaris* in France.

BETA—Continued.

- Ascochyta beticola** Prill. and Delacr. On petioles of *B. vulgaris* in France. Probably not distinct from above species.
- Bacillus** spp. Bacteriosis has been reported from a number of countries, particularly in Europe, as a serious disease of the beet (*B. vulgaris*) under various names, including *B. betae* Mig., *B. bussei* Mig., *B. lycopersi* Mig., *B. mycoides* Mig., and *B. tabificans* Delacr. The status of these species and their relation to *B. carolusporus* is most uncertain.
- Bacillus** sp. Bacterial scab of *B. vulgaris* in Korea.
- Enyoma betiphilum** Bub. Leaf smut on *B. vulgaris* in Austria. A doubtful species.
- Helicobasidium mompa** Tan. See *Morus*.
- Leptosphaeria circinans** (Fekl.) Sacc. See *Asparagus*.
- Moniliopsis aderholdi** Ruhl. See *Solanum*.
- Peronospora schachtii** Fekl. The downy mildew of beet (*B. maritima* and *B. vulgaris*) is serious in Europe, Egypt, Japan, and Argentina. The disease occurs to a limited extent in New York, Minnesota, and California. Delicate gray patches of mildew appear on lower leaf surfaces. Young central leaves are most frequently attacked and the growth of the plants checked. Seedlings may be killed outright.
- Pionnotes betae** Sacc. Yellow spots coalescing to form crustlike areas and cankers of roots of *B. vulgaris* in Great Britain.
- Pleospora putrefaciens** (Fekl.) Frank. Causes a serious disease of beets (*B. vulgaris*) in Germany.
- Ramularia betae** Kestr. Numerous subcircular grayish spots with reddish margins on both surfaces of leaves of *B. maritima* and *B. vulgaris* in Bulgaria, Belgium, Denmark, Holland, and Russia.
- Rhizoctonia napi** West. See *Brassica*.
- Septoria betae** Werd. Pale-brown spots with white centers and dark-brown margins, on leaves of *B. vulgaris* in Australia and Belgium. Reported from Indiana.
- Sorolpidium betae** Nemec. On roots of *B. vulgaris* in Europe.
- Typhula betae** Reestr. Attacks seedlings of *B. vulgaris* in Sweden, Denmark, Holland, Great Britain, and the Azores and is often destructive.
- Uredo marmorata** Speg. Rust on leaves and stems of *B. procumbens* in the Canary Islands.
- Uromyces betae** (Pers.) Léw. Yellow to brown rust pustules on leaves of *B. cicla*, *B. maritima*, and *B. vulgaris* in Argentina, Uruguay, Union of South Africa, Australia, New Zealand, and Europe. Found sparingly in California and Colorado.
- Urophysalis leproides** (Trab.) P. Maen. Forms nodular outgrowths from the upper portion of the roots, which often become as large as walnuts. These outgrowths are modified roots and remain attached by narrow bases. On *B. vulgaris* in New Zealand, Algeria, Argentina, Great Britain, France, Belgium, and Sweden. Occurs scatteringly in California and has been found in Iowa.
- Urophysalis pulposa** (Walt.) Schreit. See *Atriplex*.
- A disease characterized by orange-colored spots, irregular swellings, and distortion of the foliage of beet has been reported from northern France, the cause being unknown.
- BETULA**. BUSH. Forest and shade trees with handsome green foliage and ornamental bark.
- Didymosphaeria latebrosa** Mont. On leaves of *B. alba* in Belgium.
- Exoascus alpinus** (Johans.) Sacc. On leaves and branches of *B. nana* in Scandinavia.
- Exoascus betulae** Fekl. Fruiting areas on lower leaf surfaces of *B. alba*, *B. odorata*, *B. pubescens*, and *B. verrucosa* (*B. pendula*) in middle Europe.
- Exoascus betulinus** (Rostk.) Sadeb. Produces witches'-brooms on *B. alba*, *B. odorata* and *B. pubescens* in north and middle Europe. Reported from Wisconsin.
- Exoascus janus** Thomas. Large, circular, pale rose-colored spots on both leaf surfaces of *B. verrucosa* in Switzerland and Germany.
- Exoascus nanus** (Jehan.) Sacc. Produces witches'-brooms and yellow-green leaf spots on *B. nana* and *B. odorata* in Norway and Sweden.
- Exoascus turgidus** Sadeb. On leaves of *B. alba*, *B. odorata*, *B. pubescens*, and *B. verrucosa* in Finland, Russia, Sweden, Denmark, Switzerland, Great Britain, Austria, and Germany.
- Guignardia betulae** (Amsw.) Sacc. and Trott. On leaves of *B. verrucosa* in Denmark and Germany.
- Macrophoma sydoviana** (Bres.) Sacc. and Syd. On leaves of *B. alba* in Germany.
- Marsonia betulae** (Lib.) Sacc. Irregular, radiate, dull-brown spots on leaves of *B. alba* and *B. verrucosa* in Europe.
- Myxosporium devastans** Rostk. Causes withering and die-back of young twigs of *B. alba*, *B. verrucosa*, and *Acer pseudoplatanus* in Denmark.
- Phyllosticta betulae** Oud. On leaves of *B. alba* in Holland.
- Phyllosticta betulina** Sacc. On leaves of *B. alba* in Italy and France. Reported from New York.
- Ploeroglyphia virgultorum** Sacc. Black knot of birch. Shoots and branches from 1 to 4 years old are attacked, narrow elongated stromata forming, which are olive to black in color and carbonaceous in texture. Leaves on infected branches wilt and die and the twigs die back. A witches'-broom effect is produced finally. On *B. alba*, *B. nana*, *B. pubescens*, and *B. verrucosa* in Great Britain, Sweden, Finland, Switzerland, Russia, Denmark, and Germany. Theissen and Sydow delete this species.
- Rhytisma nervale** (Alb. and Schw.) Rehm. See *Alnus*.
- Sclerotinia betulae** Wor. Attacks the catkins of *B. alba*, *B. papyrifera*, and *B. verrucosa* in Japan, Russia, and Denmark.
- Septoria betulina** Pass. Irregular, often confluent, gray spots on leaves of *B. alba* and *B. pubescens* in Russia, Denmark, Tyrol, and Italy. Reported from Wisconsin.
- Taphrina aurea** (Pers.) Fr. Attacks seedlings of *Betula* sp. in Great Britain.
- BIGNONIA**. TRUMPET CREEPER. Woody climber.
- Ascochyta tweediana** Penz. and Sacc. On capsules of *B. tweediana* in Italy.
- Cercospora bignoniicola** Speg. Irregular, often confluent, gray spots on leaves of *Bignonia* sp. in Brazil.
- Cercospora unguis-cati** Speg. Subcircular dark-purple spots on both leaf surfaces of *B. unguis-cati* (*Doranthia unguis-cati*) in Argentina.
- Colletotrichum bignoniae-igneae** Rangel. Anthracnose on leaves of *Bignonia* sp. in Brazil.
- Phyllachora amphigenia** Speg. (*P. nitidula* Pat. and Gaill.) Black stromata on pale brown spots on leaves of *Bignonia* sp. and *Cytostoma cellistegioides* in Venezuela, Paraguay and Brazil.
- Phyllachora nitidula** Pat. Black stromata on leaves of *Bignonia* sp. in Venezuela. A doubtful species.
- Phyllachora tenuis** Speg. Tar-spot on leaves of *Bignonia* sp. in Argentina. This species should be renamed, since *P. tenuis* (B. and C.) Sacc. has priority.
- Phyllosticta tweediana** Penz. and Sacc. On capsules of *B. tweediana* in Italy.
- Pleospora briosiana** Luigi. Large hazel-colored blotches, with black borders, on leaves of *B. buccinatoria* (*Phaedranthus buccinatorius*) in Italy.
- Puccinia appendiculatoides** P. Henn. Yellow rust pustules on round or effuse circular leaf spots on leaves of *Bignonia* sp. in Brazil.

BIGNONIA—Continued.

Puccinia bignoniacarum Speg. Brown rust pustules on leaves of *Bignonia* sp. in Brazil.

Uredo cuticulosa E. and E. Brown rust pustules on leaves of *Bignonia* sp. in Nicaragua.

Cropixis rickiana P. Magn. Irregular, woody galls, on which brown rust pustules occur on trunks of *Bignonia* sp. in Brazil.

BIHAL. See *Heliconia*.

BIKUKULLA. See *Dicentra*.

BISCHOFFIA. Ornamental tropical trees.

Phyllachora bischoffiae Syd. Irregular, black, shining stromata on leaves of *B. jamaica* (*P. trifoliata*) in India.

BIXA. ANATTO. Ornamental tropical trees. Red coloring matter produced from the pulp surrounding the seeds.

Cercospora bixae Alesch. Large, irregular, brown spots, with surrounding yellow zones, on leaves of *B. orellana* in Brazil.

Ovularia bixae Rac. On leaves of *B. orellana* in Java and Ceylon.

Phyllosticta bixina Young. Subcircular to irregular pale yellow spots, surrounded by purplish-black borders, on leaves of *B. orellana* in Porto Rico.

Rhizoctonia lamellifera Small. See *Grevillea*.

Uredo bixae Arth. Brown powdery rust pustules on leaves of *B. orellana* in Porto Rico.

BLECHNUM. Ferns.

Aphelenchus olesistis Ritz. Bos. See *Begonia*.

Milesina blechni Syd. Brown rust pustules on leaves of *B. occidentale*, *B. spicatum*, and *Lycopodium polymorphum* in Colombia, Trinidad, Great Britain, Denmark, France, Spain, Switzerland, Austria, and Germany.

Polystomella pulchella (Speg.) Theiss. See *Alsophila*.

Uredinopsis mayriana Diet. Rust on leaves of *B. blechnoides* in Colombia.

Uredo blechnicola P. Henn. Powdery yellow rust pustules on leaves of *B. colubifolia* in Brazil.

Uredo orientalis Racib. Brown, erumpent, rust pustules deforming the leaves of *B. orientalis* in Java.

BLETTIA. Terrestrial orchids. See *Orchidaceae*.

BLUMENBACHIA. SALOA Ag. STING LILY. Ornamental annuals.

Septoria blumenbachiae Speg. Indefinite light-brown spots on leaves of *B. arens* in Argentina.

BOCCONIA. PLUME POPPY. Tall garden herbs.

Puccinia bocconiae Mayr. Rust on leaf blades and petioles of *B. frutescens* in Colombia.

BOEHMERIA. Ramie. Herbs, shrubs, or trees. Tropical fiber and ornamental plants.

Cercospora boehmeriana Wor. On leaves of *B. nirex* in Russia.

Colletotrichum boehmeriae K. Saw. Scattered ashen-brown circular spots on leaves and fusiform cankers on stems of *B. nirex* in Japan and Formosa.

Puccinia boehmeriae P. Henn. Yellow-brown rust pustules on leaves of *Boehmeria* sp. in Brazil.

Pucciniastrum boehmeriae Syd. Yellow-brown rust pustules on leaves of *B. blanda*, *B. grandiflora*, *B. japonica*, *B. longifolia*, *B. sieboldiana*, *B. spicata*, and *B. triacuspis* in Japan and the Philippines.

Ramularia boehmeriae Fujii. Causes a root and rhizome rot of *B. nirex* in Japan.

Septoria boerhaviae Pat. On leaves of *B. verticillata* in Abyssinia.

Uredo boehmeriae Diet. Rust on leaves of *B. platyphylla* in Ceylon.

BOLTONIA. Asterlike herbs.

Colosporium asterum (Diet.) Syd. See *Aster*.

BOMAREA. Twining, herbaceous plants of the *Amariyllidaceae*.

Aecidium bomareae Mayr. Yellow rust pustules on leaves and stems of *B. californiana* and *B. polacoensis* in Colombia, South America.

Heterosporium alli-bomareae Pat. On leaves of *Bomarea* sp. in Ecuador.

Puccinia bomareae P. Henn. (*Uredo bomareae* Lagerh.) This rust produces elongated lemon-yellow pustules, which later become dark-brown, on lower surfaces of leaves of *B. acutifolia*, *B. edulis*, *B. ovala*, and other species in Ecuador, Mexico, and Brazil.

Puccinia pallor Arth. and Holw. Yellow to brown powdery rust pustules on lower surfaces of leaves of *B. acutifolia* in Costa Rica and Guatemala.

Urocystis bomareae Diet. and Neg. Black powdery spore masses form in swellings on the stems of *B. salicilla* in Chile.

Uromyces bomareae P. Henn. Bright yellow, then brown, rust pustules on the lower surfaces of the leaves of *Bomarea* sp. in Brazil.

BOMBAX. Tropical shrubs and trees, supplying a useful floss from the cottony contents of the pods. Sometimes known as silk cotton tree.

Chrysomyxa bombacis Petch. Rust on leaves of *B. malabaricum* in Ceylon.

Cladotrichum foliolica (Niessl.) Ferro. Effuse black areas on leaves of *B. malabaricum* in India.

Fomes lignosus Klotzsch. See *Hevea*.

Gloeosporium bombacis Putte. Irregular, then confluent, dull-brown areas on leaves of *B. insignis* in Brazil.

Puccinia bombacis Diet. Brown rust pustules on leaves of *Bombax* sp. in Colombia.

Uredo bombacis Petch. Rust on leaves of *B. malabaricum* in Ceylon.

BORAGO. BORAGE. Coarse, annual herbs, sometimes cultivated for culinary purposes.

Cercospora sp. Leaf spot on *B. officinalis* in Cuba.

Cylindrosporium myosotidis Sacc. See *Myosotis*.

Entyloma serotinum Schroet. Smut sori in numerous white, then brown, circular leaf spots on *B. officinalis*, *Pulmonaria officinalis*, *Symphytum officinalis*, and *S. tuberosum* in Denmark, France, Austria, and Germany. Ciferri has described the form on *Borago* as *E. borraginis* Cif.

BORASSUS. Palmyra palm. See *Palmae*.

BORONIA. Ornamental shrubs.

Puccinia boroniae P. Henn. Dark cinnamon-colored rust pustules on branches of *B. spinosa* in Australia.

BOTOC. See *Psophocarpus*.

BOUGAINVILLEA. EUGENVILLEA Ag. Tropical woody climbers with brilliant colored bracts.

Puccinia bougainvilleae (Speg.) Schroet. Yellow to dark-brown rust pustules, on circular yellow spots, on leaves and branches of *B. frondosa* and *B. spectabilis* in Argentina.

BOUSSINGAULTIA. Madeira vine. Herbaceous perennial vines.

Cercospora boussingaultiae Roum. On leaves of *B. baselloides* in France.

Phyllosticta boussingaultiae Speg. Small whitish spots, with purple borders, on leaves of *B. baselloides* in Argentina.

BOUARDIA. Small flowering shrubs.

Uromyces bouvardiae Syd. Yellow and brown to black rust pustules on leaves of *B. hirtella*, *B. leiantha*, *B. versicolor* and *Bouvardia* sp. in Mexico and Guatemala.

BRACHYCHITON. BOTTLE TREE. Australian trees.

Phyllosticta sterculiace brachychiti Wint. On leaves of *B. diversifolium* and *B. populineum* in Portugal.
Septoria ochracea-maculans Theum. Irregular, dull yellow spots, with narrow dark purple borders, on leaves of *B. diversifolium* and *B. populineum* in Portugal.

BRACHYCOME. SWAN RIVER DAISY. Australian herbs.

Puccinia brachycomae McAlp. Yellow and black rust pustules on swollen and distorted portions of leaves, and on stems and branches of *B. alberti*, *B. diversifolia*, *B. pathophora* and *B. wrightii* in Australia.

BRASSAVOLA. Epiphytic orchids. See *Orchidaceae*.**BRASSICA.** Cabbage, Mustard, Turnip, Rutabaga, Cauliflower, Kale, Kohlrabi, Brussels sprouts. Annual and perennial herbs.

Ascochyta brassicae Tulas. Large, irregular, indefinite, yellow-gray spots on leaves of *B. oleracea capitata* and other varieties in Europe.

Bacillus brassicaevorus Del. Bacteriosis of *B. oleracea capitata* and *borealis* in France. Probably not distinct from *B. carotovorus*.

Bacillus sp. (?) Bacterial leaf spot of *B. oleracea capitata* and *B. rapa* in Guam.

Cercospora brassicae campestris Rangel. On leaves of *B. campestris* in Brazil.

Cercospora brassicae P. Henn. Circular, numerous pale spots, with well-defined surrounding zones, on leaves of *B. oleracea* and *B. pekinensis* in Japan and the Philippines.

Colletotrichum brassicae Sacc. and Holw. Anthracnose on stems of *B. oleracea* in Yugoslavia.

Cylindrosporium brassicae Sacc. and Henn. Numerous, large, irregular, dark water spots on leaves of *B. campestris* and *B. rapa* in Yugoslavia and France. Reported from Missouri and Georgia.

Gloeosporium concentricum Grav. B. and Br. Concentric black spots on leaves of *B. capitata*, *B. oleracea* and *B. rapa* in Japan, Great Britain, Denmark, and Germany.

Monilopsis adnholdii Ruhl. See *Solanum*.

Mycosphaerella brassicae Durr. and Loh. *Phyllosticta brassicae* [Curr.] West. The "ring-spot" disease of *B. oleracea* var. *capitata* in Canada, and also occurs in Europe, Australia, New Zealand, Argentina, Uruguay, and California. Numerous distinct circular, light to dark gray spots occur on the leaves, surrounded by olive-green or blue-green borders. Affected leaves have little value. The disease develops rapidly after harvesting and is often serious enough to cause heavy losses in marketing.

Ophiaster radialis (De Willd.) Pascher. See *Linum*.

Oidium brassicae Wor. Attacks the roots of seedlings of *B. oleracea* var. *capitata* and *Brassica* spp. causing wilting-off. In Russia, Finland, Sweden and Belgium. Reported from Wisconsin.

Oidium radicleolum De Willd. On roots of *B. oleracea* and *Brassica truspastensis* in Russia and Belgium.

Oularia brassicae Bras. and Allesch. Subcircular or irregular, often confluent, white spots on leaves of *B. napus* and *B. oleracea* in Germany.

Phoma napae-brassicae Rostr. Causes a dry rot of roots of *B. napae* and *B. oleracea* in New Zealand, Great Britain, Denmark, and Holland. Reported from Canada.

Phyllosticta brassicae Sacc. Circular to angular, dull, greenish-gray spots on leaves of *B. oleracea* in Italy and Malta.

Phyllosticta napi Sacc. Indefinite spots on leaves of *B. napae* and *B. oleracea* in France and Italy.

Pseudomonas Bacterium destructans Potter. Causes a white rot of *B. napae* and *B. oleracea* in Great Britain and *Rapionovirus* in Japan. Probably not distinct from *B. carotovorus* Jones.

Ramularia rapae Pm. Irregular spots on leaves of *B. napae* in Great Britain.

Rhizoctonia napi West. Covers stems, pods, leaves, and roots with white mycelium in which sclerotia develop, white at first, turning black on the outside. Causes a dry rot of infested areas, seriously diseased plants breaking over at the ground level. The recognized hosts are *Aceranthus tristis*, *Argemone mexicana*, *Artemisia sativa*, *Beta betulae*, *Brassica campestris*, *Calamagrostis*, *Cannabis sativa*, *Chenopodium album*, *Cicuta oviformis*, *Cnicus arvensis*, *Fumaria parviflora*, *Geranium vulgare*, *Lactuca sativa*, *Limonium*, *Leucaena*, *Lupinus*, *Lycium vestitissimum*, *Medicago lupulina*, *Pisum sativum*, *Sisymbrium*, *Sisymbrium*, *Triticum vulgare*, and *Vicia sativa*. Shaw states that this species should be transferred to *Botrytis*. Other workers consider it synonymous with *Sclerotinia sclerotiorum* Mass.

Sporidesmium brassicae Mass. Indefinite olive-gray or green spots, on leaves and pods of *B. campestris* in India and China.

Sporidesmium exitiosum Kuehn. On stems and pods of *B. napae*, *B. oleracea*, *B. rapa*, and *B. cretensis* in Japan, Sweden, Denmark, and Germany. Also reported from Canada. *Leptospharia napi* (Fekl.) Sacc. is said to be the perfect stage of this species.

Trocytis coralloides Rostr. Affected plants are stunted and galls up to one and one-half inches in diameter form on the roots. On *Brassica* sp. (mustard) in India.

Uromyces brassicae Nussl. On *B. oleracea* in France. A very doubtful species.

BRICKELLIA. COLEOSANTHUS Ag. Herbs or small shrubs.

Puccinia decora Diet. Dull-brown rust pustules on leaves of *Brickellia* sp. in Mexico.

Puccinia praemorsa Diet. and Holw. Rust on leaves of *B. americana* in Mexico.

Pucciniosira brickelliae Diet. and Holw. Brown rust pustules on leaf blades and petioles and stems of *B. alamosarpa*, *B. caranilloi*, *B. secundiflora*, and *Coleosanthus caranilloi* in Mexico and Guatemala.

BRIZA. QUAKING GRASS. Annual or perennial grasses.

Dilophia graminis (Fekl.) Sacc. See *Triticum*.

Septoria crispiniana Gz. Frag. On leaves of *B. maxima* in Spain.

Tilletia brizae Ue. Brown striations on leaves which finally rupture exposing powdery black masses of smut spores. On *B. media* in Germany.

BROMELIA. Bushy leafy Bromeliads.

Perisporium bromeliae Stevens. Forms smoky patches on leaves of *B. pinguin* in Porto Rico.

Phyllosticta bromeliae Alm. and Cam. Whitish spots on leaves of *B. acanga* in Portugal.

BROMUS. BROMEGRASS.

Exobasidium graminicolum Bres. See *Arrhenatherum*.

Helminthosporium fragosii Bub. On leaves of *B. stertilis* in Spain.

Hypochnus fuciformis (Berk.) McAlp. See *Lolium*.

Leptosphaeria culmicola Fr. Karst. On culms of *B. strictus*, *Festuca arundinacea*, *F. sylvatica*, *Phalaris arundinacea*, *Sesale cereale*, and *Triticum sativum* in Denmark, Italy, Sweden, France, and Germany.

Mycosphaerella longissima Fekl. On leaves of *B. asper* and *B. maritimus* in Italy, Switzerland, Denmark, and Germany.

Mycosphaerella recutita Fr. Johans. See *Aira*.

Ophiobolus cariceti (B. and Br.) Sacc. See *Triticum*.

Phyllachora bromi Fekl. Tar spot on leaves and leaf sheaths of *B. japonicus*, *B. unioloides*, *Brachypodium* sp., and *Dactylis* sp. in Argentina, Japan, Algeria, Italy, France, and Germany.

BROMUS—Continued.

- Phyllosticta bromi** Poteb. On leaves of *B. patulus* in Russia.
- Puccinia brachypus** Speg. Linear, brown, rust pustules on leaves of *B. caudatus*, *Lolium perenne*, and *Triticum sativum* in Argentina.
- Puccinia bromina** Erikss. Yellow and brown to black rust pustules on leaf blades and petioles of *Palmatoria montana*, *Symphytum officinale*, and nearly 50 species of *Bromus* in Argentina, Australia, Asia Minor, Japan, and all of Europe.
- Puccinia bromi-japonica** S. Ito. Yellow-brown to black linear rust pustules on leaves of *B. japonicus* in Japan.
- Septoria affinis** Sacc. Linear whitish spots with rufous margins, on leaves of *Brachypodium cineratum*, *Bromus inermis*, and *B. mollis* in Italy and Russia.
- Septoria bromicola** Speg. On leaves of *B. unioloides* in Chile.
- Septoria bromivora** Speg. On leaves of *Bromus* sp. in Argentina.
- Tilletia belgradensis** P. Magn. Smut in inflorescences of *B. scabellus* in Yugoslavia.
- Tilletia holci** (West.) Rostr. See *Holcus*.
- Tilletia velenovskii** Bub. Smut sori in ovaries of *B. cernuus* and *B. scabellus* in Bulgaria. Said to be the same as *T. guineensis* Har. which is a limited occurrence in the western United States.
- Uredo bromi-pauciflorae** S. Ito. Yellow-brown rust pustules on leaves of *B. pauciflorus* in Japan.
- BROUSSONETIA PAPIRIUS** Ag. PAPER MULBERRY. Ornamental trees or shrubs. Bark used for paper making.
- Gibberella moricola** (Ces. and De N.) Sacc. See *Morus*.
- Phyllosticta broussonetiae** Trav. and Mighl. Irregular whitish areas, often involving entire leaves of *B. papyrifera* in Italy.
- BROWNEA**. Small, tropical evergreen trees.
- Fomes lamaeensis** Murr. See *Hevea*.
- Phyllachora juruensis** P. Henn. Black stromata on small brown spots scattered irregularly over leaves of *Brownea* sp. in Brazil.
- BRUNELLA**. **PRUNELLA** Ag. Salt-tolerant. Hardly herbaceous perennials. Also called rock-rose.
- Ophiobolus rostrupii** Berk. and Winge. On leaves of *B. vulgaris* in the Mexican States.
- Puccinia brunellarum-moliniae** Coudan. Rust with yellow anther stage on leaves of *B. grandiflora* and *B. vulgaris*, and brown antheral and telial stages on *Melvinia cerulea* in Europe.
- Ramularia brunellae** E. and E. *pyrenaica* Gz. Frag. On leaves of *B. byrtica* in Spain.
- Ramularia harloti** Sacc. More or less circular, grayish white spots on both leaf surfaces of *B. vulgaris* in France.
- Septoria greschikii** Bres. On *B. grandiflora* in Europe.
- Septoria trulliana** Sacc. and var. *italica* Ferr. On leaves of *B. vulgaris* in Yugoslavia, Italy, Russia, and Great Britain.
- BRUNFELSIA**. Ornamental tropical shrubs.
- Fomes lamaeensis** Murr. See *Hevea*.
- BRUNSVIGIA**. Summer or autumn flowering bulbs.
- Aecidium brunsvigiae** P. Henn. Leaf rust on *Brunsvigia* sp. in the Union of South Africa.
- Tilletia sydownii** Sacc. and Trent. Smut sori 0.5 mm long in place of ovaries, protected at first by ashen-colored membranes which rupture to expose powdery black spore masses. On *B. (Arumochloa) subcaulis* in Algeria.
- BRYONIA**. BAYONET. Herbaceous perennial climbers.
- Asochyta bryoniae** Bub. and Kab. Irregular, grayish-yellow or brown spots, often confluent, on leaves of *B. alba* in Bohemia.
- Asochyta tirolensis** Bub. Small circular to oblong whitish spots with irregular surrounding zones, on leaves of *B. dioica* in Tyrol.
- Ramularia bryoniae** Fautr. and Roum. Large, irregular, brown spots on leaves of *B. dioica* in France.
- BRYOPHYLLUM**. Succulent foliage plants.
- Stilbella flavida** (Cke.) Lind. See *Coffea*.
- BUDDEIA**. BETTERLY BUSH. Ornamental shrubs or trees cultivated for their showy flowers.
- Diplodia buddleiae** Pat. On leaves of *Buddleia* sp. in Ecuador.
- Phyllosticta auriculata** Kalch. and Cke. Circular pale-brown spots, with purple margins, on leaves of *B. auriculata* and *B. globosa* in the Union of South Africa and Denmark.
- Phyllosticta montemartini** Ciferri. Irregular, yellow-brown spots on leaves of *B. variabilis* (*B. davidii*) in Italy.
- Septoria buddleiae** Kalch. and Cke. Irregular, dark brown spots on leaves of *B. salicifolia* in the Union of South Africa.
- Septoria merrilli** Syd. On leaves of *B. asiatica* in the Philippines.
- Septoria thimienii** Sacc. Red-brown spots on leaves of *B. auriculata* in the Union of South Africa.
- BUGINVILLEA**. See *Bougainvillea*.
- BULBINE**. Bulbous and nonbulbous African and Australian plants.
- Aecidium bulbines** Henn. and Erens. Brown spots on both leaf surfaces in which yellow aecia appear. On *Bulbine* sp. in the Union of South Africa.
- Uromyces bulbini** Thum. Brown rust pustules are arranged concentrically in large circles on sunken spots on both leaf surfaces of *B. aloides*, *B. bulbosa*, and *B. latifolia* in New South Wales and Victoria.
- Uromyces semibarbata** Osborn. Leaf rust on *B. semibarbata* in Australia.
- BULBOCODIUM**. CROCUSLIKE bulbous plants.
- Aecidium bulbocodium** Koen. Yellow rust pustules on leaves of *B. ternum* in Asia.
- Urocystis colchici** (Schlecht.) Rab. See *Colchicum*.
- BULBOPHYLLUM**. See *Orchidaceae*.
- BUPHTHALMUM**. OX-EYE. Perennial Asian herbs.
- Phyllosticta buphthalmi** Allesch. Irregular spots, often occupying entire leaf area, on *B. salicifolium* in Germany.
- Ramularia buphthalmi** Allesch. Subcircular or irregular, often confluent, dark-violet spots, drying out to brown, on leaves of *B. salicifolium* in Germany.
- Septoria buphthalmi** Allesch. Small, irregular, purple or olive-colored spots on leaves of *B. salicifolium* in Germany.
- BUPLEURUM**. Shrubs or herbs.
- Asochyta bupleuri** Thum. (Including var. *bupleurum-fruticosum* Gz. Frag. Gray spots on leaves of *B. falcatum*, *B. fruticosum*, and *B. obovatum* in Spain, Portugal, and Germany.
- Cercospora bupleuri** Pass. Reddish spots on stems and branches of *B. tenuisternum* in Italy.
- Didymella fruticosa** v. Hoeh. Circular or oblong purple-brown spots with black margins on leaves of *B. fruticosum* in Corsica.
- Entyloma bupleuri** Lindr. Smut sori in brown spots on leaves and stems of *B. glaucum* in France.
- Laestadia bupleuri** (Dur. and Mont.) Sacc. On leaves of *B. spinosum* in Algeria.

BUPLEURUM—Continued.

Phyllosticta asteromoides Bub. Irregular, confluent, purple-brown or black leaf spots on leaves of *B. falcatum* in Bohemia.

Phyllosticta bupleuri (Eckl.) Sacc. Gray spots on leaves of *B. falcatum*, *B. fruticosum*, and *B. gibraltarium* in Algeria and Germany.

Puccinia bupleuri-falcati (DC.) Wint. This rust in all its stages occurs on a wide range of species of *Bupleurum* in Japan, Indo-China, India, China, Siberia, Asia Minor, north Africa, and all of Europe.

Septoria amphigena Miy. On leaves of *B. falcatum* in China.

Septoria bupleuri Desm. Small subcircular whitish spots with black borders on leaves and buds of *B. frutescens*, *B. fruticosum*, and *B. semiconpositum* in Spain, Portugal, France, Italy, and Austria.

Septoria bupleuricola Sacc. Circular, dark brown spots, becoming gray with dark-brown margins on leaves of *B. aureum* and *B. longifolium* in Siberia.

Septoria bupleuri-falcata Died. Subcircular or angular spots, white above gray below, with black margins, on leaves of *B. falcatum* in Germany.

Septoria bupleurina de Lamarl. Black, then brown, circular or angular spots with narrow greenish borders on leaves of *B. longifolium* in France.

Septoria diffusa F. Tassi. Diffuse dark-brown spots on leaves of *B. fruticosum* in Italy.

Uromyces bupleuri P. Magn. Chestnut-brown rust pustules on leaves and stems of *B. frutescens* and *B. graminifolium* in Persia and Spain.

BURCHARDIA. Herbaceous perennials, allied to *Veratrum*.

Puccinia burchardiae Sacc. Yellow-brown to dark-brown rust pustules on both leaf surfaces of *B. umbellata* in Australia.

BUTOMUS. FLOWERING RUSH. Hardy perennial aquatic plants.

Doassansia niesslii De Toni. Smut sori in gray-brown spots on leaves of *B. umbellatus* in Poland, Austria, and Germany.

Physoderma butomi Schroet. On leaves of *B. umbellatus* in Finland, Denmark, Norway, and Germany.

Ramularia butomi Lind. Small, oblong or angular, scattered black spots on leaves of *B. umbellatus* in Sweden.

BUXUS. Box. Ornamental evergreen, small trees or shrubs.

Ascochyta buxina Sacc. Indefinite light-brown spots on leaves of *B. sempervirens* in France and Italy. Possibly *Phyllosticta buxina* Sacc., which occurs in the United States, is the same species.

Dothidella depazeoides (Desm.) Theiss. and Syd. Black stromata on white spots on leaves of *B. sempervirens* in France, Belgium, Switzerland, and Austria.

Gloeosporium louisiae Baumei. Anthracnose on leaves of *B. sempervirens* in Belgium and Hungary.

Guignardia buxi Desm. Yellow spots on lower surfaces of leaves of *B. sempervirens* in Europe.

Helminthosporium martinicense Thuen. Black effuse spots on leaves of *B. citrifolia* in Martinique.

Phyllosticta auerswaldii Allesch. On leaves of *B. sempervirens* in Germany. Reported from New Jersey.

Phyllosticta buxina Sacc. Variable, pale-brown leaf spots with distinct dark purple borders on *B. sempervirens* in Great Britain and Italy.

Phyllosticta phacioides (Sacc.) Allesch. On leaves of *B. sempervirens* in Yugoslavia.

Puccinia buxi DC. Brown rust pustules on leaves of *B. sempervirens* in Madeira, Persia, Dalmatia, Spain, Portugal, Greece, Russia, Italy, Switzerland, France, Great Britain, Austria, and Germany.

Septoria phacioides Desm. On leaves of *B. sempervirens* in France and Belgium.

Uromyces ambiens Cke. Yellow-brown rust pustules on leaves of *B. sempervirens* in India.

BYRSONIMA. Tropical American trees and shrubs.

Aecidium byrsonimae Kern. & Kellerm. Leaf and stem rust on *B. crassifolia* in Brazil and Guatemala.

Aecidium byrsonimaticola P. Henn. Leaf rust on *Byrsonima* sp. in Brazil.

Aecidium byrsonimatis P. Henn. Leaf and stem rust on *B. verbascifolia* and *Byrsonima* sp. in Brazil and Trinidad.

Aecidium singulare (Diet and Holw.) Arth. Leaf rust on *B. crassifolia* in Guatemala.

Cronartium byrsonimatis P. Henn. Leaf rust on *B. coccobolifolia* in Brazil.

Cronartium notatum Arth. Brown leaf rust on *B. crassifolia* in Cuba and Porto Rico.

Uredo byrsonimatis P. Henn. Leaf rust on *Byrsonima* sp. in Brazil.

Uredo uberabensis P. Henn. Leaf rust on *Byrsonima* sp. in Peru and Brazil.

CACALIA. Perennial herbs.

Coleosporium cacaliae Othth. Golden-yellow rust pustules on needles of *Pinus montana*, *P. silvestris*, and on leaves of *Adenostyles albifrons*, *A. alliariae*, *A. alpina*, *Cacalia atriplicifolia*, *C. farfaraefolia*, *C. hastata*, *C. pubescens*, and *C. suaveolens* in Japan, Siberia, Russia, Spain, Bohemia, Montenegro, Denmark, Great Britain, France, Italy, Switzerland, Austria, Hungary, and Germany.

Puccinia senecionicola Arth. See Senecio.

Puccinia tranzschelii Diet. Dark-brown rust pustules on leaves of *C. hastata* in Russia.

Rhytisma hysteroideis Fr. On stems of *C. hastata* in Siberia.

Uromyces cacaliae (DC.) Wint. See *Adenostyles*.

Uromyces veratri (DC.) Schroet. See *Veratrum*.

CACARA. See *Pachyrhizus*.**CAESALPINIA**. Ornamental tropical shrubs or trees grown for showy flowers and attractive leaves.

Some species yield tanning material and dye stuffs.

Helminthosporium bonducellae P. Henn. Leaf spot of *C. bonducella* in Brazil.

Laestadia caesalpiniae Pat. Pale-brown spots on leaves of *Caesalpinia* sp. in Java.

Ravenelia cohniiana P. Henn. Cinnamon-brown rust pustules on leaves and stems of *Caesalpinia* sp. in Brazil.

Ravenelia humphreyana P. Henn. Cinnamon-brown to dark-brown rust pustules on leaves of *C. bahamensis* and *C. pulcherrima* in Porto Rico, Jamaica, Mexico, and Guatemala.

Ravenelia inconspicua Arth. Rust on leaves of *Caesalpinia* sp. in Mexico and *C. exostemma* in Guatemala.

CAJANUS. PIGEON PEA. Sometimes called Congo pea or gandul. Cultivated for the edible peas.

Aecidium cajani Petch. Rust producing small galls on under surfaces of leaves and on petioles and stems of *C. indicus* in Ceylon. Probably should be referred to *Woroninella umbilicata*.

Cercospora cajani P. Henn. (*Velloziella cajani* Rangel.) Small circular to irregular dark-brown spots on leaves of *C. indicus* in Porto Rico, Cuba, and Brazil.

Cercospora instabilis Rangel. Small angular, then confluent, dark-brown spots on leaves, branches and pods of *C. indicus* in Brazil.

Colletotrichum cajani Rangel. Anthracnose on leaves of *C. indicus* in Brazil.

Corticium salmonicolor B. and Br. See *Citrus*.

Diplodia cacaoicola P. Henn. See *Theobroma*.

CAJANUS—Continued.

Fusarium udum Butl. This fungus produces a gradual, or at times, sudden wilting and dying of infected plants. Tissues at the base of the stems and along the main roots of diseased plants are blackened either entirely or in streaks. A serious disease in India on the pigeon pea (*C. indicus*). See also *Dianthus*.

Heterosporium lagunense Syd. On *C. indicus* in the Philippines.

Phoma cajani Rangel. On stems of *C. indicus* in Brazil.

Phyllosticta cajani Rangel. On leaves of *C. indicus* in Brazil.

Rosellinia bunodes B. and Br. See *Citrus*.

Uredo cajani Syd. Rust on leaves of *C. indicus* in India.

Woroninella umbilicata (B. and Br.) v. Hoeh. Minute, red swellings on the stems and leaves which burst, exposing orange-red powdery spore masses. On *C. indicus* in Ceylon.

CALACINUM. See *Muehlenbeckia*.

CALADIUM. Large-leaved aroids grown for their variegated foliage. Herbaceous perennials.

Cercospora caladii Cke. Circular dull-brown leaf spots on *Caladium* sp. in India.

Gloeosporium arecarum P. Henn. Circular to irregular brown spots, with darker brown surrounding zones, on leaves of *Caladium* sp. and *Philodendron bipinnatifidum* in Japan and Brazil. Hemmi in Japan has described what is probably the same species under the same name.

Helmintosporium caladii Stevens. Large (up to 1.5 to 2 centimeters), circular or oval ashen white spots, with tan-colored borders, on leaves of *C. bicolor* in Porto Rico.

Macrophoma surinamensis (B. and C.) Berl. and Vogl. On leaf blades and petioles of *Caladium* spp. in Surinam.

CALAMAGROSTIS. Hay and forage grasses.

Dilophia graminis (Fckl.) Sacc. See *Triticum*.

Leptosphaeria sparsa (Fckl.) Sacc. See *Poa*.

Metasphaeria avenae (Auersw.) Sacc. See *Avena*.

Mycosphaerella calamagrostidis Volk. On leaves of *C. varia* in Switzerland.

Mycosphaerella lincolata (Desm.) Schroet. On leaves of *Anemophila* sp., *Calamagrostis epigejos*, and *Phragmites communis* in Great Britain, Denmark, France, Italy, and Germany.

Mycosphaerella perforans Desm. On leaves of *C. arenaria* in Denmark, France, and Holland.

Mycosphaerella pusilla (Auers.) Johans. On leaves of *C. arenaria* in Finland, Denmark, and Germany.

Puccinia borealis Juel. See *Agrostis*.

Puccinia brevicornis S. Ito. Powdery golden-yellow and black rust pustules on leaves of *C. canadensis* and *C. villosa* in Japan.

Puccinia epigejos Ito. As in the preceding species on *C. arundinacea* and *C. epigejos* in Japan.

Puccinia erythropus Diet. Brown to black rust pustules on purple leaf spots on *C. sciurioides* and *Miscanthus sinensis* in Japan.

Puccinia ishikawai S. Ito. Golden-yellow and black rust pustules, on brown sunken spots, on leaves of *C. epigejos* var. *densiflora* in Japan.

Puccinia pertenuis Ito. Rust on leaves of *C. arundinacea* in Japan.

Puccinia pygmaea Erikss. Golden-yellow and dark-brown rust pustules on leaves of *C. arundinacea*, *C. epigejos*, and *C. halleriana* in Sweden, Finland, and Germany.

Puccinia rangiferina Ito. Powdery black rust pustules on leaves and sheaths of *C. arundinacea*, *C. hillebrandii*, and *Melica* sp. in Japan.

Puccinia stichosora Diet. Rust on leaves of *C. sciurioides* in Japan.

Septoria arenariae Rostr. On leaves of *C. arenaria* in Denmark.

Septoria calamagrostidis (Lib.) Sacc. White linear spots on leaves of *C. arundinacea*, *C. epigejos* and *C. sylvatica* in Russia, Denmark, and France.

Septoria epigejos Theum. On leaves of *C. epigejos* in Russia and Denmark.

Septoria nebulosa Rostr. Indefinite areas on leaves and culms of *Calamagrostis* sp., *Poa* sp., and *Trisetum* sp. in Greenland.

Septoria rhizodes Bres. and Krieg. On leaves of *C. halleriana* in Germany.

Tilletia calamagrostidis Fckl. Long linear smut pustules on the leaves which break open exposing black spore masses. On *C. arundinacea*, *C. epigejos*, *C. halleriana*, and *C. lanceolata* in Germany.

Uredo paulensis P. Henn. Rust on leaves of *Calamagrostis* sp. in Brazil.

CALAMUS. See *Palmae*.

CALANDRINIA. ROCK PURSLANE. Fleshy, trailing plants used in borders or rockeries.

Heterosporium calandriniae Mass. Irregular pale olive-colored spots on leaves and inflorescences of *C. potentilloides* in Argentina.

Peronospora calandriniae Speg. Downy mildew on leaves of *C. portulacoides* in Argentina.

Ustilago calandrinicola Speg. Forms dark violet powdery spore masses in inflorescences of *C. affinis* in Argentina.

CALATHEA. Tropical perennial foliage plants.

Niptera calathea P. Henn. On leaves of *Calathea* sp. in Peru.

CALCEOLARIA. FAGELIA. Ag. Showy-flowered herbs and shrubs.

Aphelenchus olesistis Ritz. Bos. See *Begonia*.

Entyloma calceolariae Lagerh. Smut producing sori in circular whitish spots on leaves of *Calceolaria* sp. in Ecuador.

Peronospora indica Syd. Downy mildew on leaves of *C. scabiosaefolia* in India.

Phyllosticta calceolariae Speg. Elliptical, definite whitish spots on leaves of *C. foliosa* in Argentina.

CALENDULA. POT MARIGOLD. Flower garden annuals or perennials.

Cercospora calendulae Sacc. Subcircular, dull-brown leaf spots on *C. officinalis* in Italy and Portugal.

Entyloma calendulae (Oud.) De By. Smut sori in circular, greenish or white, finally brown, spots on leaves of *Arnica montana*, *Arnoseria minima*, *Aster alpinus*, *Bellidistram michelii*, *Calendula arvensis*, *C. officinalis*, *C. pluvialis*, *Cineraria papposa*, *Crepis biennis*, *Erigeron acer*, *Hieracium murorum*, *H. pilosella*, *H. schmidtii*, *H. silvaticum*, *H. stygium*, *H. sudeticum*, *H. tridentatum*, *H. villosum*, *H. vulgatum*, *Leontodon hispidus*, and *Parthenium hysterophorus* in Argentina, Algeria, and throughout Europe. American references to this species apparently are referable to *E. arnicale* E. and E. and other species.

Puccinia calendulae McAlp. Rust on leaves, stems, and fruits of *C. officinalis* in Australia.

CALIMERIS. Daisylike border plants.

Puccinia calimeris Syd. Dark-brown rust pustules on leaves of *C. altaica* in Siberia.

CALLA. Wild calla. Bog plants.

Ascochyta pellucida Bub. Large irregular spots on leaves of *C. palustris* in Bohemia.

CALLIANDEA. Evergreen shrubs and trees.

Ravenelia affinis Syd. Yellow-brown and deep-brown rust pustules on leaves of *C. turbinata* in Brazil.

Ravenelia armata Syd. Brown leaf rust on *Calliandra* sp. in Brazil.

CALLIANDRA—Continued.

Ravenelia bizonata Arth. and Holw. Powdery brown rust pustules on leaves of *C. houstoni* and *Calliandra* sp. in Guatemala.

Ravenelia dieteliana P. Henn. Brown rust pustules on leaves of *C. macrocephala* in Formosa and Brazil.

Ravenelia echinata Lagerh. and Diet. Cinnamon-brown to black rust pustules on leaves of *Calliandra* sp. in Ecuador.

Ravenelia ectypa Arth. and Holw. Brown leaf rust on *C. gracilis* and *Calliandra* sp. in Costa Rica and Guatemala.

Ravenelia lagerheimiana Diet. Brown rust pustules on leaves of *Calliandra* sp. in Ecuador.

Ravenelia mexicana Tranzsch. Brown rust pustules on leaves of *C. grandiflora* in Mexico.

Ravenelia pazschkeana Diet. Brown to black rust pustules on leaves of *Calliandra* sp. in Brazil.

Uredo longipedis P. Henn. Dull yellow-brown rust pustules on brown subcircular leaf spots on *C. longipes* in Brazil.

CALLICARPA. BEAUTY BERRY. Ornamental woody plants cultivated chiefly for their bright-colored fruit.

Uromyces callicarpae (Petch.) Fuji. Brown rust pustules on leaves of *C. formosana*, *C. lanata*, and *C. tomentosa* in Ceylon and Japan.

CALLISTEMMA. See *Callistephus*.

CALLISTEMON. BOTTLE BRUSH. Hardy ornamental shrubs.

Phyllosticta flavidula Sacc. On leaves of *Callistemon* sp. in France.

Septoria callistemonis F. Tassi. On leaves of *Callistemon* sp. in Italy.

CALLISTEPHUS. **CALLISTEMMA** Ag. CHINA ASTER. Garden annuals.

Coleosporium asterum (Diet.) Syd. See *Aster*.

Phyllosticta asteris Bres. Subcircular to angular brown spots on leaves of *C. chinensis* in Europe.

Phytophthora sp. The "black neck" or wilt disease of China asters in Great Britain. Plants of all ages wilt and die, the lower parts of the stems becoming brown or black. The causal organism is said to be close to *P. omnivora*, but is considered distinct.

CALLUNA. HEATHER. Low evergreen shrubs.

Metasphaeria callunae Fautr. On branches of *C. vulgaris* in France.

Plowrightia polyspora (Bref.) Sacc. Small subcircular stromata on branches of *C. vulgaris* in Germany.

CALONYCTION. MOONFLOWER. Twining perennial herbs.

Septocylindrium platense Speg. Indefinite brown leaf spots on *C. aculeatum* (*C. speciosum*) in Argentina.

CALOTROPIS. MODAR-BUSH. Milkweedlike shrubs or small trees.

Ascochyta tripolitana Sacc. and Trot. Circular, definite, whitish spots with reddish margins, on leaves of *C. procera* in north Africa.

Cercospora calotropidis Speg. Indefinite yellow leaf spots on *C. procera* in Senegal. This species should be renamed, the name being occupied by the following species.

Cercospora calotropidis E. and E. Large dirty-brown spots on leaves of *C. procera* in the Bahamas and Central America.

Cercospora inconspicua Pat. and Har. Circular spots on leaves of *Calotropis* sp. in French North Africa.

Cercospora patouillardii Sacc. and D. Sacc. On leaves of *C. procera* in Guadalupe.

Napcladium calotropidis Mors. Circular, often confluent, yellow-green to olive-green spots on leaves and stems of *C. procera* in Tanganyika.

CALPUERNIA. Tropical shrubs and trees.

Ascochyta calpurniae Wint. Angular to irregular, white or dirty-gray spots, with dull purple margins, on leaves of *C. sylvatica* in the Union of South Africa.

Cercospora calpurniae Petch. Diffuse yellow areas on leaves of *C. aurea* in Ceylon.

Ravenelia glabra Kalkbr. and Cke. Brown rust pustules on leaves of *C. sylvatica* in the Union of South Africa.

CALTHA. MARSH MARGOLD. Hardy, blooming marsh plants.

Cercospora calthae Cke. Circular dull-brown leaf spots on *Caltha* sp. in Great Britain and Esthonia.

Physoderma bohemicum Sacc. Small brown galls on leaf blades and petioles of *C. palustris* in Bohemia.

Puccinia zopfi Wint. Chestnut-brown to dark-brown rust pustules on leaves of *C. leptosepala* and *C. palustris* in Argentina, Japan, Great Britain, France, Denmark, Belgium, Holland, Russia, Switzerland, Italy, Austria, and Germany.

Ramularia calthae Lindr. (*R. calthae* Gz. Frag.) Irregular whitish spots, with brown margins on leaves of *C. palustris* in Spain, Bohemia, Denmark, Finland, Austria, and Germany.

CALYCANTHUS. SWEET SHRUB. Ornamental shrubs with fragrant flowers.

Ascochyta calycanthi Sacc. and Speg. and var. **occidentalis** Fl. Tass. Leaf spots on *C. floridus* and *C. occidentalis* in Italy.

Phyllosticta calycanthi Sacc. and Speg. Brown spots, becoming white, on leaves of *C. orientalis* and *Meratia praecox* (*C. praecox*) in Denmark and Italy.

Septoria calycanthi Sacc. and Speg. Dull-yellow to brown spots on leaves of *C. floridus* and *C. laevigatus* in Italy and Portugal.

CAMELLIA. Woody plants grown for their showy white or red flowers and handsome evergreen leaves.

Ascochyta minutissima Pass. On young branches of *C. japonica* in Italy.

Ascochyta camelliae (Pass.) Fl. Tass. Oblong or irregular, grayish-white leaf spots on *C. japonica* in France.

Exobasidium assamense Syd. and Butl. Circular, then confluent and irregular, white areas on leaves of *C. drupifera* in India.

Exobasidium camelliae Shirai. See *Thea*.

Helminthosporium polyphragmum Syd. On stems of *C. japonica* in India.

Leptosphaeria camelliae Cke. and Mass. Brown indefinite spots on leaves of *Camellia* sp. in Australia.

Phyllosticta amphigena D'Alm. Brown leaf spots on *C. japonica* in Portugal.

Phyllosticta camelliae West. and var. **meranensis** Bub. On leaves of *C. japonica* in Denmark, France, and Austria, and on *Thea* (tea) in Japan.

CAMPANULA. BELLFLOWER. HAREBELL. CANTERBURY BELLS. Hardy herbaceous perennials.

Ascochyta bohemica Kab. and Bub. Circular to angular, often confluent, small yellowish or brown spots with purple margins on leaves of *C. trachelium* in Bohemia.

Ascochyta carpathica (Allesch.) Keissl. Small oblong or irregular whitish spots with brown margins on leaves of *C. carpathica* in Germany.

Clathrosorus campanulae Ferd. and Winge. Small swellings on roots of *C. rapunculoides* in Denmark.

CAMPANULA—Continued.

- Marsonia campanulae** Bres. and Allesch. Small subcircular dull-yellow leaf spots on *C. latifolia* and *C. morettiana* throughout Europe.
- Peronospora corallae** Tranzsch. Downy mildew attacking the flowers of *C. persicifolia* in Russia and Sweden.
- Phyllosticta alliariaefolia** Allesch. Large subcircular to angular ashen spots, with indefinite margins, on leaves of *C. alliariaefolia* in Germany.
- Phyllosticta campanulae** Sacc. and Speg. Yellowish-brown spots with darker margins on leaves of *C. glomeratum* and *C. trachelium* in Italy and Siberia.
- Phyllosticta campanulina** Moesz. Whitish spots with red-brown margins on leaves of *C. persicifolia* in Hungary.
- Phyllosticta michauxioides** P. Magn. Large leaf spots, yellow above, black beneath, on *C. michauxioides* in Asia Minor.
- Phyllosticta rehml** Bub. On leaves of *C. trachelium* in Hungary.
- Placosphaeria campanulae** (DC.) Bäuml. Black stromata on leaves of *C. bononiensis*, *C. rapunculoides*, and *C. trachelium* in France, Yugoslavia, Switzerland, Austria, and Germany.
- Pseudopeziza radicans** Sacc. On leaves of *C. patula*, *C. rapunculoides*, and *C. trachelium* in Great Britain, Finland, France, Belgium, and Denmark.
- Puccinia heideri** Wettst. Black rust pustules on leaves of *C. barbata* in Austria.
- Ramularia campanulae-barbatae** Jaap. and Lind. Subcircular brown spots on leaves of *C. barbata* in Switzerland.
- Ramularia campanulae-rotundifoliae** Lindr. Small gray or reddish spots on leaves of *C. rotundifolia* in Finland.
- Ramularia coleosporii** Sacc. and var. **campanulae** Car. On leaves of *C. rapunculoides*, *C. trachelium*, *Campanula* sp., *Hypericum androsaemum*, *Melampyrum* sp., *Senecio* sp., and *Telekia speciosa* in Europe.
- Ramularia macrospora** Fres. and var. **major** Lindr. Subcircular or irregular greenish-gray spots with ochraceous margins on leaves of *C. glomerata*, *C. latifolia*, *C. media*, *C. persicifolia*, *C. pyramidalis*, *C. rapunculoides*, and *C. trachelium* in Alaska and Europe.
- Ramularia michauxioides** P. Magn. On leaves of *C. michauxioides* in Asia Minor.
- Septoria campanulae** (Lév.) Sacc. On leaves of *Campanula* sp. in Persia.
- Septoria obscura** Trail. Dull brown circular or irregular leaf spots on *C. rotundifolia* in Scotland.
- Septoria trachelii** Allesch. Irregular spots on leaves of *C. trachelium* in Switzerland and Russia.
- CAMPANUMAEAE**. Twining perennial herbs.
- Coleosporium campanumaeae** Diet. Yellow rust pustules on leaves of *C. javanica* in Japan.
- Puccinia campanumaeae** Pat. Yellow-brown rust pustules on lower surfaces of leaves of *C. javanica* and *Campanumaea* sp. in China and Java.
- CAMPE**. See *Barbarea*.
- CANAVALLIA**. JACK BEAN. SWORD BEAN. Prostrate trailing herbs used for green manure crops.
- Cercospora canavaliae** Syd. Dull-red circular to irregular areas on leaves of *C. ensiformis* and *C. gladiata* in the Philippines.
- Cerotelium canavaliae** Arth. Brown rust pustules on red-brown leaf spots on *C. ensiformis*, *C. gladiata*, and *Canavalia* sp. in Porto Rico and French Congo.
- Eisinoe canavaliae** Racib. This fungus causes distortion of the leaves and stems of *C. ensiformis* and *C. gladiata* in Ceylon and the Philippines.
- Gloeosporium canavaliae** Syd. Anthracnose on stems of *C. ensiformis*, *C. gladiata* and *C. turgida* in Hawaii and the Philippines.
- Mycosphaerella canavaliae** Syd. Indefinite brown leaf spots on *C. gladiata* in the Philippines.
- Physalospora guignardioides** Sacc. On leaves of *C. ensiformis* and *C. gladiata* in the Philippines.
- Septoria canavaliae** Lyon. Small circular to irregular spots on leaves causing them to drop prematurely. The spots are dark brown to black, becoming confluent, with ashy-gray centers on *C. ensiformis* in Hawaii.
- Septoria molleriana** Bres. Circular pale-brown spots with reddish margins on leaves of *C. lineata* and *C. obtusifolia* in the Philippines and island of St. Thomas (Africa).
- CANDOLLEA**. See *Styidium*.
- CANNA**. Tall ornamental plants cultivated as bedding plants for the foliage and showy flowers. One species with edible root stocks.
- Asochyta cannae** Rangel. On leaves of *Canna* sp. (cult.) in Brazil.
- Bunchy-top**. See *Musa*.
- Puccinia cannae** (Wint.) P. Henn. Yellow to black rust pustules covering the under sides of leaves of *Maranta arundinacea*, *Thalia geniculata*, *Canna coccinea*, *C. glauca*, *C. indica*, and other cultivated species and varieties throughout Central America, the West Indies and South America. Occurs also to a limited extent in south Florida. This disease destroys the ornamental value of the canna to such an extent that its culture has been abandoned in some countries where the rust is prevalent. A bacterial stem disease of *Canna* has been reported from Ceylon.
- CANNABIS**. HEMP. Fiber plants and occasionally used as ornamentals.
- Cercospora cannabina** Wakef. Circular to elongate, often confluent, pale-brown leaf spots on *C. sativa* in Uganda.
- Dendrophoma marconii** Cav. On *C. sativa* in Italy.
- Leptosphaeria cannabina** Ferr. and Mass. On leaves of *C. sativa* in Italy.
- Peronospora cannabina** Oth. Downy mildew causing yellowing of leaves of *C. indica* and *C. sativa* in Japan, Switzerland, Russia, and Italy.
- Phyllachora cannabina** P. Henn. On *C. sativa* in Brazil. Referred to the *Clypeosphaeriaceae* by Theissen and Sydow.
- Phyllosticta cannabis** (Kirch.) Speg. Subcircular dark-brown spots on leaves of *C. indica* and *C. sativa* in Yugoslavia, Bohemia, Italy, and Russia.
- Rhizoctonia napi** West. See *Brassica*.
- Septoria cannabis** Sacc. Indefinite dirty-yellow spots on leaves of *C. sativa* in Russia, Italy, Belgium, and Germany. Probably the imperfect stage of *Leptosphaeria cannabina* Ferr. and Mass.
- Uredo kriegeriana** Syd. Brown rust pustules on leaves of *C. sativa* in Germany.
- CAPNOIDES**. See *Corydalis*.
- CAPRIOLA**. See *Cynodon*.
- CAPSCICUM**. RED PEPPER. Sometimes called Cayenne pepper. Herbs or shrubs cultivated for their fruit.
- Acrothecium capsicii** Turc. Hazel-colored circular to oval depressed areas with brown margins on fruit of *C. annuum* in Italy.
- Bacillus capsicii** Pavar. and Tarc. Bacterial fruit rot and wilt of plants of *C. annuum* in Italy.
- Phomopsis capsicii** (Magn.) Sacc. On harvested fruit of *C. annuum* in Italy and the Philippines.
- Phyllosticta capsicii** Speg. Circular grayish leaf spots with definite deep-brown margins on *C. annuum* and *Capsicum* sp. in Argentina.

CAPSICUM—Continued.

Puccinia capsici Mayor. Rust on leaves of *C. baccatum* in Colombia. This species has priority over the following, which is probably not distinct.

Puccinia capsici Avena-Sacca. Rust pustules on leaves of *C. frutescens*, *C. microcarpum*, *C. odoriferum*, and *C. pendulum* in Brazil.

Puccinia gonzalezi Mayor. Rust on leaves of *Capsicum* sp. in Colombia.

Puccinia paulsenii Rangel. (*P. capsici* Avena?). Rust on leaves and twigs of *C. annuum* in Brazil.

Rhizoctonia palida Matz. See *Saccharum*.

Rosellinia arcuata Petch. See *Thea*.

Vermicularia capsici Syd. (*Steirochaete capsici* [Syd.] Sacc.). This fungus causes the most serious disease known of pepper in India, Ceylon, China, and the Philippines. Butler describes the disease as follows: "At first the flowers droop and dry up, the disease then spreading, if weather conditions are favorable, through the flower stalks to the stem. The bark turns brown and finally white. The upper portion of the plant may be killed. Round, sunken, anthracnose spots occur on the fruit, dark at first then pale at the center and coalescing. Diseased fruits shrivel and dry up." Known hosts are *Capsicum annuum*, *C. frutescens*, *Dolichos biflorus*, *D. lablab*, *Lycopersicon esculentum*, *Solanum melongena*, and *Vigna catjang*.

CARAGANA. PEA SHRUB. Siberian pea tree. Ornamental shrubs grown chiefly for their bright-yellow flowers, also as hedge plants.

Asochyta bondarzewii P. Henn. Dull-brown spots on leaves of *C. arborescens* in Russia.

Hendersonia septem-septata Vesterg. On leaves of *C. arborescens* in Russia.

Melasma caraganae Thum. Black areas on leaves and branches of *C. arborescens* in Siberia.

Mycosphaerella jazewskii A. Pot. (*Phyllosticta borszczowii* Thum.). On leaves of *C. arborescens* in Siberia, Russia, and Germany.

Peronospora lagerheimii Güm. Downy mildew on leaves of *C. arborescens* in Norway.

Phleospora caraganae Jacz. Indefinite yellow spots on leaves of *C. arborescens* in Russia.

Phyllachora crebia Syd. Black stromata on rust-colored spots on leaves of *Caragana* sp. in India.

Phyllosticta caraganae Syd. Whitish spots with brown margins, often at the top or along the margins of leaves of *C. chamglag* in Germany.

Phyllosticta caraganae Fl. Tass. On leaves of *C. arborescens* in France. This species should be renamed, the name being occupied as above.

Phyllosticta gallarum Thum. On leaves of *C. arborescens* in Siberia.

Phyllosticta spaethiana All. and Syd. Small irregular white spots with dull-brown margins on leaves of *C. arborescens* in Russia, Austria, and Italy.

Septoria caraganae P. Henn. Yellow then brown leaf spots on *C. arborescens* and *Caragana* sp. in Russia and Germany.

Tromyces genistae-tinctoriae (Pers.) Wint. Powdery cinnamon-brown rust pustules on under leaf surfaces of *C. arborescens*, *C. chamglag*, *C. frutescens*, *C. fruticosa*, *C. pygmaea*, *Colutea arborescens*, *C. orientalis*, *Cytisus alpinus*, *C. austriacus*, *C. biflorus*, *C. capitatus*, *C. decumbens*, *C. falcatus*, *C. heuffelii*, *C. hirsutus*, *C. laburninus*, *C. liberus*, *C. linifolius*, *C. nigricans*, *C. proliferus*, *C. ramentaceus*, *C. ratisbonensis*, *C. sagittalis*, *C. sessilifolius*, *C. supinus*, *Galega officinalis*, *G. orientalis*, *Genista angelica*, *G. germanica*, *G. pilosa*, *G. sagittalis*, *G. sericea*, *G. tinctoria*, *Laburnum anagyroides*, *Sorothamnus scoparius*, and *Tetragonolobus siliquosus* in Argentina, Japan, Siberia, Canary Islands, and Europe.

CARDAMINE. BITTER CRESS. CUCKOO-FLOWER. Small leafy-stemmed perennials.

Phyllosticta cardamines Allesch. On leaf blades and petioles of *C. bellidifolia* in Greenland.

Puccinia cardamines-bellidifoliae Diet. Chestnut-brown rust pustules on leaf blades and petioles of *C. bellidifolia* in Norway and Greenland.

Puccinia cardamines-cordatae Diet. and Neg. Yellow-brown rust pustules on leaf blades and petioles of *C. cordata* and *C. reniformis* in Chile.

Puccinia cruciferarum Rud. Red-brown rust pustules on leaves of *C. alpina*, *C. asarifolia*, *C. bellidifolia*, *C. gelida*, *C. pratensis*, and *C. resedifolia* in Iceland, Spain, Switzerland, France, Belgium, Austria, and Hungary.

Ramularia cardamines Syd. Circular to irregular spots on leaves of *C. amara* and *C. pratensis* in Switzerland, Denmark, Bohemia, and Germany.

Septoria cardamines Fekl. Dark-brown effuse spots on leaves and stems of *C. amara* and *C. pratensis* in Germany.

Septoria cardamines-resedifoliae Heim. Small brown leaf spots, becoming whitish, on *C. resedifolia* in Tyrol.

Septoria cardamines-trifoliae v. Hoehn. Subcircular whitish spots with dull brown margins on both leaf surfaces of *C. trifolia* in Austria.

CARDIOSPERMUM. HEARTSEED. BALLOON VINE. Tendril climbing tropical herbs.

Aecidium cardiospermi Cke. Yellow rust pustules on yellow-brown leaf spots on *C. halicacabum* (*C. microcarpum*) in Zanzibar.

Cercospora cardiospermi Petch. On leaves of *C. halicacabum* in Ceylon.

Phyllosticta cardiospermi Speg. Circular spots on leaves of *C. retinutrum* in Argentina.

CARICA. PAPAYA. Small rapid-growing unbranched trees of the Tropics with milky juice. One species with edible fruit.

Asochyta caricae Pat. Dark linear or ovate spots on petioles becoming ashen on *C. papaya* in Ecuador.

Astericia caricarum Rehm. On leaves of *C. papaya* in Brazil.

Cercospora caricae Speg. Small, pale, indefinite spots on leaves of *C. papaya* in Brazil.

Diplodia cacaecicola P. Henn. See *Theobroma*.

Reimithosporium papayae Syd. Definite circular ochraceous concentrically zoned leaf spots on *C. papaya* in the Philippines.

Mycosphaerella caricae Syd. On leaves of *C. papaya* in the Philippines and Ceylon.

Ovulariopsis papayae Van der Bijl. Powdery mildew on lower leaf surfaces of *C. papaya* in the Union of South Africa. A powdery mildew, which may be the same, occurs in Florida and other countries where the papaya is grown.

Phyllosticta caricae-papayae Allesch. On leaves of *C. papaya* in Brazil and the Union of South Africa.

Phyllosticta papayae Sacc. Small subcircular whitish spots on fruit of *C. papaya* in Abyssinia.

Phytophthora faberi Maubl. See *Theobroma*.

Sphaerostilbe repens B. and Br. See *Hevea*.

CAEISSA. Tropical spinose shrubs cultivated for ornament and hedges as well as for the edible berrylike fruit.

Leveillina arduinae (K. and C.) Theiss. and Syd. Black irregular stromata on upper leaf surfaces of *C. arduina* in the Union of South Africa.

Phyllosticta carissae Kalchb. and Cke. Subcircular pale-brown leaf spots with brown margins on *C. arduina* in south and central Africa.

Puccinia carissae Cke. and Mass. Dark-brown rust pustules on leaves of *C. orata* in Australia.

- CARLUDOVICA.** HAT PALM. Palmlike plants of tropical America, used in making Panama hats.
Microthyrium carludovicae P. Henn. Dark-brown effuse spots on leaves of *Carludovica* sp. in Brazil.
- CARPINUS.** HORNBANE. Shade and timber trees.
Ascochyta carpinea Sacc. Subcircular or sinuous dull-yellow leaf spots on *C. betulus* in Italy.
Exoascus carpini Rostr. Produces yellowing and distortion of leaves and witches'-brooms of *C. caroliniana* and *C. betulus* in Europe.
Gloeosporium carpini (Lib.) Desm. Irregular indefinite olive-brown anthracnose spots on leaves of *C. betulus* in France, Austria, and Germany.
Melampsorium carpini (Eckl.) Diet. Yellow to yellow-brown rust pustules on lower leaf surfaces of *C. betulus*, *C. cordifolius*, and *C. yedoensis* in Japan, Russia, Italy, Switzerland, Austria, Hungary, and Germany.
Phyllosticta carpinea Sacc. Subcircular or sinuous dull-yellow leaf spots with brown margins on *C. betulus* and *C. orientalis* (*C. duinensis*) in Italy.
Phyllosticta carpini Schulz. and Sacc. On leaves of *C. betulus* in Russia.
Sclerotium carpini West. Round black sclerotia on living leaves of *Carpinus* sp. in Belgium.
Uncinula geniculata Ger. var. *carpinicola* K. Hara. Powdery mildew on leaves of *Carpinus* sp. in Japan.
- CARTHAMUS.** SAFFLOWER. Hardy annuals.
Cercospora carthami Syd. On leaves of *C. tinctorius* in Russia.
Cercosporina carthami Syd. Foliage frequently entirely destroyed and plants killed by this leaf disease of *C. tinctorius* in the Philippines. The spots are circular to irregular and gray-brown in color.
Erysiphe taurica Lévl. See Althaea.
Gloeosporium carthami Hori and Hemmi. Brown elliptic to irregular sunken areas on leaves and stems, in which pink spore masses appear. The tips of diseased plants wilt and fall over. On *C. tinctorius* in Japan.
Puccinia carthami (Hutzel.) Cda. Brown to dark-brown rust pustules on leaves of *C. oxyacanthus* and *C. tinctorius* in Japan, India, Egypt, Austria, and Germany.
- CARUM.** CARAWAY. Annual or perennial herbs, some yielding aromatic and edible garden products.
Cercospora cari Westerd. and Luyk. Sunken brown spots with dark-brown margins occur on petioles and flower stalks of *C. cari* in Holland.
Protomyces macrosporum Ung. See Coriandrum.
Puccinia dictyospora Tranzsch. The yellow aecia of this rust form elongate swellings along the veins of the leaves with brown telial pustules following on *Carum* sp. and *Scaligeria hirtula* in Turkestan. Possibly the same as the following species.
Puccinia microsphenetia Lindr. Black rust pustules on leaves and stems of *C. atosanguineum* in Turkestan.
Urophylctis kriegeriana P. Magn. Small pearl-like galls formed on the surface of leaves, stems, and floral parts. These galls are depressed at the apex and hollow. On *C. cari* and *C. persicum* in Switzerland and Germany.
- CARYOPHYLLUS.** EUGENIA. Malabar plum. Rose apple. Tropical ornamental and fruit trees.
Puccinia jambosae P. Henn. Brown rust pustules on leaves of *C. jambos* in Brazil.
Puccinia jambulana Rangel. Brown rust pustules on leaves and fruit of *C. jambos* in Brazil.
Puccinia psidii Wint. See Psidium.
Roeselia bunodes B. and Br. See Citrus.
Uredo flavidula Wint. Rust on leaves of *C. jambos* in Brazil.
- CARYOTA.** See Palmae.
- CASIMIROA.** WHITE SAPOTE. Tropical evergreen trees. One species cultivated for its fruit.
Bacterium citri Hassé. See Citrus.
Cercospora coteroides Sacc. Subcircular then confluent gray leaf spots with dark-brown margins on *C. edulis* in Mexico.
- CASTANEA.** CHESTNUT. Nut and timber trees.
Actinopelte japonica Sacc. On leaves of *C. vesca* in Japan. Reported from New Jersey.
Bacterium castanicolum Carvara. Said to cause galls at the base of stems of seedlings of *Castanea* sp. in Italy and to be distinct from *B. tumefaciens*.
Blepharospora cambivora Petri. See Ink disease.
Cylindrosporium castanicolum (Desm.) Berl. Dark-brown indefinite spots on both leaf surfaces of *C. vesca* (*C. sativa*) in Europe.
Diplodina castaneae Prill. and Delacr. Forms cankerous swellings on branches of *C. vesca* in France and Italy.
Endothia parasitica (Murr.) Ander. and Ander. This serious disease, native of Japan and China, has been introduced into the United States and Belgium. The chestnut forests of the eastern United States have been nearly exterminated, entailing a loss of many millions of dollars by this fungus, which in its native home is a comparatively harmless wound parasite. *C. mollissima*, the Chinese chestnut, is very resistant. *C. dentata*, the American chestnut, and other species of chestnut and chinquapin, are extremely susceptible. The disease appears as sunken cankers on the bark of trunks and limbs, which increase rapidly in size until girdling is effected and the limb, or the entire tree, dies. The wood is not attacked, the fungus working in the inner bark, where its presence can be noted by the characteristic fan-shaped mycelium. Diseased areas soon become covered with yellow, orange, or reddish-brown fruiting pustules. In damp weather spores are extruded in long irregular yellow or greenish horns.
Fomes korthalsii (Lévl.) Cke. Causes heart and butt rot of *C. argentea* and *C. javanica* in Java.
Guignardia diffusa (Crie.) Sacc. and Trott. On leaves of *Quercus* sp. and *Castanea* sp. in France.
Lanomyces tibodensis Güm. On leaves of *C. argentea* in Java.
Melanconis modonia Tul. (*Coryneum perniciosum* Br. and Farn.) On dead branches of *C. vesca* in France, Great Britain, Italy, and Germany, but reported as "a very destructive" parasite in the Caucasus. Associated with the so-called ink disease, (q. v.).
Monochaetia pachyspora Bub. Large circular brown leaf spots on *Castanea* sp. in Bohemia.
Mycosphaerella punctiformis (Pers.) St.-rb. (*Phyllosticta punctiformis* Sacc.) Irregular, dull, yellowish-red leaf spots on *C. vesca* in Italy and Dalmatia.
Phoma endogena Sacc. Causes a hard rot of the kernels of the nuts of *C. vesca* in Italy.
Phyllosticta nubecula Pass. On leaves of *C. vesca* in Italy.
Plowrightia noxia (Ruhl.) Sacc. On *Castanea* sp., *Fagus* sp., and *Quercus* sp. in Germany. Said to be not a *Plowrightia*, and its systematic position uncertain.
Pucciniastrum castaneae Diet. Yellow rust pustules on leaves of *Castanopsis javanica* and *Castanea vesca* in Japan, India, and the Philippines.
Sclerotinia candolleana (Lévl.) Eckl. See *Quercus*.
Scolecodothis castaneae Raeb. Brown-black stromata on the upper sides of leaves, with corresponding yellowish-red spots beneath, on *C. argentea* in Java.
Septoria castaneae Lévl. On leaves of *C. vesca* in France and Austria.
Tubercularia confuens Pers. On leaves of *C. vesca* (*C. vulgaris*) in Spain.

CASTANEA—Continued.

Uredo castaneae P. Henn. Brown rust pustules on leaves of *C. vesca* in the Philippines.

Black spot. A disease of unknown origin attacking the fruit of *Castanea vesca* in Italy. The nuts are normal in appearance until cut open, when the kernel is found dark brown or black in color.

Ink disease. The so-called ink disease of European chestnut (*C. vesca*) has caused heavy losses in France, Italy, Portugal, and Spain. A number of causes, including several fungi and various environmental influences, have been assigned by various workers. Diseased trees wilt and die, and upon close examination the roots are found dead and rotted, the bark at the base of the trunk being similarly affected. There is a black liquid exudate, rich in tannin, from the roots and trunk which gives the characteristic name to the disease. The disease spreads in irregular circles from the original infection points, trees dying in from 2 to 3 years after being attacked. Some workers have assigned *Melanconis monodia* Tul. (or its imperfect stage *Coryneum perniciosis* Br. and Farn.) as the cause through destruction of the bark. More recently a phycomycete, *Blepharospora cambitor* Petri, has been considered as the causative agent. This fungus is described as bringing about the formation of dark cankerous bands in the bark in the region of the root collar and out along the larger roots. This type of disease would be similar to the collar rot of *Citrus* (*Phytophthora terrestris* Sherb.).

A root rot, "gangrene humide," said to be due to bacteria, is reported from Portugal.

CASTANOPSIS. Chinquapin. Ornamental trees or shrubs.

Pucciniastrum castaneae Diet. See *Castanea*.

CASTILLA (CASTILLOA). MEXICAN RUBBER TREE.

Corticium salmonicolor B. and Br. See *Citrus*.

Diplodia cacaicola P. Henn. See *Theobroma*.

Fomes lamaoensis Murr. See *Hevea*.

Rosellinia bunodes B. and Br. See *Citrus*.

Uredo artocarpi B. and Br. See *Artocarpus*.

CASTILLEJA. PAINTED CUP. Herbs with showy bracts in terminal heads or spikes.

Puccinia nesodes Arth. and Holw. Brown rust pustules on leaves of *C. communis*, *C. tenuiflora* and *Lamoureauxia* sp. in Guatemala and Costa Rica.

CASUARINA. BEEFWOOD. Ornamental trees and shrubs with equisetum like leaves.

Botryosporium diffusum Cda. On branches of *C. leptoclada* in Russia.

Corticium salmonicolor B. & Br. See *Citrus*.

Rhizoctonia lamellifera Small. See *Grevillea*.

Sphaerostilbe repens B. & Br. See *Hevea*.

Trichosporium vesiculosum Butl. Causes a root disease of *C. equisetifolia* in India and possibly in Mauritius.

CATALPA. Ornamental trees cultivated for their flowers and foliage.

Ascochyta catalpae F. Tassi. Circular ashen-colored leaf spots with dull-brown margins on *C. speciosa* in Italy.

Phyllosticta bacillispora Kab. and Bub. Circular to angular brown spots becoming gray and often increasing so as to occupy entire leaf area on *C. syringifolia* in Bohemia.

Septoria catalpae Sacc. On pods of *C. bignonioides* and *C. syringifolia* in Italy.

CATASETUM. Epiphytic terrestrial orchids. See Orchidaceae.**CATTLEYA.** Epiphytic orchids. See Orchidaceae.**CEDRELA.** CIGAR-BOX CEDAR. Ornamental trees grown for their handsome foliage and as timber trees.

Cercospora sp. On leaves of *C. odorata* in Cuba.

Fomes lamaoensis Murr. See *Hevea*.

Phyllachora balansae Speg. Small black shiny stromata on leaf spots 0.5 to 2 centimeters in diameter on *C. brasiliensis*, *C. fissilis* and other species in Brazil, Argentina, and Paraguay.

Phyllosticta agnostolea Speg. Leaf spot of *Cedrela* sp. in Brazil.

Triphragmium cedrelae Hori. Yellow to black powdery pustules on both leaf surfaces of *C. chinensis* and *C. sinensis* in Japan.

Uncinula delavayi Pat. Powdery mildew on under leaf surfaces of *Ailanthus* sp. and *Cedrela sinensis* in China and Japan.

Uromyces cedrelae P. Henn. Dull-brown rust pustules on lower leaf surfaces of *C. serrata* in Java.

CEDRUS. CEDAR. CEDAR OF LEBANON. Valuable timber trees. Also grown for their handsome foliage and striking habit.

Peridermium cedri Barcl. A rust producing witches'-brooms with golden-yellow pustules on the needles of *C. deodora* in Northern India.

CEIBA. A fiber of great economic importance (silk cotton or kapok), is obtained from the pods. Also grown as shade trees.

Cercospora sp. A leaf spot on *C. pentandra* in Cuba.

CELASTRUS. BITTERSWEET. Woody climbers grown for their brightly colored fruit and handsome foliage.

Pucciniastrum celastri Syd. Brown rust pustules on leaves of *C. paniculatus* in India.

Uncinula sengokul Salmon. Powdery mildew on leaves of *C. articulatus* in Japan.

CELERI. See *Apium*.**CELOSIA.** COCKSCOMB. Garden annuals grown for their showy agglomerated flower heads, and sometimes colored foliage.

Cercospora gilbertii Speg. Circular gray leaf spots on *C. trigyna*, *Iresine celosioides*, and *I. paniculata* in Ecuador, Uruguay, and Argentina.

Phyllosticta celosiae Thum. On leaves of *C. cristata* in Italy and Portugal.

Uromyces celosiae Diet and Holw. Brown rust pustules on leaves of *C. latifolia*, *Iresine calea*, and *I. canescens* in Mexico, Cuba, Guatemala, and Costa Rica.

CELTIS. HACKBERRY. Shade trees.

Exoascus aemiliae Pass. Grayish-white spots on leaves of *C. australis* in Italy. Probably the same as the following species.

Exoascus celtidis (Sad.) Sacc. Brown spots or small excrescences on leaves of *C. australis* in middle and south Europe.

Mycosphaerella castagnel (Har. and Br.) Jaap. On leaves of *C. australis* in Dalmatia.

Napcladium celtidis Cav. Subcircular to angular, olivaceous spots on leaves of *C. latifolia* in Italy.

Phyllosticta celtidicola Bub. and Kab. Angular to irregular, dull-yellow or brown leaf spots on *C. australis* in Austria.

Phyllosticta destruens Desm. Whitish spots on leaves of *C. australis*, *Prunus laurocerasus* and *P. lusitanica* in France, Denmark, Russia, Italy, Portugal, and Austria. Reported from Wisconsin and Kansas.

Phyllosticta gel Bres. On leaves of *C. australis* in Europe.

Phyllosticta immersa Bub. Small circular sinuate greenish or yellow leaf spots on *C. australis* in Hungary.

CELTIS—Continued.

- Phyllosticta talae** Speg. Circular ashen-colored leaf spots on *C. tala* in Argentina.
- Septoria celtidis** Sacc. On leaves of *C. australis* in France and Italy.
- Sporodesmium celtidis** Syd. Small, definite circular gray leaf spots on *C. australis* in Yugoslavia.
- Uncinula kusanoi** Syd. Powdery mildew on under sides of leaves of *C. sinensis* in Japan.
- Uncinula shiraiana** P. Henn. Powdery mildew on lower leaf surfaces of *C. sinensis* in Japan.
- Uredo celtidis** Pазsch. Brown powdery rust pustules on lower leaf surfaces of *Celtis* sp. in Peru and Brazil.
- Uromyces celtidis** Diet. Rust on leaves of *Celtis* sp. in Brazil.
- CENTAUREA**. CORN FLOWER. BACHELOR'S-BUTTON. Annuals or half-hardy perennials, used for bedding, borders, etc.
- Ascochyta cyani** Cruch. On leaves of *C. cyanus* in Switzerland.
- Ascochyta schelliana** Thuem. On leaves and stems of *C. glastifolia* in Siberia.
- Bremia centaureae** Syd. Downy mildew on leaves of *C. jacea* and *C. montana* in Europe.
- Cercospora centaureae** Diet. Irregular ashen-colored spots on both leaf surfaces of *C. phrygia* in Germany.
- Cercospora centaureae** Syd. Small circular, or somewhat irregular, blackish leaf spots on *C. scabiosa* in Denmark and Germany.
- Cercospora tributiana** Sacc. and Let. Small angular leaf spots on *C. amara*, *C. nigrescens*, *C. phrygia*, and *C. pratensis* in Spain and France.
- Heterosporium centaureae** Ranof. On stems of *C. spinulosa* in Yugoslavia.
- Phyllosticta tagana** (Thuem.) Fl. Tass. On leaves of *C. tagana* in Portugal.
- Puccinia arenaricolor** Plowr. Yellow to brown rust pustules on leaves of *C. nigra* and *Carex arenaria* in Great Britain.
- Puccinia beltrani** Gz. Frag. Rust on leaves of *C. lingulata* in Spain.
- Puccinia carlicis-montanae** Ed. Fisch. Yellow rust pustules of *C. montana*, *C. nemoralis*, *C. nigra*, *C. nigrescens*, *C. plumosa*, and *C. scabiosa* in Russia, France, Switzerland, Austria, Hungary, and Germany.
- Puccinia centaureae** Mart. Brown rust pustules on many species of *Centaurea* spp. in India, Siberia, Asia Minor, Tripoli, and throughout Europe. Reported from several localities in the United States.
- Puccinia leuzeae** Syd. Cinnamon-brown powdery rust pustules on leaves of *Leuzea* (*Centaurea*) *carthamoides* in Siberia.
- Puccinia montana** Fekl. Orange-yellow, then chestnut-brown or dark-brown, rust pustules on *C. cana*, *C. montana*, and *C. phrygia* in Anatolia, Belgium, Italy, Switzerland, Yugoslavia, Austria, Hungary, and Germany.
- Puccinia obducens** Syd. Dark-brown powdery rust pustules on leaves of *C. ruthenica* in Turkestan.
- Puccinia pencana** Syd. Powdery cinnamon-brown to dark-brown rust pustules on leaves of *C. bulbosa* in Chile.
- Puccinia persica** Wettst. Yellow-brown to black powdery rust pustules on leaves of *C. belangerana*, *C. carduiformis*, and *C. kermanensis* in Persia.
- Puccinia tenuistipes** Rostr. Rust on leaves of *C. jacea* and *Carex muricata* in Yugoslavia, Sweden, Russia, Denmark, Italy, Austria, and Germany.
- Puccinia veruca** Thuem. Rust on leaves of *C. jacea*, *C. maculosa*, *C. montana*, *C. napifolia*, *C. nicaensis*, *C. orientalis*, *C. pullata*, *C. romana*, *C. scabiosa*, *C. sonchifolia* and *C. spinulifolia* in Egypt, Algeria, Corsica, Yugoslavia, Spain, Italy, Russia, Denmark, Austria, Hungary, and Germany.
- Puccinia vestita** Syd. Yellow-brown rust pustules on leaves of *C. myriocephala* in Kurdistan and Asia Minor.
- Pyrenochaete centaureae** Vogl. Indefinite gray or ashen-brown leaf spots on *C. candidissima* in Italy.
- Ramularia centaureae** Lindr. Small whitish spots, with dull-brown margins on leaves of *C. dealbata*, *C. nigra*, *C. phrygia*, *C. salicifolia*, and *C. scabiosa* in Denmark, Finland, Russia, France, Switzerland, and Germany.
- Ramularia centaureae atro-purpureae** Bub. Subcircular, dull-brown leaf spots with yellow centers on *C. atro-purpurea* in Hungary.
- Septoria aderholdi** Vogl. Indefinite, ashen-brown leaf spots on *C. candidissima* and *C. pseudophrygia* in Italy and Austria.
- Septoria centaureae** (Roum.) Sacc. Small olivaceous leaf spots on *C. jacea* and *C. nigra* in France.
- Septoria centaureicola** Brun. Circular, purple-brown leaf spots with white centers on *C. phrygia*, *C. scabiosa*, and *C. spinulifolia* in Russia, Yugoslavia, and France.
- Septoria collinae** Gz. Frag. On leaves of *C. collina* in Spain.
- Septoria cyani** Höll. Indefinite brown leaf spots on *C. cyanus* in Hungary.
- CENTURIUM**. See *Erythraea*.
- CENTRANTHUS**. JUPITER'S BEARD. Red valerian. Annual or biennial herbs.
- Aecidium centranthi** Thuem. Rust on leaves of *C. calcitrapa* in Spain, France, Switzerland, and Belgium.
- Puccinia valerianae** Carest. See *Valeriana*.
- Ramularia centranthi** Brun. Subcircular to irregular brown leaf spots on *C. ruber* in France.
- Septoria centranthicola** Brun. Angular to irregular, dirty-white leaf spots with narrow dull-green margins on *C. ruber* in France.
- CEPHALANTHUS**. BUTTONBUSH. Shrub grown for its attractive white flower heads.
- Aecidium cephalanthi-peruviani** P. Henn. Chestnut-brown rust pustules on red-brown leaf spots on *C. peruvianus* in Peru.
- Fusicladium cephalanthi** Speg. Superficial dense olivaceous fungal layers on branches, leaves, and young inflorescences, all of which as a result are more or less distorted. On *C. sarandus* in Uruguay and Argentina.
- CEPHALARIA**. Coarse annual or perennial herbs.
- Laestadia cephalariae** (Awd.) Sacc. See *Alternanthera*.
- Peronospora cephalariae** Vincens. Downy mildew on leaves of *C. alpina*, *C. leucantha*, and *C. transylvanica* in France.
- Phyllosticta cephalariae** Wint. Circular to irregular and finally confluent white leaf spots with dark purple margins on *C. attenuata* in the Union of South Africa.
- Septoria amicabilis** Boy. and Jacz. Brown leaf spots on *C. leucantha* in France.
- Septoria cephalariae-alpinae** Roum. Small, numerous, often confluent, whitish leaf spots with purple margins on *C. alpina* in France.
- Septoria scabiosicola** Desm. See *Scabiosa*.
- Venturia cephalariae** Kalchbr. and Cke. Dull-brown, circular leaf spots on *C. attenuata* in the Union of South Africa.
- CERASTIUM**. SNOW-IN-SUMMER. MOUSE-EAR CHICKWEED. Decumbent annuals or perennials, used in rockeries or for borders.
- Aecidium balearicum** Gz. Frag. Rust on leaves of *Cerastium* sp. in the Balearic Islands.
- Ascochyta alpina** Rostr. On leaves of *C. alpinum* in Norway.

CERASTIUM—Continued.

Ascochyta silenes E. and E. f. *cerastii* Sacc. On leaves of *C. arvense* in Italy.

Fabraea cerastiorum (Fr.) Rehm. Yellow spots on leaves and stems of *Cerastium* spp. and *Stellaria* sp. in Argentina, Australia, and throughout Europe.

Isariopsis alborosella (Desm.) Sacc. See *Stellaria*.

Peronospora helvetica Gäum. Downy mildew on leaves of *C. latifolium* in Switzerland.

Peronospora tornensis Gäum. Downy mildew on leaves of *C. alpinum* in Norway, Sweden, and Iceland.

Peronospora trivialis Gäum. As above on *C. arvense*, *C. semidecandrum*, and *C. triviale* in Europe and Argentina.

Septoria celanensis D. Sacc. Discoloring leaves of *C. tomentosum* in Italy.

Septoria cerasticola Rost. On leaves of *C. alpinum* in Denmark.

Septoria cerastii Desm. and Rob. Pale-yellow leaf spots, turning ashen-gray, on *C. pumilum*, *C. triviale* and *C. vulgatum* in Italy, Spain, France, Belgium, Russia, and Austria.

Septoria commersoniana Speg. On leaves of *C. humifusum* in Argentina.

Ustilago duriaana Tul. A smut producing brown powdery spore masses in ovaries of *C. arvense*, *C. brachypetalum*, *C. glomeratum*, and *Stellaria media* (*Alsine media*) in French and Italian north Africa, Great Britain, and Germany.

CERASUS. See *Prunus*.

CERATONIA. CAROB. A leguminous evergreen tree, the pods of which are used for forage and to some extent for human food.

Cercospora ceratonia Pat. and Trab. Small circular or angular brown spots on leaves of *Ceratonia* sp. in Algeria and Malta.

Phyllosticta calaritana Briosi. and Cav. Irregular marginal brown blotches on leaves of *C. siliqua* in Italy.

Phyllosticta ceratoniae Berk. Irregular black marginal spots on leaves of *C. siliqua* in Tunis, Russia, Italy, Malta, and Portugal.

Ramularia australis Sacc. Leaf spot on *C. siliqua* in Italy.

Septoria carrubi Pass. Dark-brown, irregular spots on leaves and pods of *C. siliqua* in France and Malta. Said to "menace the existence of the trees in this part of the country."

Septoria ceratoniae Pass. On leaves of *C. siliqua* in Italy and France.

CERCIDIPHYLLUM. Tree grown for its handsome foliage and habit.

Polythrincium shiraianum P. Henn. Circular to angular dull-brown leaf spots on *C. japonicum* in Japan.

CERCIS. REBUD. Judas tree. Trees or shrubs grown for their pink flowers.

Accidium lipskianum Jacz. A rust on leaves and pods of *C. siliquastrum* in Turkestan and Russia.

Ascochyta siliquastri Pass. On pods of *C. siliquastrum* in Italy.

Ovularia cercidis S. Cam. Irregular brown spots with deeper colored borders on leaves of *C. siliquastrum* in Portugal.

Phaeosphaerella japonica Shir. and Har. On leaves of *C. chinensis* in Japan.

Phyllosticta siliquastri Sacc. and Speg. Reddish leaf spots on *C. siliquastrum* in Italy and Portugal.

Septoria cercidis Fr. On leaves of *C. canadensis* and *C. siliquastrum* in Italy, France, Albania, Yugoslavia, and Russia.

CERCOCARPUS. Mountain mahogany. Small trees or shrubs cultivated for their attractive foliage and peculiar feathery-tailed achenes.

Ascochyta zonata Syd. Circular brown leaf spots (0.5 to 1 centimeter in diameter) on *C. parvifolius* in Germany.

Phyllosticta cercocarpi Syd. Small circular pale leaf spots with red margins on *C. parvifolius* in Germany.

CEREUS. Arboresecent, columnar forms of cactus.

Accidium cerei P. Henn. Rust on *Cereus* sp. in Argentina.

Cytosporella cerei Pöhl. On *Cereus* sp. in Italy.

Gloeosporium josephinae D. Sacc. Anthracnose on branches of *C. nycticalis* in Italy.

Leptothyrium parasiticum Pollacci. Large gray or ochraceous spots on stems of *C. stellatus* and *C. triangularis* in Italy.

Montagnella opuntiarum Speg. On *Cereus* sp. and *Cactus peruvianus* in Brazil and Paraguay.

Mycosphaerella cerei P. Henn. On phylloides of *Cereus macrogenus* in Brazil.

Septoria caeticola P. Henn. Pale-brown, indefinite spots, often confluent, on trunks of *C. pentagonus* in Germany.

Stagonospora assans Pass. On *Cereus* sp. and *Echinocactus* sp. in Italy.

CERINTHE. HONEYWORT. Annual or perennial herbs, with showy purple bracts.

Colosporium cerinthes Schroet. Golden-yellow rust pustules on leaves of *C. minor* in Russia and Germany.

Ramularia cerinthes Höll. Circular, black leaf spots, with dull-yellow margins on *C. minor* in Hungary.

CEROPEGIA. Tropical vines cultivated in greenhouses.

Accidium nummular Berk. Rust on leaves of *C. biflora*, *C. candelabrum*, and *C. cumingiana* in Ceylon and the Philippines.

Phyllachora ajrekari Syd. Irregular black stromata on both leaf surfaces of *C. hirsuta* in India and Japan.

CESTRUM. Ornamental tropical shrubs.

Didymosporium stromaticum Pat. On leaves of *Cestrum* sp. in Ecuador.

Phyllachora cestri Pat. Black stromata on leaves of *Cestrum* sp. in Ecuador.

Phyllosticta cestri Speg. Ashen leaf spots with reddish borders on *C. pubescens* in Argentina.

Phyllosticta humerispora Speg. On leaves of *Cestrum* sp. in Brazil.

Puccinia cestri Diet. and P. Henn. Brown rust pustules on lower leaf surfaces of *Cestrum* sp. in Brazil and Ecuador.

Septoria cestri (Mont.) Sacc. Circular white leaf spots with dull-brown margins on *C. parqui* in Chile.

Stigmatea cestri Pat. Black, circular spots on leaves of *Cestrum* sp. in Costa Rica.

Tuberculina jaffueli Speg. On leaves of *C. parqui* in Chile.

Uromyces cestri Mont. Yellow and powdery black rust pustules on leaves of *C. athroxanthum*, *C. aurantiacum*, *C. auriculatum*, *C. lanatum*, *C. laurifolium*, *C. macrophyllum*, *C. pallidum*, *C. parqui*, and *C. pseudoquina* in Chile, Peru, Bolivia, Brazil, Argentina, Uruguay, Porto Rico, Virgin Islands, Jamaica, Mexico, Costa Rica, and Guatemala.

Uromyces maculans Arth. Brown leaf rust on *C. nocturnum* in Costa Rica.

Uromyces venustus Diet. and Holw. Rust on leaves of *C. nitidum* in Mexico.

- CHAEROPHYLLUM.** Scented annual, biennial, or perennial herbs, often with tuberous roots.
- Protomyces macrosporus** Ung. See *Coriandrum*.
- Puccinia aromatica** Bub. and Lindr. Golden-yellow and deep-brown rust pustules on leaves of *C. aromaticum* in Russia, Austria, Hungary, and Germany.
- Puccinia chaerophylli** Purt. Cinnamon-brown to dark-brown rust pustules on leaves, petioles, and stems of *C. aureum*, *C. coloratum*, *C. hirsutum*, *C. temulum*, *Anthriscus sylvestris*, *A. nemerosa*, *A. tenerima*, and *Myrrhis odorata* in Europe and Siberia.
- Puccinia enormis** Fekl. Cinnamon-brown, powdery rust pustules on leaf blades, and petioles and on stems of *C. villarsium* in Switzerland and Austria.
- Puccinia pozzi** Sem. Leaf rust on *C. hirsutum* in Europe.
- Puccinia prescottii** Lindr. Dark-brown or black rust pustules on leaf blades and petioles of *C. prescottii* in Russia.
- Puccinia retifera** Lindr. Yellow and cinnamon-brown to dark-brown rust pustules on leaves of *C. bulbosum* in Russia, Hungary, and Germany.
- Septoria chaerophylli** Bres. Leaf spot on *C. aromaticum* in Germany.
- CHAETOCHLOA.** Annual grasses, mostly weeds. See also *Setaria*.
- Puccinia camellae** (Mayor) Arth. Rust on leaves of *C. setosa* in Porto Rico and Colombia.
- Sphaelotheca pamparum** (Speg.) Clint. Sori infecting the ovaries of the spikes, extending between the glumes as oblong bodies, 3 to 7 millimeters long, inclosed by a false membrane which ruptures irregularly at the apex, disclosing the powdery, olive-black spore masses. On *C. geniculata*, *C. imberbis*, and *Chaetochloa* sp. in Argentina, Cuba, and the Bahamas.
- CHALCAS.** ORANGE JASMINE. Small evergreen-leaved shrubs.
- Bacterium citri** Hassé. See Citrus.
- Corticium salmonicolor** B. and Br. See Citrus.
- CHAMAECYPARIS.** WHITE CEDAR. Trees or shrubs cultivated for their handsome evergreen foliage. Timber trees.
- Asterula chamaecyparidis** Shir. and Hara. On needles of *C. obtusa* in Japan.
- Coniothyrium pallido-fuscum** Sacc. On needles of *C. pisifera squarrosa* (*Retinospora*) and *Araucaria brasiliensis* in France.
- Gymnosporangium solenoides** (Diet.) Kern. (*G. miyabei* Yam. and Miy.). A rust with the aecial (cluster-cup) stage on leaves of *Sorbus alniifolia* and *S. aria*, and the telial stage on branches of *Chamaecyparis pisifera* (*C. plumosa*) in Japan. The telial stage is characterized by elongated reddish-brown swellings with rough surfaces, on which red-brown spore masses appear.
- Hediodasidium mompa** Tanaka. See Morus.
- Lophodermium chamaecyparidis** Shir. and Hara. On needles of *C. obtusa* in Japan.
- CHAMAEDOREA.** See Palmae.
- CHAMAEROPS.** See Palmae.
- CHEIRANTHUS.** WALLFLOWER. Flower-garden perennials.
- Ascochyta cheiranthi** Bres. Circular to oblong yellow or brown spots on leaves of *C. cheiri* in Alaska and Germany.
- Bacterium** sp. A bacterial disease of *C. annuus* is reported from Holland, characterized by a constriction of the upper part of the tap root of diseased plants, followed by yellowing and dropping of the leaves. The vascular bundles in the roots are stained brown or black.
- Cercospora cheiranthi** Sacc. White leaf spots on *C. cheiri* in Italy.
- Peronospora cheiranthi** Gäm. Downy mildew on leaves of *C. cheiri* in Europe.
- Phyllosticta cheiranthicola** Bub. and Zimm. Circular, greenish then yellowish spots on leaves of *C. cheiri* in Bohemia.
- Phyllosticta cheiranthorum** Desm. Scattered irregular white or gray-green leaf spots on *C. cheiri* in France.
- Septoria cheiranthi** Rob. and Desm. Circular or irregular leaf spots on *C. cheiri* in France.
- CHELIDONIUM.** CELANDINE POPPY. Garden herbs.
- Ascochyta chelidonii** Kab. and Bub. Irregular brown to black leaf spots on *C. major* in Bohemia.
- Didymaria chelidonii** Jacz. Indefinite brown leaf spots on *C. uniflorum* in Siberia.
- Melampsora magnusiana** G. Wagn. See *Populus*.
- Peronospora chelidonii** Miy. Downy mildew on leaves of *C. major* in Japan.
- Phyllosticta chelidonii** Bres. Irregular yellow to dull-brown leaf spots on *C. major* in Germany.
- Septoria chelidonii** Desm. Circular to angular greenish-olive or brown leaf spots with white centers on *C. major* and *Argemone mexicana* in Porto Rico, China, Japan, Siberia, and throughout Europe.
- CHENOPODIUM.** GOOSEFOOT. Quinoa. Weedy herbs.
- Ascochyta boni-henrici** Ranoj. On leaves of *C. bonus-henricus* in Yugoslavia.
- Ascochyta chenopodii** Rostr. See *Atriplex*.
- Ascochyta nebulosa** Sacc. and Berl. and f. *foliicola* Gz. Frag. On stems and leaves of *C. glaucum* in Siberia, Italy, and Spain.
- Cercospora chenopodii** Fres. On leaves of *Chenopodium* sp. in Denmark, Russia, and Japan.
- Cercospora chenopodiolica** Bres. Small subcircular pale-brown leaf spots on *C. polyspermum* in Russia and Germany.
- Cercospora macularis** (Schroet.) Magn. On leaves of *C. bonus-henricus* in Austria.
- Hypochnus peronosporoides** Speg. Thin white fungus layers on lower leaf surfaces of *C. ambrosioides* in Argentina.
- Oulariopsis erysipoides** Pat. and Har. On leaves of *C. ambrosioides* in Spain.
- Peronospora bohemia** Gäm. Downy mildew on leaves of *C. ficifolium* in Bohemia.
- Peronospora boni-henrici** Gäm. Downy mildew on leaves of *C. bonus-henricus* in Europe.
- Peronospora chenopodii-ficifolia** Saw. Circular to irregular pale-yellow spots, often covering entire leaf area of *C. ficifolium* in Japan.
- Peronospora chenopodii polyspermi** Gäm. Downy mildew on leaves of *C. polyspermum* in central and north Europe.
- Peronospora chenopodii rubri** Gäm. As above on *C. rubrum* in Holland, Yugoslavia, and Germany.
- Peronospora muralis** Gäm. Downy mildew on leaves of *C. muralis* in Europe and Argentina.
- Phragmidium ambrosioides** Thuem. Leaf rust on *C. album* in Yugoslavia.
- Phyllosticta dimorphospora** Speg. Circular pale-brown indefinite leaf spots on *C. hirtum* in Argentina.
- Ramularia chenopodii** Speg. On leaves of *C. album* in Argentina.
- Ramularia dubia** Reiss. On leaves of *C. album* and *C. patula* in Russia and Germany.
- Ramularia macularis** Schroet. Angular pale-yellow then white leaf spots on *C. bonus-henricus* in Switzerland, Poland, Austria, and Germany.
- Rhizoctonia napi** West. See Brassica.
- Septoria ambrosioides** Speg. On leaves of *C. ambrosioides* in Argentina.
- Septoria atriplicis** (West.) Fekl. See *Atriplex*.

CHENOPODIUM—Continued.

Septoria chenopodii West. Greenish-white than yellow leaf spots on *C. album*, *C. bonus-henricus*, *C. murale*, *C. rubrum*, and *C. viride* in Belgium, Portugal, Italy, and Germany.

Septoria undulipora Bub. On leaves of *C. murale* and *C. vulvaria* in Spain and Hungary.

Tolyposporium leptideum Syd. Cinnamon-brown smut sori in ovaries of *Chenopodium* sp. in Germany.

Urophlyctis pulposa (Wall.) Schroet. See *Atriplex*.

CHIOGLOTTIS. See Orchidaceae.**CHIONODOXA**. GLORY-OF-THE-SNOW. Spring bulbs.

Ustilago vaillantii Tul. See Hyacinthus.

CHIONANTHUS. FRINGE TREE. Woody plants grown for their profuse white flowers.

Aecidium chionanthi B. and Br. Rust on leaves of *Chionanthus* sp. in Ceylon.

Phyllosticta chionanthi Thuem. Irregular yellowish-white leaf spots with narrow red-brown margins on *C. virginica* in Portugal.

Septoria veleospora Sacc. On leaves of *C. virginica* in France.

CHLORIS. FINGER GRASS. Rhodes grass. Annual or perennial forage grasses. Some species grown as ornamentals.

Balanisia discoidea P. Henn. On culms of *C. distichophylla* and *Chloris* sp. in Peru and Brazil.

Heterosporium chloridis Speg. On leaves of *Chloris* sp. in Brazil.

Phyllachora chloridicola Speg. Black stromata on leaves of *C. distichophylla* and *C. radiata* in Argentina.

Puccinia chloridicola P. Henn. Dull yellow-brown to black elongate rust pustules on linear dull-brown leaf spots on *Chloris* sp. in the Congo.

Puccinia chloridina Bacc. Black linear rust pustules on leaves of *Chloris* sp. in Abyssinia.

Sorosporium chloridicola Beeli. Smut on *C. polydactyla* in the Congo Free State.

Tolyposporium chloridis P. Henn. Spikelets of *C. abyssinica* and *C. gayana* become black smutty masses in central and south Africa.

Ustilago deserticola Speg. Brown-black powdery smut sori in culms and rhizomes of *Chloris* sp. in Argentina.

Ustilago ulei P. Henn. Dark-brown or dark olivaceous spore masses covering and often distorting the leaves of *Chloris* sp. in Brazil.

CHLOROPHYTUM. Rhizomatous herbaceous plants.

Aecidium chlorophyti Har. and Pat. Brown rust pustules on large leaf spots on *Chlorophytum* sp. in African Congo.

CHORIZEMA. Small shrubs grown for their showy pea-like yellow, orange, and red flowers.

Phyllosticta chorizemae P. Henn. Brown confluent leaf spots on *Chorizema* sp. in Germany.

Septoria straussiana P. Henn. Dull-brown then paler leaf spots with red-brown margins on *Chorizema* sp. in Germany.

CHRYSANTHEMUM. Herbaceous and subshrubby plants cultivated in greenhouses and out of doors for their flowers.

Aphelenchus olesistus Ritz. Bos. See Begonia.

Aphelenchus ritzemii-bosi Schwartz. This nematode is parasitic on the leaves of *Chrysanthemum* (cult.), causing them to turn brown and fall prematurely. Blossom buds also shrivel up and the entire plant ultimately dies. The disease occurs in England and Germany.

Cercospora chrysanthemi Puttem. On leaves of *C. indicum* in Brazil. Heald and Wolf have used the same name for a species found in Texas.

Corticium chrysanthemi Plow. Said to cause a basal rot of stems of *Chrysanthemum* (cult.) in Great Britain.

Entyloma leucanthemi Syd. Small circular yellow-green leaf spots on *C. leucanthemum* in Switzerland and Austria.

Peronospora danica Gäum. Downy mildew on leaves of *C. segetum* in Denmark.

Peronospora leptosperma (De By.) Gäum. As above on *C. inodorum*, *Matricaria chamomilla*, and *M. discoidea* in Europe.

Peronospora radii De By. Downy mildew attacking leaves and deforming flower heads of *Achillea ptarmica*, *Anthemis arvensis*, *A. austriaca*, *Chrysanthemum leucanthemum*, *Matricaria chamomilla*, and *M. inodora* in Switzerland, France, Sweden, Denmark, Belgium, Italy, Finland, Bohemia, Austria, and Germany.

Phoma chrysanthemi Vogl. On stems and leaves of *Chrysanthemum* sp. in Italy, causing a wilt.

Phyllosticta leucanthemi Speg. Grayish-white circular leaf spots on *C. leucanthemum* in Italy.

Protomycopsis chrysanthemi Büren. Flat yellowish-white callosities on lower leaf surfaces of *C. alpinum* in Switzerland.

Protomycopsis leucanthemi Büren. On leaves of *C. leucanthemum* in Switzerland.

Puccinia acidil-leucanthemi Ed. Fisch. Brown to black rust pustules on leaves of *C. leucanthemum* in Europe. The asexual stage occurs on species of *Carex*.

Puccinia horiana P. Henn. Dark-brown to black rust pustules on lower leaf surfaces of *C. chinense*, *C. decussatum*, and *C. morifolium* (*C. sinense*) in Japan.

Puccinia leucanthemi Pass. Rust pustules on leaf blades and petioles of *C. leucanthemum* in Italy.

Puccinia proximella Syd. Brown rust pustules on leaves of *C. millefolium* in Russia.

Puccinia pyrethri Rabh. Pale brown to black rust pustules on leaves of *C. chinense*, *C. corymbosum*, *C. partheniifolium*, *C. parthenium*, and *C. morifolium* in Japan, Russia, Switzerland, Italy, Spain, Portugal, Rumania, Austria, Hungary, and Germany.

Septoria cercosporoides Trail. Irregular indefinite spots on leaves of *C. leucanthemum* in Scotland.

Septoria chrysanthemella Cav. Circular dull red-brown leaf spots of varying size and with yellow centers on *C. morifolium* and *Chrysanthemum* (cult.) in Denmark, Italy, and Japan. Also said to cause a damping-off of seedlings.

Septoria chrysanthemi-indiei Bub. and Kab. Circular, rarely irregular, dull-brown to dark-brown leaf spots on *C. indicum* in Bohemia.

Septoria chrysanthemi-rotundifolii Namys. Brown circular or irregular, often confluent, leaf spots on *C. rotundifolium* in Poland.

Septoria leucanthemi Sacc. and Speg. Yellow-brown circular or irregular leaf spots with white centers on *C. (Leucanthemum) maximum* in Italy and Portugal.

Septoria obesa Syd. Circular to irregular leaf spots on *C. arcticum* in Japan.

Septoria rostrupii Sacc. and Syd. Circular dark-brown leaf spots on *C. indicum* in Russia and Denmark.

Septoria socia Pass. and f. *catalaunica* Gz. Frag. Subcircular to irregular dull-brown leaf spots on *C. leucanthemum* in Italy, Spain, and Portugal.

Uredo autumnalis Diet. Yellow-brown rust pustules on leaves of *C. boreale*, *C. indicum*, and *C. sinense* in Japan.

- CHRYSOBALANUS.** COCOA PLUM. *ICACO*. Shrubs or trees with edible fruit. Sometimes planted for ornament.
- Helminthosporium chrysobalani** P. Henn. Large indefinite leaf spots on *Chrysobalanus* sp. in the Congo.
- Mycosphaerella chrysobalani** Miles. Irregular spots, generally confluent and occupying most of the leaf surface on *C. icaco* in Porto Rico.
- CHRYSOPHYLLUM.** STAR APPLE. *CAINITO*. Handsome broad-leaved evergreen trees grown for fruit and ornamental effect.
- Helminthosporium chrysophylli** P. Henn. Dark-brown circular leaf spots on *Chrysophyllum* sp. in East Africa.
- Phyllosticta chrysophylli** Syd. Indefinite gray leaf spots on *C. imperialis* in Germany.
- Septoria evansii** Syd. Small circular white leaf spots with dull-purple margins on *C. magalis-montana* in the Union of South Africa.
- Uredo chrysophyllicola** P. Henn. Brown rust pustules on dark-brown leaf spots on *Chrysophyllum* sp. in Brazil.
- CHRYSOSPLENIUM.** GOLDEN SAXIFRAGE. Semiaquatic plants.
- Entyloma chrysosplenii** (B. and Br.) Schroet. Smut sori in small circular white or yellow leaf spots on *C. alternifolium* in Great Britain, Denmark, and Germany.
- Peronospora chrysosplenii** Fckl. Downy mildew on leaves of *C. alternifolium* in central and northern Europe.
- Puccinia chrysosplenii** Grev. Brown rust pustules on leaves of *C. alternifolium*, *C. oppositifolium*, *C. trachyspermum*, and *Mitella longicaapa* in Japan, eastern Asia, Russia, Finland, Belgium, Great Britain, Austria, Hungary, and Germany.
- Septoria posoniensis** Bauml. Circular or irregular greenish-gray or ashen leaf spots with narrow brown margins on *C. alternifolium* and *C. oppositifolium* in Great Britain and Hungary.
- CHUSQUEA.** See *Bambuseae*.
- CICER.** CHICK-PEA. Garbanzo. Gram. Herbaceous legumes.
- Erysiphe taurica** Lévl. See *Althaea*.
- Phyllosticta rabiei** (Pass.) Trott. (*Phyllosticta cicerina* Prill. and Delacr.) Dull-yellow to gray leaf spots on *C. arietinum* in France, Italy, and Spain. The stems, petioles, and rachides are also attacked, diseased plants often breaking over at the lesions and dying.
- Rhizoctonia napi** West. See *Brassica*.
- Uromyces ciceris-arietini** (Grogg.) Jacz. and Boy. Small round or oval cinnamon-brown powdery rust pustules, which tend to coalesce, on both leaf surfaces of *C. arietinum* in India, East Africa, Yugoslavia, Spain, Portugal, France, and Italy.
- CICORIUM.** CHICORY. Endive. Herbs cultivated for their fleshy roots and as salad plants.
- Puccinia cichorii** (DC.) Bell. Cinnamon-brown to dark-brown rust pustules on leaves and stems of *C. endivia* and *C. intybus* in India, Argentina, and Europe. Reported from Oregon.
- Puccinia endiviae** Pass. Cinnamon-brown to dark-brown powdery rust pustules on leaves and stems of *C. endivia* in Argentina, Italy, Spain, France, Holland, and Denmark. Reported from Connecticut.
- Ramularia lam psanae** (Desm.) Sacc. On leaves of *C. intybus*, *Lampasana communis*, and *Taraxacum officinale* in Europe.
- Septoria endiviae** Thuem. On leaves of *C. endivia* in Austria and Holland.
- Septoria intybi** Pass. On branches of *C. intybus* in Italy.
- CIMICIFUGA.** BUGBANE. Tall ornamental herbaceous perennials.
- Aecidium shiraianum** Syd. Rust on leaves of *C. biternata*, *C. foetida*, *C. intermedia*, and *C. japonica* in Japan and Siberia.
- Coleosporium cimicifugatum** Thuem. Yellow or orange rust pustules on leaves of *C. foetida*, *C. intermedia*, and *C. simplex* in Siberia and Japan.
- CINCHONA.** Tropical trees cultivated for their medicinal bark, which is sometimes called Peruvian bark and from which quinine is made.
- Colletotrichum cinchonae** Koord. Anthracnose on leaves of *Cinchona* sp. in Java.
- Corticium salmonicolor** B. and Br. See *Citrus*.
- Dasycephala warburgiana** P. Henn. Causes a serious canker disease of limbs of *C. ledgeriana* in Java.
- Guignardia yersinii** Vinc. On *C. ledgeriana* and *C. succirubra* in Indo-China.
- Moniliopsis aderholdii** Ruhl. See *Solanum*.
- Phyctaena cinchonae** Vinc. Causes roughened, cankered areas on bark of branches of *Cinchona* spp. in Indo-China.
- Phyllosticta cinchonae** Koord. Irregular yellowish spots on leaves of *Cinchona* sp. in Java. The name of this species is untenable, the following having priority.
- Phyllosticta cinchonae** Pat. Angular dull-yellow leaf spots on *Cinchona* sp. in Ecuador.
- Phyllosticta cinchonaeacola** Vinc. Circular deep-brown patches with violet borders on leaves of *C. ledgeriana* and *C. succirubra* in Indo-China. On old leaves a shot hole effect is produced; on young leaves the numerous spots cause deforming. Developing shoots are also involved, stunting the plants.
- Phyllosticta honbaensis** Vinc. On leaves of *C. ledgeriana* and *C. succirubra* in Indo-China.
- Phyllosticta yersinii** Vinc. Attacks young leaves and shoots of *Cinchona* spp. in Indo-China.
- Physalospora cinchonae** Vinc. On *C. ledgeriana* and *C. succirubra* in Indo-China.
- Tulasnella cinchonae** Racib. Thin pale-rose fungus layers on lower leaf surfaces of *C. ledgeriana*, *C. officinalis*, and *C. succirubra* in Java.
- Uredo cinchonae** P. Henn. Brown rust pustules on leaves of *Cinchona* sp. in Java.
- CINEBARIA.** Herbs cultivated for their flowers. See also *Senecio*.
- Aecidium cinerariae** Rostr. Rust on leaves and stems of *C. cruenta* (*Senecio cruentus*) and *C. palustris* in Siberia, Denmark, Austria, and Germany. Said to be a stage of *Puccinia eriophori* Thuem. which occurs on *Eriophorum* spp.
- Asochyta cinerariae** Fl. Tass. On leaves of *C. maritima* in Italy.
- Asochyta fibricola** Sacc. On stems of *C. cruenta* and *C. maritima* in France.
- Coleosporium senecionis** Fr. See *Senecio*.
- Entyloma calendulae** (Oud.) De By. See *Calendula*.
- Phytophthora cryptogea** Pethybr. and Laff. See *Lycopersium*.
- Puccinia cinerariae** McAlp. Orange-yellow and chestnut-brown to black rust pustules on leaves of *Cineraria* (cult.) in Australia.
- Ramularia senecionis** (B. and Br.) Sacc. See *Senecio*.
- CINNAMOMUM.** CINNAMON. CAMPHOR TREE. CASSIA BARK TREE. Evergreen trees and shrubs with aromatic leaves and wood, yielding aromatic and medicinal products.
- Aecidium cinnamomi** Rac. Rust on leaves, petioles, and stems of *C. iners* and *C. zeylanicum* in Java.
- Asochyta camphorae** Turc. Whitish spots with purple borders on leaves of *C. glandulifera* in Italy.
- Atichia millardetii** Racib. Black, superficial, radiate fruiting bodies on leaves of *Anacardium occidentale*, *Cinnamomum zeylanicum*, *Coffea arabica*, *Styrax benzoin*, and *Syzygium jambolanum* in Java and the Malay Peninsula.

CINNAMOMUM—Continued

- Colletotrichum briosii** Turc. Anthracnose on leaves *C. burmanni* in Italy.
- Corticium salmonicolor** B. and Br. See *Citrus*.
- Cytospora cinnamomi** Turc. Circular to irregular white leaf spots with dull-brown margins on *C. burmanni* in Italy.
- Desmotaseus cinnamomi** Petch. Attacks the trunks and branches of *C. zeylanicum* in Ceylon.
- Diplodia cacaicola** P. Henn. See *Theobroma*.
- Exobasidium cinnamomi** Petch. Rough excrescences up to 8 centimeters long and of pale-brown color on branches of *C. cassia* and *C. zeylanicum* in Ceylon.
- Fomes carophylli** (Rac.) Bres. Causes a root rot and wood rot of *C. zeylanicum*, *Citrus* spp., *Dalbergia latifolia*, *Mesua ferrea*, and other economic plants in Java, Ceylon, Mauritius, Japan, and Central Africa.
- Fomes lamaensis** Murr. See *Hevea*.
- Fomes lignosus** Klotzsch. See *Hevea*.
- Gleosporium camphorae** Sacc. Anthracnose on leaves of *C. camphora* (*C. officinarum*) in Italy. Reported from Texas and Alabama.
- Glomerella cinnamomi** Yosh. (*Guignardia cinnamomi* [Yosh.] Hara.) Anthracnose on leaf blades and petioles, buds, and young shoots of nursery stock of *C. camphora* in Japan and Formosa. The diseased spots are circular, elliptical or fusiform, reddish-brown at first and finally fading out to a light brown. The stems are girdled by a brown rot, the plants dying.
- Hypochnus cucumeris** Frank. See *Cucumis*.
- Hypochnus sasakii** Shirai. Leaf blight of *C. camphora* in Japan.
- Leptosphaeria cinnamomi** Shir. and Hara. Branch canker and leaf spot of *C. camphora* in Japan.
- Macrophoma cinnamomi-glanduliferi** Luigi. Large irregular blotches on leaves of *C. glanduliferum* in Italy.
- Phyllosticta aromatophila** Sacc. Small gray leaf spots on *C. camphora* (*C. officinarum*) in Italy.
- Phyllosticta cinnamomi** Delacr. Irregular dull-yellow leaf spots on *C. zeylanicum* in India.
- Phyllosticta cinnamomi-glanduliferi** P. Henn. Circular, finally confluent, dull-brown leaf spots on *C. camphora* and *C. glanduliferum* in Russia and Germany.
- Phytophthora cinnamomi** Rands. Irregular vertical stripes of dead bark (1 to 5 centimeters wide) appear on the trees, extending upward from the ground for a distance of 1 to 10 meters. At the upper end of each canker drops of wine or amber colored exudate appear. The host is *C. burmanni* in Sumatra.
- Rosellinia arcuata** Petch. See *Thea*.
- Rosellinia bunodes** B. and Br. See *Citrus*.
- Scolecotrichum cinnamomi** Rac. Circular gray leaf spots from 1 to 4 centimeters in diameter on *C. zeylanicum* in Java.
- CISSAMPELOS**. Twining herbs or shrubs.
- Cercospora pareirae** Speg. Circular dull-brown leaf spots on *C. pareira* in Argentina.
- Phyllosticta cissampeli** Speg. Small brown leaf spots with white centers on *C. pareira* in Argentina.
- Uromyces cissampelidis** Diet. Powdery cinnamon-brown to dark-brown rust pustules on leaves of *Cissampelos* sp. in Colombia.
- CISSUS**. Tropical woody climbers.
- Aecidium cissigenum** Welw. Leaf rust on *C. adenanchus* in Abyssinia.
- Cercospora riachuelii** Speg. On leaves of *C. palmata* in Argentina.
- Cornartium wilsonianum** Arthur. Straw-colored to dark-brown powdery rust pustules on leaves of *C. rhombifolia* in Cuba, Costa Rica, and Ecuador.
- Endophyllum guttatum** (Kunz.) Syd. (*Endophyllum circumscriptum* [Schw.] W. and O.) Yellow rust pustules on large swollen areas on leaves of *C. quadrangularis*, *C. rhombifolia*, and *C. sicyoides* in Italian north Africa, Central and South America, Trinidad, and the West Indies.
- Entyloma cissigenum** P. Henn. Smut sori in small, circular, black, often confluent, leaf spots on *Cissus* sp. in tropical Africa.
- Phyllosticta cissicola** Speg. Small, circular, whitish leaf spots with brown borders on *C. sicyoides* in Argentina and Porto Rico.
- Schizonella colemani** Iyen and Nar. This smut causes abnormal growths resembling witches' brooms at the nodes of *C. quadrangularis* in India. The diseased branches are pale-green in color and contain dark-green elongate swellings within which the spores form as black powdery masses.
- Schroeteria arabica** P. Henn. Smut on *C. quadrangularis* in Arabia and India. Probably the same as the following species.
- Schroeteria cissi** (DC.) De T. (*Mykosyrinx cissi* [DC.] G. Beck.) Smut sori in the peduncles and pedicels of inflorescences distorting the infected parts somewhat and filling them with a dusty-purple spore mass. On *C. acida*, *C. erosa*, *C. quadrangularis*, and *C. sicyoides* in Uganda, Cameroon, Mexico, South America, and the West Indies. Also reported from Florida.
- Uromyces cladomanes** Traverso. Rust causing coralloidal formations on *Cissus* sp. in Somalia.
- CISTUS**. ROCKROSE. Low shrubs grown for their red or white flowers.
- Ovariopsis cisti** Jaap. On leaves of *C. albidus*, *C. mouspeltensis*, and *C. salvifolius* in Dalmatia.
- Phyllosticta cistina** Thuem. Small circular purplish black, finally ashen-colored leaf spots with purple borders on *C. crispus*, *C. laurifolius*, and *C. salvifolius* in Greece, France, and Portugal.
- Uredo helianthemii** Rbh. See *Helianthemum*.
- CITHAREXYLUM**. Shrubs or trees cultivated for their ornamental effect.
- Phyllachora citharexyli** (Rehm.) v. Hoeh. Black stromata on leaves of *C. suberosum* and *Citharexylum* sp. in Brazil.
- CITROPSIS**. AFRICAN CHERRY ORANGE. Trees allied to *Citrus*.
- Bacterium citri** Hasse. See *Citrus*.
- CITRULLUS**. WATERMELON. Colocynth. Annual or perennial vines.
- Hypochnus cucumeris** Frank. See *Cucumis*.
- Puccinia citrulli** Syd. and Butl. Powdery red-brown rust pustules on leaves of *C. colocynthis* in India.
- Scolecotrichum melophthorum** Prill. and Delacr. See *Cucumis*.
- Septoria citrullicola** Poteb. Small circular white leaf spots on *C. vulgaris* in Russia.
- CITRUS**. ORANGE. LEMON. GRAPEFRUIT. CITRON. LIME. MANDARIN ORANGE. TANGERINE. Tropical and subtropical fruit trees.
- Ascochyta citri** Penz. Nearly circular dark-brown, finally grayish, leaf spots with definite margins on *C. limonia* and *C. sinensis* in Italy and Brazil.
- Ascochyta corticola** McAlp. This "bark-blotch" disease attacks the lemon (*C. limonia*) and orange (*C. sinensis*) in Australia and New Zealand. The disease begins as dark-brown spots on the lower portion of the trunk and spreads rapidly until the trunk is girdled, death of the tree resulting.
- Ascochyta destruens** (McAlp.) F. Tass. Minute black pustules on yellowish-brown scab-like patches with red-brown margins on leaves of *Citrus sinensis* in Australia. Diseased leaves wither and the scabby areas fall out.

CITRUS—Continued.

Bacillus citrimaculans Doidge. Bacterial spot. This bacterial disease produces circular to irregular slightly sunken spots on the fruit which are yellowish at first but soon become brown, varying in shade, and often with red margins. The skin only may be affected, or the disease may penetrate into the pulp with a peculiar accompanying odor. Under moist conditions yellow viscid drops appear. Branches are attacked, infection taking place at points of attachment of the leaves, water-soaked areas of 6 to 10 millimeters in diameter appearing which increase in size. Leaves and twigs within these areas wither and die. On the larger branches irregular brown areas occur. Practically all *Citrus* spp. are attacked, including orange, lemon, citron, mandarin, tangerine, lime, and grapefruit in the Union of South Africa.

Bacterium citri Hassé. Citrus canker attacks all parts of the tree above ground. On the leaves the disease first appears as minute yellow-brown translucent spots, which soon become visible on both sides. These spots increase rapidly in size, and the tissues at the center soon rupture, giving the characteristic corky appearance, white or pinkish at first, later brown. The corky raised areas are surrounded by a yellowish-brown, oily, translucent area, and are inclosed by yellowish-green zones which shade into the normal green. The spots vary in size from a quarter to half an inch at maturity and are circular in shape, occurring singly or in groups. Similar corky areas occur on the twigs, thorns, branches, and fruit, tending to be more irregular and more prominent, and with a tendency to develop cracks or fissures. The fruit is attacked only while green, and may ripen prematurely and fall. While the canker spots do not penetrate deeply, the cracking of the spots opens a way for decay organisms.

This disease occurs in China, Japan, the Philippines, Hawaii, Mauritius, Ceylon, Siam, Dutch East Indies, Borneo, and Formosa. It has been introduced into the Union of South Africa, northern territory of Australia, and the Gulf States of the United States, where eradication campaigns are in progress. Practically all species and varieties are susceptible, although to a greatly varying degree, the grapefruit (*C. grandis*), *Citrus* (*Poncirus*) *trifoliata*, lime (*C. aurantifolia*) and round orange (*C. sinensis*), being especially so, and the Satsuma, King orange and kumquat rather resistant. In addition to the large number of natural hosts of the genus *Citrus* and its segregates, many *Citrus* relatives have been found susceptible to a greater or less extent through artificial inoculations. Among these are *Aegle marmelos*, *Alatalania citroides*, *A. ceylonica*, *A. disticha*, *Casimiroa edulis*, *Chaetosperrum glutinosum*, *Chalcas exotica*, *Citropsis schweinfurthii*, *Claucaena lamsium*, *Eremocitrus glauca*, *Erodia ridleyi*, *E. latifolia*, *Feronia limonia*, *F. lucida*, *Hesperethusa crenulata*, *Melicope triphylla*, *Microcitrus australasica*, *M. garroayi*, *M. australis*, *Paramignya longipedunculata*, *Poncirus trifoliata*, and *Toddalia asiatica*.

Bacterium gummi Comes. This bacterium is reported from Italy, Paraguay, and Brazil as the cause of a gumming disease of twigs, branches, and fruit of orange (*C. sinensis*) and lemon (*C. limonia*). In Brazil the type of disease known as mal-di-gomma or foot rot is also attributed to this organism. The disease is apparently the same as the foot rot and gummosis of Florida, now attributed to other causes.

Bark rot. This disease attacks the calomondin (*Citrus mitis*), grapefruit (*C. grandis*), mandarin (*C. nobilis* var.), and orange (*C. sinensis*) in the Philippine Islands and China, its cause being as yet unknown. It is marked by the oozing out of sap from the bark on the trunk and branches, a putrid sore from 0.5 to 3.5 centimeters in diameter forming. These sores run together and finally girdle the infected tree. The attempts of the tree to heal the spots causes a rough, scaly appearance. Gum pockets form between the wood and cambium. Any fruit present matures prematurely and falls.

Botrytis citricola Briozzi. Oranges (*C. sinensis*) and lemons (*C. limonia*) are rotted and mummified by this fungus in Italy and Brazil.

Cercospora sp. Circular dark-brown leaf spots on *C. sinensis* in Japan.

Cercospora fumosa Penz. Gray irregular areas on the lower leaf surfaces of *Citrus* spp. in Algeria, Argentina, and Italy.

Cercospora penzigii Sacc. On leaves of *C. limonia* in Italy.

Cercospora undulata (Bern.) Sacc. See *Aegle*.

Cladosporium brunneo-atrum McAlp. Minute brown, generally circular, raised spots on fruit, leaves, and young shoots of *C. sinensis* characterize the false melanose of Australia. On the fruit these spots usually run together into long, irregular curves and become black.

Cladosporium elegans Penz. Dry, bleached, subcircular to oval spots on orange (*C. sinensis*) leaves in Italy and Australia. This species has been confused in the literature with *C. citri* Mass.

Cladosporium furfuraceum McAlp. The lemon-scurf disease, forming dingy-white patches on the surface of lemon (*C. limonia*) fruits in Australia. The discolored areas peel off in small flakes.

Cladosporium subfusoidium McAlp. Forms gray-brown scabs on lemons (*C. limonia*), which crack and become covered with darker fruiting patches. Occurs in Australia and the Philippines.

Coniothecium citri McAlp. Forms numerous small brown patches with raised brown or black margins on the upper sides of leaves of the orange (*C. sinensis*) in Australia.

Coniothecium scabrum McAlp. Causes "black scurf" of citrus (*Citrus* spp.) fruits in Australia and the Union of South Africa. Small black depressions, surrounded by yellowish-green areas appear, which enlarge to form large, sooty-black patches. The black masses become detached in flakes, leaving dirty-gray scurfy depressed areas.

Coniothorium cervinum McAlp. Irregular gray to fawn-colored leaf spots on *C. limonia* in New South Wales.

Coniothorium citricolum McAlp. Grayish-white leaf spots with red-brown margins on *C. sinensis* in New South Wales.

Corticium koleroga (Cke.) v. Hoeb. See *Coffea*.

Corticium salmonicolor B. and Br. This fungus produces the well-known and widespread "pink disease" of the Tropics, which attacks practically all woody economic plants. It occurs on the trunk and branches, first being noted as dirty-white to pinkish pustules, which push through the dead bark. Following this stage, a layer of mycelium spreads out over the bark, turning to a characteristic pink color, constituting the fruiting areas of the fungus. This layer which may extend for several feet, and even girdle the trunk and one or more limbs, is at first smooth and velvety, but with age becomes roughened and breaks into patches, which fade out to a dirty-white. The bark beneath is at first watery and somewhat darker than normal, soon, however, becoming dried out, and is finally reduced to a dry mass of fiber. The outermost wood layers are also penetrated and killed. The disease has been reported from the following countries: West Indies, including Porto Rico, Trinidad, British Guiana, Surinam, Brazil, Caucasus, Ceylon, India, Java, Guam, Philippines, Borneo, Malaya, Sumatra, Burma, Mauritius, Fiji, Formosa, west Africa, Portuguese India, and the Cameroons. Among the known hosts are *Amherstia nobilis*, *Annona muricata*, *A. squamosa*, *Artocarpus integrifolia*, *Averrhoa* sp., *Bauhinia* sp., *Cajanus indicus*, *Castilla elastica*, *Casuarina equisetifolia*, *Chalcas exotica*, *Cinchona ledgeriana*, *C. succirubra*, *Cinnamomum camphora*, *C. zeylanicum*, *Citrus aurantifolia*, *C. grandis*, *C. hystrix*, *C. limonia*, *C. medica*, *C. micrantha*, *C. nobilis*, *C. sinensis*, *C. webberi*, *Clitoria cajanifolia*, *Coffea arabica*, *C. liberica*, *C. robusta*, *Corchorus* sp., *Duranta*

CITRUS—Continued.

Corticium salmonicolor—Continued.

plumieri, *Durio zibethinus*, *Eriobotrya japonica*, *Erythroxylon coca*, *Eucalyptus* sp., *Feronia elephantum*, *Feroniella lucida*, *Ficus* spp., *Gardenia* sp., *Gliricidia maculata*, *Hevea brasiliensis*, *Hibiscus* sp., *Indigofera arrecta*, *Lansium domesticum*, *Mangifera indica*, *Myristica fragrans*, *Photinia japonica*, *Piper* sp., *Pithecolobium dulce*, *Prunus persica*, *Randia macrantha*, *Shorea* sp., *Strobilanthes* sp., *Thea* sp., and *Theobroma cacao*.

Cytospora citri May. On *Citrus* sp. in Italy.

Cytosporina citriperda Camp. Circular depressed brownish-red, then black, patches on fruit of *C. sinensis* in Italy.

Dendrophoma valispora Penz. On lemon (*C. limonia*) leaves in Italy.

Diaporthe citricola Rehm. Found in connection with *Citrus* die-back in the Philippines.

Didymella citri Noack. Produces a canker on branches and twigs of *C. sinensis* (orange) in Brazil.

Diplodia cacaicola P. Henn. See *Theobroma*.

Diplodia citricola McAlp. On fruit, leaves, twigs, and branches of *Citrus* spp. in Australia.

Diplodia destruens McAlp. Said to cause scabby, dirty-gray patches with red-brown margins on leaves of *C. sinensis* in Australia.

Dry rot. See *Nematospora* sp.

Eutypella citricola Speg. On branches of *C. sinensis* in Brazil and Argentina.

Exobasidium citri Jacc. On young fruit of *C. nobilis* in the Caucasus.

Fusarium sp. A blossom end-rot of orange (*C. sinensis*) is produced by this fungus in Porto Rico, marked by a brown staining of the blossom ends and an internal pink rot. Affected fruits turn a deep abnormal yellow and fall prematurely.

Gloeosporium aurantium West. Anthracnose on *Citrus* spp. in Belgium and Argentina.

Gloeosporium citri Mass. This species, reported on leaves of *C. sinensis* from Trinidad and Australia, does not seem to be other than *Colletotrichum gloeosporioides* Penz.

Gloeosporium citricolum Hori. Irregular indefinite brown patches on leaves, causing defoliation. Young twigs are also attacked and caused to die-back. Brown anthracnose spots appear on the fruit. Attacks *C. nobilis* var. and *C. sinensis* in Japan. *G. citricolum* Massee is referred to *Colletotrichum gloeosporioides* Penz.

Gloeosporium foliicola Nish. Rapidly spreading brown areas around the blossom end of fruit of Satsuma and Mandarin (*C. nobilis* var.) in Japan. These spots become grayish and sunken at the center with dark-brown surrounding areas.

Gloeosporium hesperidearum Catt. Anthracnose on leaves of *Citrus* spp. in Italy and Argentina. Referred by Grove to *G. aurantium* West.

Gloeosporium intermedium Sacc. and var. **subramulosum** Sacc. Anthracnose on *Citrus* spp., *Corynocarpus laevis*, and *Hoya carnea* in Italy, France, and Australia. Also referred to *G. aurantium* West.

Gloeosporium spagazzini Sacc. Anthracnose on *C. limonia* and *C. sinensis* in Brazil and Argentina. Like most of the other *Gloeosporium* of *Citrus* not definitely distinct from *Colletotrichum gloeosporioides*.

Hainesia aurantii P. Henn. On leaves of *C. sinensis* in Costa Rica.

Hendersonia citri McAlp. On leaves of *C. aurantium* in Australia.

Hendersonia socia McAlp. On leaves and bark of *C. limonia* in Australia.

Laestadia socia Penz. On leaves of *C. limonia* in Italy.

Lasiodiplodia citri Avena. Produces a fruit rot, die-back of twigs, and yellow to brown leaf spots of *C. sinensis* and *C. limonia* in Brazil. Said to be one of the worst of Brazilian *Citrus* diseases. The fungus is apparently not distinct from *D. cacaicola* P. Henn.

Leaf drop. This disease, due to an unknown cause, defoliates Satsumas (*C. nobilis* var. *unshiu*) in Japan, and is considered serious.

Leptosphaeria citricola Penz. On *Citrus* sp. in Italy.

Leptosphaeria coniothyrium Sacc. On twigs of *C. limonia* and *C. sinensis* in Brazil.

Leptosphaeria diana Sacc. and Speg. On leaves of *C. aurantium* in Italy.

Macrophoma aurantii Scallia. On leaves of *C. sinensis* in Italy.

Macrophoma mantegazziana (Penz.) Berl. and Vogl. On leaves of *Citrus* spp. in Italy and Ceylon.

Meliola butleri Syd. Circular black superficial fungus patches on leaves of *C. medica* and *C. grandis* in India.

Meliola citricola Syd. Superficial black fungus patches on leaves of *C. grandis*, *C. nobilis*, and *C. sinensis* in China.

Microthyrium citri Penz. On leaves of *C. limonia* in Italy.

Mycosphaerella citricola McAlp. On leaves of *C. limonia* in Ceylon and Australia.

Mycosphaerella gibelliana Pass. Subcircular, brown leaf spots on *Citrus* spp. in Italy, Portugal, Spain, and Austria.

Mycosphaerella horii K. Hara. Circular to irregular definite reddish-brown to gray leaf spots with raised blackish-brown margins on *C. sinensis* and *C. nobilis* in Japan. The "small round brown spot disease."

Mycosphaerella inflata Penz. On living branches of *C. aurantium* in Italy.

Mycosphaerella loefgreni Noack. Small, circular to irregular, brown to dirty-white spots on leaves, fruit, and twigs of *C. sinensis* in Brazil. The spots on the twigs are raised, but sunken on leaves and fruit. The disease may cause premature fall of leaves and fruit.

Nematospora sp. This fungus is reported as the cause of the "dry-rot" disease of fruit of *C. sinensis*, *C. aurantium*, and *C. nobilis* in the Philippine Islands, China, Japan, Formosa, and possibly Barbados. There are no external symptoms. Internally infected fruits are more or less dry and free of juice, the sections becoming atrophied and falling apart. The pulp at first has an unpleasant sour taste, but soon becomes dry and tasteless.

Nothopateella lecanidium (Speg.) Sacc. On trunks and branches of *C. sinensis* in Brazil.

Oidium sp. Forms a white powdery layer on leaves and twigs, causing yellowing and premature fall of leaves and die-back of twigs of *Citrus* spp. in India and Ceylon. In the latter country said to be so serious as to prevent the growing of *Citrus* on any scale. *O. tingitaninum* Carter, described from California, is probably the same species.

Oospora aurantii Petch. Causes a rotting of fruit of *C. sinensis* in Ceylon.

Ovularia aurantii McAlp. Eruptent, effused, grayish-green patches on fruit of *C. sinensis* in Australia and Ceylon.

Ovularia citri Br. and Farn. Attacks orange fruits (*C. sinensis*) in Italy causing a disease known as "white rust."

Phoma aurantium (Rabenh.) Sacc. On branches of *Citrus* sp. in middle Europe.

Phoma citricarpa McAlp. The black-spot disease causes a spotting of fruit of *C. limonia*, *C. maxima*, *C. mitis*, *C. nobilis*, and *C. sinensis* in China, Australia, and New Zealand. The spots are small, irregular, from 1 to 9 millimeters in diameter, reddish-brown at first, later much darker and sunken.

CITRUS—Continued.

Phoma flaccida McAlp. Brown, apical leaf spots on *C. sinensis* in Australia.

Phoma omnivora McAlp. This fungus forms irregular gray to black scabby patches on the fruit and leaves. Leaves and twigs die-back, gray blotches appearing on the latter. Dark-brown cankers occur on the roots. The hosts are *C. limonia*, *C. medica*, and *C. sinensis* in Australia and Ceylon.

Phoma stigmaea Dur. and Mont. On leaves of *C. aurantium* in Algeria.

Phomopsis caribaea Horne. Causes a stem-end rot of *C. grandis* in the Isle of Pines. Probably not distinct from *P. citri* Fawc.

Phyllosticta arethusa Bub. On leaves of *C. aurantium* in Austria.

Phyllosticta beltranii Penz. On leaves of *C. limonia* in Italy.

Phyllosticta circumsepta Sacc. On fruit of *C. nobilis* in the Philippines.

Phyllosticta citricola Hori. Circular to irregular brown leaf spots on *C. nobilis* and *C. sinensis* in Japan.

Phyllosticta delicosa Pass. Brown leaf spots with raised margins on *Citrus* sp. in Italy.

Phyllosticta disciformis Penz. and var. **brasiliensis** Speg. Circular to irregular ashen-white leaf spots with dull-brown margins on *Citrus* spp. and *Atalantia buxifolia* in Russia, Ceylon, Italy, and Brazil.

Phyllosticta fuliginosa Massal. Irregular olivaceous to dull-brown leaf spots on *Citrus* sp. in Italy.

Phyllosticta hesperidearum Penz. Small, irregular, brown, then yellow, and finally ashen leaf spots, and dull-brown corky areas on twigs of *C. limonia* and *C. sinensis* in Brazil, Algeria, and Australia.

Phyllosticta lenticularis Pass. Leaf spot on *Citrus* sp. in Portugal.

Phyllosticta longispora McAlp. Brown to gray apical and marginal leaf spots on *C. sinensis* in Australia.

Phyllosticta marginalis Penz. Irregular white spots with brown margins on leaves of *C. medica* in Italy.

Phyllosticta micrococcoides Penz. Dark-brown, irregular spots on young leaves of *Citrus* sp. in Italy.

Phyllosticta scabiosa McAlp. On leaves of *C. limonia* in Australia.

Phytophthora sp. Causes a rot of *Citrus* fruits in Porto Rico.

Phytophthora citri Ven. Causes a soft rot of fruit of *C. media* in India.

Pleospora batumensis Naou. On petioles of *C. sinensis* in south Russia.

Pleospora citrorum Sacc. On *Citrus* sp. in Italy and Argentina.

Pleospora disrupta McAlp. Disorganizes the tissues of orange (*C. sinensis*) leaves in Australia.

Pleospora hesperidearum Catt. Causing a scabbing of fruit of *Citrus* sp. in Italy and Uruguay.

Polyporus caryophylli Racib. See *Cinnamomum*.

Pyrenochaeta destructiva McAlp. Circular to oval gray leaf spots with brown margins on *C. aurantium* and *C. grandis* in Australia.

Ramularia citri Penz. On leaves of *C. sinensis* in Italy.

Ramularia scabiosa McAlp. and Tryon. Circular definite light-brown, then nearly black, raised spots on leaf blades and petioles of *C. limonia* and *C. sinensis* in Australia.

Ramularia undulata Bern. On leaves of *Citrus* sp. and *Aglaia odorata* in Java.

Rhabdospora flexuosa (Penz.) Sacc. On bark of *C. sinensis* in Italy and Brazil.

Rhynchosdiploia citri Br. and Farn. This fungus, reported as the cause of a serious lemon (*C. limonia*) disease in Italy and Brazil, is probably the same again as *D. cacaicola* P. Henn.

Rosellinia sp. An undetermined species of *Rosellinia* causes a root rot of *Citrus* spp., *Erythrina* sp., *Theobroma cacao*, and possibly other economic plants in the West Indies. Diseased trees die-back slowly, suckering freely, and finally dying. The rot in the roots and crowns resembles that produced by *R. pepo*. A *Rosellinia* sp. is reported as the cause of a root disease of *C. sinensis* in Spain and the Balearic Islands.

Rosellinia bunodes B. and Br. The "black root disease" attacks and destroys a wide range of tropical plants, including *Acalypha* sp., *Artocarpus incisa*, *Cajanus indicus*, *Castilla elastica*, *Cinnamomum camphora*, *Citrus* spp., *Coffea arabica*, *Dryobalanops aromatica*, *Eugenia jambos*, *Ficus dubia*, *Ilexa brasiliensis*, *Hibiscus rosa-sinensis*, *Mangifera indica*, *Maranta arundinacea*, *Miconia* sp., *Panax plumatum*, *Persea gratissima*, *Peltiera alliacea*, *Piper nigrum*, *Sterculia caribaea*, and *Theobroma cacao* in the West Indies, Trinidad, India, Malaya, Java, and Ceylon. The disease is apparently most prevalent and most serious in the West Indies, the *Rosellinia* root disease of tea and other woody plants in the oriental tropics being attributed for the most part to other species.

The fungus attacks the roots and crowns of the hosts, working through the bark to the wood. Black fungus strands or rhizomorphs appear on the surface and may grow together to form a dense layer. From the lower surface black strands, with a white core, grow out, penetrating the bark and finally the wood. The fungus fruits on the surface layer which extends up the trunk of the host a short distance, a conical stage of black bristle-like stalks (*Graphium*) appears first, followed by the black globose carbonaceous perithecia. Infected plants die back gradually as the disease progresses, until girdled, when wilting and death occurs.

What is thought to be the same species has been found in the British West Indies, causing a root disease and rot of tubers or rhizomes of arrowroot (*Maranta*), *Colocasia*, *Dioscorea*, and other herbaceous plants. The external rhizomorphs are lacking, but the internal conditions are similar to those of woody plants attacked by *R. bunodes*.

Rosellinia pepo Pat. The root disease attributed to this species is called white root rot and attacks *Cajanus indicus*, *Citrus* spp., *Coffea arabica*, *Hymenaea courbaril*, *Manihot utilisima*, *Theobroma cacao*, and other woody plants in Porto Rico, Martinique, Guadeloupe, Trinidad, and the British Antilles. Herbaceous plants, such as dasheens (*Colocasia*), *Musa*, and *Canavalia*, are also attacked at times.

The fungus forms an irregular gray, then black, coating over infected roots. White strands grow out from this layer through the bark and form another much-branched white fungus layer or "fan" between the wood and the bark. The white strands penetrate the wood along the medullary rays. The fungus is black only when exposed. The effect on diseased plants is similar to that produced by *R. bunodes*.

Septobasidium acaciae Saw. See *Acacia*.

Septoria arethusa Penz. Ochraceous leaf and fruit spots on *C. limonia* and *Citrus* sp. in Brazil, Argentina, Italy, France, and Algeria. *Podocarpus* sp. has also been given as a host, but this reference seems doubtful.

Septoria cattanei Thum. Small ochraceous leaf spots on *C. medica* in India and the Canary Islands.

Septoria cinereascens (Dur. and Mont.) Sacc. On leaves of *C. aurantium* in Algeria.

Septoria citri Pass. On leaves of *Citrus* sp. in Italy, Spain, Portugal, and France.

Septoria depressa McAlp. Brown to black circular to irregular sunken spots on fruit of *C. limonia*, *C. medica*, and *C. sinensis* in Australia.

Septoria flaccescens McAlp. Apical brown areas on leaves of *C. sinensis* in Australia.

Septoria glaucescens Trab. "Spot disease" of mandarin (*C. nobilis* var.) in French North Africa.

CITRUS—Continued.

- Septoria limonum** Pass. On leaves and fruit of *Citrus* spp. in Italy.
- Septoria sicula** Penz. On leaves of *C. limonia* in Italy.
- Septoria tibia** Penz. Brown leaf spots on *C. limonia* in Italy.
- Septoria westraliensis** McAlp. Circular to irregular ashen-gray leaf spots with brown margins on *C. sinensis* in Australia.
- Septoriopsis citri** Paul. and Gz. Frag. On fruit of *C. sinensis* in Spain.
- Sphaeropsis pseudo-diplodia** (Fckl.) Delacr. On twigs and branches of *Citrus* sp. in Brazil. Apparently not distinct from *Diplodia cacaotola*.
- Sphaeropsis tumefaciens** Hedges. Knots or galls are produced on the twigs and branches, varying from 0.37 to 0.66 inch in diameter. A witches'-broom effect may appear, but ultimately the portion above the gall dies. The disease occurs on *C. aurantifolia* and *C. sinensis* in Cuba, British Guiana, and Jamaica.
- Sphaerostilbe repens** B. and Br. See Hevea.
- Sporidesmium griseum** McAlp. Dirty-gray scabby patches, cracking and breaking into irregular areas on fruit and leaves of *C. limonia* and *C. sinensis* in Australia and New Zealand.
- Trichoseptoria alpei** Cav. Causes a spotting of fruit of *C. limonia* in Italy.
- CLADRASTIS.** YELLOWWOOD. Trees grown for their white flowers and handsome foliage.
- Ascochyta cladrastidis** Kab. and Bub. On *C. lutea* in Bohemia.
- Cercospora cladrastidis** Jacz. Subcircular brown leaf spots with red margins on *C. (Maackia) amurensis* in Japan and Siberia.
- Uromyces amurensis** Kom. Yellow and cinnamon-brown to black rust pustules on leaves of *C. (Maackia) amurensis* in Siberia.
- Uromyces cladrastidis** Kusano. Black rust sori on leaves of *C. shikokiana* in Japan.
- Uromyces shikokianus** Kusano. Yellow-brown to black rust sori on leaves of *C. shikokiana* in Japan.
- CLARKIA.** Flower-garden annuals.
- Phytophthora parasitica** Dastur. See Ricinus.
- CLAYTONIA.** SPRING BEAUTY. Succulent herbs cultivated for their flowers.
- Puccinia claytoniae** Thuem. Rust on leaves of *C. arctica* in Siberia.
- LEMATIS.** Climbing or erect perennial garden plants with handsome, showy flowers.
- Aecidium englerianum** P. Henn. and Lind. Leaf rust on *C. wightiana* in Abyssinia and Tanganyika.
- Aecidium orbiculare** Barcl. Rust causing hypertrophy of stems of *C. grata*, *C. orientalis*, and *C. puberula* in India.
- Aecidium otagensis** Limds. A rust deforming the leaves, peduncles, and flowers of *C. colensoi*, *C. hexapetala*, *C. indivisa*, and *C. marata* in New Zealand. Possibly a stage of *Puccinia clavata*.
- Ascochyta indusiata** Bres. Subcircular to ovoid dull-brown leaf spots on *C. recta* in France and Germany.
- Ascochyta vitalba** Br. and Har. On leaves and branches of *C. vitalba* in France.
- Coleosporium clematidis** Barcl. Yellow to orange rust pustules on leaves of *Clematis* spp. in China, Japan, Siberia, India, Australia, Russia, Abyssinia, central and south Africa, and Australia.
- Coleosporium clematidis-apifoliae** Diet. Yellow rust pustules on leaves of *C. epifolia* and *C. parvifolia* in Japan and China.
- Coleosporium elongatum** Syd. Leaf rust on *C. hedysarifolia* in Japan.
- Erysiphe taurica** Lévl. See Althaea.
- Marsonia clematidis** All. Large irregular indefinite dull-brown zoned leaf spots on *C. vitalba* and *Thalictrum minus* in Germany.
- Phyllosticta bacteriosperma** Pass. Irregular angular dull-brown leaf spots on *C. vitalba* in Italy.
- Phyllosticta beguinotiana** Sacc. Small subcircular light-brown, then white, leaf spots on *C. flammula* in Italy.
- Phyllosticta corrodens** Pass. Indefinite gray leaf spots with dull-brown margins on *C. vitalba* in Italy and Denmark.
- Phyllosticta intermedia** Allesch. Small subcircular to angular whitish leaf spots with brown margins on *C. recta* in France and Germany.
- Phyllosticta vitalba** Cke. Leaf spots on *Clematis* sp. in Great Britain.
- Phyalospora disseminata** Sacc. On leaves of *C. glauca* in Siberia.
- Puccinia clavata** Syd. Brown rust pustules on small sunken brown leaf spots on *C. hexapetala* in New Zealand.
- Puccinia exhausta** Diet. Brown powdery rust pustules on leaves of *C. heracleaeifolia*, *C. javanica*, *C. talwaniana*, and *C. tuberosa* in Java, Japan, and the Philippines.
- Puccinia insidiaria** Berk. Leaf rust on *C. nutans* in India.
- Puccinia wattiana** Barcl. Brown rust pustules on sunken black leaf spots on *C. buchaniana*, *C. gouriana*, and *C. puberula* in India.
- Septoria aecidicola** Pat. On leaves of *C. cirrhosa* in Tunis.
- Septoria cirrhosae** Sacc. Circular dull-yellow leaf spots with dark-purple margins on *C. cirrhosa* in Sicily.
- Septoria clematidis** Rob. and Desm. Circular to angular grayish leaf spots with dull-brown margins on *C. flammula*, *C. glauca*, and *C. vitalba* in Ecuador, Siberia, and Europe.
- Septoria clematis-flammulae** Roum. Small subcircular gray, then white, leaf spots with brown margins on *C. flammula* in Dalmatia, France, and Italy.
- Septoria clematis-rectae** Sacc. Grayish-white leaf spots on *C. recta* in Italy.
- Septoria viticellae** Pass. Leaf spots on *C. viticella* in Italy.
- CLEOME.** Spiderflower. Mostly weedy herbs.
- **Cercospora conspicua** Earle. Leaf spot on *C. pentaphylla* in Porto Rico.
- CLERODENDRUM.** GLORY BOWER. Shrubs, trees, or climbers grown for their showy flowers.
- Aecidium clerodendri** P. Henn. Leaf rust on *C. calamatosum*, *C. cyrtophyllum*, *C. fragrans*, *C. intermedium*, and *C. paniculatum* in Java, Japan, and the Philippines.
- Aecidium clerodendricola** P. Henn. Leaf rust on *C. buchnerum* in Tanganyika.
- Aecidium clerodendronis** P. Henn. Leaf rust on *Clerodendrum* sp. in Indo-China.
- Cercospora bakeri** Syd. On leaves of *C. intermedium* in the Philippines.
- Cercospora clerodendri** Miy. Leaf spot on *Clerodendrum* sp. in China.
- Chrysomyxa peregrina** Syd. and Butl. Yellow-brown rust pustules on leaves of *Clerodendrum* sp. in India.
- Coleosporium clerodendri** Diet. Yellow to orange rust pustules on lower leaf surfaces of *C. cyrtophyllum*, *C. fragrans*, and *C. trichotomum* in Java, Japan, and Formosa.
- Hemileia scholzii** Syd. Yellow powdery rust pustules on leaves of *Clerodendrum* sp. in central and south Africa and Portuguese East Africa.
- Phyalospora clerodendri** Syd. On leaves of *Clerodendrum* sp. in the Congo.
- Puccinia erebia** Syd. Leaf rust on *C. commersonii* and *C. minahassae* in the Philippines.
- Synchytrium collapsum** Syd. and Butl. Small swellings on both leaf surfaces of *Clerodendrum* sp. in India.
- Uredo clerodendricola** P. Henn. Leaf rust on *C. inerme* in Java, Ceylon, and China.

- CLETHRA.** WHITE ALDER. Shrubs or small trees grown for their spikes of white fragrant flowers.
Phyllachora rubefaciens Rehm. Shiny black stromata on red-brown leaf spots on *C. laevigata* in Brazil.
Phyllosticta clethrae Syd. On leaves of *C. scabra* in Austria.
Pucciniastrum kusanoi Diet. Leaf rust on *C. barbinervis* in Japan.
- CLIANTHUS.** PARROTBEAK. Sometimes called Glory pea. Tender half trailing shrubs with large showy flowers.
Ascochyta clianthi Tassi. Dull-brown to whitish irregular leaf spots on *C. arboreus* in Italy.
- CLINTONIA.** Low-growing liliaceous herbs.
Puccinia clintoniae-udensis Bub. Cinnamon-colored rust pustules on the upper surfaces of leaves of *C. udensis* in Amur Province, Siberia.
- CLITORIA.** BUTTERFLY PEA. Hardy perennial vines with pealike flowers.
Cercospora pantoleuca Syd. Dark-brown blotches on leaves of *C. ternatea* in India and the Philippines.
Cercospora ternatea Petch. Circular black spots with gray centers, which become gray with yellow or pale-greenish margins, on leaves and pods of *C. ternatea* in Ceylon.
Corticium salmonicolor B. and Br. See *Citrus*.
Melasmia juruana P. Henn. Black shiny stromata on leaves of *Clitoria* sp. in Brazil.
Uromyces clitoriae Arth. Powdery brown rust pustules on lower leaf surfaces of *C. mexicana* in Mexico.
Uromyces neurocarpi Diet. Powdery brown rust pustules on leaves of *C. cajanifolia* and *C. rubiginosa* in Porto Rico, Cuba, Jamaica, Trinidad, and Brazil.
Uromyces yurimaguasensis P. Henn. Brown rust pustules in circular dull-brown leaf spots on *C. arborens* and *Clitoria* sp. in Peru and Panama.
- CLIVIA (Himantophyllum).** KAFIR LILY. Tender bulbous plants with evergreen foliage and showy red or red and yellow flowers.
Ascochyta cliviae Magnaghi. On leaves of *C. nobilis* in Italy.
Colletotrichum cliviae Oud. Small irregular red to brown leaf spots on *C. miniata* and *Clivia* sp. in Bohemia, Holland, and France.
Physalospora himantophylli Gz. Frag. On leaves of *Himantophyllum (Clivia) mimata* in Spain.
- CLUSIA.** Tropical trees used to some extent for shade.
Coccomyces clusiae (Lév.) Sacc. Black fruiting bodies on leaves of *Clusia* sp. in Costa Rica and northern South America.
Phyllosticta clusiae Allesch. On leaves of *Clusia* sp. in Brazil.
Septogloeum clusiae Karst. and Har. On leaves of *Clusia* sp. in Colombia.
Uredo clusiae Arth. Dark-brown rust pustules on leaves of *C. rosea* in Porto Rico.
- CLYTOSTOMA.** Ornamental vines, grown for their beautiful flowers.
Phyllachora amphigena Speg. See *Bignonia*.
- COBAEA.** Climbers grown for their large bell-shaped flowers.
Phyllosticta cobaeae Tassi. Leaf spot on *C. scandens* in Italy.
Septoria oligocarpa Nannizzi. Small, subcircular leaf spots with narrow purple margins on *C. scandens* in Italy.
- COCCOLOBA.** SEA GRAPE. Tropical shrubs and trees, some species grown for their fruit or foliage.
Cercospora bicolor Wint. Leaf spots on *C. sagittata* in Portugal.
Fabraea cocolobae P. Henn. Effuse dull-brown areas on leaves of *Coccoloba* sp. in Brazil.
Phyllachora cocolobae Speg. Black stromata on leaves of *Coccoloba* sp. in Argentina.
Phyllachora simplex Starb. Small scattered black stromata on leaves of *C. laurifolia* and *Coccoloba* sp. in Paraguay and Porto Rico.
Phyllosticta cocolobae E. and E. Circular, purplish leaf spots, 3 to 5 millimeters in diameter on *C. unifera* in Porto Rico and the Bahamas.
Physalospora cocolobae Rehm. Circular yellowish leaf spots on *Coccoloba* sp. in Brazil.
Uredo cocolobae P. Henn. Powdery light-brown rust pustules on leaves of *C. populifolia* and *C. unifera* in Brazil, Porto Rico, and Cuba. Also reported from Florida.
- COCCOTHRINAX.** See *Palmae*.
- COCCULUS.** SNAIL SEED. Shrubs grown for their handsome foliage and ornamental fruit.
Phomopsis cocculi Luigi. Light-ashen spots with irregular black borders on leaves of *C. laurifolius* in Italy.
Phyllosticta cocculi Speg. Leaf spot on *C. laurifolius* in Argentina.
Phyllosticta thunbergii Wint. Large, indefinite, irregular, pale-yellow leaf spots on *C. thunbergii* in Japan.
- COCHLEARIA.** Small fleshy seaside herbs.
Puccinia cochleariae Lindr. Powdery black rust pustules on leaf blades and petioles of *C. fenestrata*, *C. groenlandica*, and *C. pyrenaica* in France and Greenland.
- COCOS.** Coconut. See *Palmae*.
- CODIAEUM (PHYLLAUREA Ag.).** Tropical shrubs grown for their variegated foliage.
Ascochyta banosensis Syd. On leaves of *C. variegatum* in the Philippines.
Corticium koleroga (Cke.) v. Hoeh. See *Coffea*.
Corticium stevensii Burt. Leaf blight of *C. variegatum* in Trinidad.
Fomes lamaensis Murr. See *Hevea*.
Gloeosporium sorauerianum Allesch. Large irregular indefinite yellow to ashen leaf spots on *C. variegatum* in France, Holland, and Germany.
Laestadia minuscula (Lév.) Sacc. On leaves of *C. variegatum* in Java.
Phyllosticta codiae Diet. On leaves of *Codiaeum* sp. in the Philippines.
Phyllosticta reysei (Sacc.) Yates. On leaves of *C. variegatum* in the Philippines.
- CODONOPSIS.** Twining or decumbent perennials with showy flowers.
Coleosporium horianum P. Henn. Golden-yellow to brown rust pustules on leaves of *C. lanceolata* in Japan.
- CODONORCHIS.** See *Orchidaceae*.
- COELOGLOSSUM.** See *Orchidaceae*.
- COELOGYNE.** See *Orchidaceae*.
- COFFEA.** COFFEE. Tropical shrubs or small trees.
Anthostomella coffeae Delacr. Found in connection with die-back of twigs of *C. arabica* in Mexico.
Ascochyta coffeae P. Henn. Subcircular brown leaf spots with darker margins on *C. arabica* in Brazil.
Ascospora coffeae Faber. A wound parasite destroying large areas of bark of *C. robusta* trees in Java. Leaves of infected plants turn yellow and fall.
Atichia millardeti Raeb. See *Cinnamomum*.
Cephalosporium sp. Concentrically ringed brown leaf spots on *C. arabica* in Porto Rico.

COFFEA—Continued.

- Cercospora coffeae** Zimm. Subcircular dark-brown leaf spots on *C. arabica*, *C. laurina*, *C. robusta*, and *C. stenophylla* in Malaya and Tanganyika.
- Cercospora coffeicola** B. and C. Circular brown leaf spots, 6 to 10 millimeters in diameter, sometimes causing defoliation. Young twigs are also attacked to some extent. The disease is serious on the fruit, appearing first as small brown spots which enlarge to cover about half the fruit, generally the upper half. The diseased pulp turns black and dries down, adhering to the parchment, making the process of preparation more difficult, and lowering the grade of the final product. The disease occurs in practically all coffee-growing countries, including Porto Rico, Cuba, Jamaica, Trinidad, Surinam, Guatemala, Costa Rica, Brazil, Hawaii, Dutch East Indies, India, Java, New Caledonia, Queensland, the Philippines, and central Africa. The hosts are *C. arabica* and *C. buxobensis*.
- Cercospora herrerana** Farn. Circular chestnut-brown leaf spots with dark-purple margins on leaves and fruit of *C. arabica* in Mexico. Probably the same as the preceding species.
- Cobweb disease.** A Javan disease similar to the "koleroga" disease of coffee (*C. arabica*), but due to a different fungus. White, thick, branching fungus threads grow along the surface of twigs and leaves. Infected leaves turn yellow, wilt, and die, accompanied by die-back of the twigs.
- Colletotrichum coffeanum** Noack. Circular to irregular, large, brown, and finally gray spots on leaves of *C. arabica* in Porto Rico, Costa Rica, Brazil, central Africa, Madagascar, Reunion, Java, and India. A die-back of twigs is often associated with this or closely related species. The berries are also attacked, the spotting reducing the grade of infected material. *Colletotrichum incarnatum* Zimm. from Ceylon is probably not distinct.
- Cortium koleroga** (Cke.) v. Hoeh. (*Pellicularia koleroga* Cke.) The "koleroga" disease of coffee (*C. arabica*) occurs in Porto Rico, Cuba, Jamaica, Surinam, Venezuela, Guatemala, India, Java, Malaya, Queensland, Congo, and possibly in Brazil. The fungus threa is grow up along the undersides of the twigs and out onto the lower leaf surfaces. The leaves turn black, wither up or drop, or remain suspended hanging by the fungus filaments. These strands are at first white and finally deep brown. Diseased twigs may die back for considerable distances. The berries are attacked to some extent, infected berries showing blackened grains. Additional hosts are *Citrus* spp., *Codiaeum variegatum*, *Cucumis anguria*, *Garcinia mangostana*, *Hevea brasiliensis*, and *Luffa aegyptiaca*.
- Corticium salmonicolor** B. and Br. See *Citrus*.
- Dictyothyriella mucosa** Syd. Small, black, superficial fruiting bodies on *C. liberica* in Indo-China and the Philippines.
- Diplodia cacaicola** P. Henn. See *Theobroma*.
- Euryachora coffeicola** Avena. Black stromata on leaves of *C. arabica* in Brazil.
- Euryachora liberica** Oud. A doubtful species on branches of *C. arabica* and *C. liberica* in Java.
- Fomes lamaensis** Murr. See *Hevea*.
- Fomes lignosus** Klotzsch. See *Hevea*.
- Fusarium coffeicola** P. Henn. Found in connection with die back of *C. arabica* and *C. liberica* in central Africa.
- Gloeosporium coffeanum** Delacr. Anthracnose on leaves of *C. arabica* in Reunion and Ceylon. Probably the same as *Colletotrichum coffeanum* Noack.
- Gloeosporium coffecolum** Tass. On *C. arabica* in Italy. Not distinct from *G. cingulata*.
- Helmintosporium coffeae** Mass. On leaves of *C. liberica* in Gold Coast Colony.
- Helmintosporium ubangense** P. Henn. On leaves of *Coffea* sp. in tropical Africa.
- Hemileia vastatrix** B. and Br. This very destructive rust has destroyed the coffee industry of some countries and made it unprofitable in others. The characteristic yellow rust pustules develop during wet weather on the lower leaf surfaces, and more rarely on fruits and tips of young branches. Small yellow spots first appear, which increase in size, merge, and finally involve the entire leaf blade, which becomes brown and falls. Plants of all ages are attacked, defoliated, and killed within a season or two. *C. arabica* is very susceptible. Other species (*C. bengalensis*, *C. laurina*, *C. liberica*, *C. robusta*, *C. traxan-korensis*) vary in their resistance, some varieties being practically immune. The rust occurs in Burma, Ceylon, India, Indo-China, China, Malaya, Philippines, Union of South Africa, east Africa, Malacca, Sumatra, Java, Fiji, Samoa, Mauritius, Reunion, New Hebrides, New Caledonia, Gold Coast, and Madagascar. The disease does not occur in the western Tropics.
- Laestadia coffeicola** Speg. Circular whitish leaf spots on *C. arabica* in Costa Rica.
- Leptosphaeria coffeigena** (B. and C.) Sacc. On leaves of *C. arabica* in Brazil.
- Leptosphaeria tonduzi** Speg. On leaves of *Coffea* sp. in Costa Rica.
- Metasphaeria bifoveolata** Speg. On *C. arabica* in Costa Rica.
- Mosaic** (*Phloem necrosis*). This disease, apparently of the group of infectious mosaics, attacks *C. liberica*, and occasionally *C. arabica*, in Surinam. In the acute form the leaves wilt rapidly and hang down. The roots become black and the entire plant is dead within a few weeks. In other cases the disease progresses more slowly. The leaves turn pale-green, then yellow, and finally fall, being replaced by small abnormal ones.
- Muratella coffeae** Bunting. Attacks *C. liberica* on Gold Coast Colony.
- Mycosphaerella coffeae** Noack. On leaves of *C. arabica* in Brazil.
- Mycosphaerella coffeicola** Cke. Subcircular, brown leaf spots with narrow gray margins on *C. arabica* in Jamaica, Ecuador, Venezuela, Brazil, Uganda, and India.
- Ophionectria follicola** A. Zimm. On leaves of *C. liberica* in Java.
- Phoma coffeicola** Tassi. On branches and twigs of *C. arabica* in Italy.
- Phyllosticta camoensis** Delacr. White thickened leaf spots on *C. camoensis* in tropical Africa.
- Phyllosticta coffeicola** Delacr. Circular whitish leaf spots on *C. arabica* in Malaya, Java, Mexico, Costa Rica, and Brazil. Probably synonymous with the Spegazzini species, which has priority.
- Phyllosticta coffeicola** Speg. On leaves of *Coffea* sp. in Costa Rica and Brazil.
- Phyllosticta usteri** Speg. Circular, then confluent, leaf spots on *C. arabica* and *C. robusta* in Ceylon and Brazil.
- Placosphaeria citricola** Berl. and Roum. On leaves of *C. arabica* in Indo-China.
- Pleonectria coffeicola** A. Zimm. On leaves of *C. liberica* in Java.
- Ramularia goldiana** Sacc. Subcircular zonate spots on leaves and stems of *C. arabica* in Brazil.
- Rhizoctonia lamellifera** Small. See *Grevillea*.
- Rosellinia bunodes** B. and Br. See *Citrus*.
- Rosellinia pevo** Pat. See *Citrus*.
- Rostrella coffeae** Zimm. On *C. arabica* in Guatemala.
- Saccardinula tahitensis** Pat. On leaves of *Coffea* sp. in Tahiti.
- Sclerotium** sp. Causes a damping-off of coffee (*C. arabica*) seedlings in the Philippines.
- Sclerotium coffecolum** Stahl. The "coremium" or sclerotia disease of *C. abocuta*, *C. arabica*, *C. excelsa*, and *C. liberica* in Surinam is characterized by dark-brown concentric spots on leaves and fruits. Small, greenish-brown or orange-brown sclerotia develop on the diseased areas.
- Septoria berkeleyi** Sacc. and Trav. Subcircular pale-brown leaf spots on *C. arabica* in South America.

COFFEA—Continued.

- Septoria coffeae** Wakef. On leaves of *C. arabica* in Uganda.
- Septoria coffeicola** P. Henn. Circular white leaf spots with dark-brown margins on *C. liberica* in Kamerun.
- Stibella flavida** (Cke.) Lind. Circular dark-brown and finally light-brown leaf spots on *C. arabica*, *C. excelsa*, *C. perrieri*, and *C. robusta* in Porto Rico, Jamaica, Cuba, British West Indies, Central America, Mexico, Trinidad, Ecuador, Venezuela, Colombia, British Guiana, Dutch Guiana, and Java. Other plants, including *Commelina* sp., *Bryophyllum pinnatum*, *Inga vera*, and *Mangifera indica*, growing under or adjacent to coffee are attacked. The spots are from 6 to 12 millimeters in diameter, and are often so numerous as to destroy the leaves. On the upper surfaces of the spots and, to some extent, on the lower surfaces yellow, hairlike projections, 1 to 4 millimeters long, with a small red terminal spore mass occur. Young twigs and the fruit are also attacked to some extent. The disease may be sufficiently serious to reduce yields as much as 75 per cent.
- Trachysphaera fructigena** Tabor. and Bunting. The "mealy pod" disease attacks *C. liberica* and *Theobroma cacao* in Gold Coast Colony. Coffee fruits of all ages are attacked, but the young berries are most subject to the disease. Their development is stopped and they shrivel up and become hard, a dark purplish-brown discoloration appearing, which is later covered by a whitish or pinkish-brown mealy incrustation. On cacao pods small brown areas appear which increase rapidly until the entire fruit is involved. The same mealy layer ultimately appears.
- Venturia coffeicola** Avena-Sacca. On leaves of *C. arabica* in Brazil.
- COIX.** JOB'S-TEARS. Tall, broad-leaved grasses with beadlike inflorescences.
- Phyllachora coicis** P. Henn. Shiny black elongate stromata on lower leaf surfaces of *C. agrestis* and *C. lachryma-jobi* in India, Indo-China, and the Philippines.
- Sclerospora maydis** (Rac.) Butl. See Zea.
- Uredo operta** Syd. and Butl. Straw-colored rust sori on leaves of *C. lachryma-jobi* in India and Ceylon.
- Ustilago coicis** Bref. Grain transformed into a black mass of smut spores, surrounded by a membrane and confined within the glumes. On *C. agrestis* and *C. lachryma-jobi* in India, Java, and Japan.
- COLCHICUM.** AUTUMN CROCUS. Autumn-flowering bulbous plants.
- Ascochyta juehii** Bub. Round or elliptical leaf spots on *C. autumnale* in Austria.
- Cladochytrium maurii** Hariot. Small brown spots on leaves of *C. autumnale* in France.
- Cylindrosporium colchici** Pass. On leaves of *C. alpinum* and *C. officinale* in France and Holland.
- Perisporium colchici** Rbh. Black superficial fungus on leaves of *C. autumnale* in Germany.
- Phyllosticta latemarensis** Kab. and Bub. Small irregular black, often confluent, leaf spots on *C. autumnale* in Italy.
- Septoria colchici** Pass. Long light-brown spots on leaves of *C. alpinum* and *C. officinale* in France, Italy, and Germany.
- Septoria gallica** Sacc. and Syd. Oval to elongate, often confluent, spots on leaves of *C. autumnale* and *C. officinale* in France and Austria.
- Uromyces colchici** (Schlecht) Rab. Long rows of black, powdery streaks occur on the leaves of *C. autumnale*, *Allium cepa*, *A. magicum*, *A. rotundum*, *Bulbocodium vernum*, *Muscari comosum*, *M. racemosum*, *Narcissus* spp., *Paris quadrifolia*, *Scilla bifolia*, and *Tulipa* spp., in England, Yugoslavia, Spain, and Dalmatia. A smut referred to this species occurs to a limited extent in the United States on *Polygonatum* spp. and *Vagnera* spp., but if the European and American forms are conspecific different strains are most certainly involved.
- Uromyces colchici** Mass. Brown powdery rust pustules on both leaf surfaces of *C. autumnale*, *C. barbaricum*, *C. speciosum*, and *C. spectabilis* in Great Britain.
- COLEOSANTHUS.** See Brickellia.
- COLEUS.** Showy-leaved herbs used as bedding plants.
- Aphelenchus olesistis** Ritz. Bos. See Begonia.
- Meliola cavitiesis** Yates. Black superficial fungus patches on leaves of *Coleus* sp. in the Philippines.
- COLLETIA.** Spiny shrubs grown in warm regions or greenhouses.
- Dothidella colletiae** (P. Henn. and Lind.) Theiss. and Syd. On branches and cladophylls of *C. spinosa* in Chile.
- Montagnella curumamuel** Speg. Black circular stromata on leaves of *C. cruciata* and *C. ferox* in Argentina.
- Piaostroma lorentzianum** (Speg.) Theiss. and Syd. On branches of *C. ferox* in Argentina.
- COLLOMIA.** See Gilia.
- COLOCASIA.** DASHEEN. ELEPHANT'S-EAR. Yautia. Eddoe. Taro. Tropical aroids grown for their ornamental leaves (bedding plants) and for edible roots.
- Cercospora caladii** Cke. var. **colocasiae** v. Hoeh. Subcircular brown zoned leaf spots on *Colocasia* sp. in Samoa.
- Heterosporium colocasiae** Mass. On leaves of *Colocasia* sp. in Jamaica.
- Mycesphaerella colocasiae** K. Hara. Circular, then confluent, brown, zoned, more or less sunken leaf spots with dark-brown margins on *C. antiquorum* in Japan. The disease greatly reduces the yields of tubers.
- Peronospora trichotoma** Mass. This supposed species of downy mildew, said to cause a tuber rot of *C. esculenta* in the West Indies, is now considered a hyphomycete overgrowing *Phytophthora colocasiae* Rac.
- Phyllosticta colocasiae** v. Hoeh. Circular brown zoned leaf spots on *Colocasia* sp. and *Dieffenbachia* sp. in Samoa and Porto Rico.
- Phyllosticta colocasicola** v. Hoeh. Probably not distinct from the preceding species.
- Phytophthora colocasiae** Rac. This downy mildew causes a serious blight of *C. antiquorum* in India, Java, Ceylon, Formosa, Philippines, and Hawaii. Circular to oval dark-brown spots appear on the leaves and, becoming irregular, may occupy most of the leaf surface. They become yellow-brown in color, and are often marginally zoned in shades of brown, green, and yellow. The petioles may be soft rotted, so that the leaves fall over. Inflorescences and the corms are attacked in turn, although if the plants are attacked while young no corms are formed. Where corms form they are destroyed by a soft rot.
- COLQUHOUNIA.** Tender plants grown for their whorls of scarlet and yellow flowers.
- Puccinia leucophaea** Syd. and Butl. Leaf rust on *C. coccinea* in India.
- COLUTEA.** BLADDER SENNA. Shrubs grown for their attractive flowers and ornamental bladder-like pods.
- Ascochyta coluteae** Lambr. and Fautr. On leaves of *C. orientalis* in Yugoslavia.
- Ovularia vogeliana** Sacc. and Syd. White fungus layers over leaf surfaces of *C. arborescens* in Germany.
- Uromyces genistae-tinctoriae** (Pers.) Wint. See Caragana.

- COMARUM.** Decumbent herbs. The genus is usually placed in *Potentilla*.
Mycosphaerella innumerella (Karst.) Starb. On leaves of *C. palustre* in Europe.
Phyllosticta argentiniae Desm. See *Potentilla*.
Physoderma comari (Berk. and White.) Lagh. Black smut sori in leaves of *C. palustre* in Great Britain and Scandinavia.
Septogloeum comari Allesch. and Bres. Dull yellow leaf spots on *C. palustre* in Sweden, Denmark, Estonia, and Germany.
Septoria comari Lasch. On leaves of *C. palustre* in Germany.
Venturia palustris Sacc. On leaves of *C. palustre* in Belgium and Germany.
- COMBRETUM.** Tropical shrubs and trees.
Hyphaster kutuensis P. Henn. Superficial black stromata on leaves of *C. baumii* in the Union of South Africa.
Phyllosticta combreticola P. Henn. Irregular ashen leaf spots on *C. argenteum* in Germany.
Uredo kampuluvensis P. Henn. See *Baphia*.
- COMMELINA.** DAYFLOWER. Perennial or annual herbs.
Cercospora commelinae Kalchbr. and Cke. Circular, dull-brown leaf spots on *C. benghalensis* in the Union of South Africa.
Cylindrosporium kilimandscharicum Allesch. Small, definite, brown, often confluent leaf spots on *Commelina* sp. in Tropical Africa.
Ilosporium commelinae Stevens. Circular dull-yellow leaf spots on *C. longicaulis* and *C. virginica* in Porto Rico.
Kordyana celebensis Racib. Causes leaf spots on *C. benghalensis* in Java.
Phyllosticta commelinicola Young. Indefinite leaf spots on *C. nudiflora* in Porto Rico.
Puccinia commelinae Holw. Brown rust pustules on brown irregular leaf spots on *Commelina* sp. in Mexico.
Stilbella flavida (Cke.) Lind. See *Coffea*.
Tilletia (?) *commelinae* Kom. Doubtful species of smut in ovaries of *C. communis* in Manchuria and Japan.
Uredo commelyneae Kalchbr. Leaf rust on *C. elegans* and *C. virginica* in Porto Rico, St. Croix, and Trinidad.
Uredo ochracea Diet. Brown rust pustules on leaves of *C. benghalensis*, *C. communis*, *C. longicaulis*, and *C. obliqua* in Brazil, India, Japan, Ceylon, and the Philippines.
Uredo pedicellata Rac. Rust on leaves and bracts of *Commelina* sp. in Java.
Uromyces commelinae Cke. Leaf rust on *C. benghalensis*, *C. communis*, *C. forskalii*, *C. longicaulis*, *C. nudiflora*, *C. subulata*, and *Tradescantia cumanensis* in the Union of South Africa, India, Japan, Portuguese East Africa, Uganda, Porto Rico, Cuba, Jamaica, Guatemala, and South America.
Uromyces tosenis P. Henn. Brown rust pustules on leaves of *C. communis* in Japan.
- CONVALLARIA.** LILY OF THE VALLEY. Herbaceous perennials cultivated for their racemes of fragrant flowers.
Ascochyta majalis Massal. Causes red-brown oblong to irregular spots on leaves of *C. majalis* in Italy.
Dendrophoma convallariae Cav. Dark-red elongated spots are produced on the leaves of *C. majalis* in Great Britain, Denmark, Italy, and Germany.
Gloeosporium convallariae Allesch. On leaves of *C. majalis* in Germany.
Heterosporium ornithogali Klotzsch. Oval to irregular brown spots on the leaves of *Convallaria* sp. and other species of the family *Liliaceae* in New Zealand.
Lophodermium herbarum (Fr.) Fekl. On leaves of *C. majalis* in France, Belgium, Switzerland, Sweden, Finland, and Germany.
Mycosphaerella asteroma (Fr.) Lind. On leaves of *Convallaria* sp., *Maianthemum* sp. and *Polygonatum* sp. in Europe.
Mycosphaerella brunneola (Fr.) All. and Schnab. On leaves of *C. majalis* in Denmark, Belgium, and Austria.
Phyllachora melanoplaca Sacc. Forms black, shiny, slightly concave spots on the leaves of *Convallaria majalis*, *Veratrum album*, *V. nigrum*, and *V. viride* in Switzerland and Austria.
Sclerotium convallariae West. On leaves of *Polygonatum verticillatum* (*C. verticillata*) in Belgium.
Septoria brunneola (Fr.) Niessl. (*S. convallariae* West.) Causes brown leaf spots on *C. majalis* and *Polygonatum* sp. in Siberia, Great Britain, Belgium, Sweden, Italy, and Moravia. Probably the imperfect stage of *Mycosphaerella brunneola*.
Septoria majalis Aderh. Large brown spots without definite margins, often involving entire leaves of *C. majalis* in Germany.
Septoria subradians (Fr.) Karst. A leaf-spot disease of *C. majalis* in Denmark.
- CONVOLVULUS.** BINDWEED. Annual and perennial climbing herbs, mostly weeds.
Aecidium congoanum P. Henn. Leaf rust on *Convolvulus* sp. in the Kongo.
Aecidium convolvulinum Speg. Leaf rust on *Convolvulus* sp. in Argentina.
Cercospora sorokinii Sacc. On leaves of *Convolvulus* sp. in central Asia.
Entyoma convolvuli Bres. Smut sori in circular to oblong brown leaf spots, on *C. soldanella* in Portugal.
Puccinia macrocephala Speg. See *Ipomoea*.
Septoria calystegiae West. Small subcircular brown leaf spots on *C. arvensis* and *C. soldanella* in Europe.
Septoria convolvulina Speg. Circular white leaf spots with reddish margins on *C. arvensis* in Argentina.
Septoria longispora A. Bond. Small circular brown to gray leaf spots on *C. arvensis* and *C. lineatus* in Russia.
Septoria obesipora Oud. Leaf spot on *C. sepium* in Holland.
Thecaphora hyalina Fingerh. A smut reducing ovaries of *C. arvensis*, *C. sepium*, and *Soldanella* sp. to brown, powdery, spore masses in Europe. What is apparently this species has been collected in California.
Uromyces gemmatus B. and C. Leaf rust on *Convolvulus* sp. in Cuba.
- COPAIFERA.** Brazilian trees, producing a balsam called *copaiba*.
Phyllachora copaiferae Speg. Black stromata on irregular brown leaf spots on *C. langsdorffii* in Paraguay and Brazil.
Uredo copaiferae P. Henn. Leaf rust on *Copaifera* sp. in Brazil.
- COPERNICIA.** See *Palmae*.
- COPROSMA.** Shrubs or small trees grown for their ornamental fruit and variegated leaves.
Phyllosticta coprosmae McAlp. Ashen leaf spots, with narrow, raised, dark-red margins on *Coprosma* sp. in Australia.
Puccinia coprosmae Cke. Rusty-brown rust pustules on irregular leaf spots on *C. billardieri*, *C. hirtella*, and *C. lucida* in New Zealand.
Septoria coprosmae Cke. On leaves of *Coprosma* sp. in New Zealand.

CORCHORUS. Jute. Shrubs or herbs, some species yielding fiber.

Corticium salmonicolor B. and Br. See Citrus.

Diplodia corchori Syd. The "black band" disease of jute (*C. capsularis* and *C. olitorius*) is serious in India. Dense black bands form on the stems of infected plants up to the height of 2 or 3 feet, the bark splits, and finally only the brown, dry fibers are left, the tissue between having decayed away. The leaves of infected plants turn yellow and drop.

Hymenula nigra Saw. On *Corchorus* sp. in Formosa.

Hypochnus centrifugus (Lév.) Ful. On *Corchorus* sp. in Formosa.

Macrophoma corchori Saw. Indefinite spots and cankers on stems of *Corchorus* sp. (jute) in Formosa, causing die-back. Considered serious.

Phyllosticta corchori Saw. On leaves of *Corchorus* sp. in Formosa.

Rhizoctonia sp. See Vigna.

CORDIA. Tropical shrubs and trees, some species cultivated for fruit and ornament.

Aecidium brasiliensis Diet. Leaf rust on large round yellow spots on *C. cylindrostachya*, *C. macrophylla*, *C. nodosa*, and *C. rothii* in Brazil, Trinidad, and India.

Aecidium cordiae P. Henn. Leaf rust on *C. bullata* and *C. urticifolia* in the Dominican Republic.

Alveolaria andina Lagh. Chestnut-brown rust pustules on yellow-brown circular leaf spots of *Cordia* sp. in Ecuador.

Alveolaria cordiae Lagh. Rust on sunken spots on leaf blades and petioles of *C. cylindrostachya*, *C. ferruginea*, *C. laziflora*, and *C. riparia* in Costa Rica, Guatemala, Jamaica, Ecuador, and Colombia.

Dimerium stevensii Garm. Indefinite yellow spots on leaves of *C. corymbosa* in Porto Rico.

Guignardia javanica Koord. Indefinite dark-brown leaf spots on *C. suaveolens* in Java.

Helminthosporium cordiae Welw. and Curr. On leaves of *Cordia* sp. in west Africa.

Phyllachora caffra Syd. Black stromata on rust-brown indefinite leaf spots on *C. caffra* in the Union of South Africa.

Placosphaeria cordiae Allesch. Subcircular black stromata on leaves of *Cordia* sp. in Brazil.

Puccinia cordiae Arth. Brown rust pustules on leaves of *C. alliodora* and *C. gerascanthus* in Porto Rico and Guatemala.

Puccinia corticola Arth. and Rorer. Rust attacking leaves, twigs, branches, and trunks of *C. gerascanthus* in Trinidad. Large lesions are formed on trunks and branches.

Uredo cordiae P. Henn. A rust causing gall-like swellings on leaves and shoots and witches'-broom effect on *Cordia* sp. in Peru.

Uromyces cordiae P. Henn. Brown rust pustules on leaves of *Cordia* sp. in Brazil.

CORDYLINE. See Dracaena.

COREOPSIS. Annual or perennial herbs or shrubs, some species cultivated in flower gardens.

Puccinia coreopsidis Jacks. and Holw. Brown rust pustules on leaves of *C. mutica* (*C. mexicana*) in Guatemala.

Puccinia coreopsidis Wakef. Leaf rust on *Coreopsis* sp. in Uganda. The preceding species has priority.

CORIANDRUM. CORIANDER. Strong-smelling umbelliferous herbs the seeds of which are used in seasoning.

Protomyces macrosporus Ung. Galls up to half an inch long by an eighth of an inch wide are produced on flower stalks, petioles, and young stems of *Aegopodium podagraria*, *Carum carvi*, *Chaerophyllum hirsutum*, *Coriandrum sativum*, *Daucus carota*, *Heracleum sphondylium*, *Meum athamanticum*, *M. mutellina*, and *Pastinaca sativa* in Europe, north Africa, India, Japan, and Australia.

CORIARIA. Shrubs or herbs grown for their ornamental fruit.

Phyllosticta coriariicola Speg. Angular, then confluent, dull-brown leaf spots with purple margins on *C. ruscifolia* in Chile.

Pucciniastrum coriariae Diet. Yellow to brown rust pustules on leaves of *C. japonica* and *C. nepalensis* in India and Japan.

Septoria coriariae Pass. Irregular brown leaf spots on *C. myrtifolia* in Spain and Italy.

CORNUS. DOGWOOD. Woody plants grown for their attractive flowers and fruit.

Coryneum corni-albi (Roum.) Sacc. On leaves of *C. alba* and *C. mas* in France, Bohemia, Austria, and Germany.

Erysiphe tortilis (Wallr.) Fr. Powdery mildew on leaves of *C. alba* and *C. sanguinea* in Europe.

Helminthosporium phylophilum Karst. On leaves of *C. alba* in Finland.

Phoma thallina Sacc. On stems of *C. sanguinea* in Europe.

Pucciniastrum corni Diet. Yellow to brown rust pustules on leaves of *C. kousa* and *C. officinalis* in Japan.

Septoria corni-maritima Sacc. On leaves of *C. sanguinea* in northern Europe.

Sphaeropsis suspecta Vestreg. Stem canker of *C. sanguinea* in Russia.

COROKIA. Evergreen shrub.

Septoria corokeae P. Henn. Dull-brown spots, occupying large portions of leaf surfaces of *C. Ludleyoides* in Germany.

CORONILLA. Crown vetch. Shrubs and herbs grown in gardens for their flowers.

Ascochyta emeri Sacc. Dirty-white, often marginal, leaf spots with red margins on *C. emerus* in Italy.

Cercospora ratuensis C. Mass. Small subcircular pale-gray leaf spots with rufous margins on *C. varia* in Italy.

Helminthosporium bornmuelleri P. Magn. On leaves of *C. coronata*, *C. montana*, *C. scorpioides*, and *C. vaginatis* in Spain, Dalmatia, and Austria.

Mycosphaerella ariadna (Sacc.) Lind. On leaves of *C. emerus* in Italy.

Peronospora coronillae Güm. Downy mildew on leaves of *C. scorpioides* and *C. varia* in Dalmatia and Silesia.

Puccinia coronillae Wor. Leaf rust on *C. cappadocica* in Russia.

Ramularia coronillae Bres. Small pale-brown leaf spots on *C. varia* in Yugoslavia and Austria.

Septoria emeri Sacc. White leaf spots with dull-brown margins on *C. emerus* in Italy.

Uromyces anthyllidis (Grev.) Schroet. See Anthyllis.

CORREA. Australian shrubs.

Phyllosticta correae McAlp. Elongate brown leaf spots on *C. speciosa* in Australia.

Puccinia correae McAlp. Brown rust pustules on leaves of *C. lawrenciana* in Tasmania.

CORYDALIS. CAPNOIDES Ag. Hardy erect or prostrate herbaceous perennials.

Entyloma corydalis De By. Smut sori in white, then brown, leaf spots on *C. cava* and *C. solida* in Russia, France, Denmark, and Germany.

Entyloma corydalis-luteae Vogl. Smut sori in circular yellow-brown spots on leaves and stems of *C. lutea* in Switzerland and Italy.

Melampsora magnusiana G. Wagn. See Populus.

Peronospora bulbocapni Beck. Downy mildew on leaves of *C. ambigua*, *C. cava*, *C. fabacea*, and *C. laxa* in Japan and Europe.

Urocystis corydalis Niessl. Smut sori in brown circular leaf spots on *C. cava* and *C. solida* in Bohemia.

CORYLUS. HAZELNUT. FILBERT. Shrubs grown for nuts and foliage.

Ascochyta coryli Sacc. and Speg. Subcircular to irregular white leaf spots on *C. avellana* in Italy.

Bacterium sp. A bacterial disease of *Corylus* is reported as serious in Oregon. It causes a blighting of buds in new shoots, a brown speckling of the leaves, a girdling and dying-back of smaller branches, and formation of large cankers on larger branches and trunks.

Bacterium coryli Breda. This bacterium is reported as the cause of a canker disease of *C. avellana* and *C. colurna* in central Europe.

Cryptosporrella anomala Pk. Sacc. The disease due to this fungus, known as "filbert blight," is serious in the eastern part of the United States on *C. avellana*, and an effort is being made to prevent its spread to the Pacific coast. The fungus may be on *C. americana*. Sunken cankers of varying size and shape occur on branches and trunks, girdling the stems and soon killing outright many plants, particularly those of the European varieties.

Labrella coryli (Desm. and Rob.) Sacc. Irregular rufous leaf spots on *C. avellana* in Great Britain and France.

Mamiania coryli (Batsch.) Ces. and De N. On leaves of *C. avellana*, *C. heterophylla*, and *C. rostrata* in Japan, Denmark, and Austria.

Nematosporea coryli Pegl. Causes malformation of nuts of *Corylus* sp. in Italy.

Phyllosticta coryliaria (Sacc.) Tass. On leaves of *C. avellana* in Italy.

Phytophthora syringae Kleb. See *Syringa*.

Pucciniastrum coryli Kom. Ochraceous rust pustules on leaves of *C. heterophylla* and *C. rostrata* in Russia, Manchuria, and Japan.

Septoria avellanae B. and Br. On leaves of *C. avellana* in Great Britain and Italy.

CORYNOCARPUS. Ornamental evergreen trees.

Gloeosporium intermedium Sacc. See *Citrus*.

Phyllosticta corynocarpi Alm. and Cam. On leaves of *C. laurigata* in Portugal.

Septoria corynocarpi Thum. Circular to elliptical white leaf spots on *C. laurigata* in Portugal.

CORYPHA. See *Palmae*.**COSMOS.** Annual or perennial herbs grown in flower gardens.

Entyloma holwayi Syd. Smut sori in circular to irregular yellow-brown, then deep-brown, leaf spots on *C. sulphureus* in Mexico.

Uromyces bidentis Lagerh. Brown rust pustules on leaves of *C. coudatus* and *bidens* spp. in Porto Rico, Jamaica, and northern South America.

COSTUS. Perennial thick-rooted tropical herbs.

Ceroospora costina Syd. Leaf spot on *C. speciosus* in the Philippines.

Dactylaria costi Saw. Leaf spot on *C. speciosus* in Japan.

Puccinia costi P. Henn. Syd. Powdery brown rust pustules on indefinite sunken yellow leaf spots on *C. pumilus* in Brazil.

Uredo costina Syd. Leaf rust on *C. speciosus* in the Philippines.

COTONEASTER. Shrubs grown for their ornamental red or pink fruits and for the foliage.

Aecidium cunninghamianum Bardi. Leaf rust on *C. baccifer* in India.

Coleopuccinia sinensis Pat. Brown rust pustules on leaves of *Cotoneaster* sp. and *Aucubacter* sp. in China.

Entomosporium mespili (DC.) Sacc. Brown leaf spots on *C. nigra*, *Eriobotrya japonica*, *Mespilus germanica*, and *Pyrus sinensis* in Europe and Argentina. Probably not distinct from *E. maculatum* Lev.

Fusicladium pyracanthae (Oth.) Rostr. On fruit and peduncles of *C. pyracantha* in Denmark and Russia. Probably the same as *F. pirinum*.

Gymnosporangium mespili (DC.) Kern. See *Juniperus*.

Phyllosticta cotoneastri Allesch. Small subcircular, often confluent, brown, then gray, leaf spots with red-brown margins on *Cotoneaster* sp. in Germany.

COTYLEDON. Succulent herbs or shrubs grown mostly for their oddity.

Aecidium umbilici Trotter. Leaf rust on *C. umbilicus* in Portugal.

Ascochyta cotyledonis H. Zimm. Large gray leaf spots with purple-black margins on *C. gibbiflora* in Austria.

Microsphaera umbilici Kom. Powdery mildew on leaves of *C. semenovii* in Turkestan.

Septoria dalmatica Jaap. Circular gray leaf spots on *C. chierantha* in Dalmatia.

Septoria zimmermanni-hugonis Bub. Circular to elliptical brown leaf spots with gray margins on *C. desmetiana*, *C. gibbiflora*, and *C. pachyphytum* in Bohemia.

CRACCA. See *Tephrosia*.**CRASSULA.** Flesh-leaved greenhouse shrubs and herbs grown for the grotesque appearance of some species.

Aphelenchus olesistis Ritz. Bos. See *Begonia*.

Puccinia exanthematica MeO. Leaf rust on *C. cordata* and *C. spathulata* in the Union of South Africa.

CRATAEGUS. Hawthorn. Trees grown for their flowers and decorative fruit.

Aecidium patulum Syd. Rust on leaf blades, petioles, and fruit of *C. coccinea* in India.

Ascochyta crataegi Fekl. On leaves of *C. oxyacantha* in Russia, Italy, and Germany.

Ascochyta crataegicola Allesch. On leaves of *C. oxyacantha* in Russia and Germany.

Ascochyta crus-galli P. Brun. Circular to subcircular white leaf spots with brown margins on *C. crus-galli* in France.

Ascochyta misera Oud. On leaves of *C. monogyna* in Holland.

Ceroospora crataegi Sacc. and C. Mass. Small angular yellow leaf spots on *C. oxyacantha* in Italy.

Coryneum foliicolum Fekl. Ochraceous leaf spots on *C. oxyacantha*, *Quercus* sp., *Rubus discolor*, and *R. fruticosus* in Argentina, Russia, Italy, and Germany. Reported from Indiana.

Exoascus crataegi (Sad.) Sacc. Causes a crumpling and red spotting of the leaves and sometimes a deformation of twigs of *C. monogyna*, *C. oxyacantha*, and *C. sanguinea* in Europe.

Fusicladium crataegi Aderh. (*Venturia crataegi* Aderh.) Black spots on fruit of *C. monogyna* and *C. oxyacantha* in Denmark and Germany.

Gymnosporangium mespili (D. C.) Kern. See *Juniperus*.

Gymnosporangium orientale Syd. Aelia on fruit and branches of *C. ararella* and *C. insignia* in Russia, Greece, Thessaly, Asia Minor, and Persia.

Monilia crataegi Died. Effuse brownish leaf spots on *C. oxyacantha* in Germany.

Myosphaerella crataegi (Fekl.) Oud. On leaves of *C. monogyna* in Great Britain, Denmark, and Germany.

Myosphaerella crataegicola (Fekl.) Bond. and Tranz. On leaves of *C. monogyna* and *C. sanguinea* in Russia.

Myosphaerella oxyacanthae Jaap. On leaves of *C. oxyacantha* in Switzerland, Austria, and Germany.

Phloeospora oxyacanthae (Kunze. and Schm.) Walb. Crowded yellowish leaf spots on *C. oxyacantha* in Russia, Great Britain, Italy, and Germany. "Hawthorn leaf scorch."

Phyllosticta casinalbensis D. Sacc. White leaf spots on *C. azarolus* in Italy.

CRATAEGUS—Continued.

- Phyllosticta michailorskensis* Elenn. and Ohl. Leaf spots on *C. sanguinea* in Russia.
Phyllosticta monogyna Allesch. On leaves of *C. monogyna* and *C. oxyacantha* in Russia.
Phyllosticta phaea Sacc. On leaves of *C. oxyacantha* in Moravia.
Phytophthora syringae Kleb. See *Syringa*.
Septoria crataegicola Bond. and Tranz. On leaves of *Crataegus* sp. in Russia.
Septoria crataegophila Ranaj. On *C. monogyna* in Yugoslavia.
Stereum purpureum Pers. See *Prunus*.
Trichoseptoria fructigena Maubl. See *Cydonia*.
Uncinula prunastri (DC.) Sacc. See *Prunus*.

CRATAEVA. Tropical trees and shrubs.

- Aecidium crataeae* Syd. Rust pustules on large, circular to subcircular, sunken, yellow-brown leaf spots on *C. religiosa* in India.
Fusidium crataeae B. and Br. On leaves of *C. razburghii* in Ceylon.
Napadiadium crataeae Syd. On leaves of *C. religiosa* in India.

CREPIS. Annual, biennial, or perennial herbs, some species grown in flower gardens.

- Bremia ovata* Saw. Downy mildew on leaves of *C. japonica* in Japan and Formosa.
Entyloma calendulae (Oud.) De By. See *Calendula*.
Entyloma crepidicola Trot. and var. *crepidis-rubrae* Jaap. A smut forming galls on the fibrous roots and rhizomes of *C. bulbosa* and *C. rubra* in Italy and Dalmatia. The variety has been described as a new species under the name *Tolyposporium crepidis-rubri* (Jaap) Cif.
Phragmidium eximia Bub. Leaf rust on *C. grandiflora* in Yugoslavia.
Phyllosticta crepidis-paludosae Petr. Subcircular to irregular, ochre-brown leaf spots on *C. paludosa* in Austria.
Phyllosticta eximia Bub. Angular dark-brown to black leaf spots on *C. tiscidula* in Yugoslavia.
Protomyces crepidicola Buren. On leaves of *C. biennis* in central Europe.
Protomyces crepidis (Jaap.) Sacc. and Trott. Small circular dirty yellow-gray then yellow-brown pustules on *C. aurea*, *C. biennis*, *C. incarnata*, and *C. montana* in Switzerland and Austria.
Protomyces crepidis-paludosae Buren. On leaves of *C. paludosa* in Switzerland.
Protomyces inouei P. Henn. Effuse, elongate sori in peduncles of *Crepis* sp. in Japan.
Puccinia alpestris Syd. Yellow and brown rust pustules on leaves of *C. alpestris* in Switzerland and Austria.
Puccinia aschersoniana P. Henn. Brown rust pustules on leaves of *C. rupestris* in Arabia.
Puccinia barkhousiae-rhoeadifolia Bub. Yellow and dark brown to black rust pustules on leaves of *C. foetida* and *C. rhoeadifolia* in Bohemia and Spain.
Puccinia crepidicola Syd. Brown powdery rust pustules on leaves of *Crepis* spp. in Europe and Asia Minor.
Puccinia crepidis Schroet. Leaf rust on *Crepis* spp. in Europe.
Puccinia crepidis-aureae Syd. Leaf rust on *C. aurea* in Switzerland and Austria.
Puccinia crepidis-blattarioides Kasl. Leaf rust on *C. albidula*, *C. alpestris*, *C. blattarioides*, *C. tectus*, and *C. virens* in Switzerland and Spain.
Puccinia crepidis-grandiflorae Hasl. Rust on leaves of *C. bellidifolia*, *C. grandiflora*, *C. nicotensis*, and *C. tectus* in Switzerland.
Puccinia crepidis-japonicae Diet. Powdery, chestnut-brown rust pustules on leaves of *C. japonica* in Japan.
Puccinia crepidis-leontodontoides R. Maire. Powdery, cinnamon-brown to dark-brown rust pustules on leaves of *C. leontodontoides* in Corsica.
Puccinia crepidis-pygmaeae Gaill. Dark-brown rust pustules on leaf blades and petioles of *C. pygmaea* in France and Austria.
Puccinia crepidis-sibiricae Lindr. Brown powdery rust pustules on leaves of *C. sibirica* in Siberia, Russia, and Finland.
Puccinia crucheti Hasl. Rust on leaves of *C. succisaefolia* in Switzerland.
Puccinia intybi (Juel.) Syd. Leaf rust on *C. praemorsa* in Great Britain, Sweden, Estonia, Denmark, and Germany.
Puccinia major Diet. Cinnamon to dark-brown rust pustules on leaves of *C. conyzifolia*, *C. grandiflora*, and *C. paludosa* in Europe.
Puccinia praecox Bub. Powdery yellow and brown rust pustules on leaves of *C. biennis*, *C. foetida*, and *C. rubra* in Russia, Yugoslavia, Switzerland, Bohemia, Denmark, and Austria.
Puccinia scaliana Syd. Leaf rust on *C. biennis* and *C. bursifolia* in Spain and Sicily.
Puccinia silvatica Schroet. See *Taraxacum*.
Ramularia eximia Bub. Angular to irregular yellow leaf spots on *C. grandiflora* and *C. tiscidula* in Yugoslavia.
Septoria crepidis Vestergr. Irregular, often confluent, leaf spots on *C. biennis* and *C. tectorum* in Russia, Estonia, and Sweden.
Stagonospora crepidis Höll. Circular, dull-brown leaf spots with black margins on *C. biennis* in Hungary.
Taphrina rhaetica Volk. On leaves of *C. blattarioides* in Switzerland.
Uredo crepidis-integrae Lindr. Leaf rust on *C. integra*, *C. japonica*, *C. keiskeana*, and *C. platyphylla* in Japan and Australia.
Uredo crepidis-japonicae Lindr. Leaf rust on *C. japonica* in Ceylon and Australia.

CRINUM. Large, showy, flowering bulbs.

- Aecidium crini* Kalchbr. Leaf rust on *C. longifolium* (*C. capense*) and *C. cooperi* in the Union of South Africa.
Aecidium mangaranga P. Henn. Rust on large, effuse, yellow leaf spots on *Crinum* sp. in the Union of South Africa.
Gloeosporium crini Sacc. Leaf anthracnose of *Crinum* sp. in Italy.
Mycosphaerella crini Siem. On leaves of *Crinum* sp. in the Caucasus.
Phyllosticta crini Sacc. Subcircular yellow leaf spots on *Crinum* sp. in Italy.
Phyllosticta crinicola Siem. On leaves of *Crinum* sp. in southern Russia.
Septoria crini Tassi. On leaves of *C. asiaticum* in Italy.

CROCUS. Low, spring and autumn flowering bulbs.

- Bacillus croci* Miz. This bacterium causes a serious rot of *Crocus* spp. in Japan, attacking bulbs, roots, sheaths, leaves, and flower stalks. *Narcissus* sp., *Hyacinthus* sp., and *Allium* spp. have been infected by inoculation. The lower leaves become rotten, turn yellow above, and are readily detached from the corm, which is in turn soft-rotted.
Perisporium crocophila (Mont.) Sacc. Brown spots appear on corms of *C. sativus* in France, increasing in size, and finally bring about rot of the corm. A black fungus layer covers the rotted areas.
Sclerotinia tulparum (Wakk.) Rehm. See *Tulipa*.
Uromyces croci Pass. Elongate, powdery brown rust pustules on both leaf surfaces of *C. biflorus* and *C. susianus* in Russia, Italy, France, and Austria.

CROTALARIA. Rattlebox. Sunn-hemp. Annual leguminous herbs and shrubs, some species grown as green-manure crops.

Aecidium crotalariae P. Henn. Leaf rust on *C. engleri*, *C. grandibracteata*, and *Crotalaria* sp. in Uganda and west Africa.

Aecidium crotalariae P. Henn. Rust on dull-brown leaf spots on *Crotalaria* sp. in Brazil.

Aecidium dielsii P. Henn. Ochraceous rust pustules on leaves of *Crotalaria* sp. in Tanganyika.

Cercospora crotalariae Sacc. Leaf spots on *C. striata* in Ceylon.

Colletotrichum crotalariae Petch. Anthracnose on *C. striata* in Ceylon.

Mycosphaerella crotalariae Petch. Pale-brown zoned leaf spots on *C. striata* in Ceylon.

Phyllosticta crotalariae Sacc. On leaves of *C. striata* and *Crotalaria* sp. in Uganda and Ceylon.

Porla hypobrunnea Petch. See Hevea.

Uredo crotalariae Diet. Ochraceous rust pustules on leaves of *C. incana* and *C. vitellina* in Brazil.

Uredo crotalariae P. Henn. Leaf rust on *C. lachnoclada* in central Africa.

Uredo harmsiana P. Henn. Leaf rust on *C. lanceolata* in the Union of South Africa.

Uredo thesiae Neger. Brown rust pustules on leaves of *C. anagyroides* in Colombia.

Uromyces decoratus Syd. Powdery brown to black rust sori on yellow leaf spots on *C. albida*, *C. ferruginea*, *C. juncea*, and *C. vitellina* in India, Ceylon, and Costa Rica.

Uropyxis crotalariae Arth. Brown rust pustules on leaves of *C. maypurensis* and *Crotalaria* sp. in Guatemala.

CROTON. Shrubs and trees, a few species of economic importance.

Cercospora crotonophila Speg. On leaves of *Croton* sp. in Paraguay.

Cercospora manaensis P. Henn. On leaves of *Croton* sp. in Brazil.

Cercospora tigilii P. Henn. Circular, black leaf spots on *C. tigilium* in the Philippines.

Cercospora crotonis P. Henn. Leaf spots, often marginal, causing curling on *Croton* sp. in Brazil.

Gloeosporium crotonis Delacr. Anthracnose of *Croton* sp. in France.

Phakopsora crotonis (Cke.) Arth. Leaf rust on *C. cossypifolius* and *C. hirtus* in Trinidad.

Phyllachora crotonis (Cke.) Sacc. "Tar spot" on leaves of *C. arborescens*, *C. chamaedryfolius*, *C. floribundus*, and *C. silvaticus* in South America and the Union of South Africa.

Phyllachora globispora Speg. Black stromata on leaves of *Croton* sp. in Argentina.

Phyllachora fragiae (B. and C.) Sacc. Circular black stromata on leaves of *C. arborescens*, *C. chamaedryfolium*, *C. flavens*, and *C. lucidus* in Brazil, Porto Rico, French Guiana, Argentina, and Brazil.

Phyllosticta portoricensis Young. Circular yellowish leaf spots on *C. lucidus* in Porto Rico.

Schroeteriaster argentinensis (Speg.) Syd. Rust on leaf blades, petioles, and stems of *C. hirtus* in Argentina.

Schroeteriaster mexicanus (Arth.) Syd. Cinnamon-brown rust pustules on leaves of *C. calvescens* in Mexico.

Schroeteriaster stratosus (Cke.) Syd. Powdery brown rust pustules on leaves of *C. oxypetalus* and *C. sylvaticus* in the Union of South Africa and the Congo.

Septoria crotonis Bres. On leaves of *C. macrostachya* and *Croton* sp. in Denmark and Abyssinia.

Uredo crotonicola P. Henn. Leaf rust on *C. glandulosus* and *Croton* sp. in Brazil and Argentina.

Uredo crotonis P. Henn. Leaf rust on *Croton* sp. in Brazil.

CRUCIANELLA. Crosswort. Hardy rock plants.

Puccinia crucianellae Desm. Brown rust pustules on leaves and stems of *C. angustifolia*, *C. herbacea*, and *C. maritima* in Tripoli, Palestine, Portugal, Spain, Italy, and France.

Puccinia monopora Lindr. Rust on leaves and stems of *C. (Asperula) glomerata* in Greece.

Puccinia syriaca Syd. Powdery dark-brown rust pustules on leaves and stems of *C. macrostachya* and *C. syriaca* in Syria.

Uredo mediterraneae Lindr. Rust on stems of *C. maritima* in France and Portugal.

CRYPTOGRAMMA. ROCK BRAKE. Hardy subalpine ferns.

► **Hyalopora cryptogrammes** Diet. Brown rust pustules on leaves of *C. japonica* in Japan.

CRYPTOMERIA. Ornamental coniferous trees.

Fusarium blasticola Rostr. See Pinus.

Helicobasidium mompa Tanak. See Morus.

Valsa cryptomeriae Kitajima. The "blister" disease of seedlings of *C. japonica* is characterized by blackish-brown or reddish-brown areas on the stems which become sunken and cracked. These cankers cause hypertrophy and bending of diseased seedlings. The disease occurs in Japan.

CRYPTOSTEGIA. RUBBER VINE. Tropical climbers.

Ramularia cryptostegiae Pim. On leaves and fallen seeds of *Cryptostegia* sp. in Ireland.

CRYPTOSTEMMA. Composite herbs.

Ascochyta cryptostemmatis McAlp. Irregular, deep brown leaf spots on *C. calendulaceum* in Australia.

Septoria perforans McAlp. Circular, then confluent, pale gray-brown leaf spots on *C. calendulaceum* in Australia.

CUCUMIS. MELON. CUCUMBER. Herbaceous vines cultivated for their fruit.

Ascochyta melonis Poteb. On fruit, stems, and leaf blades and petioles of *C. melo* and *C. sativus* in Russia.

Corticium koleroga (Cke.) v. Hoeh. See Coffea.

Corynespora melonis (Cke.) Sacc. (*Cercospora melonis* Cke.). Small pale-green translucent spots on leaves and young fruits, becoming gray, on *C. melo* and *C. sativus* in Great Britain, Sweden, Russia, Denmark, and Germany.

Helminthosporium cucumerinum Jarb. Pale-yellow leaf spots up to 1.5 cm. in diameter, with definite margins, on *C. sativus* in south Russia. Infected plants are so weakened that death finally ensues.

Hypochnus cucumeris Frank. This fungus attacks its hosts at the ground level, rotting the stems and forming a thin, grayish film of hyphae. Diseased plants turn yellow and collapse. The hosts are *Amorpha phyllis*, *konja*, *Arachis hypogaea*, *Cinnamomum camphora*, *Citrus edulis*, *C. vulgaris*, *Cucumis melo*, *C. sativus*, *Daucus carota*, *Morus alba*, *Phaseolus mungo*, *P. vulgaris*, *Sesamum indicum*, *Sisya maz*, *Trifolium pratense*, *Vicia faba*, and *Vigna stensis* in Japan, Great Britain, Denmark, and Germany.

Puccinia cucumeris P. Henn. Yellow and black rust pustules on leaves of *C. ficifolius* in Abyssinia.

Rhizoctonia sp. See Vigna.

Scolecotrichum melophthorum Prill. and Delacr. Sunken brown spots on leaves and young stems, causing a rot of *C. melo*, *C. sativus*, and *Cucurbita vulgaris* in Great Britain, Italy, France, Belgium, Holland, and Russia.

Ustilago cucumeris Graff. A doubtful species of root smut on *C. sativus* in Great Britain.

CUCURBITA. PUMPKIN. SQUASH. GOURD. Cucurbitaceous vines.

Cercospora cucurbiticola P. Henn. Circular brown leaf spots on *Cucurbita* sp. in Brazil.

Scolecotrichum melophthorum Prill. and Delacr. See Cucumis.

- CUDRANIA.** Woody foliage plants used for hedges.
Hymenopsis cudraniae Mass. On leaves of *C. javanensis* in Queensland, India, and the Philippines.
Melasma cudraniae (Mass.) v. Hoeh. On *C. javanensis* in the Philippines.
Physopella sinense Syd. Leaf rust on *Cudrania* sp. in China.
Rhytisma hypoxanthum B. and Br. Black, irregular stromata on leaves of *C. javanensis* in Queensland.
Uredo cudraniae Petch. Leaf rust on *C. javanensis* in Ceylon.
- CUMMINGIA.** Tender summer-blooming bulbs.
Phyllosticta santiaguina Speg. Linear, red leaf spots on *C. campanulata* in Chile.
- CUNILA.** STONE MINT. Low-tufted hardy perennials.
Puccinia cunilae Diet. Brown rust pustules on leaves of *C. angustifolia* in Brazil.
Puccinia fuscata Arth. and Holw. Leaf rust on *C. leucantha* in Guatemala.
- CUNNINGHAMIA.** BELIS AG. CHINA FIR. Coniferous trees cultivated for their handsome foliage.
Mycosphaerella cunninghamiae Wor. On leaves of *C. sinensis* in Russia.
- CUNONIA.** Trees and shrubs.
Cercospora capensis (Thuem.) Sacc. Subcircular, depressed, black leaf spots on *C. capensis* in the Union of South Africa.
UPHEA. **PARSONSIA** Ag. Tropical and subtropical herbs and shrubs.
Didymariopsis cuphaeicola Speg. Circular, then irregular, leaf spots on *C. nesostemum* in Argentina.
Puccinia cuphaeae Holw. Dark-brown rust pustules on leaves of *C. acquipetala*, *C. cyanea*, *C. hookeriana*, *C. nitidula*, *C. procumbens*, and *C. squamuligera* in Mexico and Guatemala.
Puccinia jaliscensis Holw. Brown leaf rust on *C. cyanea*, *C. hookeriana*, *C. nitidula*, and *C. squamuligera* in Mexico.
Septoria cuphaeae Tassi. Brown leaf spots with wide rosy margins on *C. ilavea* in Italy.
Uredo cuphaeae P. Henn. Brown rust pustules on confluent, violet-brown leaf spots on *C. micrantha*, *C. parsonsia*, and *C. serphyllifolia* in Porto Rico, Colombia, and Brazil.
- CYPRINUS.** CYPRINUS. Ornamental evergreens and timber trees.
Gymnosporangium cunninghamianum Barel. Brown, slightly raised areas on smaller branches of *C. torulosa* in India. The aecial stage of this rust occurs on leaves of *Pyrus variolosa*.
- CURATELLA.** Small trees or scandent herbs.
Phyllosticta curatellae P. Henn. Circular red-brown to dull-brown leaf spots on *C. americana* in Peru.
- CURCULIGO.** Stemless herbs of palmlike habit.
Puccinia curcugonis Raeb. Circular to irregular dark-brown spots, in which appear brown powdery rust pustules, on both leaf surfaces of *C. latifolia*, *C. orchaioides* and *C. recurvata* in Java, India, and the Philippines.
- CURCUMA.** Turmeric. Tropical gingerlike plants.
Taphrina maculans Buit. Numerous dirty-yellow to bay, often confluent, spots on both leaf surfaces of *C. amada*, *C. angustifolia*, *C. longa*, *Hedychium* sp., *Zingiber casumunar*, *Z. mioga*, and *Z. zerumbet* in Japan and India.
- CYANTHUS.** Creeping or ascending woody herbs much like *Tradescantia*.
Uredo dactyloides Syd. Powdery brown rust pustules on leaves of *C. tuberosa* and *C. zeylanica* in India, Ceylon, and the Philippines.
Uromyces commelinae Cke. See Commelina.
- CYATHEA.** TREE FERN.
Griggsia cyathea Stev. Numerous irregular slightly elevated black spots on leaves of *C. arborea* in Porto Rico.
Pachypatella alsophilae (Rac.) Theiss. and Syd. On leaves of *C. caudata* and *Alsophila contaminans* in Java and the Philippines.
Phyllachora hieronymi P. Henn. Shiny black stromata on leaves of *C. dregii* in Nyassa and the Union of South Africa.
- CYBELE.** See Stenocarpus.
- CYCAS.** FERN PALM. SAGO PALM. Cycads.
Cladosporium cycadis Marc. Small circular yellow leaf spots with dark margins and with a greenish mold forming beneath, on *C. revoluta* in Italy.
Dendrophoma clypeata D. Sacc. Irregular, brown leaf spots on *C. revoluta* in Italy.
Hendersonia togniniana Pöhl. On leaves of *C. revoluta* in Italy.
Pestalotzia cycadis Allesch. On leaves of *C. revoluta* in Monaco.
Phyllosticta cycadina Pass. Brown leaf spots on *C. revoluta* in Italy.
Septoria montemartini Pöhl. On leaves and petioles of *C. revoluta* in Italy.
- CYCLAMEN.** Herbaceous plants with flat tubers or corms, grown under glass for their flowers.
Phyllosticta cyclaminella Bub. On leaves of *C. neapolitanum* in Yugoslavia.
Phyllosticta cyclaminis Brun. Circular brown spots finally destroying plants of *C. europaeum* and *C. persicum* in France and Italy.
Septoria cyclaminis Dur. and Mont. Large irregular smoky leaf spots with rufous margins on *C. europaeum*, *C. hederifolium*, and *C. repandum* in Algeria, Dalmatia, and Italy.
- CYCLANTHERA.** Annual and perennial herbs.
Cercospora cucubitina Speg. Small, circular to irregular, then confluent, white leaf spots on *Cyclanthera* in Brazil.
- CYDISTA.** Ornamental vines cultivated for their flowers. (See also Bignonia).
Puccinia adenocalymnatis (P. Henn.) Arth. Leaf rust on *C. aequinoctialis* in Cuba and Trinidad.
Puccinia aequinoctialis Holw. Cinnamon-brown rust pustules on circular, yellow leaf spots on *C. aequinoctialis* in Cuba.
Puccinia cuticulosa (E. and E.) Arth. Brown leaf rust on *C. aequinoctialis* in Porto Rico, Cuba, Brazil, and Nicaragua.
- CYDONIA.** QUINCE. Fruit trees.
Cercospora cydoniae Rangel. On leaves of *C. oblonga* (*C. vulgaris*) in Brazil.
Cercospora tomenticola (Thuem.) Sacc. On leaves of *C. vulgaris* in Brazil and Italy.
Eroasacus bullatus (B. and Br.) Fekl. See *Pyrus*.
Gymnosporangium koreaense (P. Henn.) Jacks. See *Juniperus*.
Gymnosporangium mespilii (DC.) Kern. See *Juniperus*.
Gymnosporangium photiniae (P. Henn.) Syd. See *Juniperus*.
Gymnosporangium spiniferum Syd. Rust on leaves of *C. vulgaris* in Japan.
Helicobasidium mompa Tan. See *Morus*.
Mycosphaerella pomacearum Sacc. See *Malus*.
Orularia necans (Pass.) Sacc. Large spots causing premature leaf fall of *C. vulgaris* and *Mespilus germanica* in Italy and France.

CYDONIA—Continued.

- Phylosticta cydoniae** (Desm.) Sacc. Circular to irregular brown leaf spots on *Chaenomelea japonica* (*Cydonia japonica*) and *C. vulgaris* in Italy, France, and Denmark.
- Phylosticta cydoniaeicola** Allesch. Large irregular leaf spots on *C. (Chaenomelea) japonica* in Italy and Germany.
- Phylosticta cydoniicola** P. Henn. Circular, then confluent, dull-brown leaf spots on *C. vulgaris* in Brazil.
- Phylosticta velata** Bub. Circular to ellipsoid, often confluent, yellow-brown to red-brown leaf spots on *C. vulgaris* in Hungary.
- Ramularia tenuior** Fautr. and Brun. On leaves of *C. vulgaris* in France.
- Sclerotinia linhartiana** Prill. and Delacr. Brown leaf spots, twig die-back, and brown rot of fruit of *C. vulgaris* in Europe.
- Sclerotinia mespill** Woron. See Mespillus.
- Septoria cydoniae** Fekl. On leaves of *Chaenomelea japonica* (*Cydonia japonica*) and *C. vulgaris* in Italy, Austria, and Germany.
- Septoria cydonicola** Thuem. Pale-gray, irregular leaf spots on *C. vulgaris* in Italy and Malta.
- Trichoseptoria fructigena** Maubl. Causes a brown rot of fruit of *C. vulgaris*, *Crataegus* sp., and *Malus* (apple) in France and Germany.

CYMBIDIUM. See Orchidaceae.**CYMOPOGON**. CITRONELLA GRASS. LEMONGRASS. Oil-producing grasses.

- Cerebella nardi** Butl. Black fungus areas in ovaries of *C. nardus* in India.
- Ophioidictis sclerotica** (Pat.) P. Henn. Black sclerotia in inflorescences of *C. nardus* and *C. schoenanthus* in India and Indo-China.
- Phyllachora andropogonis** Karst. and Har. See Andropogon.
- Puccinia cesatii** Schroet. See Andropogon.
- Puccinia cymopogonis** Mass. Leaf rust on *C. citratus* in Uganda.
- Puccinia nakanishikii** Diet. See Andropogon.
- Uredo cymopogonis-polyneuri** Petch. Leaf rust on *C. polyneuros* in Ceylon.
- Uromyces schoenanthi** Syd. Yellow to brown rust pustules on *C. schoenanthus* in India.
- Ustilago bengalensis** Syd. and Butl. Linear dark-brown smut sori in ovaries of *C. pendulus* in India.
- Ustilago mildbraedii** Syd. Powdery black smut masses replacing the ovaries of *C. schoenanthus* in tropical Africa.
- Ustilago nardi** Syd. Dark-brown elongate smut sori in ovaries of *C. nardus* in India.
- Ustilago schoenanthi** Syd. and Butl. Smut sori in inflorescences of *C. schoenanthus* in India.
- Ustilago spumioidea** B. and Br. Elongate smut sori in inflorescences of *C. martii*, *C. nardus*, and *C. venustus* in Ceylon.

CYNANCHUM. MOSQUITO TRAP. Herbaceous or semiwoody twiners.

- Cercospora beltyneckii** (West.) Sacc. On leaves of *C. acutum* and *C. vincetoxicum* in Italy, Belgium, China, and Spain.
- Cercospora mirae** Syd. On leaves of *C. caudatum* in Japan.
- Cercospora punctiformis** Sacc. and Roum. Yellow leaf spots on *C. acutum* in Russia and Algeria.
- Cronartium flaccidum** (Alb. and Schw.) Wint. See Paeonia.
- Melampsora cynanchi** Thuem. Leaf rust on *C. sibiricum* in Siberia.
- Myosphaerella abscens** Rabh. On leaves of *C. vincetoxicum* in Europe.
- Phylosticta asclepiadearum** West. and var. **minor** R. Rosai. Gray-white leaf spots on *C. vincetoxicum* and *Hoya carnosus* in Italy and Victoria.
- Phylosticta atro-maculans** Speg. Large pale-yellow areas on leaves of *C. vincetoxicum* in Italy.
- Phylosticta cynanchi** Brun. Circular to angular white leaf spots with brown margins on *C. acutum* in France.
- Phylosticta viridi-tingens** Ferr. Small subcircular leaf spots on *C. vincetoxicum* in Italy.
- Phylostictiella vincetoxici** (Sacc.) Tass. Sinuous white leaf spots with red margins on *C. vincetoxicum* in Denmark and Italy.
- Puccinia cynanchi** Lagh. Leaf rust on *C. parviflorum* and *Cynanchum* sp. in Martinique and Venezuela.
- Septoria asclepiadea** Sacc. On leaves of *C. vincetoxicum* and *Marsdenia erecta* in Italy, France, and Denmark.
- Septoria cynanchica** Sacc. On stems of *C. vincetoxicum* in Belgium.
- Septoria maculosa** Lévl. Black marginal leaf spots on *C. erectum* in France.
- Septoria rajkoffii** Bub. On leaves of *C. acutum* in Bulgaria.
- Septoria vincetoxici** (Schub.) Awd. On leaves of *C. nigrum* and *C. vincetoxicum* in Russia and Denmark.

CYNAEA. Artichoke. Cardoon. Perennial herbs.

- Ascochyta cynarac** Maffei. Circular ochraceous to gray-white leaf spots on *C. scolymus* in Italy.
- Phylosticta cynarac** West. Subcircular zoned ashen-brown leaf spots on *C. scolymus* in Argentina, Italy, and Belgium.
- Ramularia cynarac** Sacc. Circular to irregular, gray blotches on leaves of *C. scolymus* and *Silybum marianum* in Argentina, French North Africa, Denmark, Russia, Italy, France, and Portugal. The leaves are browned and destroyed by the fungus, and heads fail to form.
- Septoria cardunculi** Pass. On leaves of *C. cardunculus* in Italy.

CYNODON. CAPRIOLA Ag. BERMUDA GRASS. Lawn and pasture grasses.

- Balanisa** sp. On *C. dactylon* in the Union of South Africa.
- Cerebella cynodontis** Syd. Black, corrugated fungus masses in ovaries of *C. dactylon*, *Panicum distachyum*, and *P. prostratum* in Ceylon, India, the Philippines, and the Union of South Africa.
- Phyllachora cynodontis** (Sacc.) Niessl. (*Phyalospora cynodontis* Del.) Black stromata on leaves of *C. dactylon* and *C. incompletus* in Japan, China, the Philippines, India, French North Africa, Egypt, Union of South Africa, Madeira, Yugoslavia, Bulgaria, Italy, Spain, France, Austria, and Germany.
- Puccinia cynodontis** Desm. Rust on leaves, sheaths, and culms of *C. dactylon* in Porto Rico, Guatemala, Ceylon, China, Japan, India, Turkestan, Persia, Australia, Algeria, Asia Minor, Russia, Spain, Portugal, France, Yugoslavia, Bulgaria, Italy, Austria, and Germany. Also reported from California and Florida.
- Puccinia varians** Diet. Powdery black rust pustules on leaves of *C. dactylon* in Japan.
- Septoria cynodontis** Fekl. On leaves of *C. dactylon* in Italy and Germany.
- Ustilago cynodontis** P. Henn. Black powdery smut masses destroying the inflorescences of *C. dactylon* and *C. glabratus* in Japan, China, Australia, Uganda, French North Africa, Abyssinia, Bulgaria, Portugal, Spain, Italy, and India.
- Ustilago dregeana** Tul. Black smut sori deforming peduncles and inflorescences of *C. dactylon* in the Union of South Africa.
- Ustilago paraguayensis** Speg. Smut in culms of *C. dactylon* in France, Brazil, and Paraguay.

CYNOSURUS. CRESTED DOG-TAIL. Ornamental lawn and pasture grasses.

Phyllosticta cynosuri Gz. Frag. Leaf spots on *C. echinatus* in Spain.

Uromyces phyllachoroides P. Henn. Brown to black rust sori on leaves of *C. cristatus*, *C. echinatus*, and *C. elegans* in French North Africa.

CYPERUS. FLAT SEDGE. Papyrus. Aquatic or moist-land plants.

Asecochyta papyricola Tass. On leaves of *C. papyrus* in Italy.

Cintractia cyperi-polystachyi P. Henn. Smut in peduncles of *C. polystachyus* in the Philippines.

Cintractia javanica Rach. Black powdery smut sori in inflorescences of *Cyperus* sp. in Java.

Cintractia peribebuensis Speg. Smut in peduncles of *C. distans* in India and Ceylon.

Cintractia tangensis P. Henn. Black powdery smut masses in leaf axils of *Cyperus* sp. in Tanganyika.

Cintractia togoensis P. Henn. Smut on *Cyperus* sp. in central Africa.

Helminthosporium cyperi Sacc. On leaves of *C. dichostachys* in Abyssinia.

Kawakamia cyperi (Miy. and Ideta.) Miy. Downy mildew causing browning of leaves and stems of *C. tetigitiformis*, one of the matting sedges in Japan. The disease has also been found in Texas on imported material.

Phyllachora cyperi Rehm. Black stromata on leaves and culms of *C. giganteus* and *C. longus* in Porto Rico, Spain, and Portugal. Also reported from Wisconsin.

Puccinia abrepta Kern. Leaf rust on *C. ferax* in Costa Rica.

Puccinia antioquiensis Mayor. Leaf rust on *C. diffusus* in Panama, Trinidad, and Colombia.

Puccinia condusa Thum. Black rust pustules on leaves of *C. longus* in Portugal.

Puccinia Juncelli Diet. Leaf rust on *C. serotinus* in Japan.

Puccinia philippinensis Syd. Brown leaf rust on *C. compressus*, *C. rotundus*, and *C. polystachyus* in Japan and the Philippines.

Puccinia subcoronata P. Henn. Ochraceous to dark-brown rust pustules on leaves of *C. malaccensis* in Brazil and Japan.

Schinzia aschersoniana Magn. (*S. cypericola* [Magn.] De T.) Attacking roots of *C. flavescentis* in Algeria.

Uredo cyperi P. Henn. Rust on leaves and stems of *Cyperus* sp. in Abyssinia.

Uredo cypericola P. Henn. Leaf rust on *C. capitatus*, *C. distans*, *C. rotundus*, and *C. tuberosus* in India and tropical Africa.

Uredo philippinensis Syd. Cinnamon-brown rust pustules on leaves of *C. polystachyus* and *Cyperus* sp. in China and the Philippines.

CYPHOMANDRA. TREE TOMATO. Trees or shrubs, one species cultivated for its fruit.

Asecochyta cyphomandrae Petch. Subcircular black, then gray, leaf spots on *C. betacea* in Ceylon.

Cronartium uleanum Syd. Leaf rust on *Cyphomandra* sp. in Brazil.

CYPRIPEDIUM. LADY'S-SLIPPER. MOCCASIN FLOWER. Hardy terrestrial orchids. See Orchidaceae.

CYRTOPodium. See Orchidaceae.

CYSTOPTERIS. (*Filix*) BLADDER FERN.

Aphelenchus olesistis Ritz. Bos. See Begonia.

CYTISUS. BROOM. Shrubs cultivated for their profuse yellow or purple flowers.

Asecochyta cytisi Lib. On *C. laburnum* (*Laburnum vulgare*) in Switzerland.

Asecochyta laburni Sacc. On branches of *C. laburnum* (*Laburnum vulgare*) in France and Italy. *A. laburni* Kab. and Bub. is described on *C. laburnum* from Bohemia.

Ceratophorum setosum Kirchner. Small brown, then dark-brown to black, spots enlarging rapidly and often confluent to form large, irregular, often concentric, ringed areas on leaves of *Cytisus* sp., *Lupinus cruihshankstii*, *L. mutabilis*, *L. polyphyllus*, and other species in Japan and Europe.

Cucurbitaria laburni Pers. On twigs of *Cytisus* sp. in Europe.

Heterosporium laburni Oud. Brown leaf spots on *C. laburnum* in Denmark and Holland.

Leptosphaeria corrugans Rehm. Subcircular, oblong, or irregular yellow leaf spots with red-brown margins on *C. alpinus* (*Laburnum alpinum*) in Austria.

Marsonia carnea Vesteg. Large circular leaf spots on *C. laburnum* (*Laburnum vulgare*) in Europe.

Mycosphaerella laburni (Pass.) Lind. On leaves of *C. laburnum* and *C. ramentaceus* in Italy and Dalmatia.

Peronospora cytisi Rostr. Downy mildew, causing brown leaf spots and destroying young plants of *C. alpinus* (*Laburnum alpinum*) and *C. laburnum* in Europe.

Phyllosticta coniothyrioides Sacc. Subcircular ashen leaf spots on *C. laburnum* and *C. nigricans* in France and Denmark.

Phyllosticta cytisella Sacc. Angular white leaf spots on *C. nigricans* in Italy.

Phyllosticta cytisi Desm. Subcircular dull-brown leaf spots on *C. laburnum* in France, Great Britain, Italy, Belgium, and Austria.

Phyllosticta cytisorum Pass. On leaves of *C. laburnum* in France.

Phyllosticta laburni Oud. Irregular gray-white leaf spots on *C. laburnum* in Holland.

Phyllosticta laburnicola Sacc. On leaves of *C. laburnum* in Italy.

Septoria cytisi Desm. Numerous small subcircular white leaf spots with brown margins on *C. alpinus*, *C. capitatus*, *C. hirsutus*, *C. laburnum*, *C. nigricans*, and *Genista tinctoria* in Italy, France, Switzerland, Denmark, and Austria.

Septoria laburni Pass. On leaves of *C. sessilifolius* in Italy.

Uromyces fulgens (Haas.) Bub. Brown rust pustules on leaves and branches of *C. austriacus* and *C. hirsutus* in Yugoslavia, Bulgaria, Italy, Austria, and Germany.

Uromyces genistae-tinctoriae (Pers.) Wint. See Caragana.

Uromyces pallidus Niessl. Yellow-brown rust pustules on leaves of *C. capitatus*, *C. hirsutus*, and *C. prostratus* in Europe.

DACTYLIS. ORCHARD GRASS. Forage grasses.

Apiospora lloydii (Crouan) Sacc. See Holcus.

Aplanobacter rathayi Sm. A viscid lemon-yellow slime envelops the upper leaves, stems, and inflorescences of *D. glomerata* in Denmark and Austria, causing a dwarfing and premature drying out of infected plants.

Cladochytrium graminis Büsg. See Festuca.

Dilophia graminis (Fekl.) Sacc. See Triticum.

Leptosphaeria nigrans (Desm.) Ces. and De N. See Aira.

Mycosphaerella recutita (Fr.) Johans. See Aira.

Ovularia puchella (Ces.) Sacc. and var. *lilii-italici* Sacc. Red leaf spots on *D. glomerata* and *Lolium italicum* in France and Italy.

Phyllachora bromi Fekl. See Bromus.

Phyllosticta dactyloides Gz. Frag. On leaves of *D. glomerata* in Spain.

Puccinia dactylidina Bub. Powdery brown to black rust pustules on leaves of *D. glomerata* in India, Hungary, and Bohemia.

Septoria culmifida Lind. See Phleum.

Sphaerodothis dactylidis (Delacr.) Theiss. and Syd. Black stromata on leaves of *D. glomerata* in France and Spain.

DACTYLIS—Continued.

Uromyces dactylidis Oth. Yellow and brown to black rust pustules on leaves of *D. glomerata* and *Ranunculus* spp. in Europe and New Zealand. Reported from Virginia.

Ustilago dactylidis Maire. Smut on *D. glomerata* in French North Africa.

DAEDALACANTHUS. Tropical shrubs with blue or rose-colored flowers.

Isariopsis acanthacearum Cke. Leaf spot of *D. nervosus* and *Eranthemum* sp. in Great Britain.

Puccinia polystegia Syd. Brown rust pustules on brown sunken leaf spots on *D. nervosus* in China.

DAEMONOROPS. See *Palmae*.**DAHLIA**. Stout perennial herbs with fleshy roots and profuse autumn bloom.

Aecidium dahliae Syd. Leaf rust on *D. variabilis* in Mexico.

Bacillus dahliae Hori. and Bakis. Bacterial rot of *Dahlia* sp. in Japan.

Cercospora sp. A leaf spot of *D. variabilis* in Argentina.

Cercospora grandissima Rangel. Leaf spot on *D. variabilis* in Brazil.

Coleosporium dahliae Arth. A rust producing yellow powdery pustules on the undersides of leaves of *D. variabilis* in Mexico.

Cytospora dahliae Peyl. On stems of *Dahlia* sp. in Bohemia.

Entyloma calendulae dahliae Sternon. Minute spots appear on the leaves at flowering time. These spots are at first few in number, 5 to 7 millimeters in diameter, limited by the principal veins and visible only on the upper leaf surfaces. The spots soon enlarge, becoming confluent, brown in color with dark-brown margins, and finally involve the entire leaf area as well as the petioles. The original dead areas fall out, giving a shot-hole appearance to the leaves. The loss of all or most of the leaf area reduces or even entirely prevents the development of flowers. On *Dahlia* (cult.) in Belgium. See, also, *Calendula*.

Entyloma dahliae Syd. Smut sori in pale circular to elliptical leaf spots on *D. variabilis* in the Union of South Africa.

Ophiobolus georginae (Fckl.) Sacc. On stems of *Dahlia* sp. in Germany.

Phoma fuscata Sacc. On stems of *Dahlia* sp. in Germany.

Phyllosticta dahliae Brun. Produces large brown spots on leaves of cultivated varieties of *D. variabilis* in France, Switzerland, Italy, Austria, and Germany.

Sporotrichum dahliae Thuem. On stems of *D. variabilis* in Siberia.

Urophlyctis sp. Causes a leaf spot of *Dahlia* [cult.] in Kenya Colony.

Verticillium dahliae Kleb. Causes a wilt disease of the *Dahlia* in Germany. The fungus is said to differ from *V. albo-atrum*, the cause of a serious wilt disease of the potato and other economic plants. The symptoms of the *Dahlia* disease are typical of wilt diseases, namely, a preliminary drooping of the leaves which finally involves the entire plant. The disease can be carried in the roots.

DALBERGIA. ROSEWOOD. Tropical trees and shrubs.

Catacauma dalbergicola (P. Henn.) Theiss. and Syd. Black stromata on leaves of *D. acanthophylla*, *D. armata*, *D. ferruginea*, and *D. variabilis* in Brazil, the Union of South Africa, and the Philippines.

Endodothella natalensis Doidge. On leaves of *D. armata* in the Union of South Africa.

Fomes caryophylli (Rac.) Bres. See *Cinnamomum*.

Fomes lamaeensis Murr. See *Hevea*.

Phyllachora dalbergiae Niessl. Black stromata on leaves of *D. ferruginea*, *D. lactuca*, *D. sissoo*, and *D. variabilis* in India, the Philippines, and Brazil.

Phyllachora lactea Theiss. and Syd. Black stromata on red-brown leaf spots on *D. lactea* in Tanganyika.

Phyllachora perforans (Rehm.) Sacc. and Syd. Tar spot on leaves of *D. acanthophylla* in Brazil.

Phyllachora spissa Syd. Circular shiny black stromata on brown leaf spots on *D. sissoo* in India.

Phyllosticta dalbergiae Syd. Circular to irregular, often confluent, leaf spots with pale purple margins on *Dalbergia* sp. in Brazil.

Phyllosticta dalbergicola Syd. Circular pale-brown leaf spots on *D. variabilis* in Brazil.

Phyllosticta nivea Syd. Subcircular brown leaf spots on *Dalbergia* sp. in Brazil.

Puccinia periodica Racib. Leaf rust on *Dalbergia* sp. and *Derris* sp. in Java.

Sphaerophragmium dalbergiae Diet. Yellow to brown rust pustules on leaves of *D. amerinum* and *D. armata* in Cuba and Natal.

Uredo dalbergiae P. Henn. Ochraceous rust sori on *D. variabilis* in Brazil.

Uredo mararyensis P. Henn. Leaf rust on *Dalbergia* sp. in Brazil.

Uredo nidulans Syd. Powdery ochraceous rust pustules on *D. foliolosa* in Bolivia.

Uredo sissoo Syd. and Butl. Powdery, yellowish rust pustules on leaves of *D. sissoo* in India and Ceylon.

Uromyces achrous Syd. Leaf rust on *D. sissoo* in India.

A serious canker disease due to an as yet undetermined fungus, attacks *D. latifolia* in Java.

BALEA. PAROSELA Ag. Leguminous herbs and small shrubs, grown for their various-colored flowers.

Puccinia daeae Diet. and Holw. Cinnamon-brown to black rust pustules on leaves of *D. citradora* in Mexico.

DALECHAMPIA. Climbing tropical shrubs, cultivated for their ornamental bracts.

Aecidium cornu-cervi P. Henn. A rust, forming woody galls up to 10 centimeters in diameter on branches of *Dalechampia* sp. in Brazil.

Aecidium dalechampicola P. Henn. Leaf rust on *Dalechampia* sp. in Brazil.

Caecoma theissenii Syd. Leaf rust on small subcircular spots on *Dalechampia* sp. in Brazil.

Didymella sphaerelloides Sacc. and Syd. Indefinite pale brown sunken leaf spots on *D. ficifolia* in Brazil.

DAMMARA. See *Agathis*.**DAPHNE**. Ornamental woody plants with handsome foliage and sweet-scented flowers.

Colletotrichum daphnes-ponticae Siem. Anthracnose on *D. pontica* in Caucasasia.

Dothidea mezerei (Fr.) Theiss. and Syd. On stems of *D. alpina* in Europe.

Gloeosporium mezerei Cke. Anthracnose on leaves on *D. mezereum* in Great Britain and Australasia.

Marssonia daphne (Rob. and Desm.) Magn. Irregular greenish, then dull-brown, leaf spots on *D. mezereum* in France, Italy, Switzerland, Denmark, Austria, and Holland.

Marssonia obtusata Maubl. Small raised ochraceous, then dull dark-brown leaf spots on *D. laureola* in France.

Mycosphaerella gnidii Maire. On leaves of *D. gnidius* in Algeria.

Mycosphaerella laureolae Desm. Subcircular, yellowish leaf spots on *D. laureola* in France, Switzerland and Germany.

Phyllohendersonia daphnes (Pass.) Tass. On leaves of *D. indica* in France.

Phyllosticta carniolica Voss. On leaves of *D. blagayana* in Austria.

Phyllosticta daphnes ponticae Siem. Leaf spot on *D. pontica* in Caucasasia.

DAPHNE—Continued.

Phyllosticta laureolae Desm. Subcircular to irregular leaf spots on *D. laureola* in Denmark, France, and Portugal.

Septoria daphnes Desm. Irregular indefinite greenish leaf spots on *D. mezereum* in France and Germany.

Uredo daphnicola Diet. Brown leaf rust on *Daphne* sp. in Hongkong.

DAPHNIPHYLLUM. Broadleaved evergreen, hardy or semitropical shrubs, and small trees.

Leptothyrium yoshinagai P. Henn. On leaves of *D. glaucescens* in Japan.

Mycosphaerella daphniphylli Syd. and Hara. Circular to irregular spots on leaves of *D. glaucescens* in Japan.

DATISCA. Tall perennial herbs.

Coleosporium datiscæ Tranzsch. Golden rust pustules on leaves of *D. cannabina* in Kashmir and Caucasia.

Phyllosticta datiscæ Syd. Irregular, then confluent, often marginal, dull brown leaf spots on *D. cannabina* in Russia.

DATURA. ANGEL'S-TRUMPET. JIMSON WEED. Rank-growing annual or perennial herbs or shrubs mostly weeds.

Ascochyta daturæ Sacc. Subcircular gray-white leaf spots on *D. arborea* and *D. stramonium* in Russia, Denmark, Italy, and Portugal.

Septoria daturæ Speg. Definite circular gray-white leaf spots on *D. inermis* and *D. stramonium* in Argentina and Italy.

DAUCUS. CARROT. Annual and biennial herbs.

Aecidium carotinum Bub. Leaf rust on *D. carota* in Bohemia.

Aphelenchus modestus J. M. Attacks the roots of *D. carota* in Europe.

Cercospora pastinacæ Karst. See *Pastinaca*.

Erysiphe taurica Lév. See *Althæa*.

Helicobasidium mompa Tan. See *Morus*.

Hypochnus cucumeris Frank. See *Cucumis*.

Leptosphaeria rostrupii Lind. Gray-brown cankers on the roots, which work up into the stems of *D. carota* in Sweden and Denmark.

Mycosphaerella sagedioides (Wint.) Lind. On stems of *Daucus* sp. and *Dipsacus* sp. in Switzerland.

Phyllachora pastinacæ Rostr. (*Septoria pastinacæ* West.) Leaf spot and stem canker of *D. carota* in Great Britain and Denmark.

Protomyces macrosporus Ung. See *Coriandrum*.

Ramularia pastinacæ Bub. See *Pastinaca*.

Septoria carotæ P. Nag. On leaves of *D. carota* in Russia.

Septoria daucina Brun. Subcircular to irregular, then confluent, brown leaf spots on *D. carota* in France.

DEERINGIA. Climbing herbs or subshrubs.

Aecidium deeringiae Cke. and Mass. Leaf rust on *D. celosioides* in Australia.

Puccinia calosperma Syd. and Burt. Rust on leaf blades, petioles, stems, and flowers of *D. celosioides* in India.

Uromyces deeringiae Syd. Brown leaf rust on *D. indica* in Java, Japan, and the Philippines.

DELPHINIUM. LARKSPUR. Hardy herbs cultivated for their flowers.

Cercospora delphinii Thuem. Dull-yellow subcircular leaf spots on *D. elatum* in Siberia. Also reported from Colorado.

Coleosporium martianoffianum Syd. Golden-yellow rust pustules on leaves of *D. intermedium* in Siberia.

Phyllosticta ajacis Thuem. Irregular, often marginal, gray-white leaf spots on *D. ajacis* in Austria.

Rhizoctonia destruens Tassi. See *Solanum*.

Ramularia delphinii Jaap. Leaf spot on *D. elatum* in Switzerland.

Septoria staphysagriae Wint. Circular to irregular definite brown leaf spots on *D. staphysagria* in Portugal and Italy.

DENDROBIUM. See *Orchidaceæ*.**DENDROCALAMUS**. See *Bambuseæ*.**DENDROPANAX**. Tropical trees and shrubs.

Mycosphaerella didymopanicis Miles. Light-brown to gray, circular, sometimes confluent, leaf spots with narrow orange-brown margins on *D. arboreum* in Porto Rico.

Phyllosticta arallana Young. Leaf spot on *D. arboreum* in Porto Rico and Cuba.

DENTARIA. TOOTHWORT. Small, early-flowering herbs.

Phyllosticta dentariae Kab. and Bub. Greenish, then pale-brown to white, irregular leaf spots on *D. encephalylla* in Bohemia.

Peronospora dentariae macrophyllæ Gäum. Downy mildew on leaves of *D. macrophylla* in Japan.

Puccinia dentariae (Alb. and Schw.) Fekl. Leaf and stem rust on *Dentaria* spp. in Europe. Also reported from Oregon.

DERRIS. Tall, tropical trees or climbers.

Cercospora pumila Syd. On leaves of *Derris* sp. in the Philippines.

Dioreidium koordersi Wurtb. Leaf rust on *D. elliptica* in Java.

Dothidella derridis (P. Henn.) Theiss. On leaves of *Derris* sp. and *D. elliptica* in central Africa and the Philippines.

Fomes lignosus Klotzsch. See *Hevea*.

Haplophragmium derridis Syd. Brown rust pustules on leaves of *D. uliginosa* in Ceylon and central Africa.

Phyllachora affinis Theiss. and Syd. Black stromata on leaves of *D. philippinensis* in the Philippines.

Phyllachora derridis Syd. Black stromata on indefinite brown leaf spots on *D. glabrata* in Angola.

Phyllachora luzonensis P. Henn. See *Milletia*.

Phyllachora ramosi (P. Henn.) Theiss. and Syd. Numerous, black stromata on small, leaf spots on *Derris* sp. in the Philippines.

Phyllachora yapensis (P. Henn.) Syd. Black circular to irregular stromata on leaves of *D. diadelpa*, *D. elliptica*, and *D. philippinensis* in the Philippines, Java, and the island of Yap.

Phyllosticta derridis P. Henn. Circular or effuse, pale-brown, marginal leaf spots on *Derris* sp. in the Congo.

Placosphaeria derridis P. Henn. Circular leaf spots on *Derris* sp. in the Congo.

Placosphaeria merrillii P. Henn. Circular effused yellow or dull brown leaf spots on *Derris* sp. in the Philippines.

Puccinia periodica Racib. See *Dalbergia*.

Triphragmium pulchrum Racib. Brown rust pustules on leaves of *D. elliptica* in Java.

Uredo derridis P. Henn. Leaf rust on *D. bantamensis* in Java.

DESCHAMPSIA. See also *Aira*.

Tilletia schenckiana P. Henn. Smut sori deforming and destroying the ovaries of *D. antarctica* in Kerguelen Island.

DEUTZIA. Ornamental shrubs, grown for their showy, white or bluish flowers.

Aecidium deutziae Diet. Leaf rust on *D. gracilis*, *D. scabra*, *D. sieboldiana*, and *D. staminea* in Japan and India.

Ascochyta deutziae Bres. Irregular, dull-yellow leaf spots on *D. scabra* in Germany.

Cylindrosporum deutziae Syd. Indefinite confluent yellow-brown leaf spots on *D. scabra* in Japan.

Leptosphaeria dichroa Pass. On branches of *D. scabra* in Italy.

Mycosphaerella deutziae Syd. On leaves of *D. lemoinei* in Germany.

Phyllosticta deutzicola Petr. Large irregular brown leaf spots on *Deutzia* sp. in Bohemia and Austria.

Septoria phyllostictoides Sacc. Indefinite gray-white leaf spots on *D. scabra* in France.

DIANELLA. Tender perennial rhizomatous plants.

Uredo dianella Diet. Leaf rust on *D. ensifolia* and *D. nemorosa* in China, Hongkong, Ceylon, Java, and Japan.

DIANTHUS. PINK. CARNATION. Herbs cultivated for their showy flowers.

Ascochyta dianthi (A. and S.) Berk. Large, subcircular, brown patches on leaves of *D. caryophyllus*, *Lychnis* sp. and *Saponaria* sp. in Great Britain, Italy, Holland, and Germany.

Fusarium udum Butl. This fungus causes a wilt of seedlings of *Dianthus* sp., *Anacardium occidentale*, *Antirrhinum* sp., *Eriobotrya japonica*, *Eugenia jambos*, and *Grevillea robusta* in Uganda. See also *Cajanus*.

Gloeosporium dianthi Cke. Small circular purple confluent leaf spots with brown centers on *D. caryophyllus* in Great Britain.

Phomopsis caryophylli Grove. On calyces, peduncles, and stems of *D. caryophyllus* in Japan and Great Britain.

Phyllosticta dianthi West. Irregular gray-white blotches on leaves of *D. barbatus* in Belgium and France.

Phyllosticta dubia Sacc. On leaves of *D. chinensis* (*D. sinensis*) in Malaya.

Placosphaeria stellariae (Lib.) Sacc. See *Stellaria*.

Pseudodiscosia dianthi Host. and Laub. Large brown, then light gray spots on leaves, flower stalks and stems of *D. caryophyllus* in Germany.

Puccinia dianthi-japonici P. Henn. Brown rust pustules on circular yellow leaf spots on *D. japonicus* and *D. nipponicus* in Japan.

Puccinia fastidiosa Sacc. and De T. Powdery, ochraceous rust pustules on leaves and stems of *D. chinensis* in Siberia.

Ramularia dianthi Lind. Large leaf spots with brown margins on *D. carthusianorum* in Germany.

Septoria carthusianorum West. Irregular, dull-yellow spots on calyces of *D. carthusianorum* in Belgium.

Septoria caryophylli Scalia. Irregular elongate ochraceous spots with dull brown margins on leaves and stems of *D. caryophyllus* in Italy.

Septoria dianthicola Sacc. On leaves of *D. barbatus*, *D. caryophyllus*, and *Tunica prolifera* in Italy, Switzerland, and Portugal.

Septoria dianthophila Speg. and f. *hispanica* Gz. Frag. On leaves of *D. caryophyllus* and *D. (Tunica) prolifera* in Brazil and Spain.

Septoria sinarum Speg. Large subcircular white leaf blotches on *D. chinensis* in Italy and Great Britain.

Uredo dianthicola Har. Leaf rust on *D. caryophyllus* in France. (Doubtful species.)

Urocystis purpurea Haez. Purple masses of smut spores replacing the ovaries of *D. deltoides* and *D. (Tunica) prolifera* in Hungary.

Uromyces formosus Syd. Brown rust pustules on stems and leaves of *D. crinitus*, *D. libanotis*, and *D. macranthoides* in Persia.

DIAPENSIA. Small, compact, tufted, woody perennials.

Phyllosticta diapensiae Pat. Leaf spots on *D. lapponica* in China.

Physalospora diapensiae Rehm. On leaves of *D. lapponica* in Lapland.

Septoria diapensiae Karst. On leaves of *D. lapponica* in Lapland.

DIASIA. See *Melasmaeria*.

DICENTRA, BIKUKULIA. Ag. BLEEDINGHEART. Herbaceous perennials grown for their flowers

and cut foliage. Sometimes called *Dielytra* or *Dielytra*.

Ascochyta dicentrae Oud. On branches of *D. spectabilis* in Holland.

DICHAEA. See *Orchidaceae*.

DICHORISANDRA. Tropical perennial herbs, grown for their handsome flowers and foliage.

Colletotrichum dichorisandrae Rangel. Anthracnose on *D. thyrsiflora* in Brazil.

Uromyces dichorisandra P. Henn. Leaf rust on *Dichorisandra* sp. in Brazil.

DICHROA. Hydrangealike shrubs.

Endophyllum dichroae Rac. Rust causing hypertrophy of leaves of *D. cyanitis* in Java.

DICLIPTERA. Annual or perennial herbs or subshrubs with red or blue bracted flowers.

Puccinia diclipterae Syd. Brown rust on leaves of *D. longiflora* in Formosa.

Uromyces tweediana (Speg.) Arth. Yellow-brown to chestnut-brown rust pustules on leaves and stems of *D. maculata*, *D. squarrosa*, *D. tweediana*, and *Dicliptera* sp. in Mexico, Brazil, Argentina, Abyssinia, and India.

DICTAMNUS. GAS PLANT. Semiwoody plants.

Ascochyta nobilis Kab. and Bub. Circular to angular, often confluent, yellow-white or gray leaf spots with red margins on *D. albus* (*D. fraxinella*) in Russia and Bohemia.

Colletotrichum dictamni Höll. Circular to irregular dull-brown leaf spots on *D. albus* in Hungary.

Septoria dictamni Fekl. Discolored leaf patches on *D. albus* (*D. fraxinella*) in Russia and Italy.

DICTYOSPERMA. See *Palmae*.

DIEFFENBACHIA. TUFT ROOT. Large-leaved tropical aroids.

Gloeosporium thuenenii Sacc. See *Anthurium*.

Phyllosticta colocasiae v. Hoch. See *Colocasia*.

DIERAMA. South African cormous plants with large spikes of flowers.

Puccinia dieramae Syd. Leaf rust on *D. ensifolia* in the Union of South Africa.

DIERVILLA. BUSH HONEYSUCKLE. Ornamental deciduous shrubs with showy flowers. Certain Asiatic species sometimes placed under *Weigela*.

Ascochyta diervillae Kab. and Bub. Circular to irregular, then confluent, brown leaf spots on *D. lonicera* (*D. canadensis*) in Bohemia.

Ascochyta weigeliae Sacc. and Speg. Angular gray-white leaf spots on *D. florida* (*D. rosea*) in Italy.

Phyllosticta weigeliae Sacc. and Speg. Gray-white leaf spots on *D. florida* (*D. rosea*) in Denmark, Italy and France.

D. ERVILLA—Continued.

Phyllosticta weigelina Bub. and Kab. Leaf spots on *D. florida* in Bohemia.

Ramularia weigeliae Speg. On leaves of *D. florida* in Italy.

Septoria weigeliae Kab. and Bub. Leaf spots on *D. florida* in Bohemia.

DIGITALIS. FOXGLOVE. Hardy herbaceous perennials with long racemes of flowers. The leaves of some species used medicinally.

Ascochyta digitalis Fekl. Purplish leaf spots on *D. purpurea* in Portugal and Germany.

Gloeosporium digitalidis Rostk. Large dull-brown leaf spots on *D. purpurea* in Denmark.

Mycosphaerella mariae Sacc. and Bomm. On leaves and stems of *D. ferruginea* and *D. lutea* in Italy.

Peronospora digitalidis Gäum. Downy mildew on leaves of *D. ambigua*, *D. lutea*, and *D. purpurea* in central Europe.

Phyllosticta digitalis Bell. On leaves of *D. lutea* and *D. purpurea* in Italy and Alaska.

Ramularia variabilis Fekl. Irregular brown leaf spots starting on the basal leaves and progressing upward on *D. purpurea* in Italy, Denmark, Austria, and Germany. The medicinal value of the leaves is reduced or destroyed.

Septoria digitalis Pass. Irregular, dull-brown leaf spots on *D. ferruginea*, *D. grandiflora*, *D. lanata* and *D. lutea* in Russia, the Balkans, Italy, Great Britain, and Austria.

DILLENIA. Tall tropical trees.

Cercospora dilleniae Petch. Gray-brown leaf spots on *D. retusa* in Ceylon and India.

DINCHLOA. See Bambuseae.

DIOCLEA. Tender woody twiners.

Phyllachora diocleae P. Henn. Black stromata on circular to irregular brown leaf spots on *Dioclea* sp. in Brazil.

DIOSCOREA. YAM. AIR POTATO. Tropical climbers with edible roots.

Bagniopsis dioscoreae Wakef. Produces a reduction in length of the internodes and in size of the leaves, giving a witches'-broom effect on *D. prehensis* in Nigeria. The stromata are produced on stems and petioles, but not on the leaf blades which are, however, chlorotic.

Catacauma glaziovii (P. Henn.) Theiss. and Syd. Circular, shiny black stromata on leaves of *D. heptaneura*, *D. pachycarpa* and *D. polygonoides* in Brazil.

Cercospora brasiliensis Avena. Leaf spot on *Dioscorea* sp. in Brazil.

Cercospora carbonacea Miles. Angular, black leaf spots 1 to 1.5 centimeters in diameter on *D. alata* in Porto Rico and Cuba.

Cercospora pachyderma Syd. On leaves of *D. alata* and *D. esculenta* in the Philippines.

Cercospora ubi Rac. Circular leaf spots, yellow-brown above, gray-brown below, on *D. alata*, *D. daemona*, *D. esculenta*, and *D. glabra* in Java, China, Malaya, and the Philippines.

Cercospora contraria Syd. Gray-brown leaf spots on *Dioscorea* sp. in the Congo.

Colletotrichum dioscoreae Avena. Anthracnose on *Dioscorea* sp. in Brazil.

Cylindrosporium dioscoreae Miy. and I. Ito. Small, yellowish to brown spots on leaf blades, petioles and stems of *D. batatas* and *D. japonica* in Japan. Defoliation results.

Ellisiodothis rehmanniana Theiss. and Syd. Black stromata on stems of *D. esculenta* in the Philippines.

Gloeosporium pestis Mass. Anthracnose spots on leaves of *Dioscorea* sp. in Fiji and Russia.

Hemileia dioscoreae-aeuleatae Rac. Yellow leaf rust on *D. aculeata* and other species in Uganda and Java.

Laestadia perusta (B. and Br.) Sacc. Circular dull-brown leaf spots on *D. tomentosa* and *Dioscorea* sp. in Ceylon and India.

Mycosphaerella dioscoricola Syd. Leaf spot on *D. esculenta* in the Philippines.

Phyllachora ulei Wint. Shiny black circular stromata on leaves of *Dioscorea* sp. in Brazil.

Phyllosticta dioscoracearum Bacc. Subcircular gray leaf spots on *Dioscorea* sp. in India.

Phyllosticta dioscoreaeicola P. Brun. Irregular red-brown leaf spots on *D. batatas* in France.

Phyllosticta dioscoreae-daemona P. Henn. Circular pale-brown leaf spots with dark-brown margins on *D. daemona* in Brazil.

Phyllosticta graffiana Sacc. On leaves of *D. esculenta* and *D. pentaphylla* in the Philippines.

Puccinia valida Arth. Cinnamon-brown rust pustules on lower leaf surfaces of *D. convolvulacea* in Mexico.

Rostrupia dioscoreae (Kom.) Syd. Cinnamon-brown to black rust pustules on leaves of *D. quinqueloba* in Manchuria and Japan.

Septoria versicolor Pat. Ashen-white leaf spots with brown margins on *Dioscorea* sp. in Ecuador.

Uredo dioscoreae (B. and Br.) Petch. (*Uredo dioscoreae* P. Henn.) Leaf rust on *D. alata*, *D. bulbifera*, *D. esculenta*, *D. grandiflora*, *D. piperifolia* and *D. polygonoides* in Ceylon, the Philippines, Porto Rico, Cuba, Brazil, India, and Uganda.

Uredo dioscoreae-alatae Rac. Brown rust sori on leaf blades, petioles, and stems of *D. alata* and other species in Java and the Philippines.

Uredo dioscoreae-filiformis Rac. Rust pustules on dull-brown leaf spots on *D. filiformis* in Java.

Uredo dioscoreae-pentaphyllae Petch. Leaf rust on *D. pentaphylla* in Ceylon.

Uredo dioscoreae-quinquelobae P. Henn. Leaf rust on *D. quinqueloba* in Japan.

Uredo dioscoreae-sativa Syd. Brown rust pustules on leaves of *D. sativa* in India.

Urocystis dioscoreae Syd. Black, elongate, smut pustules on leaf blades, petioles and stems of *D. quinqueloba* and *D. tokoro* in Japan and Russia.

DIOSPYROS. PERSIMMON. Kaki. Ebony. Fruit and timber trees.

Aecidium atro-album P. Henn. Leaf rust on *Diospyros* sp. in the Union of South Africa and the Congo.

Aecidium atrocrustaceum Syd. Leaf rust on *D. discolor* in the Philippines.

Aecidium calosporum Juel. Rust on circular leaf spots on *Diospyros* sp. in Brazil.

Aecidium diospyri A. L. Sm. Leaf rust on *D. mespiliformis* in Angola.

Aecidium melaenum Syd. Leaf rust on *Diospyros* sp. in the Philippines.

Aecidium ramosii Syd. Leaf rust on *Diospyros* sp. in the Philippines.

Aecidium reyesii Syd. Leaf rust on *D. discolor* in the Philippines.

Aecidium rhytismoidem B. and Br. and var. *mabae* P. Henn. Rust on black leaf spots on *D. discolor*, *D. embryopteris*, *D. mespiliformis*, *D. ovalifolia*, *D. tomentosa*, and *Maba abyssinica* in Java, India, Ceylon, Abyssinia, and the Philippines.

Aecidium ulei P. Henn. Leaf rust on *Diospyros* sp. in Brazil.

Botrytis diospyri Brizi. A fruit rot of *D. kaki* in Italy, Russia, and Japan. The fungus also causes a leaf spot and die-back of twigs.

Colletotrichum kaki Maffei. Anthracnose on leaves of *D. kaki* in Italy.

Cylindrosporium kaki Syd. Circular dull-brown or gray leaf spots with black limiting lines on *D. kaki* in Japan.

Fusicladium diospyrae Hori. and Yosh. Circular, black spots with definite margins on leaves, fruit and buds of *D. kaki* in Japan, producing a curl of young leaves. On young shoots the spots are fusiform to oblong, dark-brown to black with sunken centers. Diseased twigs die-back and spotted fruit falls.

DIOSPYROS—Continued.

- Fusicladium kaki** Hori. and Yosh. On leaves of *D. kaki* in Japan.
- Fusicladium leverii** P. Magn. Circular brown leaf spots with black margins on *D. lotus* in Caucasia.
- Gloeosporium kaki** Ito. Small black spots, increasing in size and becoming sunken, on fruit of *D. kaki* in Japan. The spots may fuse and crack irregularly. Complete rot of the fruit ensues. Infections also occur on young shoots, showing typical blackish spots of elliptical shape, usually depressed. On stems of young plants all parts above the infected area die. Infected leaves fall, complete defoliation often resulting.
- Helicobasidium mompa** Tan. See Morus.
- Lophodermium diospyri** Pat. Long, black, superficial fruiting bodies on circular, often confluent, leaf spots on *D. ebenum* in Indo-China.
- Melasmia falcata** Syd. Black stromata on leaves of *Diospyros* sp. in Brazil.
- Myxosporium kaki** Hara. Dark-brown, more or less sunken areas, on stems, branches and rarely on fruit of *D. kaki* in Japan.
- Pestalozzia diospyri** Syd. Circular to irregular reddish-brown spots on leaves of *D. kaki* in China and Japan. Smaller brown spots, generally regular, also occur on the fruit.
- Plaocaccardinula diospyricola** P. Henn. Black flat fruiting bodies on leaves of *Diospyros* sp. in Brazil.
- Phyllosticta erythraea** Bacc. Brown leaf spots with black margins on *D. mespilifolius* in Abyssinia.
- Septoria diospyri** McAlp. On leaves of *D. cargillia* in New South Wales.
- Uredo peteloti** Pat. Brown rust pustules on circular leaf spots on *D. siamensis* in Indo-China.
- DIOTIS.** COTTONWEED. Perennial seaside plants.
- Puccinia diotidis** Pat. and Roum. Leaf rust on *D. candidissima* in France.
- DIPCADL.** Tender, bulbous, scape plants.
- Accidium dipcadi** Har. and Pat. Leaf rust on *D. udellensis* in the Congo.
- DIPHYSA.** Shrubs or trees.
- Calliospora diphsae** Arth. Powdery, black rust pustules on leaves and stems of *D. robinoides* and *D. suberosa* in Guatemala, Costa Rica, and Mexico.
- DIPLOGLOTTIS.** Australian trees.
- Uromyces diploglottidis** Cke. and Mass. Leaf rust on *D. cunninghamii* in Queensland.
- DIPLOSTEPHIUM.** Composites resembling *Aster*.
- Rhynchosoma biolleyana** Bomm. and Rouss. On leaves and stems of *D. costaricense* (*D. rupestre*) in Costa Rica.
- DIPLOTHEMIUM.** See Palmae.
- DIPSACUS.** TEASEL. Biennial or perennial herbs, used as fuller's teasels.
- Aecchyta dipsaci** Bub. Subcircular small ochraceous leaf spots with dull-brown margins on *D. pilosus* in Asia Minor.
- Mycosphaerella sagedioides** (Wint.) Lind. See Daucus.
- Peronospora violacea** Berk. Downy mildew on leaves of *Dipsacus pilosus*, *Knautia silvatica*, *Scabiosa columbaria*, and *Succisa pratensis* in Europe.
- Phyllosticta dipsaci** Br. and Fautr. Gray leaf spots on *D. pilosus* in France.
- Phyllosticta vandae** Namys. On leaves of *D. sylvestris* in Poland and Galicia.
- Ramularia silvestris** Sacc. On leaves of *D. fullonum* and *D. sylvestris* in Spain, Yugo-Slavia, Denmark, France, Bohemia, Belgium, and Germany.
- Septoria dipsaci** West. Small circular brown, then white leaf spots on *D. azureus* and *D. fullonum* in Belgium and Italy.
- Septoria fullonum** Sacc. Gray-white leaf spots on *D. fullonum* in Austria.
- Uredo involucrorum** Rabh. Leaf rust on *D. pilosus* in France, Belgium, and Switzerland.
- DISCHIDIA.** Evergreen trailing plants cultivated for their white flowers.
- Phyllachora dischidiae** Syd. Circular black shiny stromata on leaves of *D. rosea* in Chile and the Philippines.
- Uredo dischidiae** P. Henn. Yellow rust pustules on leaves of *Dischidia* sp. in New Guinea.
- DISPORUM.** FAIRY BELLS. Small perennial rhizomatous herbs.
- Accidium dispori** Diet. Leaf rust on *D. sessile* in Japan.
- DISSOTIS.** Bristly-hairy herbs or shrubs.
- Puccinia dissotidis** P. Henn. Yellow-brown rust sori on circular dull-brown leaf spots on *Dissotis* sp. in the Congo.
- Puccinosira dissotidis** Wakef. (*Accidium dissotidis* Cke.) Brown leaf rust on *D. incana* and *D. princeps* in Uganda, the Congo, and the Union of South Africa.
- Uredo dissotidis-longicaudae** P. Henn. Brown leaf rust on *D. longicaudata* in the Union of South Africa.
- DIURIS.** See Orchidaceae.
- DODONAEA.** HOP BUSH. Ornamental trees and shrubs.
- Uredo dodonaeae** Koord. Leaf rust on *D. viscosa* in Java.
- DOLICHOLUS.** See Rhynchosia.
- DOLICHOS.** HYACINTH BEAN. Tropical twiners.
- Cercospora wildemanii** Syd. Circular brown leaf spots with purple margins on *Dolichos* sp. in the Congo.
- Helminthosporium accedens** Syd. On leaves of *D. baumii* in the Union of South Africa.
- Helminthosporium dolichi** Syd. On leaves of *D. euryphyllus* in the Union of South Africa.
- Phyllachora dolichogena** (B. and Br.) Sacc. and var. *samoensis* Theiss. and Syd. Black stromata on leaves of *D. lablab* and *Dolichos* sp. in Ceylon and Samoa.
- Phyllosticta dolichi** Brun. Brown leaf spots with dark margins on *D. lablab* and *D. myoides* in Russia and France.
- Physopella concors** Arth. See Phaseolus.
- Puccinia dolichi** Arth. Powdery brown rust pustules on leaves of *D. reticulatus* in Cuba.
- Rhizoctonia** sp. See Vigna.
- Septoria lablabina** Sacc. Circular dull-brown leaf spots on *D. lablab* in tropical Africa and the Philippines.
- Septoria lablabis** P. Henn. Circular, then confluent, dull-brown spots on leaves of *D. lablab* in Tropical Africa and the Philippines.
- Uromyces dolichi** Cke. Brown rust pustules on leaves of *D. azillaris* and *D. gibbosus* in the Union of South Africa.
- Uromyces kisanuensis** P. Henn. Powdery dark-brown rust pustules on leaves of *Dolichos* sp. in the Congo.
- Vermicularia capsiei** Syd. See Capsium.
- Woronnetia dolichi** (Cke.) Syd. Orange galls on leaves of *D. lablab* and *Rhynchosia volubilis* in China and the Philippines.

- DOMBEYA. ASSONIA** Ag. Shrubs and small trees with large loose umbels of showy flowers.
Phyllachora dombeyae Syd. Black stromata on leaves of *D. rotundifolia* and *D. schimperiana* in the Union of South Africa.
- DONAX.** See *Arundo*.
- DOREMA.** Large perennial herbs yielding gum and resins.
Puccinia doremae Spechn. Leaf rust on *Dorema* sp. in Turkestan and Transcaspia.
- DORONICUM.** LEOPARD'S-BANE. Hardy herbs with yellow many-flowered heads.
Ascochyta daronici Allesch. Large ovoid subochraceous to ashen-brown leaf spots on *D. caucasicum* in Germany.
Coleosporium daronici Namysl. Yellow leaf rust on *D. austriacum* in Austria.
Mycosphaerella aronici (Fekl.) Volk. On leaves of *D. austriacum*, *D. caucasicum*, *Aronicum scorpioides*, and *Carduus defloratus* in north Africa, Switzerland, Italy, and Austria.
Phyllosticta austriaca Sacc. Brown leaf spots on *D. austriacum* in Italy.
Phyllosticta daronicella Maire. Black leaf spots on *D. thirkum* in Asia Minor.
Phyllosticta daronicigena Bub. On leaves of *D. cordatum* in Hungary.
Puccinia daronicella Syd. Brown rust pustules on leaves of *D. austriacum* and *D. cordatum* in Austria and Hungary.
Puccinia daronici Niessl. Black rust sori on leaves of *D. austriacum* and *D. macrophyllum* in Bulgaria and Austria.
- Ramularia daronici** (Sacc.) Grove. On leaves of *D. pardalianches* in Great Britain, Denmark, and France.
- Ramularia daronicum** Vogl. Subcircular to oblong, often marginal, yellow, then white, leaf spots on *D. chusium*, *D. cordatum*, and *D. scorpioides* in Italy and Spain.
- Ramularia filaris** Fres. See *Adenostyles*.
- Septoria czarnohorica** Nam. Circular to irregular and confluent brown leaf spots on *D. cordifolium* in France and Poland.
- DORSTENIA.** Tropical herbs and small shrubs.
Aecidium bertonii Speg. Leaf rust on *D. brasiliensis* in Paraguay.
Aecidium buchwaldii P. Henn. Rust on brown leaf spots on *Dorstenia* sp. in Tanganyika.
Aecidium dorsteniae-holstii P. Henn. Rust on circular dull-brown leaf spots on *D. holsti* in Tanganyika.
Puccinia dorsteniae Lagh. Leaf rust on *D. psilurus* in Angola.
Uredo consanguinea Syd. Brown leaf rust on *D. multiformis* in Brazil.
Uredo rubescens Arth. Powdery brown rust pustules on leaves of *D. contrajerva* and *D. houstoni* in Porto Rico, Guatemala, and Trinidad.
- DOVYALIS.** See *Aberia*.
- DRABA.** WHITLOW GRASS. Spring-flowering annuals.
Ascochyta drabae Oud. On leaves of *D. alpina* in Russia.
Peronospora norwegica Güm. Downy mildew on leaves of *D. hirta* in Norway.
Puccinia drabae Rud. Brown rust pustules on leaves, stems, and flowers of *Draba* spp. in Europe and Asia Minor. Reported from Utah and New York.
- DRACAENA.** (including *Cordylina*.) DRACENA. DRAGON TREE. Variegated-leaved shrubs.
Cercospora cordylines P. Henn. On leaves of *C. dracaenoides* in Argentina and Brazil.
Colletotrichum cordylines Pöhl. Anthracnose on leaves of *C. indivisa* in Italy.
Colletotrichum dracaena Allesch. On leaves of *D. latifolia* in Germany.
Colletotrichum dracaenicola Sacc. and Trott. Anthracnose on leaves of *D. fragrans* in Italy.
Gloeosporium polymorphum Trinch. Large dull-brown to ashen leaf spots on *D. fragrans* in Italy.
Gloeosporium thuenenii Sacc. See *Anthurium*.
Graphiola disticha (Ehrbg.) Lévl. Raised brown hard fruiting bodies on leaves of *D. draca* in India.
Lembosia orbicularis Pat. Elongate black fruiting bodies disfiguring the leaves of *Dracaena* sp. in Zanzibar.
Lembosia patouillardii Sacc. and Syd. On leaves of *Dracaena* sp. in east Africa.
Macrophoma bakeri Syd. Discolored areas on leaves of *Dracaena* sp. in Brazil.
Melanconium stictoides Sacc. and Paol. On leaves of *Dracaena* sp. in Malacca.
Meliola dracaenicola Pat. and Har. Superficial black fungus patches on leaves of *Dracaena* sp. in the Congo.
Meliola subdentata Pat. Circular black superficial fungus patches on leaves of *Dracaena* sp. in Indo-China.
Mycosphaerella dracaenae Tassi. On leaves of *D. reflexa* in Italy.
Myxosporium dracaenacolum B. and Br. Destroys leaves of *Dracaena* sp. in Great Britain.
Phyllachora nervisequa Wint. Black stromata on leaves of *D. cannifolia* in Queensland.
Phyllosticta cordylines Sacc. and Berl. On leaves of *D. australis* and *D. terminalis* in Queensland.
Phyllosticta dracaenae Griff. and Maubl. Large irregular leaf spots on *Dracaena* sp. in France, working injury to greenhouse plants.
Phyllosticta draconis Berk. Irregular pale-brown leaf spots with purple margins on *D. cooperi*, *D. draca*, and *D. terminalis* in Portugal, France, Great Britain, and Finland.
Physalospora amphididyma Syd. Subcircular white leaf spots with red-purple raised margins on *D. papahii* in Tanganyika.
Placasterella schweinfurthii (P. Henn.) Theiss. and Syd. On leaves of *D. ombeitis* in Abyssinia.
Rosellinia echinata Mass. See *Ficus*.
Ustilago dracaenae S. da Cam. Leaf smut on *D. draco* in Portugal.
Ustilago trabutiana Sacc. Black powdery masses of smut spores in ovaries of *D. draco* in Algeria.
- DRACOCEPHALUM. MOLDAVICA** Ag. DRAGONHEAD. Hardy herbaceous annuals or perennials.
Peronospora rossica Güm. Downy mildew on leaves of *D. thymiflorum* in Russia.
Septoria dracoccephali Thuem. On leaves of *D. peregrinum* in Siberia.
- DRIMYS.** Broad-leaved evergreen trees and shrubs.
Actinothyrium drimydii Speg. Circular pale-brown leaf spots on *D. winteri* in Chile.
Gloeosporium walteri McAlp. Anthracnose on leaves of *D. aromatica* in Australia.
Helminthosporium orbiculare Lévl. On leaves of *D. chilensis* and *D. winteri* in Chile.
Munkella drymidis (Lévl.) Speg. On leaves of *D. winteri* in Chile.
Mycosphaerella drymidis (Berk.) Sacc. Circular brown leaf spots on *D. winteri* in Chile and Brazil.
Phyllosticta drymidis Speg. Circular gray-white leaf spots with purple margins on *D. winteri* in Chile.
Phyllosticta winterii Speg. Circular brown leaf spots on *D. winteri* in Chile.
Septoria drimydicola Speg. Circular to irregular leaf spots on *Drimys* sp. in Brazil.
Septoria drymidos Mont. On leaves of *D. chilensis* in Chile.
Septoria winterii Speg. On leaves of *D. winteri* in Chile.

- DRYAS.** Dwarf, hardy, tufted evergreen shrubs.
Didymella dryadis Speg. Indefinite leaf spots on *D. octopetala* in Italy.
Guignardia rhythmoides (Berk.) Trav. On *D. octopetala* in Spain.
Mycosphaerella ootheca (Sacc.) On leaves of *D. octopetala* in Alaska.
Septoria semilunaris Johans. On peduncles of *D. octopetala* in Scandinavia.
Venturia tirolensis v. Hoeh. On leaves of *D. octopetala* in Austria.
- DRYOPTERIS.** WOOD FERNS. Wood ferns with much dissected leaves.
Hyalospora filicum Diet. See Asplenium.
Milesta columbensis Diet. Rust on fronds of *D. patens* and *Nephrolepis pendula* in Porto Rico and Colombia.
- Uredo gymnogammes** P. Henn. Rust on fronds of *Adiantum latifolium*, *D. mollis*, *D. poiteana*, *Pityrogramma calomelanos*, and *Tectaria martinicensis* in Porto Rico, Cuba, Jamaica, Trinidad, and Brazil.
- DURANTA.** SKY FLOWER. Tropical shrubs cultivated for their flowers.
Corticium salmonicolor B. and Br. See Citrus.
Phyllachora durantae Rehm. Black stromata on small yellowish to reddish leaf spots on *Duranta* sp. in Ecuador.
Phyllachora fuscicarpa Seaver. Black, irregular, slightly raised stromata on leaves of *D. repens* in Porto Rico and the Bahamas. This fungus causes a serious leaf fall.
- DURIO.** Durian. Tropical fruit trees.
Colletotrichum durionis Koord. Anthracnose on leaves of *D. zibethinus* in Java.
Corticium salmonicolor B. and Br. See Citrus.
Gloeosporium zibethinum Sacc. Anthracnose on leaves of *D. zibethinus* in Malaya.
Homostedia durionis Racib. Circular black stromata on leaves of *D. zibethinus* in Java.
Phoma durionis Petch. Attacks fruit of *D. zibethinus* in Ceylon.
Phyllachora macrospora A. Zimm. Circular black stromata on leaves of *D. zibethinus* in Java.
Phyllosticta durionis A. Zimm. Pale-brown leaf spots on *D. zibethinus* in Java.
Phyllosticta nephelii Delacr. Dull-brown indefinite leaf spots on *D. zibethinus* and *Nephelium lappaceum* in the Congo and France.
Placosphaeria durionis Syd. On leaves of *D. zibethinus* in the Philippines.
- ECHEVERIA.** Fleshy-leaved plants.
Endophyllum sempervivi (A. and S.) De B. See Sempervivum.
- ECHINOCACTUS.** A genus of the cactus family.
Stagonospora assans Pass. See Cereus.
- ECHINOCHLOA.** BARNYARD GRASS. A few species sometimes cultivated for grain or forage.
Claviceps balansoides A. Moell. Irregular black-brown sclerotia in spikes of *Echinochloa* sp. in Brazil.
- ECHINOPS.** GLOBE THISTLE. Coarse thistle-like plants with flowers in globose masses.
Puccinia echinopsis DC. Powdery brown to black rust pustules on leaves of *E. banaticus*, *E. cyaneus*, *E. ritro*, and *E. sphacrocephalus* in Europe.
Puccinia hellenica Trotter. Leaf rust on *E. microcephalus* in Greece.
Puccinia pulvinata Rabb. Brown powdery rust pustules on leaves of *E. amplexicaulis*, *E. chamaecephalus*, *E. cornigerum*, *E. echinatus*, *E. heldreichii*, *E. machrochaetus*, *E. spinosus*, and *E. viscosus* in India, Abyssinia, Uganda, Asia Minor, and French North Africa.
- ECHITES.** Tropical American twining shrubs.
Puccinia balansae Speg. Leaf rust on *E. fusiformis* in Argentina.
Septoria echitis Syd. Small, gray-white leaf spots with purple margins on *E. tweediana* in Argentina.
- ECHIMUM.** Coarse, rough herbs and shrubs with flowers in spikes.
Aecidium asperifolii Pers. Leaf rust on *E. italicum*, *E. lusitanicum*, and *E. rubrum* in Dalmatia, Russia, and Portugal.
Aecidium echii Thum. Leaf rust on *E. lusitanicum* in Portugal.
Cercospora echii Wint. Angular to irregular brown leaf spots on *E. tuberculatum* in Portugal.
Cercospora euchlora Maire. On leaves of *E. australe* in North Africa.
Ramularia anchusae Mass. See Anchusa.
Synchytrium echii Speg. Small galls on pale-brown leaf spots on *E. plantagineum* and *E. violaceum* in Argentina.
- EDGEWORTHIA.** PAPER TREE. Ornamental shrubs grown for their yellow flowers.
Helicobasidium mompia Tan. See Morus.
- EHRETIA.** Tender trees and shrubs.
Schroeteriaster ehretiae (Hir.) Syd. and Butl. Leaf rust on *E. acuminata* in Japan and Formosa.
Uredo ehretiae Barcl. Leaf rust on *E. acuminata*, *E. macrophylla*, and *E. serrata* in India and Japan.
- ELAEAGNUS.** RUSSIAN OLIVE. Shrubs and small trees grown chiefly for their handsome foliage and ornamental fruits.
Aecidium elaeagni Diet. Leaf rust on *E. glabra* and *E. pungens* in Japan.
Aecidium elaeagni-latifolia Petch. Ochraceous rust pustules on small, definite leaf spots on *E. latifolia* in Ceylon.
Aecidium elaeagni-umbellatae Diet. Rust on circular to irregular leaf spots on *E. macrophylla* and *E. umbellata* in Japan.
Aecidium minoense Syd. Leaf rust on *E. japonicus* in Japan.
Aecidium quintum Syd. Leaf rust on *E. umbellata* in Japan.
Ascochyta elaeagni Sacc. Ochraceous leaf spots on *E. argentea*, *E. multiflora* (*E. edulis*), and *E. gussonei* in Russia, Italy, and France.
Ochrospora nambuana (P. Henn.) Diet. Leaf rust on *E. multiflora* (*E. longipes*), *E. macrophylla*, and *E. umbellata* in Japan.
Puccinia achora Syd. Black rust sori on leaves of *E. macrophylla* in Japan.
Puccinia elaeagni Yosh. Leaf rust on *E. pungens* in Japan.
Septocylindrium olivascens Thum. Subcircular, olivaceous leaf spots on *E. angustifolia* and *E. rhannoides* in France and Denmark.
Septoria argyrea Sacc. Ochraceous leaf spots on *E. angustifolia* and *E. argentea* in Italy. Reported from North Dakota.
- ELAEIS.** African oil palm. See Palmae.
- ELAEODENDRON.** FALSE OLIVE. Tropical shrubs and small trees.
Septoria elaeodendri P. Henn. Circular dull-brown leaf spots on *E. xylocarpum* in Germany.
- ELETTARIA.** Cardamom. Perennial hothouse herbs.
Placostroma ellettariae (B. and Br.) Theiss. and Syd. Shiny black stromata on leaves of *E. floribunda* in Ceylon.
Schroeteriaster ellettariae Rac. Rust sori on narrow, long, gray or brown leaf spots on *Ellettaria* sp. in Java.

ELEUSINE. AFRICAN MILLET. Annual grasses, some grown as ornamentals.

Acrothecium lunatum Wakk. See Panicum.

Phyllachora eleusines Speg. Black, smooth, subcircular stromata on leaves of *E. coracana*. *E. indica*, and *E. tristachya* in Argentina, Brazil, Uganda, and the Congo.

Ustilago eleusines Kulkarni. Circular to elongate green, then chocolate-brown to black, smut sori in single grains or groups of grains of *E. coracana* in India, the sori rupture exposing the deep brown to black spore masses.

ELSOLTZIA. Herbs or shrubs cultivated for their spikes of blue or lilac flowers.

Coleosporium perillae Syd. See Perilla.

ELYMUS. WILD RYE. DUNE GRASS. Erect perennial grasses, sometimes cultivated as ornamentals or for forage.

Cercospora elymi Rostr. Oblong dull-brown leaf spots on *E. arenarius* in Denmark.

Ophiobolus cariceti (B. and Br.) Sacc. See Triticum.

Puccinia elymi-sibiricae S. Ito. Powdery, brown rust pustules on leaves of *E. sibiricus* in Japan.

Rostrupia elymi (West.) Lagh. Brown leaf rust on *E. arenarius*, *E. mollis*, and *E. sibiricus* in Japan and northern Europe.

Septoria elymi E. Rostr. On leaves of *E. arenarius* in Denmark.

Septoria elymicola Died. On leaves of *E. arenarius* in Europe.

Septoria elymi-europaei Jaap. Yellow, then brown, sunken leaf spots on *E. europaeus* in Switzerland.

Tilletia bornmülleri Magn. Brown smut sori in ovaries of *E. crinitus* in Asia Minor.

Tilletia controversa Kuehn. See Triticum.

Tilletia serbica Ranoj. Smut sori in spikelets of *E. crinitus* in Yugoslavia.

Tylenchus hordei Schoeyen. See Avena.

Uredo elymi-capitis-medusae Gz. Frag. Leaf rust on *E. caput-medusae* in Spain.

Ustilago phrygia Magn. Brown spore masses in spikes of *E. crinitus* in Asia Minor.

EMBOTHRUM. South American trees and shrubs.

Phyllosticta embotryi Speg. Indefinite dull-brown leaf spots on *E. coccineum* in Chile.

EMILIA. Annual or perennial herbs with orange or scarlet flower heads.

Aecidium emiliae Petch. Leaf rust on *E. sonchifolia* in Ceylon.

Aecidium formosanum Syd. Leaf rust on *E. sonchifolia* in Formosa.

Puccinia syndrellae P. Henn. Brown rust sori on lower-leaf surfaces of *E. sagittata*, *E. sonchifolia*, and *Synedrella nodiflora* in the West Indies and Trinidad.

Uredo emiliae-zeylanicae Petch. Leaf rust on *E. zeylanica* in Ceylon.

EMPETRUM. CROWBERRY. Low shrubs, sometimes grown for their evergreen foliage and attractive fruit.

Melasmia empetri Magn. Destroys young shoots of *E. nigrum* in Russia.

Metasphaeria empetri (Fr.) Sacc. On leaves of *E. nigrum* in Europe and Alaska.

Physalospora empetri Rostr. On leaves of *E. nigrum* in Denmark.

Rhytisma empetri White. Black smooth stromata on stems of *E. nigrum* in Great Britain, Italy, Denmark, and Sweden.

ENCELIA. Herbs or subshrubs.

Aecidium enceliae Diet. and Holw. Leaf rust on *E. canescens* in Peru.

Puccinia enceliae Diet. and Holw. Powdery dark-brown rust sori on leaves of *Simsia amplexicaulis* (*E. mexicana*) in Mexico.

Uredo enceliae-tomentosae Maire. Leaf rust on *E. tomentosa* in Chile.

ENDYMION. See Scilla.

ENKIANTHUS. Shrubs grown for their handsome flowers.

Aecidium enkianthi Diet. Leaf rust on *E. japonicus* in Japan.

ENTADA. Tropical shrubs.

Ravenelia entadae Lagh. and Diet. Brown rust pustules on leaves of *E. polystachya* in Panama and Guatemala.

Ravenelia schweinfurthii Syd. Brown leaf rust on *E. sudanica* in central Africa.

ENTEROLOBIUM. Tropical trees.

Phyllachora enterolobii Speg. Circular black stromata on leaves of *E. timbouva* in Brazil.

Ravenelia hassleri Speg. Leaf rust on *E. timbouva* in Paraguay.

Ravenelia oligotheles Speg. Dull-brown rust pustules on leaves of *E. timbouva* in Argentina.

EPHEDRA. JOINT PIR. Shrubs.

Aecidium ephedrae Speg. Rust on leaves and branches of *E. tweediana* in Argentina.

Epicium nigerianum Sacc and Syd. Black, crust-like areas on twigs of *E. andina* in Chile.

EPIDENDRUM. See Orchidaceae.

EPILOBIUM. WILLOW WEED. Herbs with willow-like foliage and large showy flower spikes.

Ascochyta epilobii Oud. Elongate dull-brown leaf spots on *E. angustifolium* in Holland.

Cercospora epilobii Schn. On leaves of *E. hirsutum* and *E. montanum* in Europe.

Cylindrosporium epilobianum Sacc. and Faut. Confluent leaf spots on *E. hirsutum* in France.

Fusicladium heterosporum V. Hoehn. Brown, often confluent, leaf spots on *E. parviflorum* in Austria.

Gloeosporium epilobii Pass. Brown leaf spots on *E. angustifolium* in France.

Marsonia chamaenerii (Rostr.) P. Magn. and var. *germanica* Syd. Irregular ochraceous brown leaf spots on *E. angustifolium* and *E. hirsutum* in Greenland and Germany.

Mycosphaerella adusta Fekl. On *E. boreale*, *E. bongardi*, and *Epilobium* sp. in Alaska.

Mycosphaerella microspila B. and Br. On leaves of *E. dodonaei*, *E. montanum*, and *E. palustre* in Denmark, Great Britain, Belgium, and Italy.

Ovularia epilobiana Sacc. and Fautr. Pale-brown leaf spots on *E. hirsutum* in France.

Ovularia epilobii Lindr. Large, irregular, confluent gray or gray-brown leaf spots on *E. palustre* in Finland.

Phyllosticta chamaenerii Allesch. Small angular to irregular, ochraceous then white, leaf spots with red margins on *E. angustifolium* in Germany.

Phyllosticta epilobii Brun. Small circular gray-white leaf spots with brown margins on *E. hirsutum* in France.

Phyllosticta epilobii-rosei Krieg. Circular to irregular, often confluent, brown leaf spots on *E. roseum* in Germany.

Plasmopara epilobii (Rh.) Schroet. Downy mildew on leaves of *E. hirsutum*, *E. latifolium*, and *E. parviflorum* in Alaska, Denmark, and Germany.

Puccinia epilobii DC. Brown rust sori on lower leaf surfaces of *Epilobium* spp. in Australia and Europe.

Puccinia epilobii-fleischeri Ed. Fisch. Yellow to brown rust pustules on leaves of *E. fleischeri* in Switzerland.

Puccinia gigantea Karst. Rust on brown sunken spots on leaves and stems of *E. angustifolium* in Switzerland and Scandinavia.

EPILOBIUM—Continued.

- Puccinia krookii** P. Henn. Powdery, dark-brown rust sori on leaves of *E. flavescens* in the Union of South Africa.
- Ramularia enecans** Magn. Gray-white to yellow leaf spots on *E. angustifolium* in Germany.
- Ramularia epilobii** All. On leaves of *E. palustre* in Germany.
- Ramularia epilobii-parviflori** Lindr. Subcircular to oblong gray-brown leaf spots with purple margins on *E. parviflorum* in Denmark and Finland.
- Ramularia epilobii-rosei** Lind. Subcircular or oblong grayish-green to pale-brown leaf spots on *E. roseum* in Denmark and Austria.
- Ramularia hornemanni** Lindr. Subcircular brown, yellow, or red leaf spots on *E. hornemanni* in Lapland.
- Ramularia punctiformis** (Schlechtld.) V. Hoeh. Leaf spot on *Epilobium* spp. in Europe. Reported from Wisconsin.
- Septoria alpicola** Sacc. On leaves of *E. alpinum* in Italy.
- Venturia maculaciformis** (Desm.) Sacc. On leaves of *E. hirsutum* in Austria.
- EPIMEDIUM**. Herbs used in rock gardens.
- Cercospora epimedii** Jach. On leaves of *E. pinnatum* in Russia.
- Mycosphaerella epimedii** (Sacc.) Jaap. On leaves of *E. alpinum* in Dalmatia, Italy, and Switzerland.
- Phyllosticta epimedii** Sacc. Irregular ochraceous leaf spots with red margins on *E. alpinum* in Italy.
- Puccinia epimedii** (Henn. and Shir.) Miy. and Ito. Rust sori on circular yellow-brown leaf spots on *E. macranthum* in Japan.
- EPIPACTIS**. Hardy terrestrial orchids. See Orchidaceae.
- ERAGROSTIS**. LOVE GRASS. Teff. Annual or perennial grasses.
- Epichloe eragrostis** Pole-Evans. Attacks *E. plana* in the Union of South Africa.
- Helminthosporium eragrostidis** P. Henn. Black, velvety fungus patches in the glumes of *Eragrostis* sp. in the Congo.
- Sorosporium turneri** McAlp. Dense black masses of smut spores in the ovaries of *E. nigra* in Australia.
- Sphaeclotheca kusanoana** P. Henn. (*Ustilago kusanoana*, P. Henn.) Smut in inflorescences of *E. ferruginea* in Japan.
- Uromyces pedicellata** Pole-Evans. Yellow-brown to black rust sori on leaves and culms of *E. abyssinica* and *E. curvula* in the Union of South Africa.
- Ustilago egenula** Syd. and Butl. Olivaceous smut sori replacing the ovaries of *E. nutans* in India.
- ERANTHEMUM**. Tropical shrubs, some species cultivated for the foliage or flowers.
- Isariopsis acanthacearum** Cke. See Daedalacanthus.
- EEANTHIS**. WINTER ACONITE. Low perennial herbs.
- Peronospora eranthidis** (Pass.) Fekl. Downy mildew on leaves of *E. hyemalis* in Switzerland.
- Septoria cajadensis** Speg. Ashen white or ochraceous circular to angular, often confluent, leaf spots on *E. hyemalis* in Italy.
- Tubercinia eranthidis** (Pass.) Liro. Smut on *E. hyemalis* in southern Europe.
- EREMOCITRUS**. Small trees allied to *Citrus*.
- Bacterium citri** Hassé. See Citrus.
- EREMURUS**. DESERT CANDLE. Hardy liliaceous desert plants with tall flowering stalks.
- Puccinia eremuri** Kom. Dark-brown rust pustules on both leaf surfaces of *E. iberiensis*, *E. kaufmanni*, *E. robustus*, and *E. spectabilis* in Asia Minor and Persia.
- Rhabdospora eremuri** Ohl. Dark-brown spots on stems of *Eremurus* sp. in Russia.
- ERIA**. See Orchidaceae.
- ERIANTHUS**. RAVENNA GRASS. Tall, reedlike, perennial grasses with large woolly plumelike inflorescences.
- Cintractia pulverulenta** Cke. and Mass. Black smut sori in ovaries of *Erianthus* sp. in India.
- Puccinia damlofi** Bub. Brown to black powdery rust pustules on leaves of *E. hostium* in Yugoslavia.
- Uredo ravennae** Maire. Leaf rust on *E. ravenna* in French north Africa and Spain.
- Ustilago erianthi** Syd. Smut sori destroying the ovaries and forming dark-olive spore masses on *E. ravenna* in Russia.
- Ustilago sacchari** Rabh. See Saccharum.
- ERICA**. HEATH. Ornamental shrubs.
- Gibbera satiburgensis** Niessl. On leaves of *E. carnea* in Germany.
- Phyllosticta ericae** Allesch. Red-brown patches on leaves of *E. carnea* in Germany.
- Stemphylium ericetonum** B. and De B. On *Erica* spp. in Europe.
- Venturia straussii** Sacc. On branches and leaves of *E. scoparia* in Italy and France.
- ERIGERON**. FLEABANE. Mostly weeds, a few species cultivated in border plantings like *Aster*.
- Acididium spegazzinii** Pat. Leaf rust on *E. albus*, *E. bonariensis* and *E. maximus* in South America.
- Coleosporium erigerontis** Syd. Powdery golden rust pustules on leaves of *E. linifolius* in Formosa.
- Entyloma calendulae** (Oud.) De B. See Calendula.
- Puccinia doloris** Speg. Brown rust pustules on small brown sunken leaf spots of *E. bonariensis*, *E. decum*, and *Erigeron* sp. in Argentina, Chile, Colombia, Costa, Rica and Guatemala.
- Puccinia dorensis** Blytt. Dark-brown rust pustules on leaves of *E. alpinus* and *E. uniflorus* in Norway, Switzerland, and India.
- Septoria chanousii** Ferr. Irregular brown leaf spots on *E. uniflorus* in Italy.
- Septoria erigeronata** Thuem. On leaves of *E. elongatus* in Siberia.
- Venturia bonariensis** Speg. Brown indefinite leaf spots on *E. bonariensis* in Argentina.
- Venturia tucumanensis** Speg. On leaves of *E. canadense* in Argentina.
- ERIOBOTRYA**. LOQUAT. Small trees grown for ornament and edible fruit.
- Ascochyta eriobotryae** Vogl. Circular to oblong dull-brown to white leaf spots on *E. japonica* in Italy.
- Coleopuccinia simplex** Diet. Leaf rust on *E. japonica* in Japan.
- Corticium salmonicolor** B. and Br. See Citrus.
- Coryneum eriobotryae** Scalia. Circular, gray-white leaf spots with dark-purple margins on *E. japonica* in Sicily.
- Entomosporium** sp. Raised shiny spots on fruit and leaves surrounded by yellowish rings and never confluent on *E. japonica* in the Union of South Africa.
- Entomosporium mespili** (DC.) Sacc. See Cotoneaster.
- Fomes lamaensis** Murr. See Hevea.
- Fusicladium** sp. Circular brown, then velvety dark olive-green, often confluent, spots on fruit and leaves of *E. japonica* in the Union of South Africa. Probably the same as the following species. The leaves are often distorted.
- Fusicladium eriobotryae** Cav. Scabby areas on leaves, stems and fruit of *E. japonica* in Australia, Russia, and Italy. Reported from California.

ERIOBOTRYA—Continued.

Fusarium udum Butl. See *Dianthus*.

Gloeosporium eriototryae Speg. Circular, then confluent, ashen leaf spots on *E. japonica* in Argentina.

Leptosphaeria eriototryae Syd. and Butl. On leaves of *E. japonica* in India.

Phyllosticta eriototryae Thuem. Subcircular to irregular and confluent gray-brown leaf spots on *E. japonica* in Argentina, Brazil, Japan, Formosa, Australia, Russia, and Italy.

Phyllosticta uleana Syd. Large circular, rarely confluent, dull-yellow to brown leaf spots with purple margins on *E. japonica* in Brazil.

Septoria eriototryae Maffei. Irregular, black leaf spots on *E. japonica* in Italy.

ERIOSTEMON. Australian evergreen shrubs.

Puccinia eriostemonis McAlp. Dark-orange rust pustules on leaves of *E. myoporoides* in Australia.

ERLANGEA. Vernonia-like shrubs.

Mycosphaerella erlangae Bacc. On leaves of *E. abyssinica* in Abyssinia.

Puccinia erlangae Grove. Leaf rust on *E. tomentosa* in British east Africa.

ERODIUM. HERON'S-BILL. Afilaria. Filaree. Annual or perennial herbs grown in flower gardens.

Gloeosporium schweinfurthianum Thuem. On leaves of *E. glaucophyllum* in Egypt.

Peronospora erodii Fekl. Downy mildew on leaves of *E. cicutum*, *E. cicutarium*, and *E. malachoides* in Sardinia, Denmark, Russia, Austria, and Germany.

Phyllosticta erodii Speg. Subcircular brown leaf spots with broad purple margins on *E. malachoides* in Argentina.

Ramularia erodii Bres. Small irregular leaf spots on *E. cicutarium* in Spain, Denmark, and Germany.

Ramularia geranii (West.) Fekl. See *Geranium*.

Synchytrium papillatum Farl. Small galls on dull-purple leaf spots on *E. cicutarium* in Guadeloupe, Canary Islands, St. Croix, and the Union of South Africa. Also in California on imported plants.

ERYNGIUM. ERYNGIO. Sea holly. Annual and perennial herbs valued for the steel-blue cast of their stems.

Aphelenchus olesistis Ritz. Bos. See *Begonia*.

Ascochyta phomoides Sacc. On stems of *Eryngium* sp. in France.

Entyloma eryngii (Cda.) De By. Smut pustules on circular to irregular, often confluent, brown leaf spots on *E. campestre* and *E. planum* in France, Italy, Austria, and Germany. Reported from Iowa.

Erysiphe taurica Lévy. See *Althaea*.

Leptosphaeria woodrowi wilsoni Garb. On leaves of *E. campestre* in Russia.

Mycosphaerella eryngii (Fr.) Lind. and var. *libanotis* Fekl. On leaves of *E. campestre*, *E. maritimum*, and *Libanotis montana* in Europe.

Phyllosticta eryngiana Sacc. and Fautr. Dull yellow-brown leaf spots on *E. campestre* in France.

Phyllosticta eryngiella Bub. Gray-white angular leaf spots on *E. campestre* in Hungary.

Phyllosticta eryngii Syd. Large indefinite black, often confluent, leaf spots on *E. maritimum* and *E. paniculatum* in Denmark and Germany.

Phyllosticta eryngiella Bub. On leaves of *E. campestre* in Hungary.

Puccinia eryngii DC. Leaf rust on *E. bovei*, *E. campestre*, *E. congestum*, *E. creticum*, *E. glomeratum*, and *E. virens* in Europe, Egypt, French North Africa, and Asia Minor.

Ramularia eryngii Jachw. On leaves of *E. planum* in Russia.

Septoria eryngicola Oud. and Sacc. On leaves of *E. campestre* in Italy and Spain.

Septoria eryngii West. Angular brown to gray-white leaf spots on *E. maritimum* in Denmark and Belgium.

ERYSIMUM. BLISTER CRESS. Early-blooming hardy annuals with yellow and orange flowers.

Septoria erysimi Niessl. Pale-brown leaf spots on *E. cheiranthoides* in Bohemia.

Septoria repanda Bub. Circular to irregular, yellow or white leaf spots on *E. banaticum* and *E. repandum* in Yugoslavia and Bohemia.

ERYTHRAEA. CENTAURIUM. Ag. Low, annual, biennial or perennial herbs.

Cercospora erythraea Höll. Dull-yellow areas on leaves and stems of *E. linariaefolia* in Hungary.

Peronospora erythraea (Kuehn) Güm. Downy mildew on leaves, peduncles and stems of *E. centaurium*, *E. linariaefolia*, *E. litoralis*, *E. pulchella*, and *E. ramosissima* in central and north Europe.

Phyllosticta erythraea Sacc. and Speg. On leaves of *E. centaurium* in Italy.

Pycnospezia plantaginifolia Fekl. var. *erythraea* Pat. Brown spots on lower leaf surfaces of *E. centaurium* in Tunis and Algeria.

Septoria gentianae Thuem. var. *erythraea* Gz. Frag. On leaves of *E. chlooides* in Spain.

ERYTHRINA. CORAL TREE. Herbs, shrubs, and trees cultivated for ornament and as coffee shade.

Coleosporium erythrinae Petch. Leaf rust on *E. lithosperma* in Ceylon.

Colletotrichum erythrinae Koord. Anthracnose on leaves of *E. lithosperma* in Ceylon and Java.

Dichelirinia binata (Berk.) Arth. (*Uredo cabreriana* Kern. and Kellerm.) Powdery deep-brown rust pustules on lower leaf surfaces of *E. glauca* and *E. umbrosa* in Porto Rico, Cuba, Guatemala, and Trinidad.

Diplodia cacaoicola P. Henn. See *Theobroma*.

Fomes lamaeensis Murr. See *Hevea*.

Fomes lignosus Klotzsch. See *Hevea*.

Helminthosporium extensum Petch. On leaves of *E. lithosperma* in Ceylon.

Helminthosporium inversum Sacc. On leaves of *E. indica* in the Philippines.

Mycosphaerella erythrinae Koord. Circular gray-white leaf spots on *E. lithosperma* and *E. ovalifolia* in Java and Ceylon.

Phyllosticta australis Speg. Indefinite grayish-white leaf spots on *E. crista-galli* in Argentina.

Phyllosticta erythrinae Petch. Spots on branches and leaves of *E. lithosperma* in Ceylon.

Phyllosticta erythrinicola Young. Numerous small, circular, sordid-white leaf spots on *E. microphylla* in Porto Rico.

Ravenella erythrinae Güm. Leaf rust on *E. microcarpa* and *E. velutina* in Java.

Ravenella platensis Speg. Powdery brown rust sori on pyriform galls on leaf blades and petioles and on branches of *E. crista-galli* in Argentina and Uruguay.

Rosellinia arcuata Petch. See *Thea*.

Rosellinia pepo Pat. See *Citrus*.

Septoria bonanseaana Sacc. Numerous subcircular gray-white leaf spots with ochraceous margins on *E. breviflora* in Mexico.

Sphaerostilbe repens B. and Br. See *Hevea*.

Telimena erythrinae Rach. Subcircular yellow-green, then yellow, finally brown, leaf spots on *E. lithosperma* in Java.

Trabutia erythrina Rick. Circular black, often confluent, stomata on *E. crista-galli* in Brazil.

ERYTHRINA—Continued.

Uredo erythrinae P. Henn. Rust pustules on circular, then confluent, yellow-brown leaf spots on *E. indica*, *E. tomentosa* and *Erythrina* sp. in the Philippines, the Congo, and Uganda.

Uredo erythrinae-ovalifoliae Petch. Leaf rust on *E. ovalifolia* and *E. velutina* in Ceylon.

Uromyces erythrinae Lagh. Brown rust pustules on leaves of *Erythrina* sp. in Ecuador.

ERYTHRONIUM. TROUT LILY ADDER'S-TONGUE. Small, spring-flowering, hardy bulbous plants.

Ascochyta erythronii Sacc. and Speg. Indeterminate white leaf spots on *E. dens-canis* in Italy.

Septoria erythronii Sacc. and Speg. Subcircular to irregular white spots on leaves of *E. dens-canis* in Italy.

Septoria fatiscipora Bub. Circular to elongate dirty-white spots without definite margins on leaves of *E. dens-canis* in Montenegro.

Uromyces erythronii (DC.) Pass. Yellow rust pustules, which later become brown, on both sides of the leaves and on the petioles of *E. dens-canis* in France, Spain, Italy, Switzerland, Bulgaria, Yugoslavia, Dalmatia, Austria, Hungary, Japan, Siberia, and Germany.

ERYTHROXYLON. COCAINE TREE. Woody medicinal plants.

Corticium sp. Black rot of *E. coca* in Ceylon. Similar to *Corticium theae* on *Thea* (q. v.).

Corticium salmonicolor B. and Br. See Citrus.

Cronartium gilgianum P. Henn. Red-brown rust pustules on leaves of *Erythroxylon* sp. and *Euclea* sp. in central and east Africa.

Fomes lamaensis Murr. See Hevea.

Phyllachora asteriana Speg. Shiny black stromata on leaves of *E. suberosum* in Brazil.

Phyllosticta erythroxyl Graz. Small circular to irregular gray-white leaf spots with yellow to dull-brown margins on *E. coca* in Bolivia and Peru.

Uredo erythroxylonis Graz. Leaf rust on *E. areolatum* and *E. coca* in Porto Rico, Cuba, Peru, Bolivia, and Brazil.

ESCALLONIA. South American evergreen shrubs or small trees.

Phyllachora escalloniae Pat. Subcircular black stromata on leaves of *Escallonia* sp. in Ecuador.

Phyalospora escalloniae P. Henn. Black spots on leaves of *E. chlorophylla* in Brazil.

Trabutia escalloniae (P. Henn.) Theiss. and Syd. Black stromata on leaves of *E. rubra* in Argentina.

EUCALYPTUS. Valuable ornamental and timber trees, natives of Australia, many species of which are called gum trees, e. g., blue gum, spotted gum.

Aulographium eucalypti Cke. and Mass. On leaves of *E. obliqua* in Australia and Tasmania.

Camarosporium eucalypti Wint. Angular to irregular pale-brown to ashen leaf spots on *Eucalyptus* sp. in Australia.

Cercospora epicoccoides Cke. and Mass. Small purple, then confluent, leaf spots on *Eucalyptus* sp. in Australia.

Cercospora eucalypti Cke. and Mass. Subcircular, then confluent, pale-brown leaf spots with rosy margins on *Eucalyptus* sp. in Australia.

Cercospora theae Petch. See Thea.

Coniothecium eucalypti Thuem. On branches of *E. globulus* and *Eucalyptus* sp. in Spain, Portugal, and Ecuador.

Corticium salmonicolor B. and Br. See Citrus.

Coryneum eucalypti d'Alm. and da Cam. Small irregular white spots with brown margins on *E. globulus* in Portugal.

Cylindrosporium eucalypti McAlp. Circular definite gray leaf spots with red-brown margins on *E. melioidora* in Victoria.

Diaporthe cubensis Bruner. Produces a canker of trunks and limbs of *Eucalyptus* spp. in Cuba. The bark is destroyed, often with a gummy exudation, and lesions are formed which permit the entry of other fungi.

Endothia havanensis Bruner. On bark of twigs and branches of *E. botryoides*, *E. microphylla*, *E. occidentalis*, *E. robusta*, *E. rostrata*, *Mangifera indica*, *Persea gratissima*, and *Spondias mombin* in Cuba.

Gloeosporium eucalypti McAlp. Leaf anthracnose on *E. corynocalyx* in Australia.

Gloeosporium nigricans Cke. and Mass. Anthracnose on leaves of *E. pauciflora* in Australia.

Gloeosporium rhapidium Speg. Anthracnose on leaves of *E. globulus* in Uruguay and Argentina.

Harknessia uromycoides Speg. On leaves of *Eucalyptus* sp. and *E. globulus* in west Australia, Portugal, and California.

Hendersonia grandispora McAlp. Attacks leaves and twigs of *Eucalyptus* sp. in Victoria.

Hypoaspila eucalypti Wakef. Red-brown spots on leaves of *Eucalyptus* sp. in Queensland.

Lembosia orbicularis Wint. Black elongate superficial perithecia on subcircular leaf spots on *E. pilularis* in Australia.

Leptostromella eucalypti Cke. and Mass. Subcircular red-brown leaf spots on *Eucalyptus* sp. in Australia.

Melasma eucalypti Cke. and Mass. On *Eucalyptus* leaves in Queensland.

Mycosphaerella cryptica Cke. On leaves of *Eucalyptus* sp. in Australia.

Mycosphaerella molleriana Thuem. and var. **megalospora** da Cam. Irregular dull-brown leaf spots on *E. ficifolia*, *E. globulus* and *Eucalyptus* sp. in Brazil, Algeria, and Portugal.

Mycosphaerella nubilosa Cke. On leaves of *Eucalyptus* sp. in Australia.

Oidium sp. Powdery mildew causing crinkling and distortion of leaves on young shoots of *Eucalyptus* spp. in Brazil.

Oidium eucalypti Rostr. Powdery mildew on leaves of *Eucalyptus* seedlings in Denmark.

Phoma australis Cke. Elliptical, then confluent, dull-brown leaf spots on *Eucalyptus* sp. in Australia.

Phoma eucalyptidea Thuem. On leaves of *E. globulus* in Victoria and Portugal.

Phyllachora eucalypti (Cke. and Mass.) Theiss. and Syd. Black stromata on leaves of *E. viminialis* in Tasmania.

Phyllachora maculata Cke. Shiny black stromata on upper leaf surfaces of *Eucalyptus* sp. in Australia.

Phyllosticta eucalypti Thuem. Large irregular dull-brown, then gray-white, leaf spots on *E. giganteus* and *E. globulus* in Russia, Denmark, Spain, Portugal, Algeria, and Australia.

Phyllosticta eucalyptina Pat. On leaves of *E. globulus* in Tunis.

Phyllosticta globuli Pass. On leaves of *E. globulus* in Italy.

Phyllostictella phomatoides (Cke. and Mass.) Tass. On leaves of *Eucalyptus* sp. in Victoria.

Phyalospora latitans Sacc. Attacks leaves and twigs, causing a die-back of *E. colosseae* and *E. rostrata* in Portugal and Brazil. Reported from California.

Plocastroma inaequale (Cke.) Theiss. and Syd. Black shiny stromata on brown leaf spots on *Eucalyptus* sp. in Victoria.

Readeriella mirabilis Syd. Stromata on circular, sunken, dull-brown leaf spots on *E. capitellata* in Victoria.

EUCALYPTUS—Continued.

- Rhytisma eucalypti** P. Henn. Black opaque stromata on dull-brown, circular leaf spots on *E. santalifolia* in Australia.
- Schizothyrium eucalyptorum** Cke. and Mass. Circular reddish leaf spots on *E. obliqua* in Australia.
- Septoria eucalypti** Wint. and Roum. On leaves of *E. amygdalina* in Algeria.
- Septoria mortolensis** Penz. and Sacc. See *Acacia*.
- Sphaerostilbe repens** B. and Br. See *Hevea*.
- Stictus emarginata** Cke. and Mass. On leaves of *Eucalyptus* spp. in Ceylon.
- Ustilago vriesiana** Vuill. A smut said to cause galls on stems and roots of seedlings from which witches'-broomlike growths develop. On *E. amygdalina* in France and Holland.
- Canker**. An undetermined fungus attacks *Eucalyptus* spp., *Mangifera indica*, *Persea gratissima*, and other trees in Hawaii and possibly in Brazil. It works inward from the leaves and young shoots, as well as through the bark, forming cankers and ultimately killing infected trees.
- EUCHARIS**. AMAZON LILY. Bulbous plants.
- Accidium delicatulum** Arth. Leaf rust on *Eucharis* sp. in Trinidad.
- Uredo eucharidis** P. Henn. Brown leaf rust on *E. candida* in Peru and Ecuador.
- EUCHLAENA**. TEOSINTE. Stout annual or perennial grasses grown for forage and ornament.
- Helminthosporium euchlaenae** Zimm. Elongate brown leaf spots on *E. mexicana* in Tanganyika.
- Phyllosticta euchlaena** Sacc. Turns leaves of *E. luxurians* white, destroying their forage value. In the Philippines.
- Sclerospora maydis** (Rac.) Butl. See *Zea*.
- Sclerospora philippinensis** Weston. See *Zea*.
- Sclerospora sacchari** Miy. See *Saccharum*.
- Sclerospora spontanea** Weston. See *Zea*.
- Ustilago kellermanii** Clint. Smut sori involving the terminal and nodal growths of *E. luxurians* in Guatemala. The brown-black dusty spore masses are inclosed by the leaf sheaths.
- EUGENIA**. SURINAM CHERRY. CLOVE TREE. A large group of tropical trees and shrubs, many species cultivated for ornament or fruit.
- Actinothecium callicola** Speg. Small circular pale-brown leaf spots on *Eugenia* sp. in Brazil.
- Campotruncum eugeniae** Pat. Circular dull-brown leaf spots on *Eugenia* sp. in Indo-China.
- Catacauma peglerae** Doidge. Black stromata on leaves of *E. capensis* in the Union of South Africa.
- Coniothyrium trigonicolum** Rangel. On leaves of *E. uniflora* in Brazil.
- Entyloma eugeniaram** Cke. and Mass. Dark-brown smut sori on circular to angular leaf spots on *Eugenia* sp. in Queensland.
- Entyloma paradoxum** Syd. Leaf smut on *E. aquea* in Java.
- Fusarium udum** Butl. See *Dianthus*.
- Helminthosporium asterinoides** Sacc. and Syd. Black superficial fungus layers on lower leaf surfaces of *Eugenia* sp. in Brazil.
- Lasmenia balansae** Speg. Small black irregular stromata on leaves of *Eugenia* sp. in Paraguay and Brazil.
- Melasmia pulchella** Speg. On leaves of *Eugenia* spp. in Paraguay and Brazil.
- Metasphaeria incompleta** Rehm. On *Eugenia* sp. in the Philippines.
- Microcyclus labens** Sacc. and Syd. On leaves of *Eugenia* sp. in Chile.
- Mycosphaerella eugeniae** Rehm. On leaves of *E. uniflora* (*E. michellii*) in Brazil.
- Myiocopron valdivianum** Speg. On leaves of *Eugenia* sp. in Chile.
- Napicadium fumago** Speg. Thin black superficial fungus layers on leaves of *Eugenia* sp. and *Myrtus* sp. in Chile.
- Napicadium myrtacearum** Speg. As above on *Eugenia* sp. in Brazil.
- Phaeophleospora eugeniae** Rangel. On leaves of *E. uniflora* in Brazil.
- Phyllachora ambigua** Syd. Black irregular stromata on leaves of *E. jambolana* in India.
- Phyllachora angustispora** Speg. Black stromata on leaves of *Eugenia* sp. in Argentina.
- Phyllachora biareolata** Speg. Black stromata on circular brown leaf spots on *E. guabiju* in Argentina.
- Phyllachora phylloplaca** (Kze.) Theiss. and Syd. Shiny black stromata on leaves of *Eugenia* sp. in Brazil.
- Phyllachora vimulosa** Speg. Black stromata on leaves of *Eugenia* sp. in Costa Rica.
- Phyllachora wetzelii** Chardon. Small black circular stromata on brown leaf spots on *Eugenia* sp. in Porto Rico.
- Phyllosticta eugeniae** Young. Circular dark-brown leaf spots with raised light-brown margins on *E. buxifolia* in Porto Rico.
- Phyllosticta icarabhyensis** Rangel. Leaf spots on *E. uniflora* in Brazil.
- Phyllosticta myrticola** Speg. Circular white leaf spots with definite narrow dull-red margins on *E. smithii* and *Eugenia* sp. in Brazil and Victoria.
- Polystomella granulosa** (Kl.) Theiss. and Syd. Dull-black stromata on leaves of *Eugenia* sp. in Chile.
- Puccinia barbacensis** Rangel. Leaf rust on *Eugenia* sp. in Brazil.
- Puccinia eugeniae** Rangel. Leaf rust on *E. grandis* in Brazil.
- Puccinia grumixamal** Rangel. Leaf rust on *E. brasiliensis* in Brazil.
- Septoria eugeniaram** P. Henn. On leaves of *Eugenia* sp. in Argentina.
- Septoria eugenicola** Speg. Sanguineous leaf spots with white centers on *Eugenia* sp. in Argentina.
- Sphaerodothis balansae** (Tass.) V. Hoeh. Black stromata on lower leaf surfaces of *Eugenia* sp. in Paraguay.
- Uncinula australis** Speg. Powdery mildew on leaves of *Eugenia* sp. in Paraguay.
- Uredo eugeniaram** P. Henn. Leaf and stem rust on *E. uaiaba* and *Eugenia* sp. in Brazil.
- Uredo goldiana** P. Henn. Yellow to golden rust pustules on fruit of *Eugenia* sp. in Brazil.
- Uredo myrtacearum** Pasch. Yellow-brown rust pustules on leaves of *E. grandis* in Brazil.
- EULOPHIA**. Terrestrial orchids. See *Orchidaceae*.
- EUONYMUS**. BURNING BUSH. WINTER CREEPER. WAHOO. Deciduous or evergreen shrubs and small trees.
- Ascochyta evonymi** Oud. Circular gray-white leaf spots with brown margins on *E. europaea*, *E. japonica*, and *E. vulgaris* in Bohemia and Holland.
- Ascochyta evonymicola** Allesch. Irregular red-brown, then gray-white, spots on leaves of *E. europaea* in Italy and Germany.
- Cercospora evonymi** Erikss. Subcircular to angular brown leaf spots with purple margins on *E. europaea* in Scandinavia.
- Gloeosporium euonymicolum** Hemmi. Anthracnose causing leaf fall of *E. japonica* and *E. radicans* in Japan.
- Gloeosporium evonymi** Br. and Cav. Leaf anthracnose on *E. japonica* in Italy.

EUONYMUS—Continued.

Gloeosporium frigidum Sacc. Anthracnose on leaves of *E. japonica* in Italy.

Macrophoma cylindrospora (Desm.) Sacc. See *Hedera*.

Marssonina thomasi Sacc. Subcircular dull-red leaf spots on *E. europaea* and *E. latifolia* in Europe. Reported from Wisconsin.

Melampsora evonymi-capraearum Kleb. See *Salix*.

Microspora evonymi (DC.) Sacc. Powdery mildew on leaves of *E. europaea* and *E. verrucosus* in Europe.

Mycosphaerella evonymi (Kunze.) Schroet. On leaves of *E. europaea* in Europe.

Oidium evonymi-japonici (Arcang.) Salm. Dense white patches of mildew on leaves of *E. japonica* and *E. radicans* in Japan, Argentina, and Europe. Also known from Louisiana.

Phyllosticta aliena (Fr.) Sacc. Elongate spots on branches of *E. europaea* in France.

Phyllosticta bolleana Sacc. On leaves of *E. japonica* in Italy and Austria.

Phyllosticta destructiva Desm. See *Althaea*.

Phyllosticta evonymella Sacc. Angular olive leaf spots on *E. europaea*, *E. japonica*, and *E. latifolia* in Spain, Italy, France, and Germany.

Phyllosticta evonymicola Togn. Gray-white leaf spots with ochraceous margins on *E. europaea* in Italy.

Phyllosticta nemoralis Sacc. Indefinite pale-brown leaf spots on *E. europaea* in Spain and France.

Phyllosticta pustulosa S. and R. Rufous gray-white leaf spots on *E. japonica* in Algeria.

Phyllosticta sardoa Pass. Large gray-white leaf spots of *E. japonica* in Italy.

Septoria evonymella Pass. On leaves of *E. japonica* in Italy and Austria. Reported from South Carolina.

Septoria evonymi Rbh. Large pale-brown leaf spots on *E. europaea* in Italy, Russia, and Germany. Reported from Virginia.

Septoria evonymi-japonicae Pass. On leaves of *E. japonica* in Italy.

Septoria evonymina Sever. On *E. japonica* in Italy.

Septoria japonicae Oud. Large pale-brown leaf spots on *E. japonica* in Holland.

Septoria semicircularis Sacc. and Scalia. On leaves of *E. fimbriatus* in Portugal.

EUPATORIUM. THOROUGHWORT. BONESET. MIST FLOWER. Perennial herbs, the tropical species shrubby.

Aecidium ampliistrum Jacks. and Holw. Leaf rust on *Eupatorium* sp. in Costa Rica.

Aecidium eupatorii Diet. Rust pustules on large yellow circular leaf spots on *Eupatorium* sp. in Brazil.

Aecidium heteromorphum Speg. Leaf rust on *Eupatorium* sp. in Argentina.

Aecidium paramense Mayor. Rust on leaf blades and petioles and stems of *E. obscurifolium* in Colombia.

Aecidium roseum Diet. and Holw. Golden rust pustules on red or purple leaf spots on *Eupatorium* sp. in Mexico.

Baeodromus eupatorii Arth. Brown rust pustules on leaves and stems of *E. aschenbornianum* and *E. pazcuarensis* in Mexico and Guatemala.

Cionothrix praelonga (Wint.) Arth. Yellow rust pustules on leaves of *Eupatorium* spp. in Mexico, Guatemala, Costa Rica, Trinidad, Colombia, Ecuador, and Brazil.

Coleosporium eupatorii Arth. Golden rust pustules on leaves of *E. chinense*, *E. collinum*, *E. macrophyllum*, and *E. oerstedianum* in Porto Rico, Cuba, Guatemala, Costa Rica, Nicaragua, Colombia, and Formosa.

Cronartium andinum Lagh. Brown rust pustules on leaves of *E. glutinosum* in Ecuador.

Cronartium eupatorium Speg. Leaf rust on *Eupatorium* sp. in Argentina.

Mycosphaerella eupatoriicola Petr. On leaves of *E. cannabinum* in Bohemia and Silesia.

Phyllosticta eupatoriicola Kab. and Bub. Irregular, often confluent, dull-brown leaf spots on *E. cannabinum* and *E. odoratum* in Porto Rico and Bohemia.

Phyllosticta eupatorii Allesch. Small subcircular dull-brown to ashen, often confluent, leaf spots with dull-purple margins on *E. cannabinum* in Germany.

Phylllosticta eupatorina Thum. Subcircular ashen leaf spots with purple borders on *E. cannabinum* in Italy and Portugal. Reported from Illinois and New Jersey.

Puccinia aegopogonis Arth. and Holw. Brown leaf rust on *E. mairetianum*, *Aegopodium cenchroides*, and *A. tenellus* in Guatemala.

Puccinia basi-petala Jacks. and Holw. Leaf rust on *E. mairetianum* and *E. rafaelense* in Guatemala.

Puccinia caeuteutensis Speg. Brown rust pustules on leaves and branches of *E. patens* in Argentina.

Puccinia espinosarum Diet. and Holw. Brown to black rust pustules on leaves of *E. espinosarum*, *E. smithii*, *E. sordidum*, and *E. spiraeae-folium* in Mexico.

Puccinia eupatorii Diet. Leaf rust on *E. ballotaefolium*, *E. iresinoides*, *E. macrocephalum*, and *E. viridiflorum* in Trinidad.

Puccinia eupatoriicola Mayor. Leaf rust on *E. pycnocephalum* and *E. schiedeannum* in Colombia.

Puccinia eupatorii-columbiani Mayor. Leaf rust on *E. columbianum* and *E. inulaefolium* in Colombia and Trinidad.

Puccinia hodgeana Kern. Leaf rust on *E. phoenicolepis* and *E. schultzei* in Guatemala.

Puccinia horrida Lagh. Cinnamon-brown rust pustules on leaves of *Eupatorium* sp. in Ecuador.

Puccinia inanipes Diet. and Holw. Brown to black rust pustules on leaves of *E. brevipedis*, *E. hirsutum*, and *E. tubiflorum* in Mexico.

Puccinia inermis Jacks. and Holw. Powdery brown rust pustules on leaves of *Eupatorium* sp. in Costa Rica.

Puccinia noackii Syd. Dark-brown rust pustules on leaves of *Eupatorium* sp. in Brazil.

Puccinia pachyspora Diet. Black rust pustules on leaves and stems of *E. oblongifolium* in Brazil.

Puccinia rosea Roth. See *Ageratum*.

Puccinia solidipes Jacks. and Holw. Brown leaf rust on *E. tubiflorum* in Guatemala and Mexico.

Puccinia tinctoria Speg. Brown rust pustules on leaves of *E. prasilfolium* and *E. tinctorium* in Argentina and Paraguay.

Puccinosira eupatorii Lagerh. Leaf rust on *E. aschenbornianum* and *Eupatorium* sp. in Guatemala and Ecuador.

Septoria albo-maculans Syd. Circular to irregular leaf spots with raised margins on *E. nubigenum* and *E. pomaderiffolium* in Colombia and Guatemala.

Septoria eupatorii Rob. and Desm. Numerous small subcircular gray-white leaf spots on *E. cannabinum* in Italy, France, and Germany. Reported from the lower Mississippi Valley.

Synarpella castagnei (Speg.) Theiss. and Syd. Black stromata on brown, indistinct leaf spots on *E. tinctorium* in Brazil.

Uredo eupatoriicola P. Henn. Brown leaf rust on *Eupatorium* sp. in Brazil.

Uredo scopigena P. Henn. Rust on leaves and stems of *Eupatorium* sp. in Brazil.

Uredo suspecta Jacks. and Holw. Brown rust pustules on leaves of *E. daleoides* in Costa Rica.

- EUPHORBIA.** SPURGE. Poinsettia. Plants with milky juice and very diverse habit, from succulent cactus-like plants to low or prostrate herbaceous weeds.
- Cercospora euphorbiae** Pat. Circular gray leaf spots on *Euphorbia* sp. in Ecuador.
- Colletotrichum euehroum** Syd. Anthracnose, destroying the leaves of *E. antiquorum* and *E. neritifolia* in China and the Philippines.
- Endophyllum euphorbiae-silvaticae** (DC.) Wint. Yellow rust pustules on leaves of *E. amygdaloides* and *E. silvatica* in Europe.
- Erysiphe taurica** Lévl. See Althaea.
- Melampsora euphorbiae** (Schub.) Cast. Yellow and brown to black rust pustules on leaves and stems of *Euphorbia* spp. in Europe, Egypt, and Tripoli. Sparingly introduced into the United States on *E. cyparissias*.
- Melampsora euphorbiae-amygdaloides** Muell. Leaf rust on *E. amygdaloides* in Europe.
- Melampsora euphorbiae-dulcis** Othb. Golden and brown to dark-brown rust pustules on leaves and stems of *Euphorbia* spp. in Japan and Europe.
- Melampsora euphorbiae-engleri** P. Henn. Leaf rust on *E. engleri* in Tanganyika.
- Melampsora euphorbiae-gerardianae** W. Muell. Rust on stems and leaves of *Euphorbia* spp. in Asia Minor, Tripoli, Persia, India, and Europe. Also known from Indiana.
- Melampsora gelmii** Bres. Yellow and dark-brown rust pustules on leaves of *Euphorbia* spp. in Persia, north Africa, and Europe.
- Melampsora helioscopiae** Wint. Yellow and dark-brown to black rust pustules on leaves and stems of *Euphorbia* spp. (over 30 reported species) in Europe, Egypt, Abyssinia, Italian North Africa, Tunis, Asia Minor, India, Japan, and Ceylon.
- Mycosphaerella canariensis** Tass. On stems of *E. canariensis* in Italy.
- Ovulariopsis erysiphoides** Pat. and Har. White erysiphoid spots on leaves of *E. balsamifera* in central Africa.
- Peronospora andina** Speg. Downy mildew deforming the leaves of *E. rhytisperma* in Argentina.
- Peronospora cyparissiae** De By. Downy mildew on *E. cyparissias* in central Europe.
- Peronospora esulae** Gäum. Downy mildew on leaves of *E. esula* in central Europe.
- Peronospora valesiaca** Gäum. Downy mildew on leaves of *E. gerardiana* in Switzerland.
- Phoma baldratii** Bacc. On stems of *E. thurcilli* in Eritrea.
- Phragmodothis asperata** Syd. On *Euphorbia* sp. in the Union of South Africa.
- Phyllosticta euphorbiaecola** P. Brun. Subcircular brown leaf spots on *E. amygdaloides* in France.
- Phyllostictella euphorbiae** (Roum.) Tass. On leaves of *E. silvatica* in France.
- Puccinia euphorbiae** P. Henn. and var. **longipes** Syd. Leaf rust on *E. agowensis* and *E. eritrea* in Abyssinia and *E. calyculata*, *E. colinifolia*, and *Euphorbia* sp. in Mexico. Reported from Minnesota.
- Puccinia intumescens** (Syd.) Holw. Yellow aecial and black powdery telial sori on leaves of *E. calyculata* in Mexico.
- Ramularia euphorbiae** (Cast.) Sacc. Whitish areas on leaves of *E. peplus* in France.
- Septoria bractearum** Mont. Rufous spots on bracts on *E. serrata* in France and Spain.
- Septoria euphorbiae** Guep. Small subcircular olivaceous leaf spots on *E. amygdaloides*, *E. angulata*, *E. esula*, and *E. peplus* in Russia and France.
- Septoria euphorbiaicola** Höll. Circular olive-brown leaf spots on *E. procera* in Hungary.
- Septoria hartiottiana** Sacc. White leaf spots with narrow dark-purple margins on *E. palustris* and *E. pilosa* in France and Spain.
- Septoria kalthbrenneri** Sacc. On leaves of *E. aspera*, *E. palustris*, and *E. silvatica* in Austria and Germany.
- Septoria media** Sacc. and Brun. Subcircular, then confluent, brown leaf spots with gray-white centers on *E. amygdaloides* and *E. palustris* in France and Yugoslavia.
- Septoria thuemeniana** Pass. Attacks leaves, stems, and twigs of *E. exigua* in Denmark and Italy.
- Uredo cornu Har.** Brown leaf rust on *Euphorbia* sp. in Ecuador.
- Uredo euphorbiae-nudiflorae** P. Henn. Dull-brown rust pustules on yellow leaf spots on *E. nudiflora* in Jamaica.
- Uredo velata** E. and E. Brown leaf rust on *E. cordata* in Hawaii.
- Uromyces alpestris** Tranzsch. Brown to black rust pustules on leaves and stems of *E. cyparissias* in Europe.
- Uromyces andinus** P. Magn. Powdery chestnut-brown rust pustules on leaves of *E. chilensis*, *E. collina*, and *E. portulacoides* in Chile and Argentina. Reported from Utah.
- Uromyces astragali** (Opiz.) Sacc. See Astragalus.
- Uromyces bresadolae** Tranzsch. Brown rust pustules on leaves of *E. angulata* in Austria.
- Uromyces cristulatus** Tranzsch. Dark-brown to black rust pustules on leaves of *E. girardiana* and *E. petrophila* in Bohemia.
- Uromyces excavatus** (DC.) Lévl. Leaf rust on *E. angulata*, *E. dulcis*, *E. gloriosa*, *E. pubescens*, and *E. verrucosa* in Europe.
- Uromyces fischeri-eduardi** P. Magn. See Vicia.
- Uromyces haussknechtii** Tranzsch. Brown rust pustules on leaves of *E. spinosa* and *E. thamnoides* in Syria, Italy, Dalmatia, and France.
- Uromyces hermonis** P. Magn. Leaf rust on *E. caudiculosa* in Asia Minor.
- Uromyces kalmusii** Sacc. Chestnut-brown rust pustules on leaves of *E. cyparissias* and *E. esula* in Manchuria and central Europe.
- Uromyces kawakamii** Syd. Leaf rust on *E. serrulata* in Formosa.
- Uromyces laevis** Koernicke. Dark-brown rust sori on leaves of *E. gerardiana* in Russia, France, Switzerland, Holland, Austria, and Germany.
- Uromyces loti** Blytt. See Lotus.
- Uromyces mayorii** Tranzsch. Leaf rust on *E. orbiculata* in Colombia.
- Uromyces monspessulanus** Tranzsch. Brown rust pustules on leaves of *E. serrata* in France and Spain.
- Uromyces natalensis** P. Magn. Powdery brown rust pustules on leaves of *E. gucinzii* in the Union of South Africa.
- Uromyces pisi** (Pers.) Wint. See Pisum.
- Uromyces scutellatus** (Schränk.) Lévl. Leaf rust on *Euphorbia* spp. in Europe. Reported from Colorado.
- Uromyces striatellus** Tranzsch. Brown leaf rust on *E. hebecarpa* in Persia.
- Uromyces striolatus** Tranzsch. Leaf rust on *E. cyparissias* and *Euphorbia* sp. in Turkestan, Italy, France, and Switzerland.
- Uromyces sublevis** Tranzsch. Dark-brown to black rust pustules on leaves of *Euphorbia* spp. in Caucasia, Syria, and Europe.
- Uromyces tinctoricola** P. Magn. Leaf rust on *Euphorbia* spp. in Kurdistan, Persia, Syria, Anatolia, Spain, France, Russia, Macedonia, Austria, and Germany.

EUPHORBIA—Continued.

- Uromyces torridensis** Speg. Powdery brown rust pustules on leaves of *E. ovalifolia* and *E. serpens* in Argentina and Uruguay.
- Uromyces tuberculatus** Fekl. Cinnamon-brown to black rust pustules on leaves and stems of *E. exigua*, *E. hissarica*, and *E. platyphyllos* in Turkestan, Spain, France, Switzerland, and Germany.
- Uromyces uleanus** Diet. Brown, powdery, rust pustules on leaves of *E. stenophylla* in Brazil.
- Uromyces undulatus** Tranzsch. Leaf rust on *Euphorbia* sp. in Turkestan.
- Uromyces winteri** Wettst. Black rust pustules on leaves of *E. falcata* and *Euphorbia* sp. in Tripoli, Asia Minor, and Turkestan.

EURYA. Shrubs.

- Aulographum euryae** Syd. Large irregular leaf spots on *E. sinensis* in Japan.
- Catacauma euryae** (Racib.) Theiss. and Syd. Black stromata on circular, yellow-green leaf spots on *E. acuminata* and *E. japonica* in the Philippines and Java.
- Exobasidium euryae** Syd. and Butl. Causes hypertrophy of flowers and fruit of *E. acuminata* in India.
- Phyllachora transiens** Syd. and Butl. Black stromata on minute indefinite yellow leaf spots on *E. acuminata* in India.
- Septoria euryae** P. Henn. Dull-brown, then white leaf spots on *E. canescens* in Ecuador.

EVELYNA. See Orchidaceae.**EVODIA**. Ornamental shrubs grown for their handsome foliage.

- Bacterium citri** Hasse. See Citrus.
- Cercospora evodiae** Syd. Small yellow leaf spots on *E. meliaefolia* in Formosa.
- Coleosporium evodiae** Diet. Powdery golden rust pustules on leaves of *E. meliaefolia* in Japan and Formosa.
- Phyllosticta evodiae** Cke. Leaf spots on *E. accedens* in Australia.

EVOLVULUS. Annual or perennial prostrate or erect herbs.

- Cercospora balansae** Speg. On leaves of *Evolvulus* sp. in Brazil.
- Helminthosporium balansae** Speg. Indefinite, olivaceous spots, often involving entire leaf blades of *Evolvulus* sp. in Brazil.
- Puccinia desertorum** Syd. Yellow-brown rust pustules on leaves and stems of *E. alsinoides* in tropical Africa.
- Puccinia enecta** Speg. Dark-brown rust pustules on leaves and stems of *Evolvulus* sp. in Argentina.
- Puccinia tuyutensis** Speg. Brown rust pustules on leaves and stems of *E. falcatus* in South America.
- Uredo evolvuli** Speg. Powdery pale-brown rust pustules on leaves and stems of *E. nummularius* and *E. sericeus* in Argentina and Venezuela.

EXACUM. Herbs, cultivated for their blue, white, or lilac flowers.

- Coleosporium exaci** Syd. Golden rust pustules on leaves of *E. tetragonum* in the Philippines.

FABA. See Vicia.**FAGELIA**. See Calceolaria.**FAGOPYRUM**. BUCKWHEAT.

- Ascochyta fagopyri** Theum. and var. *italica* Trav. On leaves of *F. vulgare* (*F. esculentum*) in Russia, Italy, and Germany. Reported from Vermont.
- Fusicladium fagopyri** Oud. On leaves of *F. vulgare* (*F. esculentum*) in Holland.
- Phyllosticta polygonorum** Sacc. See Polygonum.
- Phytophthora parasitica** Dast. See Ricinus.
- Puccinia fagopyri** Barcel. Brown to black rust pustules on leaves of *F. vulgare* (*F. esculentum*) in India.
- Ramularia curvula** Fautr. On leaves of *F. vulgare* in Europe.
- Sclerotinia fagopyri** Hori. Sclerotia form in diseased seed of *F. vulgare* in Japan.
- Sphaelotheca fagopyri** Syd. and Butl. Smut forming dark-violet sori in ovaries of *F. vulgare* in India.

FAGUS. BEECH. Timber and ornamental trees.

- Actinonema fagicola** Allesch. Subcircular spots causing premature leaf-fall and swelling of buds of *F. sylvatica* in Germany.
- Ascochyta fagi** Wor. On *Fagus* sp. in Russia.
- Clasterosporium epiphyllum** (Lév.) Sacc. On leaves and fruit of *F. sylvatica* in France.
- Cryptostictus niesslii** Oud. On leaves of *F. sylvatica* in Holland.
- Cyttaria berteri** Berk. Yellow to golden obovate or turbinate fruiting bodies on branches of *F. obliqua* in Patagonia.
- Cyttaria darwinii** Berk. Causes a wood rot of branches of *F. antarctica* and *F. betuloides* in Chile.
- Cyttaria gunnii** Berk. Rotting branches of *F. cunninghamii* in Tasmania and Victoria and fruiting in dense clusters.
- Cyttaria harioti** Fisch. Subglobose brown fruiting bodies on rotted branches of *F. antarctica* and *F. betuloides* in Chile.
- Cyttaria hookeri** Berk. Yellowish to cinnamon-colored fruiting bodies on rotted branches of *F. antarctica* and *F. obliqua* in Patagonia.
- Discozia maculiformis** Syd. On *F. sylvatica* in Japan.
- Laestadia faginea** (Oke. and Plowr.) Sacc. On leaves of *F. sylvatica* in Great Britain.
- Leptosphaeria valdoblai** Ferr. On leaves of *F. sylvatica* in Italy.
- Linospora faginea** Sacc. Discolored areas on leaves of *F. sylvatica* in Italy and France.
- Melasmia antarctica** Speg. On branches of *F. antarctica* in Patagonia.
- Mikronegeria fagi** Diet. and Neg. Yellow to rufous rust pustules on leaves of *F. obliqua* and *F. procera* in Chile.
- Mycosphaerella fagi** (Awd.) Lind. On leaves of *F. sylvatica* in Europe.
- Phyllosticta fagi** Oud. Pale-brown, often confluent, leaf spots on *F. sylvatica* in Holland.
- Phyllosticta faginea** Bres. Small subcircular dull-brown leaf spots on *F. sylvatica* in Germany.
- Phytophthora fagi** R. Hartig. Downy mildew attacking the cotyledons of seedlings causing a serious "damping-off" disease in Europe. Among the hosts are *F. grandifolia*, *F. sylvatica*, *Abies* spp., *Acer palmatum*, *Acer* sp., *Fraxinus* spp., *Larix* sp., *Picea* sp., *Pinus montana*, *P. sylvestris*, and *Robinia pseudoacacia*. The cotyledons and basal portions of diseased stems turn brown.
- Plowrightia noxia** (Ruhl.) Sacc. See Castanea.
- Septoria fagi** Awd. and f. *amphigena* D. Sacc. On leaves of *F. sylvatica* in Italy, Germany, and Austria.
- Septoria fagicola** Speg. On leaves of *F. betuloides* in Patagonia.
- Sphaeronema fagi** Oud. On leaves of *F. sylvatica* in Holland.
- Sphaeronema fagineum** Oud. On leaves of *F. sylvatica* in Holland.
- Uncinula curvispora** K. Hara. Powdery mildew on leaves of *F. sylvatica* in Japan.

FARADAYA. Climbing shrubs allied to *Clerodendron*.

- Phyllosticta faradayae** Sacc. On leaves of *F. papuana* in Malaya.

- FATSIA.** Shrubs or small trees. Some species are put under *Tetrapanax*.
Aecidium fatsiae Syd. Leaf rust on *Tetrapanax* (*F. papyrifera*) in Formosa.
- FEDIA.** See *Patrinia*.
- FEIJOA.** Tropical fruit trees.
Catacauma feijoe (Rehm.) Theiss. and Syd. Shiny black stromata on leaves of *Feijoa* sp. in Brazil.
Schizothyrium hypodermoides Rehm. On leaves of *Feijoa* sp. in Brazil.
- FERONIA.** Spiny deciduous trees with hard-shelled fruits, related to *Citrus*.
Bacterium citri Hasse. See *Citrus*.
Corticium salmonicolor B. & Br. See *Citrus*.
- FERONIELLA.** Small spiny trees related to *Feronia*.
Bacterium citri Hasse. See *Citrus*.
Corticium salmonicolor B. & Br. See *Citrus*.
- FERULA.** GIANT FENNEL. Hardy herbs.
Aecidium ferulae Rouss. and Dur. Rust pustules on leaves and petioles of *F. communis* and *F. tingitana* in French North Africa, Sardinia, France, Switzerland and Belgium.
Puccinia elliptica Lindr. Leaf rust on *F. longifolia* in Russia.
Puccinia ferulae Rud. Powdery yellow and dark-brown rust pustules on leaves and stems of *F. communis* and *Ferulago galbanifera* in Italy, France, and Austria.
Puccinia libani P. Magn. Rust on leaves and stems of *F. nodiflora*, *Cachrys gonicarpa*, *C. odonto calyx*, and *Ferulago* spp. in Asia Minor, Persia, Turkestan, Russia, Finland, Italy, and Greece.
Puccinia sogdiana Kom. Brown rust pustules on leaves of *F. joeschkeana* and *F. rigidula* in Turkestan.
Uromyces ferulae Juel. Yellow and powdery brown rust pustules on leaf blades, petioles, and stems of *F. assoi* and *F. communis* in Algeria, Sicily, Corsica, and Spain.
- FESTUCA.** FESCUE. Annual or perennial grasses, some species cultivated for ornament or as pasture grass.
Cladochytrium graminis Busgen. This fungus attacks and destroys the roots of *F. ovina*, *Dactylis glomerata*, *Panicum* sp., *Poa annua*, and other grasses in Great Britain and Germany. The disease kills out the grass in patches, which are small and yellow at first, finally becoming of considerable size, irregular in shape, and brown in color.
Dilophia graminis (Pckl.) Sacc. See *Triticum*.
Hypochnus fuciformis (Berk.) McAlp. See *Lolium*.
Leptosphaeria culmicola (Fr.) Karst. See *Bromus*.
Leptosphaeria kerguelensis P. Henn. On leaves of *F. kerguelensis* in Kerguelen Island.
Ophiobolus cariceti (B. & Br.) Sacc. See *Triticum*.
Phyllachora sylvatica Sacc. Ovate to oblong dark-brown stromata on leaves of *F. duriuscula* in Spain and Italy.
Phyllosticta paleicola P. Henn. On leaves and palea of *F. erecta* in Kerguelen Island.
Physalospora festucae (Lib.) Sacc. See *Andropogon*.
Puccinia gibberosa Lagerh. Dark-brown to black rust pustules on leaves of *F. sylvatica* in Switzerland and Germany.
Puccinia mellea Diet. and Neg. Brown leaf rust on *F. muralis* in Chile.
Sclerospora macrospora Sacc. See *Triticum*.
Septoria festucae Died. On leaves of *F. gigantea* in Central Europe.
Septoria festucae-sylvaticae Died. On leaves of *F. sylvatica* in Central Europe.
Tilletia sterilis Ule. Leaf smut on *F. ovina* and *Koeleria cristata* in Germany.
Uredo abscondita Pautr. Brown linear rust pustules on leaves of *F. heterophylla* in France.
Uredo kerguelensis P. Henn. Brown rust pustules on yellow leaf spots on *F. kerguelensis* in Kerguelen Island.
Uromyces festucae Ule. Smut on culms and leaves of *F. ovina* in Germany.
Uromyces cuspidatus Wint. Elongate dark-brown rust sori on leaves of *F. commersonii* in Chile.
Uromyces festucae Syd. Black rust pustules on leaves of *F. arundinacea*, *F. ovina*, and *F. rubra* in Spain, Bohemia, Russia, Sweden, and Germany. The aecial stage occurs on *Ranunculus bulbosus* and *R. illyricus*.
Uromyces fuegianus Speg. Leaf rust on *F. purpurascens* in Chile.
Uromyces ranunculi-festucae Jaap. Powdery golden to brown rust pustules on leaves of *F. ovina* in Germany. The aecial stage occurs on *Ranunculus bulbosus*.
Ustilago festucae-tenellae P. Henn. Rachides and flowers deformed and converted into olivaceous-black powdery masses of smut spores. On *F. tenella* in Mexico.
Ustilago sphaerocarpa Syd. Black powdery smut sori replacing ovaries of *F. amplissima* in Mexico.
- FICARIA.** See *Ranunculus*.
- FICUS.** FIG. Banyan. A large genus of tropical trees, shrubs, and climbers. Many species of economic importance for fruit, ornament, rubber, and other products.
Acrotheciella javanica Koord. On leaves of *F. elastica* in Java.
Alternaria fici Parneti. Causes atrophy of fruit of *F. carica* in Italy.
Anisochora topographica (Speg.) Theiss. and Syd. Black irregular stromata on leaves of *Ficus* sp. in Brazil.
Aphelchus olesistus Ritz. Bos. See *Begonia*.
Ascochyta caricae Rabb. Red-brown leaf spots on *F. carica* in France and Austria.
Atichia treubii V. Hoeh. Superficial fruiting bodies on leaves of *F. elastica* in Java.
Bacterium fici Cav. This bacterium is said to cause a browning of the tissues of twigs of *Ficus* sp. in Italy.
Catacauma apoense Syd. Scattered irregular dull-black stromata on leaves of *F. apoensis*, *F. nervosa*, and *F. sibulanensis* in the Philippines.
Catacauma aspidium (Br.) Theiss. and Syd. and ff. **fici-albae** and **spinifera** Theiss. and Syd. Black stromata on leaves of *F. alba*, *F. banahaensis*, *F. fastigiata*, *F. fulva*, *F. heterophylla*, *F. minahassae*, *F. odorata*, *F. repens*, *F. riedeli*, *F. scandens*, *F. spinifera*, *F. tomentosa*, *F. ulmifolia*, and *F. validicaudata* in Java, India, Ceylon, Malaya, the Philippines, and the Union of South Africa.
Catacauma circinata Syd. Circular black stromata on leaves of *F. chrysolepis* and *F. odorata* in the Philippines.
Catacauma decalnearum (Lév.) Theiss. and Syd. Black stromata on leaves of *F. laeta* in Timor.
Catacauma elmeri Syd. Small scattered black stromata on leaves of *F. blepharostoma*, *F. manilensis*, *F. minahassae*, and *F. ulmifolia* in the Philippines.
Catacauma fici-fulvae Koord. Black stromata on leaves of *F. sinuosa* and *F. validicaudata* in the Philippines.
Catacauma fici-obscurae (Koord.) Theiss. and Syd. Shiny-black stromata on brown leaf spots on *F. obscura* in Java.

FICUS—Continued.

- Catacauma garciae** Theiss. and Syd. Black stromata on small greenish leaf spots on *F. garciae* in the Philippines.
- Catacauma grammicum** (P. Henn.) Theiss. and Syd. Black linear stromata on yellow-brown leaf spots on *F. capensis* in the Congo and the Union of South Africa.
- Catacauma infectorium** (Cke.) Theiss. and Syd. Shiny-black irregular stromata on leaves of *F. infectoria*, *F. religiosa*, and *F. ulmifolia* in India and the Philippines.
- Catacauma irregulare** (W. and C.) Theiss. and Syd. Black stromata on leaves of *Ficus* sp. in Angola.
- Catacauma karnbachii** (P. Henn.) Theiss. and Syd. Dull-black stromata on circular brown leaf spots on *F. baviocortica*, *F. heterophylla*, and *F. ulmifolia* in New Guinea and the Philippines.
- Catacauma lagunense** Syd. Shiny-black stromata on leaves of *F. hauili* in the Philippines.
- Catacauma microcentum** (B. and Br.) Theiss. and Syd. and var. **graphica** Theiss. and Syd. Circular black stromata on leaves of *F. mysorensis*, *Ficus* sp., and *Artocarpus* sp. in Ceylon, India, and the Philippines.
- Catacauma merrillii** Syd. Black stromata on leaves of *F. lucbanensis* in the Philippines.
- Catacauma microplacum** Syd. Tar-spot disease of leaves of *Ficus* sp. in Amboina.
- Catacauma nipponicum** Syd. Black circular shiny stromata on leaves of *F. nipponica* in Japan.
- Catacauma repens** (Cd.) Theiss. and Syd. Shiny-black stromata on leaves of *F. gossypina* and *F. religiosa* in Uganda, the Congo, and India.
- Catacauma robinsonii** Syd. Tar-spot on leaves of *Ficus* sp. in Amboina.
- Catacauma sanguineum** Theiss. and Syd. Small black stromata on reddish-yellow leaf spots on *F. heterophylla* and *F. odorata* in the Philippines.
- Catacauma schweinfurthii** (P. Henn.) Theiss. and Syd. Black stromata on leaves of *F. palmata* in Abyssinia.
- Catacauma ulcerata** (Mass.) Theiss. and Syd. Dull-black stromata on leaves of *F. ovata* in Uganda.
- Catacauma urophyllum** (V. Hoch.) Theiss. and Syd. Linear black stromata on leaves of *F. urophylla* in Java.
- Catacauma valsiforme** (Rehm.) Theiss. and Syd. Black stromata on brown leaf spots on *F. crassitorta* in the Philippines.
- Cercospora annulata** Cke. Circular, concentrically zoned, pale-brown leaf spots on *F. hispida* in India.
- Cercospora elastica** A. Zimm. Gray-brown leaf spots on *F. elastica* in Java.
- Cercospora rubro-cincta** Pat. On leaves of *Ficus* sp. in Indo-China.
- Cercospora rufula** Syd. Large circular rufous leaf spots on *Ficus* sp. in British North Borneo.
- Cercospora urostigmatis** P. Henn. Circular to angular brown leaf spots on *Ficus* (*Urostigma*) sp. in Brazil.
- Cladosporium scophilum** Farnetl. On fruit of *F. carica* in Italy.
- Clasterosporium elasticae** Koord. On bark of *F. elastica* in Java.
- Clasterosporium javanicum** Koord. On leaves of *F. elastica* in Java.
- Coccodotthella placida** Syd. Black stromata on upper leaf surfaces on *F. orcdryadum* in Tanganyika.
- Coleroa elasticae** Koord. On leaves and branches of *F. elastica* in Java.
- Corticium salmonicolor** B. and Br. See Citrus.
- Diplodia caecicola** P. Henn. See Theobroma.
- Eutypa erumpens** Mass. See Theobroma.
- Fomes lamaoensis** Murr. See Hevea.
- Fomes lignosus** Klotzsch. See Hevea.
- Fusicadium elasticae** Koord. On leaves of *F. elastica* in Java.
- Fusicadium fici** Bacc. Circular brown leaf spots with red-brown borders on *Ficus* sp. in Abyssinia.
- Gloeosporium pluggarii** Speg. On leaves of *Ficus* (*Urostigma*) sp. in Brazil.
- Gloeosporium sycophilum** Trinch. Anthracnose on leaves of *F. elastica* in Italy.
- Helminthosporium elastica** Koord. On leaves of *F. elastica* in Java.
- Helminthosporium ficinum** Sacc. On *F. wassa* and *Ficus* sp. in the Philippines and Amboina.
- Helminthosporium ficum** Rostr. Ashen leaf spots with dull-brown margins on *F. retusa* and *Ficus* sp. in Siam.
- Helminthosporium ficum** Yates. On leaves of *F. caudatifolia* and *F. ulmifolia* in the Philippines. This species, if distinct, should be renamed, the name being occupied.
- Mycosphaerella elasticae** Koord. On leaves of *F. elastica* in Java.
- Napicladium elasticae** Koord. On leaves of *F. elastica* in Java.
- Phaeosaccardinula ficiola** P. Henn. Effuse dull-brown leaf spots on *Ficus* sp. in Tanganyika.
- Phytactena ficum** P. Henn. Circular yellow leaf spots on *Ficus* sp. in Brazil.
- Phomopsis cinerescens** (Sacc.) Trav. Large cankers occur on the branches, rotting away the bark and sapwood and girdling and destroying involved limbs of *F. carica* in Great Britain, France, and Italy.
- Phyllachora amamensis** P. Henn. Circular to irregular groups of dull-black stromata on leaves of *F. capensis* and *Ficus* sp. in Tanganyika and the Union of South Africa.
- Phyllachora aspidioides** Sacc. and Berl. Dull-black stromata on leaves of *Ficus* sp. in Brazil.
- Phyllachora catervaria** (Br.) Sacc. Round black stromata on leaves of *F. hispida*, *F. merrillii*, and *F. oppositifolia* in India, Indo-China, Java, the Philippines, and Amboina.
- Phyllachora cayennensis** (DC.) Theiss. and Syd. A doubtful species on leaves of *F. roxburghi* and *Psidium guajava* in Brazil, Surinam, and India.
- Phyllachora dawel** Mass. Tar-spot on leaves of *Ficus* sp. in Uganda.
- Phyllachora devriesii** Koord. Black stromata on red-brown leaf spots on *F. leucantatoma* in Java.
- Phyllachora effigurata** Syd. Dull-black stromata on leaves of *Ficus* sp. in Brazil.
- Phyllachora ficiola** Allesch. and P. Henn. Round to elliptical shiny-black stromata on leaves of *Ficus* sp. in Brazil.
- Phyllachora pseudos** Rehm. Black stromata on leaves of *F. minahassae* and *F. nota* in the Philippines.
- Phyllachora vinosa** Speg. Smooth black stromata on violet leaf spots on *Ficus* sp. in Brazil.
- Phyllosticta ambigua** Sacc. Large irregular gray-white leaf spots with narrow rufous margins on *F. rubiginosa* in Italy.
- Phyllosticta caricae** C. Mass. On leaves of *F. carica* in Italy.
- Phyllosticta elasticae** Koord. On leaves of *F. elastica* in Java.
- Phyllosticta fici** Bres. Circular gray-white leaf spots with dull-brown margins on *F. macrophylla* in Portuguese St. Thomas.
- Phyllosticta ficiola** Pat. Circular gray-white leaf spots on *F. carica* in Tunis and Italy.

FIGUS—Continued.

- Phyllosticta roberti** Boy. and Jacz. Pale-brown leaf spots on *F. aurea* and *F. elastica* in France and the Bahamas. Reported from Florida.
- Phyllosticta sycina** Trav. On leaves of *F. heterophylla* in Italy.
- Phyllosticta sycophila** Thum. Large brown leaf spots on *F. carica* in Victoria and Italy.
- Physalospora atractina** Syd. Small gray-brown leaf spots on *Ficus* sp. in Brazil.
- Physalospora elasticae** Koord. Attacks branches of *F. elastica* in Java as a wound parasite.
- Phytophthora carica** (Hara) Hori. A downy mildew on fruit of *F. carica* in Japan. Small, sunken areas appear which soon involve the entire fruit in a soft brown rot. A thick cottony mycelium covers the surface of rotted fruits, accompanied by a disagreeable odor. Fruits remaining on the trees mummify.
- Phytophthora fici** Ven. Causes a soft rot of fruit of *F. carica* in India.
- Puccinia seputa** B. and C. Black leaf rust on *Ficus* sp. in Nicaragua.
- Ramularia sycina** Sacc. and D. Sacc. On leaves of *Ficus* sp. in Italy.
- Rosellinia bunodes** Sacc. See Citrus.
- Rosellinia echinata** Mass. Causes a serious decay of roots, bringing about death of infected trees of *F. dubia* in Malacca and Malaya.
- Schizochora elmeri** Syd. Circular to elliptical dull-black stromata on lower leaf surfaces of *F. guyeri* in the Philippines.
- Septogloeum elasticae** Koord. Subcircular light-brown leaf spots on *F. elastica* in Java.
- Septoria arcuata** Cke. White circular leaf spots with purple margins on *F. benghalensis* and *Ficus* sp. in India and Abyssinia.
- Septoria brachyspora** Sacc. Pale-brown leaf spots with ochraceous margins on *F. elastica* in Denmark and Italy.
- Septoria elastica** Koord. On leaves of *F. elastica* in Java.
- Septoria pipulae** Cke. Irregular purple-brown leaf spots on *F. religiosa* in India.
- Septoria pirottae** Tassi. Brown leaf spots with rufous margins on *F. pumila* (*F. repens*) in Italy.
- Septoriella conformis** Sacc. On leaves of *F. alba* in Malaya.
- Sporodesmium sieyuanum** Thum. Spots on leaves and stems of *F. carica* in Italy and Germany.
- Trabutia abyssinica** (P. Henn.) Theiss. and Syd. Irregular shiny-black stromata on leaves of *F. sycamorus* and *Ficus* sp. in Abyssinia and India.
- Trabutia amboensis** Syd. On leaves of *F. hasskarlii* in Amboina.
- Trabutia bengetensis** Yates. On leaves of *F. benguetensis* in the Philippines.
- Trabutia butleri** Theiss. and Syd. Black stromata on leaves of *Ficus* sp. in India.
- Trabutia chinense** Yates. Black stromata on leaves of *Ficus* sp. in the Philippines.
- Trabutia elmeri** Theiss. and Syd. Scattered irregular black stromata on leaves of *F. banahaensis* in the Philippines.
- Trabutia evansii** Theiss. and Syd. Small circular black stromata on yellow or brown leaf spots on *Ficus* sp. in Portuguese East Africa.
- Trabutia fici-dekdekeneae** Bacc. Black stromata on leaves of *F. dekdekene* in Abyssinia.
- Trabutia fici-hochstetteri** Bacc. Tar spot on leaves of *F. hochstetteri* in Abyssinia.
- Trabutia ficuum** (Niessl.) Theiss. and Syd. Shiny-black irregular stromata on leaves of *F. foveolata*, *F. garciae*, *F. infectoria*, *F. mysorensis*, *F. nipponica*, and *F. riedelii* in Japan, India, Portuguese East Africa, and the Union of South Africa.
- Trabutia incrustans** Raob. Black irregular stromata on leaves of *Ficus* sp. in Java.
- Trabutia nervisequens** (Lingl.) Theiss. and Syd. Black stromata on yellow leaf spots on *F. hochstetteri* and *Ficus* sp. in Abyssinia and the Union of South Africa.
- Trabutia novoguineensis** Theiss. and Syd. Black stromata on rust-brown leaf spots on *Ficus* sp. in New Guinea.
- Trabutia vernicosa** Theiss. and Syd. Irregular black stromata on leaves of *F. heterophylla* and *F. ulmifolia* in the Philippines.
- Trabutiella microthyroides** (P. Henn.) Theiss. and Syd. Black stromata on leaves of *F. erecta* in Japan.
- Uncinula pirottiana** Bacc. Powdery mildew on leaves of *Ficus* sp. in Abyssinia.
- Vol utella** sp. Causes a trunk and branch canker of *F. elastica* in Italy.
- FILIPENDULA**. MEADOWSWERT. Hardy herbs grown for their showy panicles of varicolored flowers.
- Ascochyta obducens** Fekl. Large brown irregular spots destroying the leaves of *F. ulmaria* in Siberia and Germany.
- Colletotrichum volutella** Sacc. and Malbr. On stems of *F. ulmaria* in France.
- Cylindrosporium filipendulae** Thum. Pale-yellow areas on leaves of *F. ulmaria* and *Spiraea salicifolia* in Denmark and Austria. Reported from Washington.
- Mycosphaerella maculans** Sacc. and Roum. On leaves of *F. ulmaria* in Russia, France, Belgium, and Germany.
- Phyllosticta filipendulae** Sacc. Subcircular gray-white leaf spots on *F. hexapetala* in Italy.
- Phyllosticta filipendulina** Sacc. and Syd. and var. **ulmariae** Sacc. Subcircular dull-brown leaf spots on *F. ulmaria* in Italy and France.
- Phyllosticta ulmariae** Thum. Small irregular gray-white leaf spots on *F. ulmaria* in Scotland and Siberia.
- Ramularia ulmariae** Cke. Subcircular white spots on leaves of *F. hexapetala* and *F. palmata* in France, Denmark, and Germany.
- Septoria quevillensis** Sacc. Small dark-red leaf spots with pale centers on *S. ulmaria* in France.
- Triphragmium anomalum** Trenz. Leaf rust on *F. palmata* in Russia.
- Triphragmium filipendulae** (Lasch.) Pass. Orange-yellow rust pustules on leaf blades and petioles of *F. hexapetala* and *Ulmaria pentaphylla* in central and northern Europe.
- Urocystis filipendulae** Fekl. Black powdery smut sori on leaf petioles and nerves of *F. hexapetala* in Estonia, Sweden, Denmark, and Germany.
- FLACOURTIA**. Tropical shrubs, one species (Governor plum) cultivated for its fruit.
- Uredo uguressae** Petch. Rust on fruit of *F. ramontchii* in Ceylon.
- FLEMINGIA**. Shrubs used as ornamentals.
- Uromyces flemmingiae** P. Henn. Brown powdery rust pustules on leaves of *Flemingia* sp. in the Congo.
- FLUGGEEA**. Tropical shrubs, sometimes cultivated.
- Nothoravenelia japonica** Diet. See Securinega.
- FOENICULUM**. FENNEL. Annual, biennial, or perennial herbs, sometimes cultivated.
- Ascochyta foeniculina** McAlp. On fruit of *F. vulgare* in Australia.
- Ascospora foeniculi** P. Magn. On leaves of *F. officinale* in Austria.
- Erysiphe taurica** Lév. See Althaea.
- Mycosphaerella himantia** (Pass.) Died. Leaf spot of *Foeniculum* sp. in Dalmatia.

- FORSYTHIA.** GOLDEN BELL. Shrubs grown for their abundance of yellow spring flowers.
- Marsonia forsythiae** Lind. Large ashen leaf spots with dull brown margins on *F. suspensa* (*F. fortunei*) in Denmark.
- Phyllosticta forsythiae** Sacc. Subcircular ochraceous leaf spots on *F. suspensa* in Denmark, Spain, France, and Italy.
- Phytophthora syringae** Kleb. See *Syringa*.
- FORTUNELLA.** KUMQUAT. Small trees, related to *Citrus*, grown for their yellow ornamental and edible fruit.
- Bacterium citri** Hasse. See *Citrus*.
- FRAGARIA.** STRAWBERRY.
- Aphelenchus fragariae** Ritz. Bos. This nematode causes a fasciation of *Fragaria* sp. in Europe known as the "brush" or "cauliflower" disease. Fleishy malformations take the place of leaves and flowers, or the flowers may appear free of the deformed leaves and be in turn caused to assume odd shapes. The diseased areas assume a deep blood-red color. Infested plants are stunted and fail to form fruit.
- Aphelenchus olesistis** Ritz. Bos. See *Begonia*.
- Peronospora fragariae** Roze. and Cornu. Downy mildew on leaves of *F. vesca* in France, Switzerland, Italy, and Russia. Also reported from Iowa.
- Pileospora fragariae** (Br. and Har.) Petr. On leaves of *F. vesca* and *Potentilla caulescens* in Europe.
- Phragmidium fragariastrum** (DC.) Schroet. See *Potentilla*.
- Phyllosticta grandimaculans** Bub. and Krieg. Brown leaf spots on *F. vesca* in Denmark and Germany.
- Puccinia fragariae** Barcl. Yellow to brown rust pustules on leaves of *F. vesca* in India.
- Septoria fragariae** Desm. Large circular brown leaf spots with red margins on *Fragaria* spp. in Brazil, Chile, Siberia, and Europe. Possibly only another name for the imperfect stage of *Mycosphaerella fragariae*.
- Thecaphora pallescens** Fingerh. Doubtful species of leaf smut on *F. collina* in Germany.
- Tylenchus dipsaci** Kuehn. See *Narcissus*.
- FRANKENIA.** Low perennial evergreen shrubs.
- Puccinia frankeniae** Link. Leaf rust on *F. hirsuta*, *F. laevis*, and *F. pulverulenta* in Egypt, Tunis, Portugal, France, Belgium, Switzerland, and Russia.
- FRAXINUS.** ASH. Ornamental, shade, and timber trees.
- Ascochyta fraxini** Kab. and Bub. (*A. fraxini* Oud.) On branches of *F. excelsior* and *F. ornus* in Europe.
- Ascochyta fraxinicola** Brun. Circular to oblong brown spots on twigs of *F. excelsior* in France.
- Ascochyta metullicola** B. and Br. Circular brown leaf spots on *Fraxinus* sp. in Scotland.
- Ascochyta orni** Sacc. and Speg. On leaves of *F. ornus* in Italy.
- Cercospora fraxini** (DC.) Sacc. Large brown areas on leaves of *F. excelsior* and *F. ornus* in Russia and Italy.
- Cercospora lumbricoides** Turc. and Maffei. Subcircular to oblong brown leaf spots with red-brown margins on *Fraxinus* sp. in Italy.
- Laestadia fraxini** Fautr. On leaves of *Fraxinus* sp. in France.
- Phyllosticta ambigua** Scalia. On leaves of *F. ornus* in Italy.
- Phyllosticta diedickii** Bub. and Syd. On leaves of *F. excelsior* in Germany.
- Phyllosticta fraxinicola** Carr. Subcircular to irregular dull-brown leaf spots on *F. excelsior* in France, Russia, and Germany. Reported from several localities in the United States.
- Phyllosticta orni** Bub. Circular dull-yellow leaf spots with narrow brown margins on *F. ornus* in Hungary.
- Phyllosticta osteospora** Sacc. See *Morus*.
- Phyllosticta trappenii** Oud. Large ochraceous leaf spots on *F. juglandifolia* in Holland.
- Phytophthora fagi** R. Htg. See *Fagus*.
- Puccinia fraxini** Kom. Powdery dark-brown rust pustules on leaves of *F. longicuspis* and *F. rhynchophylla* in Manchuria and Japan.
- Sclerotium scutellatum** A. S. See *Acer*.
- Scolecotrichum fraxini** Pass. Attacks the leaves of *F. ornus* in Italy.
- Septoria elaeospora** Sacc. Indefinite ochraceous leaf spots with brown margins on *F. excelsior*, *F. ornus*, and *F. pubescens* in Italy and Portugal.
- Septoria orni** Pass. Dull-brown effuse leaf spots on *F. excelsior* and *F. ornus* in Denmark, France, and Italy.
- Uncinula fraxini** Miy. Powdery mildew on leaves of *F. longicuspis* in Japan.
- Uncinula salmonei** Syd. Powdery mildew on leaves of *F. bungeana* in Japan.
- Uncinula sengokui** Salm. Powdery mildew on leaves of *F. bungeana* in Japan.
- Venturia fraxini** (Fr.) Aderh. On leaves of *F. excelsior* in northern Europe.
- FREESIA.** Cormous plants, cultivated for their flowers.
- Phoma bulbicola** Tassi. On bulbs of *F. odorata* in Italy.
- Uromyces ecklonii** Bub. Yellow-brown to black rust pustules on leaves of *F. refracta* in the Union of South Africa.
- Uromyces freesiae** Bub. Brown leaf rust on *F. odorata* in the Union of South Africa.
- FREY CINETIA.** Climbing shrubs.
- Sphaerodothis merrillii** (P. Henn.) Theiss. and Syd. Circular black stromata, shiny above, dull below, on leaves of *F. ensifolia* and *F. williamsii* in the Philippines.
- Uredo freycinetiae** Rac. Rust on yellow, sunken leaf spots on *F. imbricata* in Java.
- FRITILLARIA.** FRITILLARY. Crown-imperial. Guinea-hen flower. Hardy, low-growing, spring-blooming bulbous plants.
- Uromyces acidiformis** (Schlecht.) Thuen. See *Lilium*.
- Uromyces fritillariae** (Schlecht.) Thuen. Yellow-brown to chestnut-brown rust pustules on leaf blades and petioles and stems of *F. meleagris* in Europe.
- Uromyces lili** (Link.) Fekl. See *Lilium*.
- Uromyces miurae** Syd. Leaf rust on *F. kamschatensis* in Japan.
- Uromyces mogianensis** Bub. Brown powdery rust pustules in sunken yellow leaf spots on *F. bucharica* and *F. quicquardia* in Greece, Morocco, and Turkestan.
- FUCHSIA.** Shrubs or small trees, cultivated for their flowers.
- Coleosporium fuchsiae** Cke. Yellow rust pustules on leaves of *F. excorticata* in New Zealand.
- Endomyces meliolineola** Rehm. On leaves of *Fuchsia* sp. in Brazil.
- Phomopsis fuchsiae** (Brun.) Sacc. On stems of *Fuchsia* sp. in Bohemia.
- Phyllosticta fuchsicola** Speg. Gray-white circular leaf spots with broad purple margins on *F. coccinea* in Chile.
- Puccinia fuchsiae** Syd. and Holw. Leaf rust on *F. microphylla*, *F. thymifolia*, and *Lopezia hirsuta* in Guatemala, Mexico, Costa Rica, and Panama.
- Septoria fuchsiae** Roum. Irregular pale-brown leaf spots on *F. coccinea* in France.

FUCHSIA—Continued.

Septoria fuchsicola Syd. Subcircular to irregular brown leaf spots with reddish-purple raised margins on *F. coccinea* in Germany.

Uredo fuchsiae Arth. and Holw. Leaf rust on *F. splendens* in Guatemala.

FUMARIA. FUMITORY. Annuals cultivated for their flowers.

Ascochyta fumariae Höll. Indefinite brown areas on leaves of *F. schleicheri* in Hungary.

Peronospora affinis Rossm. Downy mildew on leaves of *F. capreolata*, *F. macrocarpa*, *F. officinalis*, *F. parviflora*, and *F. vaillantii* in Tunis and Europe.

Ramularia fumariae Speg. Small circular ashen leaf spots with violet borders on *F. capreolata* in Argentina.

Rhizoctonia napi West. See Brassica.

FUNKIA. See Hosta.**FUNTUMIA**. Tropical trees, one species cultivated for rubber.

Colletotrichum funtumiae Petch. Anthracnose on leaves of *F. elastica* in Ceylon.

Fomes lamaensis Murr. See Hevea.

Nectria funtumiae Mass. Said to cause cankers on trunks of *F. elastica* in Uganda.

FURCRAEA. Succulent desert plants resembling *Agave*.

Physalospora fourcroyae P. Henn. On leaves of *F. gigantea* in Tanganyika.

GAGEA. Liliaceous herbs with white, yellow, or rose flowers.

Heterosporium ornithogali Klotzsch. See Ornithogalum.

Puccinia megatherium Syd. Cinnamon-colored rust pustules on both leaf surfaces of *G. reticulata* in the Caucasus.

Puccinia pachyderma Wettst. Powdery cinnamon-colored rust pustules on both leaf surfaces of *G. caucasia*, *G. damascena*, *G. graeca*, *G. lutea*, *G. persica*, and *G. pusilla* in Assyria, Russia, and Karpatos Island (Asiatic Turkey).

Septoria commutata Bub. Indefinite yellow, then grayish, leaf spots on *G. lutea* and *G. pratensis* in Bohemia and Galicia.

Synchytrium laetum Schroet. (*Pycnochytrium laetum* Schroet.) Forms small sulphur to golden-yellow punctiform galls on leaves, sepals and peduncles of *G. arvensis*, *G. saxatilis*, *G. lutea*, *G. silvatica*, *G. minima*, *G. pratensis*, and *Tulipa sylvestris* in Japan, Norway, Denmark, Russia, Austria, and Germany.

Synchytrium punctatum Schroeter. Forms small dirty-white to yellowish swellings or gall-like structures on leaves of *G. pratensis* in Austria and Germany.

Uromyces gageae G. Beck. Brown powdery rust pustules on both surfaces of leaves of *G. lutea*, *G. pratensis*, and *G. silvatica* in Europe.

Uromyces ornithogali Lév. Linear brown to dark-brown rust pustules on leaves of *G. arvensis*, *G. bohémica*, *G. lutea*, *G. minima*, *G. pratensis*, *G. pusilla*, *G. saxatilis*, *G. stenopetala*, *Muscari racemosum*, *Ornithogalum nanum*, *O. nutans*, and *O. umbellatum* in France, Great Britain, Spain, Portugal, Hungary, Persia, Austria, and Germany.

Ustilago ornithogali (Schm. and Kze.) Kuehn. Olive-brown, powdery smut pustules on leaves and scapes of *G. arvensis*, *G. bohémica*, *G. fibrosa*, *G. liotardi*, *G. pratensis*, *G. lutea*, *G. minima*, *G. pusilla*, *G. saxatilis*, *G. spathacea*, and *G. stenopetala* in Denmark and Italy. **U. ornithogali-erythronii** Clint. has been reported on *Erythronium* sp. in the United States.

GAILLARDIA. Showy annual and perennial herbs grown in flower gardens and borders.

Septoria gaillardiae Speg. Ashen leaf spots with dull-brown margins on *G. doniana* in Argentina.

GALACTIA. Prostrate or twining leguminous, perennial herbs or shrubs, mostly tropical.

Phyllachora galacteae Earle. Irregular black stromata on leaves of *G. rudolphoides*, *G. striata*, and *G. tenuiflora* in the Bahamas and Porto Rico.

GALANTHUS. SNOWDROP. Spring-blooming bulbs with solitary white flowers.

Botrytis galanthina (B. and Br.) Sudw. Attacks the host plants as they appear above ground, distorting and finally rotting the leaves and flowers. A gray mold followed by black sclerotia forms over the diseased areas. The host is *G. nivalis* in England, Denmark, Holland, and Germany.

The species of *Botrytis* and *Sclerotium* attacking the various ornamental bulbous plants have never been carefully studied as to their interrelationships and it is not clear whether few or many species are involved, or whether one or several occur in the United States. With this fact in mind, importations of bulbs should be carefully watched.

Melampsora galanthi-fragilis Kleb. A rust characterized by sunken yellow spots up to 1.5 cm. in diameter on both surfaces of the leaves of *G. nivalis*, *Salix fragilis* and other *Salix* spp. in Russia, Bulgaria, Spain, Switzerland, Austria, Hungary, and Germany.

Puccinia galanthi Unger. Rust sori, powdery, round to elliptical, up to 6 mm. in diameter, brown to black in color. On *G. nivalis* in Austria, Hungary, and Germany.

Urocystis galanthi Pape. Smut sori in blue-gray linear spots on leaf blades, sheaths, and flower stalks of *G. nivalis* in Germany.

GALEGA. GOAT'S- RUE. Bushy perennial leguminous plants.

Ascochyta galegae Höll. Irregular dull-brown leaf spots on *G. officinalis* in Hungary.

Cercospora galegae Sacc. Oblong white leaf spots on *G. officinalis* in Italy.

Ramularia galegae Sacc. Subcircular sordid-white leaf spots on *G. officinalis* in Italy and France.

Uromyces galegae (Opiz.) Sacc. Leaf rust on *G. officinalis* in Italy and the Balkans.

Uromyces genistae-tinctoriae (Pers.) Wint. See Caragana.

GALEOPSISIS. Annual weeds of the mint family.

Phyllosticta galeopsisidis Sacc. Ochraceous leaf spots on *G. versicolor* in Italy.

Septoria cotylea Pat. and Har. Indefinite brown leaf spots on *G. ceratonica* and *G. tetrahit* in Spain, France, and Denmark.

GALIUM. BEDSTRAW. Weak, slender herbs with square stems and whorled leaves.

Aecidium molluginis Wurtz. Leaf rust on *G. mollugo* in Switzerland.

Calloria quitensis Pat. On *Galium* sp. in Ecuador.

Fusicladium ruthenicum Petr. On leaves of *G. schultesii* in Europe.

Peronospora galii Fckl. Downy mildew on leaves of *G. mollugo*, *G. palustre*, and *G. uliginosum* in central and north Europe.

Peronospora galii veri Güm. As above on *G. verum*.

Peronospora silvatica Güm. Downy mildew on leaves of *G. schultesii* and *G. silvaticum* in Switzerland, Hungary, Austria, and Germany.

Phakopsora punctiformis (Barcl. and Diet.) Diet. Yellow-brown rust pustules on leaves of *G. aparine* in India.

Phyllosticta decipiens C. Mass. On leaves of *G. mollugo* in Italy.

Physalospora molluginis (Oth.) Sacc. On leaves and stems of *G. mollugo* in Switzerland.

Placosphaeria galii Sacc. On stems of *Galium* spp. in Europe.

Pseudopeziza autumnalis (Fr.) Karst. Black spots on leaves and stems of *G. boreale*, *G. mollugo*, *G. uliginosum*, *Asperula odorata*, and *Rubia tinctorum* in Siberia, Great Britain, France, Belgium, and Germany.

GALIUM—Continued

- Puccinia celakovskiana** Bub. Powdery brown rust pustules on leaves and stems of *G. cruciata* in Europe.
Puccinia eximia Arth. and Holw. Brown leaf rust on *G. mexicanum* in Guatemala.
Puccinia gali-elliptici Maire. Powdery brown rust pustules on leaves of *G. ellipticum* in Corsica.
Puccinia gali-silvatici Wurth. Leaf rust on *G. sylvaticum* in Switzerland.
Puccinia lagerheimii Lindr. Dark-brown rust pustules on leaves and stems of *G. sylvestre* in Switzerland.
Puccinia pallidifaciens Lindr. Dark-brown rust pustules on leaves and stems of *G. boreale* in Russia.
Septoria cruciatae Rob. and Desm. Dull-brown leaf spots with brown borders on *G. boreale*, *G. chersonense*, *G. cruciata*, *G. pedemontanum*, and *G. rotundifolium* in Europe.
Septoria gali-borealis P. Henn. (*S. gali-borealis* Bub. and Kab.) On leaves of *G. boreale* in Japan and Bohemia.
Septoria relicta Bub. On leaves of *G. relicta* in Bohemia.
Septoria urens Pass. On leaves of *G. tricornis* in Italy.
Synchytrium globosum Schroet. See *Potentilla*.
Thekopsora guttata (Schroet.) Syd. See *Asperula*.
Uromyces gali Diet. Rust on leaves and stems of *G. aparine* in Japan.

GALPHIMIA. Shrubs.

- Uromyces galphimiae** Diet. and Holw. Powdery cinnamon-brown to black rust pustules on *G. humboldtiana* in Mexico.

GARCINIA. MANGOSTEEN. GAMBOGE. Tropical fruit trees.

- Ceuthospora garciniae** Syd. On leaves of *G. mangostana* in the Philippines.
Corticium koleroga (Cke.) V. Hoeh. See *Coffea*.
Gamboge disease. A fruit disease due to an unknown cause is reported on *G. mangostana* in Ceylon.
Gloeosporium garciniae Koord. Irregular dull-brown leaf spots on *G. dulcis* in Java.
Helmintosporium garciniae Petch. On leaves of *G. mangostana* in Ceylon.
Micropeltis garciniae P. Henn. Black radiate perithecia on leaves of *Garcinia* sp. in Tanganyika.
Niptera garciniae P. Henn. On leaves of *Garcinia* sp. in Tanganyika.
Pirostoma garciniae P. Henn. On leaves of *Garcinia* sp. in Tanganyika.
Zignoella garciniae P. Henn. Forms cankers on woody portions in which appear superficial, black, carbonous perithecia. Attacks *G. mangostana* in Malaya and Tanganyika, death of infected plants often resulting.

GARDENIA. WARNERIA. Ag. CAPE JASMINE. Tropical shrubs or small trees cultivated for foliage and flowers.

- Corticium salmonicolor** B. and Br. See *Citrus*.
Hemileia woodii Kalkbr. and Cke. Yellow rust sori as broad patches on leaves of *G. edulis*, *G. jasminoides*, and *G. radicans* in Java, China, and Queensland.
Phyllosticta gardeniae Tassi. Subcircular to irregular pale-yellow leaf spots on *G. jasminoides* (*G. florida*) in Italy.
Ramularia gardeniae C. Mass. On leaves of *G. grandiflora* in Italy.
Septoria gardeniae Sacc. On leaves of *Gardenia* sp. in Italy.
Uredo gardeniae-thunbergiae P. Henn. Rust pustules on yellow effuse leaf spots on *G. thunbergia* in the Union of South Africa.

GARRYA. SILK-TASSEL BUSH. Ornamental shrubs.

- Ascochyta garryae** Sacc. Brown leaf spots with dark borders on *G. elliptica* in France.
Phyllosticta garryaeicola Pass. Subcircular brown leaf spots on *G. elliptica* in France.
Septoria garryae Roum. Subcircular to irregular, often marginal, brown, then ashen-gray, leaf spots on *G. elliptica* in France.
Septoria thuretii Brun. Pale-brown leaf spots with dull-brown margins on *G. thuretii* in France.

GARUGA. Tropical fruit trees.

- Fomes lamaensis** Murr. See *Hevea*.
Kuehneola garugae Syd. Powdery yellow rust pustules on leaves of *G. abito* in the Philippines.

GASTROLOBUM. Australian shrubs.

- Achorella gastrolobii** (P. Henn.) Theiss. and Syd. On *G. spinosum* in Australia.
Aecidium stowardii Har. Leaf rust on *G. calycinum* in Australia.
Puccinia gastrolobii Diet. Cinnamon-brown rust pustules on leaves of *G. calycinum* in Australia.
Puccinia stowardii Har. Rust on leaves and branches of *G. calycinum* in Australia.
Uredo stowardii Har. Leaf rust on *G. calycinum* in Australia.

GAULTHERIA. WINTERGREEN. SALAL. Evergreen, erect or procumbent plants cultivated for their leaves, flowers, and fruit.

- Pucciniastrum gaultheriae** Syd. Rust on leaves of *G. nummularioides* in India.
GAYLUSSACIA. HUCKLEBERRY. Shrubs grown for their fruit. See also *Vaccinium*.
Cercospora gaylussacii Speg. Angular spots on leaves of *Gaylussacia* sp. in Brazil.
Exobasidium gaylussaciae P. Henn. Irregular fleshy swellings occur on leaf blades and petioles and stems of *G. brasiliensis* in Brazil.
Phaeochora densa (B. and Br.) Theiss. and Syd. Irregular black stromata on leaves of *Gaylussacia* sp. in Brazil and Costa Rica.

GEISSORRHIZA. Half-hardy lili-like bulbous plants.

- Uromyces geissorrhizae** P. Henn. Yellow rust pustules, becoming dark-brown, on both leaf surfaces of *G. rupestris*, *G. secundata*, and *Moraea ramosa* in the Union of South Africa.

GEITONOPLESIS. Woody twiners in Australia.

- Uredo geitonoplesii** McAlp. Brown rust sori on leaves of *G. cymosum* in Australia.

GENIPA. West Indian shrubs and trees, one species grown for its fruit.

- Phylachora genipae** Arth. and D.L. Black stromata on leaves of *G. americana* in Porto Rico.

GENISTA. Broom. Ornamental shrubs grown for their handsome flowers.

- Diaporthe inaequalis** (Curr.) Nke. On *Genista* sp. in Europe.
Gloeosporium gargaricum Sacc. and D. Sacc. On branches of *G. michelii* in Italy.
Septoria cytisi Desm. See *Laburnum*.
Septoria nitida Ferr. On leaves of *G. germanica* in Italy.
Uromyces genistae-tinctoriae (Pass.) Fekl. See *Caragana*.
Ustilago albidia Bub. Smut in anthers of *G. spathulata* in Yugoslavia.

- Venturia genista** (Fekl.) Wint. On branches of *G. germanica* in Austria and Hungary.

GENTIANA. GENTIAN. Herbs, cultivated for their blue flowers.

- Cronartium delawayi** Pat. Yellow to brown rust pustules on leaves of *G. picta* and *G. yunnanensis* in China.
Cronartium gentianeum Thuem. Brown rust pustules on leaves and stems of *G. asclepiadea* in Russia, Bulgaria, and Austria.
Mycosphaella gentianae Niessl. On stems of *G. asclepiadea* in China and Austria. Reported from Colorado.

GENTIANA—Continued.

- Peronospora carnioleica** G&um. Downy mildew on leaves of *G. amarellum* in Austria.
- Phyllosticta gentianellae** Mass. Brown leaf spots with yellow margins on *G. asclepiadea* in Italy.
- Ramularia evanida** (Kuehn.) Sacc. Yellow-brown leaf spots on *G. asclepiadea* in Switzerland, Bulgaria, Yugoslavia, and Germany.
- Septoria gentianae** Thuem. Dull-brown, irregular leaf spots on *G. ascendens*, *G. macrantha*, and *G. scabra* in Japan, Siberia, and Russia.
- Septoria microspora** Speg. Large brown leaf spots on *G. asclepiadea* and *G. nivalis* in Italy and Japan.
- Septoria montana** Trav. Subcircular ochraceous leaf spots on *G. acaulis* in Italy.
- Septoria tosevi** Bub. Circular to oblong, often confluent, dull-yellow leaf spots on *G. cruciata* in Bulgaria and Austria. Reported from Colorado.
- GERANIUM**. CRANE'S BILL. Annual, biennial, or perennial herbs, some species cultivated.
- Aecidium infrequens** Barcl. Leaf rust on *G. nepalense* in Japan.
- Cercospora geranii-sanguinei** P. Henn. Dull-brown spots at tips of leaves of *G. sanguineum* in Norway.
- Cercospora magnusiana** Allesch. Ochraceous-olivaceous leaf spots on *G. sylvaticum* in Switzerland, Austria, and Germany.
- Coleosporium geranii** Pat. Brown rust pustules on leaves of *Geranium* sp. in China.
- Euryachora geranii** (Fr.) Schroet. Fungus of uncertain position on leaves of *G. sylvaticum* in Germany.
- Graphium geranii** Vogl. Circular to elliptical, often confluent, dull-brown leaf spots on *G. molle* in Italy.
- Peronospora conglomerata** Fekl. Downy mildew on leaves of *G. columbinum*, *G. dissectum*, *G. molle*, *G. phaeum*, *G. pusillum*, and *G. robertianum* in French North Africa and Europe.
- Phyllosticta trallii** Sacc. Brown leaf spots on *G. sylvaticum* in Norway.
- Plasmopara pusilla** (De By.) Schroet. Downy mildew on leaves of *G. palustre*, *G. phaeum*, *G. pratense*, *G. pusillum*, and *G. sylvaticum* in Europe.
- Puccinia callaquinensis** Neger. Rust pustules on leaves and stems of *G. berterianum* in Chile.
- Puccinia cuneata** Diet. Leaf rust on *Geranium* sp. in Japan.
- Puccinia escharoides** Syd. Powdery dark-brown rust pustules on leaf blades and petioles of *G. sinense* in central Africa.
- Puccinia geranii-pilosi** McAlp. Brown to black rust pustules on leaf blades and petioles of *G. pilosum* in Australia.
- Puccinia morthieri** Koern. Brown-black rust pustules on circular yellow leaf spots on *G. macrorrhizum*, *G. pyrenaicum*, *G. robertianum*, and *G. sylvaticum* in Europe and Siberia.
- Puccinia saniniensis** P. Magn. Powdery dark-brown rust pustules on leaves of *G. crenophilum* in Syria.
- Ramularia dolomitica** Kab. and Bub. On leaves of *G. phaeum* in Finland.
- Ramularia geranii** (West.) Fekl. Subcircular dull-brown leaf spots on *Erodium cicutarium* and *Geranium* spp. in Europe and Siberia. Reported from Louisiana.
- Ramularia geranii-phaei** (Mass.) Magn. On leaves of *G. phaeum* in Austria and Bohemia.
- Ramularia geranii-sanguinei** Massal. Rufous leaf spots on *G. sanguineum* in Italy.
- Ramularia geranii-silvatici** Vesterg. Angular-brown leaf spots on *G. sanguineum* and *G. sylvaticum* in Sweden and Estonia.
- Septoria daniloi** Bub. On leaves of *G. columbinum*, *G. lucidum*, and *G. pallens* in Spain, Yugoslavia, and Russia.
- Septoria expansa** Niessl. Indefinite areas on leaves of *G. dissectum*, *G. palustre*, and *G. sanguineum* in France, Russia, and Bohemia.
- Septoria geranii** Rob. and Desm. Irregular olivaceous-brown leaf spots with purple margins on *G. molle*, *G. pusillum*, *G. robertianum*, *G. sanguineum*, *G. sylvaticum*, and *Pelargonium* sp. in Spain, France, Italy, Tunis, the Union of South Africa, and Germany.
- Septoria geranii-nodosi** Massal. Brown leaf spots on *G. nodosum* in Italy.
- Uromyces geranii** (DC.) Outh. and Wartm. Brown rust pustules on leaf blades and petioles of *Geranium* spp. (about 30 species) in Colombia, Europe, Japan, and the Union of South Africa. Several doubtful American references.
- Uromyces kabatianus** Bub. Yellow to brown rust pustules on circular yellow leaf spots on *G. molle* and *G. pyrenaicum* in Denmark, Sweden, Estonia, Austria, and Germany.
- Uromyces scarious** Berk. Brown leaf rust on *G. dissectum* and *G. potentilloides* in New Zealand.
- Uromyces trunculatus** Trott. Yellow-brown rust pustules on leaves of *G. striatum* in Italy.
- Venturia circinans** (Fr.) Sacc. On leaves of *G. molle*, *G. rotundifolium*, and *Geranium* sp. in Australia, Italy, France, and Germany.
- GERARDIA**. See *Stenandrium*.
- GERBERA**. **PERIDIUM** Ag. BARBERTON DAISY. Perennial herbs grown for their yellow or pink flower heads.
- Aecidium erythrum** Kalch. and Cke. Leaf rust on *G. lanuginosa* in India and the Union of South Africa.
- Ascochyta gerberae** Maffei. Circular spots, enlarging to occupy entire area of leaves of *G. jamesonii* in Italy.
- Septoria gerberae** Syd. Large purplish-black blotches on leaves of *G. jamesonii* and other species in the Union of South Africa. The leaves are often destroyed.
- GESNERIA**. Shrubs grown for their showy tubular flowers.
- Cocconia gesneraceae** P. Henn. Circular black leaf spots on *Gesneria* sp. in Brazil.
- Puccinia gesneriacearum** Diet. Leaf rust on *Gesneria* sp. in Brazil.
- Rostrischiella nervicola** (Ithm.) Fitzp. Black superficial rough fruiting bodies on yellow leaf spots on *G. albiflora* in Porto Rico.
- GEUM**. **AVENS**. Herbs, sometimes cultivated for their yellow flowers or long plummy fruit.
- Cercospora gei** Fekl. On leaves of *G. rivale* in Austria.
- Depazea geicola** Fr. On leaves of *G. strictum* in Siberia.
- Gloeosporium gei** Trail. On stems of *G. urbanum* in Scotland.
- Phragmidium circumvallatum** P. Magn. Yellow rust sori on leaves of *G. heterocarpum* and *G. kokanicum* in Asia Minor, Turkestan, Algeria, Spain, and Russia.
- Phyllosticta gei** Thuem. On leaves of *G. urbanum* and *Geum* sp. in Siberia, Bohemia, Austria, and Germany.
- Puccinia gei** McAlp. Brown rust pustules on yellow-green sunken leaf spots on *G. renisfolium* in Tasmania.
- Ramularia gei** (Elias.) V. Hoeh. Subangular ochraceous to gray-brown spots on leaves of *G. rivale* and *G. urbanum* in Europe. Reported from Missouri.
- Ramularia trotteriana** Sacc. Small subcircular gray leaf spots with dark purple margins on *G. montanum* and *G. urbanum* in Italy.

- GIGANTOCHLOA.** See Bambuseae.
- GILIA.** Annual, biennial, and perennial herbs.
Phytophthora parasitica Dastur. See Ricinus.
Puccinia patagonica Speg. Rust pustules on leaves, stems, and flowers of *G. gracilis* in Argentina and Chile.
- Urocystis gilliae** Speg. Smut on roots of *G. foetida* in Argentina.
- GINKGO.** MAIDENHAIR TREE. Ornamental trees.
Phyllosticta ginkgo Brun. On leaves of *G. biloba* in France and Russia. Reported from the District of Columbia.
- Phyllosticta salisburiae** Tassi. Large irregular brown leaf spots on *G. biloba* in Italy.
- GLADIOLUS.** Summer and autumn flowering cormous plants.
Ascochyta gladioli Trav. and Spess. Dead areas at base of stems of *Gladiolus* sp. in Portugal.
Bacterium gladioli Sever. (*Pseudomonas gladioli* Sever.) This bacterium is reported from Italy and Holland as the cause of a soft rot of *Gladiolus* corms. The disease is first noted as a dying of the tops, beginning with the leaf tips, but soon involving the entire leaf which droops and finally falls over. *G. colvillei* is given as the host.
Mycosphaerella fusca Pass. On *Gladiolus* sp. in Italy.
Puccinia gladioli Cast. Produces brown, linear spots, limited by the veins, on both surfaces of leaves in which brown or black rust pustules appear. The host species are *G. aleppicus*, *G. byzantinus*, *G. communis*, *G. eckloni*, *G. illyricus*, *G. reuteri*, and *G. segetum*. The disease occurs in Italy, France, Spain, Portugal, Dalmatia, Algeria, and Asia Minor.
Uredo gladioli-buettneri Bub. Brown rust pustules on leaves of *G. buettneri* in Togo.
Uredo pulchra Syd. Rust pustules on leaves of *Gladiolus* sp. in the Himalaya Mountains of India.
Urocystis gladioli (Req.) Sm. This smut fungus attacks and destroys the corms of *Gladiolus* (*G. communis*, *G. imbricatus*, and *G. segetum*), a black spore mass forming within. It occurs in England, France, Holland, and Germany.
Uromyces gladioli P. Henn. Yellow, finally brown, rust pustules on both surfaces of leaves of *G. angustus*, *G. blandus*, *G. maculatus*, *G. orchidiflorus*, and *G. quartianus* in the Union of South Africa.
Uromyces nyikensis Syd. Dark-brown to black rust pustules are produced on both surfaces of leaves of *G. nyikensis* on the Nyika Plateau, Africa. The rust greatly disfigures the leaves.
Uromyces transversalis Thuem. Round to elongate dead spots on both surfaces of the leaves in which appear brown to deep-brown or black rust pustules. The hosts are *G. psittacinus*, *G. saundersii*, *Tritonia* (*Montbretia*) *lineata*, *T. securigera*, and *T. squarida* in the Union of South Africa.
- GLAUCIUM.** HORN POPPY. Perennial herbs with large poppylike flowers and glaucous blue foliage.
Entyloma fuscum Schroet. Smut sori in gray-white, then brown, leaf spots on *G. flavum*, *G. luteum*, *Papaver somniferum*, and *Rhoeas* sp. in France, Hungary, and Germany. Also known from Maine and Iowa.
Entyloma glaucii Dang. Smut sori in subcircular leaf spots on *G. flavum* and *G. luteum* in France, Denmark, and Germany.
- GLEDITSIA.** HONEY LOCUST. Trees grown for their finely divided foliage and fragrant flowers.
Phyllosticta triacanthi Sacc. Irregular buff leaf spots with narrow brown margins on *G. triacanthos* in Italy.
- GLIRICIDIA.** Small, ornamental, leguminous trees.
Cercospora gliricidae Syd. On leaves of *G. sepium* in the Philippines.
Colletotrichum gliricidae Syd. Anthracnose on *G. sepium* in the Philippines.
Corticium salmonicolor B. and Br. See Citrus.
- GLOBULARIA.** GLOBE DAISY. Herbs and shrubs with small blue flowers in heads.
Phyllosticta globulariae West. Subcircular dull-brown leaf spots on *G. vulgaris* in Belgium.
Puccinia grisea (Strauss.) Wint. Rust sori in sunken brown leaf spots on *G. cordifolia*, *G. nudicaulis*, *G. vulgaris*, and *G. wilkommii* in Europe.
Septoria globulariae Sacc. Subcircular gray-white leaf spots on *G. nudicaulis* and *G. vulgaris* in Italy.
- GLOXINIA.** Stemless cormous plants with showy bell-like flowers.
Aphelenchus oleisistis Ritz. Bos. See Begonia.
- GLYCERIA.** PANICULARIA Ag. Perennial grasses, sometimes grown for ornament.
Entyloma spectabile Karst. Smut sori in elongate yellow to brown leaf spots on *G. spectabilis* in Finland.
Fusieladium hartiottianum Sacc. On leaves of *G. borrieri* in France.
Mycosphaerella recutita (Fr.) Johans. See Aira.
Physoderma gerhardti Schroet. Brown areas on leaves and sheaths of *G. aquatica*, *G. fluitans*, *Alopecurus pratensis*, and *Phalaris arundinacea* in Germany.
Puccinia glyceriae Ito. Powdery golden to brown rust pustules on leaves of *G. aquatica* and *G. tonglensis* in Japan.
Ustilago grammica B. and Br. See Aira.
- GLYCINE.** See Soja.
- GLYCYRRHIZA.** LICORICE. Perennial herbs with fleshy roots.
Cercospora cavarae Sacc. and D. Sacc. Brown indefinite leaf spots on *G. glabra* in Italy.
Phyllosticta glycyrrhizae Brun. Brown leaf spots on *G. glabra* in France.
Uromyces ellipticus Diet. and Neg. Brown rust pustules on leaves and stems of *G. astragalina* in Argentina and Chile.
- GOETHEA.** Evergreen shrubs.
Phyllosticta goetheae Magn. Irregular gray-white leaf spots with dull-brown borders on *G. cauliflora* (*G. strictiflora*) in Italy.
- GOMESA.** See Orchidaceae.
- GOMPHOCARPUS.** Perennial herbs or subshrubs.
Puccinia schlechteri P. Henn. Leaf rust on *G. schinzianus* in the Union of South Africa.
Septoria gomphocarpi P. Henn. Circular brown leaf spots on *G. fruticosus* in Italy and Australia.
- GOMPHOLBIUM.** Australian shrubs.
Cronartium jacksoniae P. Henn. See Jacksonia.
- GOMPHRENA.** GLOBE AMARANTH. Erect or prostrate herbs.
Cercospora gomphrenicola Speg. Indefinite pale-yellow leaf spots on *G. glauca* in Argentina.
Phyllosticta gomphrenae Sacc. and Speg. Large indefinite yellow leaf spots with dull-brown margins on *G. glauca* in Italy.
Puccinia stuckerti Speg. Black rust pustules on leaves of *Gomphrena* sp. in Argentina.
Septoria gomphrenae Sacc. and D. Sacc. Subcircular to angular gray-white leaf spots with reddish margins on *G. globosa* in Italy.
Uromyces bonariensis Speg. Leaf rust on *G. elegans* and *G. gracilis* in Argentina.
- GONGORA.** See Orchidaceae.

- GONOLOBUS. VINCETOXICUM** Ag. Trailing or climbing herbs or shrubs.
Septoria gonolobicola P. Henn. Circular white leaf spots on *G. stephanotrichus* in Germany.
- GOODENIA.** Australian herbs and shrubs.
Puccinia saccardoi Ludw. Golden to black rust pustules on leaves of *G. albiflora*, *G. amplexans*, *G. geniculata*, *G. glauca*, *G. hederacea*, *G. ovata*, and *G. pinnatifida* in Australia.
Synchytrium succisae De B. and Wor. On leaf blades and petioles of *Goodenia* sp. and *Succisa pratensis* in Australia, Russia, Scandinavia, Bohemia, and Germany.
- GOODIA.** Australian shrubs with pea-like flowers.
Aecidium soleniforme Berk. Leaf rust on *G. latifolia* in Australia.
- GOODYERA.** See Orchidaceae.
- GOSSYPIMUM. COTTON.** Herbs and shrubs.
Alternaria macrospora Zimm. Leaf spot on *Gossypium* sp. in India.
Ascochyta gossypii Syd. On *Gossypium* sp. in Russia. Also reported from Arkansas.
Bacterial. A bacterial disease of bolls of *G. hirsutum*, said to be distinct from that caused by *B. malvacearum*, is reported from India. The disease is insect carried, black shiny spots appearing on the surface of the bolls where the punctures are made. Immature bolls when infected turn black and are shed. In all cases infected bolls, seed, and lint are discolored, and often a slimy soft rot is present.
Cercospora gossypii Syd. On *Gossypium* sp. in Uganda and Brazil.
Club leaf (Cytosis). This disease, attacking cultivated *Gossypium* in China, is apparently one of the group of virus diseases. It produces a witches'-broom effect due to a shortening of the nodes and development of additional shoots. The leaves are small, crinkled, and often colored different shades of green, yellow, and red.
Doassansia gossypii Lagerh. Leaf smut of *Gossypium* sp. in Ecuador.
Fomes lamaeensis Murr. See Hevea.
Gibberella gossypina Avena. Causes a boll rot of *Gossypium* sp. in Brazil.
Internal boll disease. The disease is characterized by a brown staining of the lint in green, unopened bolls and is due to the growth of a number of species of fungi which are transmitted by sucking insects. Young bolls when infected drop prematurely, and in older bolls the lint is more or less completely rotted. The disease is serious in the British West Indies and Trinidad. Similar diseases occur in Egypt and other cotton-growing countries, including probably the United States.
Kuehneola gossypii (Lagerh.) Arth. (*K. desmum* B. and Br.) Powdery cinnamon-brown rust pustules on purplish or brown leaf spots on *G. acuminatum*, *G. arboreum*, *G. brasiliense*, *G. herbaceum*, *G. hirsutum*, *G. mexicanum*, *G. microcarpum*, and *G. peruvianum* in Cuba, Porto Rico, South America, the Philippines, India, Java, Ceylon, India, Indo-China, New Guinea, west Africa, and Trinidad. Also known from Florida.
Ovulariopsis gossypii Wakef. White powdery mildew on yellow or red irregular patches on leaves of *G. barbadense* in Barbados.
Peronospora gossypina Avena. Downy mildew causing rot of bolls of *Gossypium* sp. in Brazil.
Pestalotzia gossypii Hori. Ochraceous to brown blotches on leaves of *G. herbaceum* in Japan.
Phoma roumii Fron. On stems and lateral branches of *Gossypium* sp. in Dahomey and west Africa, causing the leaves to wither.
Phyllosticta malkoffii Bub. Large brown to gray-white leaf spots with brown margins on *G. herbaceum* in Bulgaria.
Phytophthora sp. Causes a soft rot of bolls of *Gossypium* (cult.) in the British West Indies. Several species are involved, including *P. faberi*.
Rhizoctonia sp. See Vigna.
Sporidesmium longipedicellatum Reich. Causes a leaf spot of *Gossypium* (cult.) in Egypt.
Stilbum nanum Mass. f. *gossypina* Avena. Causes a boll rot of *Gossypium* sp. in Brazil.
- GOUANIA.** Tropical shrubs.
Puccinia gouaniae Holw. Brown leaf rust on *G. lupuloides* and *G. polygama* in Porto Rico, Cuba, Costa Rica, Trinidad, Panama, Guatemala, Santo Domingo, and Uganda.
Puccinia invaginata Arth. and Johnst. Leaf rust on *G. lupuloides* and *G. polygama* in Porto Rico, Cuba, Guatemala, St. Croix, Trinidad, and South America.
Puccinia paraensis Diet. Cinnamon-brown rust pustules on leaves of *G. pyrifolia* in Brazil.
Uredo gouaniae Ell. and Kels. Leaf rust on *G. lupuloides* and *G. polygama* in St. Croix and Porto Rico.
- Uromyces gouaniae* Kern. Brown leaf rust on *G. lupuloides* in Guatemala.
- GRABOWSKIA.** Spiny tropical shrubs.
Puccinia paradoxopoda Speg. Dark-brown rust pustules on leaves of *G. obtusa* in Argentina.
Puccinia penningtonii Syd. Black leaf rust on *G. duplicata* in Argentina.
- GRAPTOPHYLLUM.** Tropical shrubs with variegated foliage.
Rosellinia bunodes B. and Br. See Citrus.
- GRATIOLA.** Low herbs.
Phyllosticta gratiolae Höll. Ochraceous leaf spots on *G. officinalis* in Hungary and Russia.
Septoria gratiolae Sacc. and Speg. Dull-brown leaf spots on *G. officinalis* in Italy.
- GREVILLEA.** SILK OAK. Timber and ornamental trees.
Dimerium orbiculatum McAlp. Numerous circular, then confluent, brown to black leaf spots on *G. victori* in Australia.
Diplodia cacaicola P. Henn. See Theobroma.
Fomes lamaeensis Murr. See Hevea.
Fusarium udum Butl. See Dianthus.
Phyllosticta grevilleae (Lév.) Sacc. Circular black stromata on leaves of *G. buxifolia* and *G. occidentalis* in Australia.
Rhizoctonia lamellifera Small. Infected trees show a browning and wilting of the foliage followed by a stag-headed condition and ultimate death. Blackened areas occur on the lower portions of the trunks from which there are resinous or gummy exudates. Diseased tissues are brown and filled with gum. The roots rot and may be partly covered with thin superficial fungus layers. Numerous small black sclerotia occur in the rotted wood, cortex, and bark, together with sheets of black, irregular fungus tissue. The disease attacks *G. robusta* in Uganda as well as *Cosuarina equisetifolia*, *Bixa orellana*, *Coffea robusta*, and *Thea* to a lesser extent.
- Rosellinia arcuata* Petch. See Thea.
- GREWIA.** Shrubs.
Aecidium warneckeianum P. Henn. Galls up to 20 centimeters long deforming branches of *G. carpinifolia* in Tanganyika.
Helminthosporium grewiae P. Henn. On leaves of *Grewia* sp. in the Congo.
Ravenelia atrides Syd. Yellow to black rust pustules on leaves of *G. caffra* and *G. occidentalis* in the Union of South Africa.
Rhytisma grewiae Kalchbr. Black applanate stromata on leaves of *G. occidentalis* in the Union of South Africa.

GREWIA—Continued.

Stigmatea grewiae P. Henn. Circular brown leaf spots on *G. caffra* in the Union of South Africa.

Uredo grewiae Pat. and Har. Leaf rust on *G. ferruginea* in Senegal.

Ustilago grewiae (Pass.) P. Henn. Smut sori in branches of *G. columnaris*, *G. microcarpa*, and *G. mollis* in Abyssinia, tropical Africa, and Ceylon.

GRIFFITHIA. Rubiaceae shrubs.

Endophyllum griffithiae (P. Henn.) Racib. Leaf rust on *G. fragrans*, *G. latifolia*, and *Randia scandens* in Java.

GRISLINIA. Shrubs with large, glossy laurel-like foliage.

Puccinia griseliniae Pazschke. Dark-brown rust pustules on small circular sunken leaf spots on *G. ruscifolia* in Brazil.

GROSSULARIA. GOOSEBERRY. See Ribes.**GUAZUMA**. Tropical trees.

Trabutia guazumae Chardon. (*Phyllachora guazumae* P. Henn.) Black circular stromata on leaves of *G. ulmifolia* in Brazil and Porto Rico.

GUETTARDA. Tropical and subtropical shrubs.

Septoria guettardae Garman. Large irregular dark-red leaf spots with white centers on *G. ocalifolia* in Porto Rico.

GUNNERA. Perennial herbs with large leaves used for ornamental effects.

Ovularia gunnerae Speg. Circular to angular purplish leaf spots on *G. chilensis* in Chile.

Phyllosticta gunnerae Speg. On leaves of *G. magellanica* in Patagonia.

GUZMANIA. Tropical American bromeliads.

Lembosia bromeliacearum Rehm. On leaves of *G. roezlii* in Brazil.

GYMNADENIA. See Orchidaceae.**GYMNOLOMIA**. Yellow-flowered herbs or shrubs.

Aecidium gymnomolmiae Mayor. Leaf rust on *G. quitensis* in Colombia.

Puccinia gymnomolmiae Arth. Powdery brown leaf rust on *G. brachypoda*, *G. patens*, *G. subflexuosa*, and *Hymenostephium microcephalum* in Mexico, Guatemala, and Costa Rica.

Puccinia semota Jack. and Holw. Dark-brown leaf rust on *G. subflexuosa* in Guatemala.

GYMNOSPORIA. Shrubs and small trees, often spiny.

Englerulaster gymnosporiae (P. Henn.) Th. On leaves of *Gymnosporia* sp. in the Union of South Africa.

Septoria gymnosporiae Syd. Small dark-brown leaf spots on *G. deflexa* in the Union of South Africa.

GYMNOSTACHYUM. White-flowered herbs from Ceylon.

Puccinia leochroma Syd. Leaf rust on *G. subcordatum* in the Philippines.

GYNERIUM. PAMPAS GRASS. Perennial grass.

Ustilago quitensis Lagh. Powdery dark olivaceous smut sori destroying panicles of *Cortaderia argentea* (*G. argenteum*) in Ecuador.

GYNURA. Tropical herbs with showy foliage.

Aecidium gynurae Petch. Leaf rust on *G. lycopersicifolia* in Ceylon.

Gyropsora gynurae Petch. Leaf rust on *G. lycopersicifolia* in Ceylon.

GYPSOPHILA. BABY'S BREATH. Herbs with a profusion of small flowers used in borders.

Mycosphaerella gypsophilae Fckl. On leaves and stems of *G. gmelini*, *G. muralis*, and *G. scleranthus* in Siberia and Germany.

Septoria lychnidis Lasch. See Lychnis.

Uromyces gypsophilae Cke. Black powdery rust pustules on leaves and stems of *G. anatolica*, *G. haussknechtii*, *G. paniculata*, *G. polyclada*, *G. pulchra*, and *Saponaria vaccaria* in Persia, Kurdistan, and Russia.

HABENARIA. Terrestrial leafy orchids. See Orchidaceae.**HABRANTHUS**. Showy bulbous plants. Generally included with *Hippeastrum*.

Puccinia habranthii Diet. and Neg. Rust sori, round to irregular, yellow, becoming dark-brown or black in circular spots with purple margins. On both surfaces of leaves of *H. andicola* in Chile.

Puccinia reichelii Diet. and Neg. Yellow to brown rust pustules on both surfaces of leaves of *H. chilensis* in Chile.

HAEMODORUM. Perennial Australian herbs.

Puccinia haemodori P. Henn. Black rust pustules on leaves of *Haemodorum* sp. in Australia.

HAEMANTHUS. BLOOD LILY. African bulbous plants.

Cercospora haemanthi Kalch. Large, elliptical brown spots on leaves of *H. punicatus* in the Union of South Africa.

HAKEA. Australian ornamental evergreen shrubs.

Phyllachora hakeae P. Henn. Black stromata on both leaf surfaces of *H. myrtoides* in Australia.

Uredo angiosperma Thuem. Powdery brown rust pustules on leaves of *Hakea* sp. in Australia.

HALLERIA. Shrubs or small trees.

Septoria halleriae P. Henn. Pale, effuse, zoned leaf spots on *H. lucida* in Italy and Germany.

Septoria hallercola Sacc. and D. Sacc. On leaves of *H. lucida* in Portugal.

HAMMAMELIS. WITCH-HAZEL. Ornamental woody plants with yellow flowers late in fall or in the winter.

Aecidium hammamelidis P. Magn. Rust pustules on purplish leaf spots on *H. japonica* in Japan.

HAMELIA. Tropical ornamental woody plants grown for their flowers and berries.

Uredo hameliae Arth. Numerous light-brown powdery rust pustules on lower leaf surfaces of *H. crecta* in Porto Rico and Costa Rica.

HARDENBERGIA. Vines grown for their handsome flowers. See also Kennedya.

Phyllosticta hardenbergiae Cke. and Mass. Irregular dull-brown leaf spots on *Hardenbergia* sp. in Australia.

Septoria hardenbergiae Sacc. Pale leaf spots with dull-brown margins on *Kennedya monophylla* (*H. monophylla*) in Australia.

Uromyces hardenbergiae McAlp. Brown rust pustules on lower leaf surfaces of *H. monophylla* in Australia.

HEDERA. Ivy. Ornamental woody vines, grown for their handsome persisting foliage.

Ascochyta diplodina Berl. and Bres. Pale, irregular, confluent leaf spots on *H. colchica* and *H. helix* in Russia.

Bacillus sp. A bacterial disease on *H. helix* in France, producing brown patches on the leaves and killing-back the young growth.

Bacterium hederæ Arn. Transparent leaf spots on *H. helix* in France.

Colletotrichum hedericola Laubert. Anthracnose on leaves of *H. helix* in Austria.

Leptosphaeria hedericola (Desm.) Sacc. On leaves of *H. helix* in France.

Leptosphaeria hellicola (Desm.) Niessl. On leaves of *Hedera* sp. in France.

Macrophoma cylindrospora (Desm.) Berl. and Vogl. On leaf blades and petioles of *Euonymus japonicus*, *Hedera helix*, *Quercus ilex*, and *Vinea* sp. in France, Belgium, Italy, Austria, and Germany.

HEDERA—Continued.

- Macrophoma ulcinjensis** Bub. Large yellow to dull-gray leaf spots on *H. helix* in Yugoslavia.
- Meliola dichotoma** B. and C. Black superficial fungus patches on leaves of *Hedera* sp. in Japan.
- Meliola kusanoi** P. Henn. Circular black fungus patches on leaves of *Hedera* sp. in Japan.
- Mycosphaerella hederae-helicis** Siem. On leaves of *H. helix* in Russia.
- Mycosphaerella hedericola** (Desm.) Lind. Subcircular to irregular brown leaf spots on *H. colchica* and *H. helix* in Italy, Yugoslavia, Russia, Great Britain, Denmark, and France.
- Phyllosticta belgradensis** Bub. and Ranj. Irregular, often marginal, gray to ashen leaf spots with brown margins on *H. helix* in Yugoslavia and Spain.
- Phyllosticta destructiva-hederae** Desm. On leaves of *Hedera* sp. in France.
- Phyllosticta hederae** (Arc.) Allesch. On leaves of *H. helix* in Germany.
- Phyllosticta hederae** Sacc. and Roum. On leaves of *H. helix* in Europe.
- Phyllosticta hedericola** Dur. and Mont. Subcircular whitish leaf spots with brown margins on leaves of *H. colchica* and *H. helix* in Algeria and Europe. Reported from New Jersey.
- Phyllosticta milenae** Bub. Large irregular gray to dull-yellow leaf spots on *H. helix* in Bulgaria and Yugoslavia.
- Phyllosticta ulcinjensis** Bub. Large yellowish-white to greenish leaf spots with narrow brown margins on *H. helix* in Yugoslavia.
- Septoria desmazieri** Sacc. On leaves of *H. helix* in France.
- Septoria hederae** Desm. On leaves of *H. colchica* and *H. helix* in Europe.
- Septoria insularis** B. A. Br. Large brown leaf spots on *H. helix* in Germany.
- Septoria mortolensis** Penz. and Sacc. See *Acacia*.
- Triphragmium thwaitesii** Berk. and Br. See *Acanthopanax*.
- HEDYCHUM**. GINGER LILY. Rhizomatous herbs allied to *Zingiber*.
- Taphrina maculans** Butl. See *Curcuma*.
- HEDYSARUM**. Herbaceous perennials.
- Anthostomella sullae** Montem. On leaves of *H. coronarium* in Italy.
- Cercospora ariminensis** Cav. Circular to ovate or elliptical dull chestnut-brown leaf spots on *H. coronarium* in Italy and Brazil.
- Erysiphe taurica** Lévy. See *Althaea*.
- Placosphaeria onobrychidis** Sacc. See *Lupinus*.
- Ustilago kusanoi** Syd. Smut on *H. esculentum* in Japan.
- HELIANTHEMUM**. SUN ROSE. Ornamental woody or herbaceous plants grown for their showy flowers.
- Cercospora cistinearum** Sacc. Subcircular gray leaf spots with red-purple margins on *H. vulgare* in Italy.
- Cercospora helianthemii** Cav. Small circular gray-brown leaf spots on *H. polifolium* in Italy.
- Erysiphe taurica** Lévy. See *Althaea*.
- Peronospora alpestris** Gäum. Downy mildew on leaves of *H. alpestre* and *H. niloticum* in Algeria and Switzerland.
- Peronospora leptoclada** Sacc. Dirty-white areas on lower leaf surfaces of *H. guttatum* and *H. vulgare* in France, Italy, and Switzerland.
- Phyllosticta helianthemii** Roum. and var. **major** P. Brun. Subcircular white leaf spots on *H. vulgare* in France.
- Phyllosticta helianthemicola** Allesch. and var. **marginatum** Syd. Irregular dull-yellow to gray indefinite leaf spots on *H. vulgare* in Great Britain and Germany.
- Septoria chamaecistis** Vestergr. Brown, then dirty-white, leaf spots with dull-brown borders on *H. chamaecistus* and *H. vulgare* in Sweden.
- Septoria helianthemii** Gz. Frag. On leaves of *H. montanum* in Spain.
- Uredo helianthemii** Rbh. Yellow to brown rust pustules on leaves of *Cistus* sp. and *Helianthemum guttatum* in Italy.
- HELIANTHUS**. SUNFLOWER; JERUSALEM ARTICHOKE. Hardy herbaceous annuals and perennials.
- Mycosphaerella compositarum** (Auersw.) Schroet. On leaves of *H. annuus* in Yugoslavia.
- Puccinia minuscula** Arth. Brown leaf rust on *H. hypargyreus* in Ecuador.
- HELICHRYSUM**. EVERLASTING. Shrubs or herbs, some species grown for their flowers.
- Entyloma ascheronii** (Ule.) Wor. A smut producing swellings on the stems and roots and causing young shoots to die-back, of *Gnaphalium* sp. and *Helichrysum arenarium* in Germany.
- Puccinia kalchbrenneri** Del. Yellow to brown rust pustules on leaves of *H. globosum*, *H. nudatum*, *H. nudifolium*, and *H. undulatum* in Australia and south and central Africa.
- Puccinia mac-owani** Wint. Yellow to fuscous rust pustules on leaves of *H. petiolatum* and *Helichrysum* sp. in the Union of South Africa.
- Septoria helichrysi** Syd. Irregular, then confluent, brown leaf spots on *Helichrysum* sp. in the Union of South Africa.
- Uromyces helichrysi** Lagh. Dark-brown rust pustules in sunken purple spots on leaves and stems of *H. stoechas* in France.
- HELICONIA**. BIHAI Ag. A foliage plant allied to *Musa*.
- Pyrenobotrys heliconiae** (P. Henn.) Theiss. and Syd. Black stromata on brown to black irregular leaf spots on *Heliconia* sp. in Brazil.
- Puccinia heliconiae** (Diet.) Arth. Brown powdery rust pustules on leaves of *H. borinquena*, *H. latispatha*, *H. psittacorum*, and *Heliconia* sp. in Porto Rico, Martinique, Brazil, Panama, and Trinidad.
- HELIOPSIS**. Hardy herbaceous plants cultivated for their profuse yellow flowers.
- Aecidium heliopsisidis** Mayor. Rust on leaves of *H. euphthalmoides* in Colombia.
- Puccinia bimbergii** Mayor. Rust on leaves of *H. euphthalmoides* in Colombia.
- HELIOTROPIMUM**. HELIOTROPE. Herbaceous perennials cultivated for their fragrant flowers.
- Aecidium caspicum** Jacz. Rust pustules on large circular yellow to purplish leaf spots on *H. europaeum* in Russia.
- Aecidium guatemalense** Kern. and Kellerm. Rust pustules on discolored areas on leaves of *H. indicum* in Guatemala.
- Cercospora heliotropii-boconii** Scalia. Subcircular, then confluent and irregular, dirty-yellow to gray leaf spots on *H. boconii* in Sicily.
- Cercospora taurica** Tranzsch. Large gray leaf spots on *H. europaeum* in Russia and Yugoslavia.
- Puccinia gilva** Arth. and Holw. Leaf rust on *H. physocalycium* in Guatemala.
- Puccinia heliotropii** Kern. and Kellerm. Powdery chestnut-brown rust pustules on leaves of *H. indicum* in Guatemala.
- Puccinia heliotropicola** Speg. Powdery dull-brown rust pustules on leaves of *H. campestris* in Argentina.
- Uromyces heliotropii** Sred. Cinnamon to chestnut-brown rust pustules on leaves of *H. europaeum* in Russia, France, and Bulgaria.

- HELLEBORUS.** HELLEBORE. Christmas rose. Hardy herbaceous perennials cultivated for their flowers and attractive leaves.
- Aecidium hellebori** Ed. Fisch. Rust on leaves of *H. viridis* in Spain and Switzerland.
- Ascochyta hellebori** Sacc. Angular whitish leaf spots with dark borders on *H. viridis* in Italy.
- Coniothyrium hellebori** Cke. and Mass. Large irregular brown leaf spots without definite margins on *H. corsicus*, *H. foetidus*, *H. niger*, *H. olympicus*, and *H. viridis* in Great Britain, France, Corsica, Italy, Switzerland, Denmark, Albania, Yugoslavia, Austria, and Germany. Reported from Oregon.
- Peronospora pulveracea** Fckl. Gray mold on leaves of *H. foetidus*, *H. niger*, *H. odoratus*, and *H. viridis* in Yugoslavia, France, Switzerland, Denmark, Austria, and Germany.
- Phyllosticta atro-zonata** Voss. Circular ashen-colored leaf spots with black centers on *H. altifolius* and *H. viridis* in Austria.
- Phyllosticta helleboriana** Brun. Small circular gray leaf spots with brown margins on *H. foetidus* in France.
- Phyllosticta helleborella** Sacc. (*Mycosphaerella hermione* Sacc.) Large, black, then lighter, angular to irregular blotches without distinct margins on upper surfaces of leaves of *H. foetidus*, *H. niger*, and *H. viridis* in Great Britain, Italy, and Austria.
- Phyllosticta helleboricola** C. Mass. Pale angular leaf spots on *H. odoratus* and *H. viridis* in Spain, Italy, and Bulgaria.
- Phyllostictella olympica** (Allesch.) Tass. On leaves of *H. olympicus* in Germany.
- Ramularia hellebori** Fckl. Small circular whitish spots with black or purple margins on bracts and floral organs of *H. foetidus* and *H. viridis* throughout Europe.
- Ramularia recognita** Massal. On leaves of *H. multifidus* and *H. viridis* in Spain and Austria.
- Ramularia nigricans** (Mass.) Ferr. On leaves of *H. foetidus* in Spain.
- HELYGIA.** See *Parsonsia*.
- HEMEROCALLIS.** DAY LILY. Yellow and orange-flowered, stout-rooted perennials with basal leaves.
- Colletotrichum liliacearum** Fen. Anthracnose on leaves of *H. fulva* in Italy.
- Mycosphaerella hemerocallidis** (Pass.) Lind. On leaves of *H. fulva* in Yugoslavia.
- Puccinia hemerocallidis** Thuem. Pale-yellow rust pustules, becoming darker, on sunken yellow leaf spots on *H. aurantiaca*, *H. dumortieri*, *H. lilio-asphodelus*, (*H. flava*), *H. fulva*, and *H. midendorffii* in Siberia and Japan.
- Sporotrichum ochraceum** (Cda.) Sacc. On stems of *Hemerocallis* sp. and *Allium* sp. in Bohemia.
- HEMIGRAPHIS.** Diffuse or prostrate herbs grown for their foliage and flowers.
- Uromyces macintirianus** Barcl. Black rust pustules on leaves of *H. latebrosa* in India.
- HEPATICA.** Stemless low perennials.
- Aphelenchus olesistus** Ritz. Bos. See *Begonia*.
- Ascochyta molleriana** Wint. On leaves of *H. nobilis* (*H. triloba*) in Hungary.
- Ascochyta vodakii** Bub. On leaves of *H. nobilis* in Bohemia.
- Phyllosticta hepaticae** Brun. Leaf spots on *H. triloba* in France.
- Septoria hepaticae** Desm. Brown to black circular to irregular, then confluent, leaf spots on *H. triloba* in Belgium, France, Denmark, Russia, Switzerland, Italy, Spain, Austria, and Germany.
- HERACLEUM.** Cow PARSNIP. Perennial or biennial herbs.
- Ascochyta heraclei** Rens. Deep-brown, circular, often confluent, leaf spots on *H. sibiricum* and *H. spondylium* in Bulgaria and Germany.
- Mycosphaerella caulicola** (Karst.) Lind. On stems of *Chamaenerium angustifolium*, *Heracleum spondylium*, and *Telekia speciosa* in Yugoslavia, Denmark, and Finland.
- Protomyces macrosporus** Unger. See *Coriandrum*.
- Puccinia cortei** Ranj. Leaf rust on *H. minimum* in Europe.
- Puccinia heraclei** Grev. Chestnut-brown to deep-brown rust pustules on leaf blades and petioles of *H. minimum*, *H. sibiricum*, and *H. spondylium* in Europe.
- Septoria heracleicola** Kab. and Bub. Small circular brown leaf spots, becoming whitish, on *H. sibiricum* in Italy and Austria.
- Septoria heraclei-palmati** R. Maire. Yellow, then brown, confluent leaf spots on *H. palmatum* in France.
- Taphridium umbelliferarum** (Rostr.) Lagerh. and Juel. Irregular gray leaf spots on leaves of *H. montanum*, *H. sibiricum*, *H. spondylium*, *Oreoselinum* sp., and *Peucedanum palustre* in Europe.
- HERMANNIA.** Shrubs.
- Septoria hermanniae** F. Tassi. Brown marginal leaf spots on *H. micans* in Italy.
- HERNARIA.** BURSTWORT. Trailing small-leaved herbs.
- Peronospora herniariae** De By. Downy mildew on leaves and stems of *H. glabra* and *H. hirsuta* in Sweden, France, Russia, and Germany.
- Puccinia herniariae** Unger. Powdery red-brown to dark-brown rust pustules on leaves and stems of *H. glabra*, *H. hirsuta*, *H. lenticulata*, and *H. odorata* in Europe.
- Puccinia montagnei** Diet. Dark-brown rust pustules on leaves of *H. glabra*, *H. latifolia*, and *H. maritima* in France and Portugal.
- HERPESTIS.** Herbs, sometimes cultivated.
- Albugo quadrata** Kalch. and Cke. Small white sori (spore masses) on leaves of *H. verticillaris* in the Union of South Africa.
- HESPERETHUSA.** Naibel. Spiny trees or shrubs related to *Citrus*.
- Bacterium citri** Hassé. See *Citrus*.
- HEPERIS.** Dames rocket. Biennial herbs.
- Phyllosticta hesperidiarum** (Catt.) Pez. Circular to irregular brown, then dull-brown or ashen, leaf spots on *H. matronalis* in Italy.
- Ramularia matronalis** Sacc. On leaves of *H. matronalis* in France.
- HETEROPOPPUS.** Hardy herbaceous perennials with blue aster-like flowers.
- Uredo heteropappi** P. Henn. Brown rust pustules on circular dull-brown leaf spots on *H. hispidus* in Japan.
- HEUCHERA.** ALUMROOT. Herbaceous perennials.
- Aphelenchus olesistus** Ritz. Bos. See *Begonia*.
- Phyllosticta heucherae** P. Brun. Circular to irregular gray leaf spots on *H. glabra* and *H. sanguinea* in France and Great Britain.
- HEVEA.** PARA RUBBER TREE. Tropical trees furnishing rubber. This account of the diseases of *Hevea brasiliensis* is drawn for the most part from "The Diseases and Pests of the Rubber Tree," by T. Petch, of Ceylon.
- Apophthecia ulei** P. Henn. Yellow to dark-brown, circular or confluent, leaf spots on *Hevea* sp. in Peru.
- Ascochyta heveae** Petch. Marginal white or brownish-white leaf spots with red-brown borders on *H. brasiliensis* in Ceylon.
- Catacauma huberi** (P. Henn.) Theiss. and Syd. Tar-spot disease, characterized by shining black crusts up to 1 centimeter in diameter on the lower leaf surfaces, with corresponding pale-green patches above. On *H. brasiliensis* in Brazil.

HEVEA—Continued.

Cercospora heveae Vinc. Leaf spot on *H. brasiliensis* in Brazil.

Colletotrichum heveae Petch. Irregular spots to large patches, dry, gray brown, with purple-brown margins on old leaves of *H. brasiliensis* in Ceylon and Brazil. Diseased areas often fall out, producing a shot-hole effect.

Coniothyrium sp. Produces a canker on *H. brasiliensis* in Uganda.

Corticium koleroga (Cke.) V. Hoehn. See *Coffea*.

Corticium salmonicolor B. and Br. See *Citrus*.

Diplodia cacaicola P. Henn. See *Theobroma*.

Dothidella ulai P. Henn. (*Melanopsammopsis heveae* Stahel.) (*Fusicladium macrosporum* Kuyper.).

This disease, the so-called South American leaf disease, is the most serious rubber disease in the Western Hemisphere. It has made rubber culture unprofitable in Surinam and has caused heavy losses in British Guiana, Trinidad, Brazil, and Peru. Other species of *Hevea* (*H. confusa* and *H. guyanensis*) are attacked, in addition to *H. brasiliensis*.

The disease attacks the young leaves within a few days after their unfolding, causing translucent olive-green to blackish-green spots, which may be so numerous that the leaves blacken and shrivel. Young trees are particularly susceptible, the disease destroying the tops. In some cases the diseased areas dry up and fall out, producing a shot-hole effect. This type of injury is most prevalent in old trees. Petioles and young green twigs are also attacked, swollen areas resulting, which later on crack and form cankers. The fungus produces a velvety mycelial layer on the diseased areas. Other stages develop on old leaves. No method of controlling this disease has yet been found.

Fomes lamaoensis Murr. (*Hymenochaete noxia* Berk.). The brown root disease is the most widely spread root disease of economic plants in the eastern tropics. The principal hosts are *Albizia stipitata*, *Annona muricata*, *Artocarpus communis*, *A. integrifolia*, *Brownea grandiceps*, *Brunsfelsia Americana*, *Castilla elastica*, *Cedrela odorata*, *Cinnamomum camphora*, *C. cassia*, *C. iners*, *C. zeylanicum*, *Codiaeum variegatum*, *Coffea arabica*, *C. robusta*, *C. stenophylla*, *Cocos nucifera*, *Dalbergia latifolia*, *Eriobotrya japonica*, *Erythrina* sp., *Erythroxylon coca*, *Ficus glomerata*, *F. infectoria*, *Puntumia elastica*, *Garuga pinnata*, *Gossypium* spp., *Grevillea robusta*, *Hevea brasiliensis*, *Hura crepitans*, *Jacaranda mimosaefolia*, *Manihot* spp., *Mysticla magnifica*, *Persea gratissima*, *Poinciana regia*, *Soyndia felrifuga*, *Spondias wightii*, *Terminalia bellerica*, *Thea* (cult.), *Theobroma cacao*, and *Thespesia populnea* in India, Ceylon, Java, Sumatra, Malaya, Samoa, New Guinea, east and west Africa, British Guiana, and possibly Brazil.

The disease makes slow progress, one tree dying at a time. The roots of diseased trees are incrustated with a mass of sand, earth, and stones, intermingled with and held together by brown mycelium. The mycelium later becomes black. The bark beneath is brown, or brown mottled with white patches. The wood may be soft and friable, with a network of fine brown lines composed of hyphae, or the rot may be comparatively hard and transversely by rather broad brown bands. In advanced stages the wood is honeycombed. The hard, purple-brown sporophore rarely develops. The minute velvety, sterile brown patches sometimes found ringing the stems of diseased plants at the crown have caused the fungus to be determined as *Hymenochaete noxia*, a conclusion shown by Petch to be erroneous.

Fomes lignosus Klotzsch. This fungus causes the most serious losses and is the most widely distributed of all the root diseases of rubber. It occurs in India, Ceylon, Java, Sumatra, Borneo, west Africa, the Congo, and Brazil, attacking in addition to *H. brasiliensis*, *Azela palembanica*, *Artocarpus integrifolia*, *Berrya ammonilla*, *Bombax malabaricum*, *Cinnamomum camphora*, *Cocos nucifera*, *Coffea liberica*, *C. robusta*, *Dendrocalamus giganteus*, *Derris dalbergioides*, *Erythrina umbrosa*, *Ficus* spp., *Livistona cochin-chinensis*, *Koompassia malaccensis*, *Manihot glaziovii*, *M. utilisissima*, *Oncosperma filamentosum*, *Shorea* sp., *Thea* (cult.), *Theobroma cacao*, and many other woody plants.

The fungus spreads from dead roots and old stumps to the living roots of the various hosts, either where they come in contact or by growing out through the soil. As a rule, smooth white, or yellowish-white to reddish rhizomorphs are produced on the exterior of diseased roots. These run longitudinally along the roots, uniting here and there to form a network. The younger growing parts of the fungus consist of a fan-shaped mycelium. The fungus penetrates both bark and wood, causing a soft friable rot, without distinguishing characteristics. Occasionally the fungus develops internally only, producing a soft watery decay without external rhizomorphs. The fruiting body is red brown above with a bright-yellow margin and bright-orange lower surface. The colors become duller with age.

Fomes pseudo-ferreus Wakef. (*Poria hypolateritia* Berk.). This fungus produces a wet rot of roots of *H. brasiliensis*, *Albizia* sp., *Symplocos spicata*, *Tephrosia* sp., *Thea* (cult.), and *Theobroma cacao* in Malaya, Ceylon, and Java. A dark-red superficial mycelium is sometimes formed over diseased roots, or occasionally snow-white patches of mycelium on small roots, but no rhizomorphs are produced. Workers in Java refer this species to *Fomes* (*Ganoderma*) *ferreus* Berk.

Fusicladium sp. A die-back disease, starting on pruned branches, has been reported from Java on *H. brasiliensis*, but is considered by Petch identical with the disease caused by *Diplodia*.

Gloeosporium alborubrum Petch. This fungus is one of the causes of abnormal leaf fall of *H. brasiliensis* in Ceylon, Sumatra, Malaya, and Uganda. On young leaves large, irregular, blackish-green, watery-looking patches occur. Older leaves turn yellow, then brown, black spots with gray centers appearing on veins and midribs. Petioles turn dark brown or black and the leaves fall. Young twigs may also be affected, dying back. Diseased fruit rot and finally become greenish gray and wrinkled.

Gloeosporium heveae Petch. This anthracnose fungus attacks young plants of *H. brasiliensis* in the nurseries of Ceylon and Brazil. The leaves turn yellow green, then yellow, and fall. The fruiting pustules appear on both leaf surfaces.

Guignardia heveae Syd. Tips and margins of leaves become pale brown with purple-brown margins. On *H. brasiliensis* in Ceylon, Malaya, and the Philippines.

Helicobasidium mompia Tan. See *Morus*.

Helminthosporium heveae Petch. The "bird's-eye spot" disease of *H. brasiliensis* attacks for the most part small plants in the nurseries, but is occasionally found on older trees. The spots are at first purple and very small, then increasing in size and becoming white and semitransparent, with narrow purple-brown borders. The disease occurs in Ceylon, India, Malaya, Java, Sumatra, the Philippines, and Gold Coast.

Hymenochaete noxia Berk. See *Fomes lamaoensis* Murr.

Marasmius spp. See *Theobroma*.

Meliola heveae Vinc. Small black superficial fungus patches on leaves of *H. brasiliensis* in Brazil.

Mycolphaerella heveae Petch. Small yellow leaf spots on *H. brasiliensis* in Ceylon, which become purple or purplish brown in color and confluent.

Ophiobolus heveae P. Henn. Circular to effuse and confluent gray leaf spots on *Hevea* sp. in Brazil.

Phacophomopsis heveae Grove. Causes a bark disease of *H. brasiliensis* in Uganda.

HEVEA—Continued.

Phyllosticta heveae Zimm. Marginal or apical brown patches on leaves of *H. brasiliensis* in Java, Trinidad, Brazil, and Malaya. Found on young trees in shaded nurseries.

Phyllosticta ramicola Petch. This fungus attacks young twigs from 6 to 18 inches back from the tip. Diseased areas are dark brown, progressing up and down from the point of infection. On *H. brasiliensis* in Ceylon, Java, and Malaya.

Phytophthora faberi Maubl. This fungus attacks practically all parts of the rubber tree, causing the diseases known as "abnormal leaf fall," pod or fruit disease, die-back of shoots, and "claret-colored canker" or "patch canker." These diseases are prevalent in India, Ceylon, Java, Malaya, Fiji, and other rubber-growing countries.

Diseased fruits are first watery green, then black and soft. They remain hanging on the tree. The fungus passes from the fruit to the leaves and twigs, causing an abnormal leaf fall and dying back of young twigs. The leaves may first turn yellow or mottled, or they may fall while still green. The leaflets often separate from the petioles. In the case of the trunk canker no external symptoms are in evidence other than a slight exudation of a rusty-colored liquid. The layer beneath the outer brown bark is black, and the cortex is first yellow gray, then claret colored. The cankered areas are surrounded by black lines. The disease may spread around the trunk and so destroy the tree. See also *Theobroma* for additional hosts and geographical distribution.

Phytophthora meadii McRae. This species causes fruit rot, leaf fall, and black-thread or black-stripe disease. The fruit rot and leaf fall caused by this fungus bring about heavy reductions in latex yield, which is not the case with the preceding species. The symptoms are similar, however. The black-thread phase of this *Phytophthora* disease occurs on the tapped surface, beginning as a series of narrow, vertical, parallel black lines, which may broaden out and coalesce so as to form a continuous wound parallel to the tapping cut. These black lines may extend through to the wood. In addition to *H. brasiliensis*, *Theobroma cacao* is attacked by this species. It is reported from Ceylon, India, Java, and Malaya.

Poria hypobrunnea Petch. A root disease of *H. brasiliensis*, *Crotalaria fulva*, *Hibiscus* sp., *Panax* sp., *Tephrosia candida*, and *Thea* (cult.) in Ceylon and Java. The mycelium forms stout red strands on the exterior of the root, which may unite into a continuous red sheet. These rhizomorphs are smooth and tough, bright red to brownish red, white within, turning black with age. Diseased wood is soft, friable, and permeated with red mycelial sheets, which often follow the lines of the annual rings. In advanced cases a soft wet rot is present, with a network of white threads between the bark and wood. The fruiting body is first yellowish white or ochraceous, then red brown, and finally dark-slate color.

Rosellinia bunodes B. and Br. See Citrus.

Sclerotium sp. Destroys bark on tapped surfaces of *H. brasiliensis* in Sumatra.

Scolecotrichum heveae Vinc. Small brown spots, the centers of which drop out on *H. brasiliensis* in Brazil.

Sphaeronema sp. Attacks the tapped surface of trunks of *H. brasiliensis* in Malaya, rotting and destroying the exposed cortex tissues and making further tapping for some years impossible.

Sphaerostilbe repens B. and Br. Black or red flattened fungous strands or rhizomorphs occur under the cortex of diseased roots and the wood is stained deep blue or purple. There is often a foul smell connected with the disease. Infected plants gradually die. Fruiting bodies of the fungus are of two types, the conidial occurring as short, erect, red, hairy stalks 2 to 8 millimeters high, surmounted by a pink head of conidia. The second consists of small, dark-red perithecia at the base of the conidial stalks. In addition to *H. brasiliensis* the disease attacks *Artocarpus integrifolia*, *Carica papaya*, *Casuarina equisetifolia*, *Citrus* spp., *Erythrina* spp., *Hibiscus* sp., *Mangifera indica*, *Maranta arundinacea*, and *Thea sinensis* in Ceylon, India, Malaya, Uganda, Fiji, Gold Coast, and the British West Indies.

Thread blight. A number of species of "thread blights" have been reported from India, Ceylon, Malaya, Java, and other tropical countries on *H. brasiliensis* and other hosts, some at least of which are parasitic. They are all characterized by white strands of mycelium, which run over stems and leaves of the hosts. This type of disease is described in detail for *Coffea*. (See *Corticium koleroga*.)

Top canker. This disease of *H. brasiliensis* in Ceylon is thought to be due to *Phytophthora faberi* and attacks the upper branches and stems. The bark splits longitudinally in lines a foot or more long, forming long, narrow scales which in time split transversely. The disease may penetrate to the wood, causing an open wound. The stems ultimately become thickened in the region of the cankers.

Venturia emergens Petch. On branches of *H. brasiliensis* in Ceylon.

White stem blight. This disease occurs in Ceylon on *H. brasiliensis* and *Thea* (cult.) and is caused by an undetermined fungus. The larger branches are attacked, white patches often 6 feet in length appearing on the bark. A fine white thread runs along the whitened patch, and the upper margin of the patch may show a fringe of hyphae.

Witches'-broom. A disease characterized by globose woody galls on the branches of *H. brasiliensis* is reported from Java. The cause has not yet been determined.

Xylaria thwaitesii Cke. This fungus appears as black, flat, irregular bands or patches on the roots of *H. brasiliensis* in Ceylon. A rot of the bark and wood results and diseased trees ultimately die. The rotten wood is dark brownish gray and moist but hard at the center of a root, light brown and decayed outside of this region. Black lines may be present. The inner cortex becomes brown or yellow brown and friable.

HIBBERTIA. Shrubs or subshrubs with yellow or white showy flowers.

Puccinia hibbertiae McAlp. Reddish-brown to black rust pustules on lower leaf surfaces of *H. sericea* and *H. stricta* in Australia.

HIBISCUS. ROSE MALLOW. Shrubs or small trees cultivated for their showy flowers and ornamental foliage.

Aecidium erythrobasis B. and Br. Rust pustules on red leaf spots on *H. collinus* and *H. eriocarpus* in Ceylon and Uganda.

Aecidium garckeanum P. Henn. Rust on leaves of *H. crassinervius* and *H. micranthus* in Abyssinia and the Congo.

Bacterium hibisci Nak. and Tak. Small circular black spots appear on the cotyledons and young leaves, enlarging gradually and becoming irregular to angular, with dull white to yellow outer zones. On *Hibiscus* sp. in Korea.

Ceroospora brachypoda Speg. On leaves of *Hibiscus* sp. in Argentina. Reported from Louisiana.

Ceroospora hibisci-manihotis P. Henn. Circular to effuse dull-brown spots on leaves of *H. manihot* in Japan.

Colletotrichum hibisci Pöhl. Circular to irregular brown leaf spots on *H. palustris* in Italy.

Colletotrichum hibiscicola Rangel. On leaves of *H. tiliaceus* in Brazil.

Corticium salmonicolor B. and Br. See Citrus.

Fomes lamaeensis Murr. See Hevea.

Phyllachora hibisci Rehm. Shiny-black stromata on indefinite brown leaf spots on *H. furcatus* and *H. tiliaceus* in Ceylon, Samoa, and Brazil.

HIBISCUS—Continued.

- Phyllachora minuta** P. Henn. Black stromata on leaves of *H. tiliaceus* and *Thespesia populnea* in Porto Rico, Java, Australia, and the Philippines.
Phyalospora hibisci Racib. On leaves of *H. tiliaceus* in Java.
Phytophthora melongenae K. Saw. See Solanum.
Poria hypobrunnea Petch. See Hevea.
Puccinia exilis Syd. var. *hibisci* Grove. Brown rust pustules on leaves of *Hibiscus* sp. in Uganda.
Puccinia hibisci P. Henn. Dull-brown rust pustules on leaves of *H. rostellatus* in the Congo.
Rhizoctonia destruens F. Tassi. See Solanum.
Rosellinia bunodes B. and Br. See Citrus.
Septoria hibisci Sacc. Subcircular to irregular whitish leaf spots with dull-brown margins on *H. syriacus* in Italy.
Septoria rosae-sinensis Brun. Olive-brown leaf spots on *H. rosa-sinensis* in France.
Sphaerostilbe repens B. and Br. See Hevea.
Tubercularia hibisci Petch. On leaves of *H. sabbdariffa* in Ceylon.
Uromyces heterogenens Cke. Brown rust pustules on leaves of *H. syriacus* in India.
Xylaria sp. This fungus causes a root disease of the ornamental *Hibiscus* in Hawaii.
Wilt. A wilt disease of *Hibiscus* spp., cause undetermined, is reported from Hawaii. Possibly the same as the preceding disease.

HIERACIUM. HAWKWEED. Hardy herbaceous perennials.

- Ascochyta hieracii** Lasch. On leaves of *H. pilosella* in Germany.
Cercospora hieracii Jaap. Large subcircular to angular brown leaf spots on *H. prenanthoides* in Switzerland.
Entyloma calendulae (Oud.) De By. See Calendula.
Entyloma hieracii Syd. Leaf smut on *Hieracium* spp. in Europe. A form of *E. calendulae*.
Phyllosticta coreontica Kab. and Bub. Subcircular to angular zoned yellow-brown to brown leaf spots becoming gray on *H. alpinum* in Austria.
Phyllosticta hieracicola E. Rostr. Dark-brown leaf spots on *H. umbellatum* in Norway.
Phyllosticta hieracii Lasch. Subcircular to irregular, often confluent, zoned dull yellow-brown leaf spots on *H. pilosella* and *H. pseudocerinthe* in Spain and Germany.
Puccinia argentina Speg. Cinnamon-brown rust pustules on leaves and stems of *Hieracium* sp. and *Picrisia longifolia* in Argentina.
Ramularia conspicua Syd. Circular olive-brown leaf spots on *H. murorum* in Bohemia and Germany.
Ramularia coreontica Bub. and Kab. Subcircular to angular dull-yellow to brown leaf spots becoming gray on *H. alpinum* and varieties in Bohemia.
Ramularia helvetica Jaap. and Lind. Irregular greenish-gray or brown leaf spots on *H. albidum* in Switzerland.
Ramularia hieracii (Baum.) Jaap. On leaves of *H. boreale*, *H. schmidtii*, *H. sylvaticum*, *H. umbellatum*, and *H. vulgatum* in France, Switzerland, Austria, and Germany.
Ramularia subalpina Bub. Angular dirty-yellow leaf spots with narrow, dark-brown margins on *H. lanatum* in Yugoslavia.
Septoria guardaromica Gz. Frag. On leaves of *H. pilosella* in Spain.
Septoria mougeoti Sacc. and Roum. Large yellow leaf spots on *H. praealtum* and *Hieracium* sp. in France and Russia.
Septoria palustris (Ces.) Sacc. On leaves of *Hieracium* sp. in Italy.
HIEROGLOA. TORRESIA Ag. Fragrant perennial grasses.
Puccinia borealis Juel See Agrostis.
Puccinia hierochloae Ito. Yellow to chestnut-brown rust pustules on leaves and sheaths of *H. borealis* and *Stipa siberica* in Japan.
Septoria oxyspora Penz. and Sacc. See Anthoxanthum.
HIPPEASTRUM. Showy bulbous plants.
Mosaic. An infectious mosaic, characterized by a mottling of the leaves of *Hippeastrum* spp., has been reported from Hawaii.
HIPPOCREPIS. Herbs or subshrubs.
Cercospora hippocrepidis Jaap. Subcircular gray to whitish leaf spots with brown margins on *H. comosa* in Switzerland.
Uromyces anthyllidis (Grev.) Schroet. See Anthyllis.
Uromyces hippocrepidis E. Mayor. Brown rust pustules on leaves of *H. ciliata* in Europe.
HIPPOPHAE. SEA BUCKTHORN. Ornamental woody plants grown for their silvery-gray foliage and bright-colored berries.
Dothidella hippophaeos (Pers.) Theiss. and Syd. Circular black stromata on leaves of *H. rhamnoides* in Italy and Switzerland.
Septocylindrium olivaceus Thuem. See Elaeagnus.
Septoria hippophaes Desm. and Rob. On leaves of *H. rhamnoides* in France.
HIPPURIS. Perennial aquatic herbs.
Physoderma hippuridis Rost. On leaves of *H. vulgaris* in northern Europe, Greenland, and Iceland.

HOHERIA. Small shrubs or trees, native of New Zealand.

- Puccinia hoheriae** Wakef. Brown rust on leaves and stems of *H. populnea* in New Zealand.
HOLCUS. SORGHUM. Johnson grass. Sudan grass. Annual or perennial grasses, cultivated for production of grain, forage, sirup, and broomcorn.
Acerothecium lunatum Wakker. On leaves and heads of *H. sorghum* in Ceylon and Java.
Aecidium evansi P. Henn. Rust on leaves of *H. sorghum* in the Union of South Africa.
Apiospora lloydii (Crouan.) Sacc. On *Holcus* sp. and *Dactylis* sp. in France.
Ascochyta sorghina Sacc. Long dull-brown leaf spots with blood-red margins on *H. sorghum* in Italy.
Bacillus oemlianskii Serbinov. Said to cause a soft black rot of tops of plants of *H. sorghum* in Russia.
Beniowskia penniseti Wakef. See Pennisetum.
Botryodiplodia sorghi P. Henn. On culms of *H. sorghum* in Tanganyika.
Cercospora acerosa Dickk. and Arendt. H. Black spots on leaves of *H. sorghum* in Java.
Cerebella sorghi-vulgaris Subra. In ovaries of *H. sorghum* in India.
Colletotrichum andropogonis A. Zimm. Dark-red leaf spots on leaves of *H. sorghum* in Tanganyika.
Dilophia graminis (Fekl.) Sacc. See Triticum.
Helminthosporium caryopsidium Sacc. Attacks the grain of various varieties of *H. sorghum*, causing "wholesale destruction of seed" in China and the Philippines.
Leptosphaeria eustonia (Fr.) Sacc. See Secale.

HOLCUS—Continued.

Microbasidium sorghi (Pass.) Bub. and Ran. Elliptical red to dark-brown leaf spots on *H. sorghum* var. *saccharatum* in Yugoslavia.

Mycosphaerella cerea Sacc. Pale leaf spots with reddish margins on *H. sorghum* in Russia and Italy.

Mycosphaerella tassiana De N. On leaves of *H. lanatus* in Austria.

Ovularia holci-lanati Cer. Small dull dark-brown leaf spots on *H. (Notholcus) lanatus* in Italy.

Phyllachora sorghi V. Hoch. Black stromata on yellow to white leaf spots with red margins on *H. sorghum* in var. in Java and the Philippines.

Phyllosticta sorghina Sacc. Irregular leaf spots with red borders on *H. effusum*, *H. halepensis*, and *H. sorghum* in Russia, India, Italy, Portugal, and Argentina. Reported from Texas.

Puccinia holcina Erikss. Yellow-brown to brown rust pustules on leaves of *H. lanatus*, *H. mollis*, and *H. setigermis* in Europe.

Ramulispora andropogonis Miuri. Leaf spot on *H. sorghum* in Manchuria.

Sclerospora graminicola (Sacc.) Schroet. var. **andropogonis sorghi** Kulk. This downy mildew attacks plants of *H. sorghum* in all stages of growth in India. The leaves of diseased seedlings are narrow, pale-yellow, and covered with a fine white down. Later on white streaks appear on the upper leaves and the tissues tear along these streaks. Diseased plants remain stunted and sterile. Older plants may also be attacked, singly or in groups, long, narrow streaks appearing on the leaves at first pale-yellow, then orange, and finally dark-brown in color.

Sclerospora philippinensis Weston. See Zea.

Septoria holci Pass. Small gray subcircular leaf spots on *H. (Notholcus) lanatus* in Italy.

Sorosporium ehrenbergii Kuhn. Smut sori in the ovaries, covered at first by a false membrane. On *H. cernuus* in Egypt and Tunis.

Sorosporium simii Pole-Evans. Smut attacking and destroying inflorescences of *H. halepense* in the Union of South Africa.

Tilletia holci (West.) Rostr. (*T. ravenhoffii* Fisch de W.) Irregular, black smut sori in the ovaries of *Bromus arvensis*, *B. mollis*, *Holcus (Notholcus)*, *lanatus* and *H. mollis* in Denmark, Belgium, and Great Britain. Reported from Washington and Oregon.

Tolyposporium filiferum Busse. Individual grains of the heads of *H. sorghum* in India, Egypt, and east Africa are transformed into smut sori. The sori are longer than the normal grain and light brown in color, soon rupturing, exposing the powdery spore masses.

Tolyposporium volkensii P. Henn. Black smut sori in the ovaries of *H. sorghum* in tropical Africa.

Ustilago bulgarica Bub. Greenish-brown smut sori in the ovaries of *H. sorghum* in Bulgaria.

Ustilago sorghicola Speg. Subcylindrical smut sori in the ovaries of *H. sorghum* in Argentina and Italy. The sori project beyond the glumes, reaching a length often of 8 millimeters.

HOMERIA. Bulbous plants from south Africa.

Uredo homeriae Bub. Rust on leaves of *Homeria* sp. in the Union of South Africa.

HOMOZYNE. Hardy perennials grown for their attractive white or purple heads of flowers.

Ascochyta homogyne Ranoj. Brown leaf spots on *H. alpina* in Yugoslavia.

Puccinia conglomerata (Strauss.) Kze. and Schm. Dark-brown rust pustules on leaves of *H. alpina* in France, Switzerland, Yugoslavia, Austria, and Germany.

Ramularia cervina Speg. On leaves of *H. alpina* in Yugoslavia, Italy, and Austria.

Septoria prostrata Kab. and Bub. Subcircular brown, then gray, leaf spots with dark-brown margins on *H. alpina* in Yugoslavia and Austria.

Uromyces veratri (DC.) Schroet. See *Veratrum*.

HONKENYA. See *Arenaria*.**HORDEUM.** BARLEY. SQUIRRELTAIL GRASS. Annual grasses.

Bacillus cerealinus Gentner. Brown spots or blotches appear on the nodes, upper internodes, and leaves of *H. vulgare* (*H. sativum*), *Secale cereale*, and *Triticum* sp. in Europe. The upper leaves become prematurely yellow and dry, the heads fill out poorly, and the glumes may split.

Cercospora elymi Rostr. Oblong dull-brown leaf spots on *H. (Elymus) arenarium* in Denmark.

Fusarium hordearium Duc. Attacks the leaves and stems of *H. distichon* in France.

Fusarium nivale (Fr.) Sor. See *Secale*.

Heterosporium hordei Bub. Rhombic or elliptic-oblong, often confluent, dull-yellow leaf spots with dark-purple margins on *H. distichum* and *H. sativum* in Denmark and Yugoslavia.

Micrococcus tritici Prill. See *Triticum*.

Mycosphaerella basicola Frank. On leaves of *H. sativum* and *Secale cereale* in Sweden, Denmark, and Germany.

Mycosphaerella exitialis Morini. On culms of *H. sativum* and *Triticum* sp. in Japan, France, Denmark, Sweden, and Italy.

Mycosphaerella hordicola Hara. On leaf blades and culms of *Hordeum* sp. and *Triticum* sp. in Japan.

Ophiobolus cariceti (B. and Br.) Sacc. See *Triticum*.

Ophiobolus herpotrichus (Fr.) Sacc. See *Triticum*.

Ramularia hordei McAlp. Subcircular, then confluent, brown to dark-brown leaf spots on *Hordeum* sp. in Australia.

Rhizoctonia napi West. See *Brassica*.

Septoria elymi Rostr. On leaves of *H. arenarium* in Denmark.

Septoria halophila Speg. Large, ashen-colored blotches on leaves of *H. halophilum* in Argentina.

Tilletia hordei Koern. Dark-green smut sori in the ovaries, concealed at first by the glumes, then black, with a foetid odor. On *H. fragile* and *H. murinum* in Turkey, Algeria, Persia, and Australia.

Tilletia hordeina Ranoj. Yellow-brown spore masses in ovaries of *H. maritimum* in Yugoslavia.

Tilletia panicli Bub. and Ranoj. Dark violet-brown smut sori in the ovaries of *H. sativum* in Yugoslavia.

Tilletia trabuti Jacz. A smut forming brown spore masses in the spikelets of *H. murinum* in Algeria.

Tylenchus hordei Schöyen. See *Avena*.

Typhula graminum Karst. See *Lolium*.

HOSTA. NIOBE. AR. PLANTAIN LILY. Hardy perennials.

Acidium hostae Diet. Rust pustules on yellow, circular leaf spots on *H. sieboldiana* in Japan.

Ascochyta herreana P. Henn. On leaves of *H. ovata* in Germany.

Ascochyta hortensis Kab. and Bub. Circular to angular brown leaf spots with purple-brown margins on *H. albomarginata* in Bohemia.

Cercospora hostae Hori. On leaves of *H. sieboldiana* in China.

Fusarium versiforme Kab. and Bub. Irregular brown spots on both leaf surfaces of *H. albomarginata* in Bohemia.

Mycosphaerella hostae Syd. On leaves and petioles of *H. japonica* in Japan.

Phyllosticta funkiae Fen. Pale irregular leaf spots with brown margins on *H. ovata* in Italy.

Puccinia funkiae Diet. Black rust pustules on the lower leaf surfaces of *H. longipes*, *H. ovata*, *H. plantaginea*, and *H. sieboldiana* in Japan and other Asiatic countries.

Uredo hostae P. Henn. Brown rust pustules on yellow leaf spots on *Hosta* sp. in Japan.

HOTTONIA. FEATHERFOIL. Perennial aquatic herbs.

Doassansia hottoniae Rostr. Small pustules in rufous leaf spots on *H. palustris* in Denmark.

HOVENIA. RAISIN TREE. Ornamental shrubs, grown for their handsome foliage.

Microspheera yamadai Syd. Powdery mildew on leaves of *H. dulcis* in Japan.

HOYA. WAX PLANT. Tropical climbers or trailing evergreen shrubs.

Gloeosporium affine Sacc. Anthracnose on leaves of *Aloe hanburyana*, *Citrus aurantium*, and *Hoya carnosa* in Great Britain, Italy, Gibraltar, and the Philippines.

Gloeosporium hoyae Syd. Anthracnose on leaves of *H. luzonensis* in the Philippines.

Gloeosporium intermedium Sacc. See *Citrus*.

Guignardia albicans Rehm. On leaves of *H. luzonensis* in the Philippines.

Phyllosticta asclepiadeorum West. See *Cynanchum*.

Phyllosticta thuemii Tass. On leaves of *H. carnosa* in Austria and Gibraltar.

Physalospora hoyae v. Hoehn. On leaves of *H. luzonensis* in the Philippines.

Puccinia variiformis Pat. Dark-brown rust pustules on leaves of *Hoya* sp. in Indo-China.

Septoria hoyae Sacc. Ivory-white irregular leaf spots with dull-brown margins on *H. carnosa* in Italy.

HUMULUS. HOP. Twining vines.

Ascochyta humuli Lasch. Pale-brown leaf spots on *H. lupulus* in Germany.

Ascochyta humuli Bub. and Kab. Circular to irregular greenish, then dirty-ochraceous to ashen, leaf spots on *H. lupulus* in Bohemia. This species if not synonymous with the preceding must be renamed.

Cercospora cantariensis Salm. and Worm. Circular gray leaf spots with narrow purple-brown margins and surrounding yellow zones on *H. lupulus* in Great Britain.

Cercospora humuli Hori. Angular dark-brown spots, often coalescing, on *H. japonicus* and *H. lupulus* in Japan.

Fusoma parasiticum Salm. This fungus causes a serious wilt disease of hops in Great Britain. Above ground its presence is shown by a wilting of the stems, which break away easily or fall over, having been rotted just below the ground line. The bark and wood at this point are brown and decayed. Pink pustules of the causative fungus appear on the basal parts of diseased plants.

Mosaic. A mosaic disease of *H. lupulus* is reported from Great Britain. The stems of diseased plants grow only to a height of 4 to 6 feet, the leaves are curled, more or less mottled, green and yellow in color, and very brittle. Diseased plants are sterile.

Peroplasmopara humuli Miy. and Tak. This serious downy mildew of the hop (*H. lupulus*) occurs in Japan and Great Britain. It has also been reported from Wisconsin. Small angular dark-green spots, with a water-soaked aspect, first appear on the leaves, becoming finally dark-brown. Diseased leaves dry up and die.

Phyllosticta bractearum Oud. On bracts of *H. lupulus* in Holland.

Phyllosticta japonica Fautr. Brown irregular leaf spots on *H. japonicus* in Japan and France.

Phyllosticta lupulina Kab. and Bub. Circular to angular, confluent, ashen-colored leaf spots with purple-brown borders on *H. lupulus* in Austria and Bohemia.

Plasmidiophora humuli Kirk. Causes a disease of *H. lupulus* in New Zealand and Tasmania which resembles clubroot of the crucifers.

Septoria divergens Bub. and Kab. Small subcircular to angular ochraceous-brown leaf spots with narrow brown margins on *H. lupulus* in Bohemia.

Septoria humuli West. On leaves of *H. lupulus* in Europe.

Septoria humulina Bond. Pale to dull-brown leaf spots on *H. lupulus* in Russia.

HURA. SAND-BOX TREE. Tropical trees.

Cercospora hurae Stevens. Small brown leaf spots on *H. crepitans* in Porto Rico.

Fomes lamaeensis Murr. See *Hevea*.

Helminthosporium hurae P. Henn. Circular, then confluent, dull-brown leaf spots on *H. crepitans* in Brazil.

HUTCHINSIA. Low, annual or perennial herbs.

Puccinia hutchinsiae Rud. Dark-brown rust pustules on leaves of *H. alba* in Turkestan.

HYACINTHUS. HYACINTH. Hardy bulbous plants cultivated for their showy flowers.

Ascochyta hyacinthi Tassi. On leaves of *Agapanthus umbellatus* and *H. orientalis* in Italy and Australia.

Bacillus croci Mizus. See *Crocus*.

Bacillus hyacinthus septicus Heinz. Bulbs, leaves, and flower stalks are attacked by this disease, which starts as a soft rot of the bulb, rapidly extending into the above-ground parts of the plant. Leaves and flower buds wither and drop off. Diseased tissues soon break up, forming a foul-smelling slime. The disease has been reported on cultivated *Hyacinthus* in Great Britain and Germany. By some workers this species is considered synonymous with *B. caratovorvus*.

Bacterium hyacinthi Wakke. The disease known as Wakker's hyacinth disease, or the yellow-slime disease, is characterized by yellow stripes beginning near the apex of the leaf and extending downward rapidly, but only slowly sideways, so that finally the leaf will have a central dead stripe along its entire length with the margins still green. In the bulb the vascular bundles become yellow and full of bacterial slime. The yellow slime extends upward into the bundles of other scales and sideways slowly into the parenchyma until finally the entire bulb is destroyed. In advanced stages other bacteria enter and aid in destroying infected bulbs. All varieties of *Hyacinthus* are attacked. The disease has been reported from France, Great Britain, Holland, Italy, and Germany. There is one report of its occurrence in the United States.

Periza vesiculosa Bull. This fungus is reported as attacking entire plants, with a growth of mold externally; in Germany on *Hyacinthus* in variety.

Puccinia illiacearum Duby. See *Ornithogalum*.

Septoria podolica Bub. and Wrobl. On leaves of *H. leucophaeus* in Galicia.

Sclerotinia tuliparum (Wakk.) Rehm. See *Tulipa*.

Tylenchus dipsaci Kuehn. See *Narcissus*.

Uromyces scillarum (Grev.) Wint. A common and widespread rust which produces brown pustules on both leaf surfaces, disfiguring the affected plants. The hosts are *Bellevalia comosa*, *B. fuliginosa*, *B. mauritanica*, *Endymium rutans*, *Hyacinthus ciliatus*, *H. fastigiatus*, *H. orientalis*, *H. pratensis*, *Leopoldia boviana*, *Muscari botryoides*, *M. comosum*, *M. monstrosum*, *M. neglectum*, *M. polyanthum*, *M. racemosum*, *M. tenuiflorum*, *Ornithogalum comosum*, *Scilla autumnalis*, *S. bifolia*, *S. bertolonii*, *S. campanulata*, *S. italica*, *S. obscurifolia*, *S. peruviana*, *S. pratensis*, *S. pumila*, *S. ramburii*, *S. verna*, *Simethis bicolor*, *Urginea maritima*, and *U. scilla*. It occurs in Great Britain, France, Spain, Portugal, Italy, Switzerland, Holland, Russia, Austria, Hungary, Corsica, Montenegro, Serbia, Rumania, Greece, Morocco, Algeria, Tunis, Egypt, Dalmatia, the Union of South Africa, and Germany.

HYACINTHUS—Continued.

Ustilago vaillantii Tul. One of the smuts producing black powdery masses of spores in place of the anthers and ovaries in the flowers. It is perennial in the bulbs and hence easily carried from place to place. The flowers are disfigured and their commercial value lowered. The hosts are *Belleria romana*, *B. trifoliata*, *Chionodoxa lucitiae*, *Chionodoxa* sp., *Gagea* spp., *Hyacinthus* spp., *Muscari comosum*, *M. racemosum*, *M. schliemannii*, *M. tenuiflorum*, *Scilla anthericoides*, *S. bifolia*, *S. cernea*, *S. ilio-hyacinthus*, *S. maritima*, *S. trifolia*, *Urginea anthericoides*, and *U. maritima*, reported from Great Britain, France, Corsica, Italy, Belgium, Serbia, Egypt, Algeria, Austria, Germany, and from a few localities in the United States.

HYALIS. See *Ixia*.**HYDRANGEA.** Ornamental woody plants grown for their showy flowers.

Aecidium hydrangeae Pat. Rust on leaves of *H. davidi* in Thibet.

Aecidium hydrangeae-paniculatae Diet. Rust pustules on large, irregular, yellow, or reddish leaf spots on *H. paniculata* in Japan.

Aecidium hydrangiocolum P. Henn. Rust on leaves of *H. hirta* and *H. thunbergii* in Japan.

Ascochyta hydrangeae Arn. Circular to irregular greenish-gray spots on leaves and stems of *H. hortorum* in France. The spots may spread to include the entire area of leaves.

Cercospora hydrangeicola Speg. Circular, then confluent, purplish leaf spots on *H. hortensis* in Argentina.

Cercospora obtegans Syd. On leaves of *H. hortensis* in Japan.

Clasterosporium hydrangeae Sacc. On *H. hortensis* in Italy and Madeira.

Rhytisma sp. On leaves of *H. paniculata* in Japan.

Septoria anthophila Fl. Tass. On flowers of *H. hortensis* in Italy.

Septoria hydrangeae Bizz. Irregular, then confluent, dull-brown leaf spots with blood-red margins on *H. hortensis* in Germany. Said to cause serious damage. Reported from Ohio.

HYMENAEA. Tropical trees.

Rosellinia pepo Pat. See Citrus.

Uredo hymenaeae Mayor. Powdery light-brown rust pustules covering the under leaf surfaces of *H. courbaril* in Porto Rico, Cuba, and northern South America.

HYMENOCALLIS. SPIDER LILY. Bulbous plants.

Tubercinia javanica Koord. Large dark-brown areas on leaves of *Hymenocallis* sp. in Java.

HYMENOPHYLLUM. Filmy fern.

Micropeltis hymenophylli Pat. Small black fruiting bodies on leaves of *Hymenophyllum* sp. in Tahiti.

Sporotrichum niveum Allesch. and Henn. On leaves of *H. bridgesii* in Chile.

HYOSCYAMUS. HENBANE. Herbs cultivated for medicinal purposes.

Ascochyta hyoscyami Pat. Circular dull-brown leaf spots on *H. niger* in Tunis, Russia, Hungary, and Germany.

Ascochyta pinzolenis Kab. and Bub. Irregular and confluent brown or dull-yellow leaf spots, becoming white with brown margins on *H. niger* in Italy.

Peronospora hyoscyami De By. Downy mildew on leaves of *H. niger* in Bohemia, Silesia, and and Russia. See also *Nicotiana*.

Puccinia hyoscyami P. Magn. Dark-brown rust pustules on leaf blades and petioles of *H. senecionis* in Persia.

Septoria hyoscyami Höll. Dull-yellow leaf spots with brown margins on *H. niger* in Hungary.

Septoria pinzolenis Kab. and Bub. Circular to angular brown or dull-yellow leaf spots, becoming white with brown margins on *H. niger* in Italy.

HYOSERIS. Stemless herbs.

Entyloma aposeridis Jaap. Smut sori in small dull-brown leaf spots on *H. foetida* in Austria.

Phyllosticta aposeridis Allesch. Irregular dull-yellow to brown leaf spots on *H. foetida* in Germany.

Protomyces hyoseridis Syd. Circular yellow leaf spots on *H. baetica* in Italy.

Puccinia hyoseridis-radiatae R. Maire. Cinnamon-brown to dark-brown rust pustules on leaves of *H. radiata* and *H. scabra* in Tripoli, Spain, and Dalmatia.

Septoria hyoseridis Maire. Brown leaf spots with purple margins on *H. radiata* in tropical Africa.

HYPERANTHERA. See *Moringa*.**HYPERICUM.** St. JOHN'S-WORT. Annual herbs.

Ascochyta hyperici Lasch. Dull-brown leaf spots on *H. perforatum* in Italy and Germany.

Melampsora hypericorum Wint. Powdery orange-yellow then reddish-brown rust pustules on leaves of *Androsace* spp. and *Hypericum* spp. in Europe, Asia, and Africa.

Melampsora kusanoi Diet. Chestnut-brown, then black, rust pustules on yellow sunken leaf spots on *H. ascyron* in Japan.

Melampsora sancti-johannis Barcl. Rust on leaves of *H. cernuum* and *H. patulum* in India.

Metasphaeria cryptopila (Berk.) Sacc. On leaves of *Hypericum* sp. in New Zealand.

Mycosphaerella hyperici (Auersw.) Starb. On stems of *Hypericum* spp. in Europe.

Ovularia minutissima Syd. Circular to elliptical brown leaf spots with purple margins on *H. quadrangulum* in Germany.

Ramularia coleosporii Sacc. See Campanula.

Septoria carestiana Ferr. Pale-yellow or reddish leaf spots on *H. montanum* in Italy.

Septoria hyperici Desm. and var. **hyperici-quadranguli** C. Mass. Circular oblong, or indefinite, red-brown leaf spots with dull-yellow margins on *H. burserum*, *H. hirsutum*, *H. montanum*, *H. perforatum*, *H. pulchrum*, and *H. quadrangulum* in Europe.

Septoria hyperici-burseri R. Maire. On leaves of *H. burserum* in France.

Uredo hyperici-japonici Petch. Rust on leaves of *H. japonicum* in Ceylon.

Uredo hyperici-mysorensis Petch. Rust on leaves of *H. mysorensis* in Ceylon.

HYPOXIS. GOLD-EYE GRASS. STAR GRASS. Herbaceous plants with hard root stocks or corms.

Cylindrosporium guttatum Wint. Round to irregular dull-brown leaf spots on *Hypoxis* sp. in Brazil.

Puccinia expallens Syd. Gray-brown rust pustules on leaves of *H. aurea* in India.

Puccinia hypoxidis McAlp. Dark-brown to black rust pustules on both leaf surfaces of *H. hygrometrica* in Victoria, Australia.

Uromyces hypoxidis Cke. Brown rust pustules on both leaf surfaces of *H. oligotricha* in Natal and Uganda.

IBERIS. CANDYTUFT. Small flower-garden and border plants.

Helminthosporium iberidis Poll. Circular dull-brown leaf spots on *Iberis* sp. in Italy.

Puccinia iberidis Duby. Rust on leaves of *I. sempervirens* in France, Switzerland, and Belgium.

ICACOREA. See *Ardisia*.**ICHTHYOMETHIA.** See *Piscidia*.**IDESIA.** Ornamental trees, grown for their handsome foliage and attractive berries.

Melampsora idesiae Miy. Yellow to brown rust pustules on leaves of *I. polycarpa* in Japan.

- ILEX.** HOLLY. Maté, sometimes called Paraguay tea. Evergreen or deciduous woody plants, grown for their attractive leaves and berries. The leaves of several South American species are used for beverage purposes.
- Boydia insculpta** (Oud.) Grove. Destroys twigs of *I. aquifolium* in Great Britain and probably continental Europe.
- Cercospora illicicola** Maubl. On leaves of *I. paraguayensis* in Brazil. A doubtful report from Texas.
- Cercospora yerbae** Speg. Dull-brown circular leaf spots on *I. amara* in Argentina.
- Cercosporina mate** Speg. Circular ashen-colored leaf spots with purple margins on *I. paraguayensis* in Argentina.
- Colletotrichum yerbae** Speg. Subcircular whitish or ashen leaf spots on *I. paraguayensis* in Argentina and Brazil.
- Cylindrosporium pollacii** Ture. Circular whitish leaf spots with broad, dark-purple margins on *I. furcata* in Italy.
- Depazea illicicola** Fr. On leaves of *I. aquifolium* in France.
- Exobasidium giseckiae** Allesch. White to reddish swellings on leaves and flowers of *Giseckia pharnacioides* and *I. aquifolium* in Italy and tropical Africa.
- Gloeosporium alutaceum** Sacc. Anthracnose on leaves of *I. aquifolium* in Italy.
- Gloeosporium aquifolii** Penz. and Sacc. Anthracnose on leaves of *I. aquifolium* in Italy.
- Gloeosporium orthosporum** Sacc. White indefinite leaf spots on *I. aquifolium* in Italy.
- Laestadia illicis** Jacz. On leaves of *I. aquifolium* in Switzerland.
- Leptosphaeria paraguayensis** Maubl. On leaves of *I. paraguayensis* in Brazil.
- Pectia mate** Speg. Fruiting bodies occur on lower leaf surfaces of *I. paraguayensis* in Argentina and cause serious damage.
- Phyllosticta azevini** Torr. Brown leaf spots on *I. azevini* in Madeira.
- Phyllosticta haynaldi** Roum. Indefinite dull-red leaf spots on *I. aquifolium* in Denmark and France.
- Phyllosticta illeperda** Oud. Irregular gray leaf spots on *I. aquifolium* in Holland.
- Phyllosticta illicis** Oud. Pale irregular leaf spots on *I. aquifolium* in Holland.
- Phyllosticta mate** Speg. Circular, then confluent, silvery leaf spots on *I. paraguayensis* in Argentina and Brazil.
- Phyllosticta yerbae** Speg. Definite circular dull-brown leaf spots on *I. paraguayensis* in Argentina.
- Rhytisma himaleense** Syd. and Butl. Black, slightly raised areas on leaves of *Ilex* sp. in India.
- Rhytisma illicis-latifoliae** P. Henn. Black patches on subcircular leaf spots on *I. integra*, *I. latifolia*, *I. othera*, and *I. peduncularis* in Japan.
- Septoria orthospora** Lévl. On leaves of *I. aquifolium* in France.
- Trabutia lagerheimiana** (Rehm.) Theiss. and Syd. Elliptical shiny black stromata on leaves of *I. scopulorum* in Ecuador.
- Venturia illicifolia** Cke. On leaves of *I. aquifolium* and *I. balearica* in Europe.
- ILLICIUM.** ANISE TREE. Small trees or shrubs.
- Gloeosporium illicii** Hemml. Anthracnose on *I. anisatum* in Japan.
- Phyllosticta aromatica** F. Tassi. Circular to elliptical gray leaf spots on *I. floridanum* in Italy.
- IMPATIENS.** SNAPWEED. TOUCH-ME-NOT. BALSAM. Flower-garden annuals.
- Ascochyta impatiensis** Bres. Subcircular to angular dull-yellow to brown leaf spots on *I. parviflora* in Germany.
- Cercospora campi-silii** Speg. Circular to angular brown leaf spots on *I. nolitangere* in Denmark, Italy, and Austria.
- Cercospora impatiensis** Bauml. Circular, then irregular, gray or whitish leaf spots with dull-brown margins on *I. nolitangere* and *I. parviflora* in Hungary.
- Cronartium faecidum** (Alb. and Schw.) Wint. See *Paeconia*.
- Gloeosporium impatiensis** Petch. Soft rot of stems of *I. balsamina* in Ceylon.
- Puccinia komarovii** Tranzsch. Powdery cinnamon-brown to chestnut-brown rust pustules on leaves of *I. amphorata* and *I. parviflora* in Turkestan and India.
- Uredo balsaminiae** Cke. Rust on leaves of *I. oppositifolia* in Ceylon.
- IMPERATA.** Large perennial tropical grasses.
- Cercospora imperata** Syd. Leaf rust on *Imperata* sp. in the Philippines.
- Phyllachora cyperi** Rehm. var. *donacis* Berl. and Sacc. Tar spot on leaves of *I. arundinacea* in India.
- Phyllachora imperatae** Syd. Black stromata on leaves of *I. cylindrica* in the Philippines. Probably identical with the preceding species.
- Puccinia fragoana** Beltram. Leaf rust on *I. cylindrica* in Spain.
- Puccinia infusans** Arth. and Holw. Brown powdery leaf rust on *I. braziliensis* in Guatemala.
- Puccinia rufipes** Diet. Dull-yellow to black rust pustules on leaves of *I. arundinacea* and *I. cylindrica* in the Philippines, Japan, Ceylon, China, India, and Egypt.
- Sphaelotheca schweinfurthiana** (Thum.) Sacc. Smut sori replacing ovaries of *I. arundinacea* and *I. cylindrica* in Uganda, Egypt, Tunis, Tripoli, and Spain.
- INDIGOFEA.** INDIGO. Shrubs or perennial herbs sometimes grown for ornament. Several species furnish dyestuff.
- Corticium salmonicolor** B. and Br. See *Citrus*.
- Diplodia cacaoicola** P. Henn. See *Theobroma*.
- Gloeosporium inocarpi** Sacc. Anthracnose on leaves of *I. arrecta* in Malaya.
- Pythium indigoferae** Butl. On leaves of *I. arrecta* in India.
- Ravenelia indigoferae** Tranzsch. Powdery brown rust pustules on leaves and stems of *I. anil*, *I. conzattii*, *I. cuernavacana*, *I. mucronata*, *I. palmeri*, and *I. suffruticosa* in Mexico, Central America, northern South America, West Indies, Trinidad, and Bermuda.
- Ravenelia laevis** Diet. and Holw. Powdery brown rust pustules on leaves of *I. densiflora* and *I. jalsensis* in Mexico.
- Ravenelia schroeteriana** P. Henn. Powdery cinnamon-brown rust pustules on leaves of *Indigofera* sp. in Argentina.
- Uredo maranguensis** P. Henn. Ashen-brown rust pustules on leaves of *I. arrecta* in tropical Africa.
- Uromyces mbelensis** P. Henn. Brown rust pustules on *Indigofera* sp. in the Congo.
- Uromyces orientalis** Syd. Cinnamon-brown rust pustules on leaves of *I. cordifolia*, *I. glandulosa*, and *I. tinifolia* in India.
- Uromyces sphaeocarpi** Syd. Brown rust pustules on leaves of *I. pseudotinctoria* and *I. tinctoria* in Japan.
- INGA.** Tropical trees and shrubs, commonly used as coffee shade.
- Linospora guaranitica** Speg. Indefinite spots on leaves of *Inga* sp. in Brazil.
- Ophiodothella ingae** (P. Henn.) Theiss. and Syd. Dull-black irregular stromata on large yellow or yellow-brown leaf spots on *I. strigillosa* and *Inga* sp. in Brazil.
- Ravenelia ingae** (P. Henn.) Arth. Cinnamon-brown rust pustules on leaves and stems of *I. edulis*, *I. imicul*, *I. laurina*, and *I. vera* in Mexico, South and Central America, and the West Indies.

INGA—Continued.

Stilbella flavid (Oke.) Lind. See Coffea.

Uredo curvata Arth. Brown leaf rust on *I. vera* in Cuba.

Uromyces ingicola P. Henn. Brown rust pustules on leaves of *Inga* sp. in Brazil.

Uromyces porcensis Mayor. Rust on *I. ingoides* in Colombia.

INTSIA. See *Afzelia*.**INULA**. Herbs with yellow or orange flower heads.

Aecidium inulae-crithmoidis Pat. Rust on leaves of *I. crithmoides* in Dalmatia

Aecidium inulae-helenii Const. Rust on yellow irregular leaf spots on *I. helenium* in Italy.

Ascochyta inulicola Petr. On *I. conyzia* in Bohemia.

Colosporium inulae Rabb. Yellow rust pustules on *Pinus sylvestris*, *Inula aschersoniana*, *I. candida*, *I. coppa*, *I. ensifolia*, *I. graveolens*, *I. helenium*, *I. heterolepis*, *I. hirta*, *I. media*, *I. salicina*, *I. vaillantium*, and *I. viscosa* in Europe, Asia Minor, northern Africa, the Congo, and the Canary Islands.

Erysiphe taurica Lévl. See *Althaea*.

Ovularia inulae Sacc. and *forma major* P. Brun. Angular to irregular leaf spots with purple margins on *I. dysenterica* in France.

Phyllosticta inulae Allesch. Circular whitish to dull brown leaf spots on leaves of *I. britannica* in Germany.

Ramularia cupulariae Pass. On leaves of *I. conyzia* and *I. viscosa* in Denmark, Spain, and Germany.

Ramularia inula-britannicae Allesch. Subcircular yellow, then brown, leaf spots on *I. britannica* in Germany.

Septoria dysentericae P. Brun. Irregular to angular small white leaf spots with narrow brown margins on *I. dysenterica* in France.

Septoria inulae Sacc. and Speg. Rufous-brown leaf spots on *I. britannica* and *I. salicina* in China and Italy.

Uredo inulae-candidae Trott. Yellow rust pustules on leaves of *I. candida* in Greece and Dalmatia.

IPOMOEA. MORNING-GLORY. SWEET POTATO. Annual or perennial twining herbs.

Aecidium kaernbachii P. Henn. Rust on leaves of *Ipomoea* spp. in Japan, Indo-China, Ceylon, New Guinea, Amboina, and the Philippines.

Cercospora batatae Zimm. Irregular dark-brown leaf spots with paler centers on *I. batatas* in Tanganyika, Indo-China, Japan, Formosa, and the Philippines.

Cercospora stuckertiana Syd. Subcircular gray leaf spots on *Ipomoea* sp. in Argentina.

Cercospora timorensis Cke. Circular brown leaf spots on *I. cymosa* in Molucca.

Cylindrosporium bakeri Syd. On *Ipomoea* sp. in the Philippines.

Helicobasidium mompa Tan. See *Morus*.

Marsonia ipomoea Cke. and Mass. Brown areas on stems and leaves of *Ipomoea* sp. in England.

Phyllosticta stuckertii Speg. Small leaf spots with purple borders on *Ipomoea* sp. in Argentina.

Puccinia batatae Syd. Brown rust pustules on leaves of *I. paniculata* in Natal and the Congo.

Puccinia holosericea Cke. Brown rust pustules on leaves of *I. holosericea* in Tanganyika and Natal.

Puccinia insignis Holw. Rust on leaves of *Ipomoea* sp. in Mexico.

Puccinia macrocephala Speg. Black rust pustules on leaves of *Ipomoea* sp. and *Convolvulus* sp. in South America.

Puccinia nocticolor Holw. Yellow, then black, powdery rust pustules on leaves of *I. fistulosa*, *I. intrapilosa*, and *I. murucoides* in Mexico and Guatemala.

Puccinia rubicunda Holw. Rust on leaves of *Ipomoea* sp. in Mexico.

Puccinia superflua Holw. Black powdery rust pustules on yellow leaf spots on *I. murucoides* in Mexico.

Ramularia batatae Rac. Angular brown to dark-brown leaf spots on *I. batatas* in Java.

Sorosporium ipomoeae Spesch. Smut on *Ipomoea* sp. in Russia.

Uredo ipomoeae-pentaphyllae P. Henn. Rust on leaves of *I. pentaphylla* in Brazil.

Uredo speschnewii S. and S. Brown powdery rust pustules on leaves of *Ipomoea* sp. in Transcaucasia.

Uromyces comptus Syd. Dark-brown to black rust pustules on leaves of *I. bipinnatifida* in Tanganyika and the Union of South Africa.

Uromyces ipomoeae (Thuem.) Berk. Yellow, then black, rust pustules on leaves of *I. argyreoides* in the Union of South Africa.

Uromyces pachyceph Lagh. Rust on leaves of *Ipomoea* sp. in Ecuador.

IRESENE. BLOOD LEAF. Ornamental-leaved bedding plants.

Cercospora gilbertii Speg. See *Celosia*.

Puccinia macropoda Speg. Cinnamon-brown to black rust pustules on leaves of *I. angustifolia*, *I. celosia*, *I. celosioides*, and *I. elatior* in South America and the West Indies.

Puccinia striolata (Speg.) Arth. Powdery brown rust pustules on leaves of *I. angustifolia* and *I. celosia* in Trinidad, Porto Rico, Cuba, St. Thomas, and Argentina.

Septoria iresines Speg. Circular or elliptical definite whitish leaf spots with dull-brown margins on *I. celosioides* in Argentina.

Uromyces celosiae Diet. and Holw. See *Celosia*.

Uromyces iresines Lagh. Dull-brown rust pustules on leaves of *I. celosia*, *I. elatior*, and *I. paniculata* in Guatemala, St. Thomas, Ecuador, and Colombia.

IRIS. Perennial rhizomatous or bulbous plants.

Ascochyta iridis Oud. Leaf spot of *I. pseudacorus* in Holland.

Ascochyta pseudacori All. On leaves of *Iris* (cultivated var.) in France.

Bacterium iridis v. Hall. Infected leaves turn yellow and dry up. The underground portions are soft rotted. The disease is generally at its worst during the flowering season. It is readily carried on rhizomes or bulbs, either as small lesions or by means of the bacteria themselves in soil or debris or on the bulb scales. Soft rot of *Iris* bulbs and rhizomes is common in the United States, but the causal agents mentioned have been other than the above. *Bacillus omnivorus* is also given as the cause of an *Iris* rot in Europe. The disease occurs on many varieties and species of cultivated *Iris* in Holland, Great Britain, and France.

Cladochytrium (Physoderma) iridis De By. Causes a leaf spot and blighting of *Iris* sp. in Germany.

Heterosporium montenegrinum Bub. Forms pale spots with purple-brown margins on leaves of *I. graminea* in Montenegro. Said to be distinct from *H. gracile*.

Leptosphaeria heterospora (De Not.) Niessl. Leaf spot and lesions on *Iris* rhizomes in Europe.

Leptosphaeria iridicola Lamb. and Fautr. (*L. vectis* [B. and Br.] Ces. and De Not.) On *I. foetidissima* in France and Italy.

Macrophoma rhabdosporioides Lamb. and Fautr. Reported as the cause of a leaf spot of *I. foetidissima* in France.

Metasphaeria iridicola (Desm.) Sacc. On leaves and stems of *I. foetidissima* in Algeria.

Microdiploia iridicola Gz. Frag. On *I. filifolia*, *I. longepetala*, *I. pumila*, and *I. spuria* in Spain.

Mycosphaerella iridis (Auersv.) Schroet. Causes a leaf spot on *I. filifolia*, *I. pseudacorus*, *I. pumila*, and *I. syriacanthum* in Great Britain, Denmark, France, Italy, Spain, and Germany.

IRIS—Continued.

Mystrosporium adustum Mass. This fungus forms large black crusty patches on the outer bulb scales, the mycelium reaching to the heart of the bulb. The disease is readily carried on the bulbs of the host (*I. reticulata*). It is said "to carry off bulbs by the hundreds, making it necessary to lift the bulbs every year and sort out the diseased ones for destruction and dry and store the remainder." The disease occurs in England and Denmark.

Nectria paludosa (Eckl.) Sacc. On leaves of *Iris* sp. and *Typha* sp. in Europe.

Ophiobolus peduncularis Feltg. At the base of stems of *I. pseudacorus* in Luxemburg. *O. bactrosporus* Feltg. and *O. pseudacori* Feltg. are reported on the same host and from the same region.

Phoma agapanthi Thuen. Attacks the base of *Iris* plants in Europe.

Phoma iridis Cke. On leaves of *I. foetidissima* in Europe.

Phyllosticta iridicola Gz. Frag. Reported from Spain as the cause of leaf spot of *I. pseudacorus*.

Phyodicta tenue (Nowk.) Karst. See *Nymphaea*.

Puccinia capensis Diet. This rust fungus produces brown pustules on the leaves of *Iris* sp. in the Union of South Africa.

Puccinia caucasica Martino. Large stroma-like rust patches on leaves of *I. flavescens* in the Caucasus.

Puccinia melanopsis Syd. Black rust pustules are produced on both leaf surfaces of *I. sisyrinchium* in Asia Minor.

Ramularia rollandi Faut. On leaves of *I. pseudacorus* in France.

Schinzia (*Naegelia*) **cellulicola**. Causes swellings on the roots of *Iris* sp. in Switzerland.

Sclerotium crustiforme Rob. At the base of plants of *I. germanica* in Belgium.

Sclerotium durum Pers. At the base of plants and on rhizomes and bulbs of *Iris* spp. in Europe.

Sclerotium iridis Thuen. Attacks the rhizomes of cultivated *Iris* varieties, causing a soft rot distinct from bacterial rot. Several shipments of *Iris* rhizomes from Germany have been completely destroyed by what is probably this species. Doubtless also occurs in Holland and France. At least two other species of *Sclerotium* have been reported on *Iris* spp. in Europe, and it is not clear how many species really are involved. What has been referred to as *S. semen* has been intercepted on *I. tingitana*.

Seoleotrichum cladosporioideum R. Maire. On leaves of *I. foetidissima* in Algeria.

Seoleotrichum iridis Fautr. and Roum. On leaves of *Iris* sp. in Europe.

Septoria iridina Sacc. On leaves of *I. foetidissima* in Portugal.

Septoria iridis Massal. Circular, brown spots with light-brown margins on leaves and flower stalks of *I. florentina* and *I. germanica* (in var.) in Italy and Dalmatia.

Stagonospora iridis C. Mass. Causes a leaf spot disease on cultivated *Iris* in Great Britain and *I. florentina* in Tunis.

Trabulia atrolineata (Wint.) Theiss. and Syd. On *Iris* sp. in Portugal.

Trabulia molleriana Cava. On leaves of *Iris* sp. in Portugal.

Uredo iridis (Thuen.) Plowr. Forms narrow elliptical chestnut-colored rust pustules on both surfaces of leaves of *I. flavissima*. Said to be distinct from *Puccinia iridis*, which is present in the United States.

ISATIS. Herbs cultivated for ornament and for dyestuff.

Aecidium isatidis P. Har. Rust on leaves of *I. tinctoria* in France.

Peronospora isatidis Güm. Downy mildew on leaves of *I. tinctoria* in Switzerland.

ISCHNOSIPHON. Calathea-like perennial herbs.

Uredo ischnosiphonis P. Henn. Powdery brown rust pustules on leaves of *I. leucophaeus* in Brazil.

ISOPYRUM. Annual or perennial dwarf stemless plants.

Aecidium isopyri Schroet. Rust on leaves of *I. thalicroides* in France, Switzerland, Belgium, and Germany.

Cercospora isopyri v. Hoeh. Irregular dark-brown areas on leaf blades and petioles and on stems of *I. thalicroides* in Austria.

Peronospora parvula Schneid. Downy mildew on leaves of *I. fumarioides* in Siberia.

Triphragmium isopyri Moug. and Nestl. Black rust pustules on leaf blades and petioles and on stems of *I. thalicroides* in Europe.

IXIA. HYALIS. Ag. Spring flowering cormous plants from south Africa with grasslike foliage.

Bacillus ixiae Sever. Appearing first as a browning and dying back of the tops of the plants, followed by a soft rot of the bulbs of *I. maculata* in Italy and Holland.

Uromyces ixiae (Lév.) Wint. Small yellow rust pustules, becoming brown on leaves of *Acidantha exscapa*, *Ixia arillaris*, *I. corulescens*, *I. erecta*, *I. paniculata*, *I. patens*, *I. polystachya*, *Sparaxis grandiflora*, and *Watsonia densiflora* in south Africa and Australia.

Uromyces zeyheri Bub. Yellow to brown rust pustules on both leaf surfaces of *I. scillaris* in south Africa.

IXIOLIRIUM. Hardy spring-flowering bulbs.

Aecidium ixioliri Komarov. Rust on leaves of *I. tataricum* in central Asia.

Aecidium tataricum Rostr. Rust on leaves of *I. tataricum* in Transcaspia.

IXORA. Woody plants with evergreen leaves and showy flower clusters.

Aecidium ixorae Arth. Rust on leaves of *I. ferrea* in Cuba.

Colletotrichum ixorae Griff. and Maubl. Large, pale anthracnose spots on leaves of *I. alba* in France.

Endophyllum ixorae Güm. Leaf rust on *I. javanica* in Guam.

Phyllachora ixorae Theiss. and Syd. Shiny black stromata on leaves of *I. parvifolia* in India.

Phyllosticta ixorae Rangel. On leaves of *Ixora* sp. in Brazil.

Stagonospora ixorae Rangel. On leaves of *I. coccinea* and *Ixora* sp. (cult.) in Brazil.

JACARANDA. Trees and shrubs with handsome tubular flowers.

Aecidium circinatum Wint. Rust on leaves of *Jacaranda* sp. in Brazil.

Aecidium jacarandae P. Henn. Rust on leaves of *Jacaranda* sp. in Brazil.

Aecidium puttemansianum P. Henn. Rust on leaves of *Jacaranda* sp. in Brazil.

Fomes lamaensis Murr. See *Hevea*.

JACKSONIA. Stiff, leafless shrubs and subshrubs.

Cronartium jacksoniae P. Henn. Brown rust pustules on stems of *Aotus villosus*, *Bossiaea cinerea*, *Gompholobius latifolius*, *Jacksonia scoparia*, and *Platylobium formosum* in Australia.

Uromyces politus (Berk.) McAlp. Bright-orange rust pustules, becoming brown to black, on dark-purple, elongated patches on stems of *Muehlenbeckia* (*Jacksonia*) *cunninghami* in Australia.

JACQUEMONTIA. Tropical climbing herbs.

Aecidium jacquemontiae E. and E. Rust on leaves of *J. pentantha* in Mexico.

Uredo jacquemontiae P. Henn. Powdery dull-yellow rust pustules on leaves of *Jacquemontia* sp. in New Guinea.

Uromyces gemmatus B. and C. Rust on leaves of *J. nodiflora* in Porto Rico, Cuba, Jamaica, and St. Croix.

- JACQUINIA.** Tropical American trees and shrubs.
Phyllachora conspicua Ferd. and Winge. Dull-black stromata on leaves of *J. armillaris* in the American Virgin Islands.
Phyllachora inclusa (B. and C.) Sacc. Small black stromata on leaves of *J. aurantiaca*, *J. berterii*, and *Jaquinta* sp. in Porto Rico, Guatemala, Nicaragua, and Paraguay.
- JAMBOSA.** See *Caryophyllus*.
- JASMINUM.** JASMINE. JESSAMINE. Climbing or erect shrubs, with attractive, often fragrant, flowers.
Aecidium jasminicola P. Henn. Rust distorting leaves and branches of *J. floribundum* in Abyssinia.
Aecidium longaeense P. Henn. Rust on yellow-brown leaf spots on *J. microphyllum* in the Union of South Africa.
Blastospora butleri Syd. Brown rust pustules on leaves of *J. malabaricum* in India.
Laestadia jasminicola (Desm.) Sacc. Black sunken leaf spots on *J. officinale* in France.
Meliola jasminicola P. Henn. Black superficial fungus patches on upper leaf surfaces of *J. auriculatum* and *J. sambac* in India and Indo-China.
Mycosphaerella jasmini-officinalis Siem. On leaves of *J. officinale* in Southern Russia.
Phyllosticta jasminorum Togn. On leaves of *J. officinale* in Italy.
Phytophthora syringae Kleb. See *Syringa*.
Puccinia abyssinica (P. Henn.) Syd. Brown rust pustules on circular to irregular sunken leaf spots on *J. abyssinicum* in Abyssinia.
Puccinia chrysopogi Barel. See *Andropogon*.
Puccinia exauriens Thuem. Cinnamon-brown rust pustules on leaves and branches of *J. flexile* and *J. tortuosum* in Ceylon and the Union of South Africa.
Puccinia jasmini DC. Dark-brown rust pustules on leaves and stems of *J. fruticans* in Europe and northern Africa.
Puccinia zimmermanniana P. Henn. Brown rust pustules on leaves of *J. mauritianum* in east Africa.
Septoria aitchisoni Syd. Circular dull-brown leaf spots with raised purple margins on *J. humile* in Afghanistan.
Septoria jasmini Roum. Circular, then confluent, brown leaf spots, becoming white with brown margins, on *Jasminum* sp. in France.
Septoria sambac Pass. Subcircular whitish leaf spots on *J. sambac* in Italy.
Uromyces comedens Syd. Yellow to dark-brown rust pustules on leaves of *J. pubescens* in India.
Uromyces hobsoni Vize. Brown rust pustules on leaves, stems, and flower stalks of *J. floribundum*, *J. grandiflorum*, and *Jasminum* sp. in India, Somali, and Abyssinia.
- JATROPHA.** PHYSIC NUT. Tropical trees and shrubs with milky juice.
Cercospora ajreklari Syd. Circular yellow-brown leaf spots on *J. nana* in India.
Cercospora jatrophae Speg. Definite circular leaf spots, concentrically zoned, on *J. macrocarpa* in Argentina.
Cercospora peronosporioides Speg. Dull-yellow indefinite leaf spots on *J. anisophylla* and *J. macrocarpa* in Argentina.
Cercosporina jatrophicola Speg. Subcircular to angular ashen-colored leaf spots with dull-brown margins on *J. macrocarpa* in Argentina.
Uredo jatrophicola Arth. Brown rust pustules on leaves of *J. curcas* and *J. gossypifolia* in Porto Rico, Cuba, and Dominican Republic.
Uromyces jatrophae Diet. and Holw. Powdery cinnamon to chestnut-brown rust pustules on leaves of *J. multifida* in Mexico.
Uromyces jatrophicola P. Henn. Powdery yellow to black rust pustules on leaves of *Jatropha* sp. in Brazil.
- JEFFERSONIA.** TWINLEAF. Attractive, hardy perennial herbs.
Triphragmiopsis jeffersoniae Naumoff. Rust on leaf blades and petioles of *J. dubia* in Russia.
- JOANNESIA.** Tropical trees with milky juice, useful for medicinal, ornamental, and timber purposes.
Puccinia joannesiae P. Henn. Pale-brown to chestnut-brown rust pustules on leaves of *J. brasiliensis* in Brazil.
Uredo macleensis P. Henn. Brown rust pustules on irregular dull-brown leaf spots and on stems of *J. princeps* in Central America.
- JUBAEA.** PALM. See *Palmae*.
- JUGLANS.** WALNUT. Butternut. Nut and timber trees.
Ascochyta juglandis Bolts. Subcircular gray-brown leaf spots, the centers of which fall out. On *J. regia* in Switzerland, Yugoslavia, and Austria.
Gummosis. A gumming disease of walnut (*Juglans* spp.) trunks and branches has been reported from France and other parts of Europe. Several distinct diseases are probably included, some due to nonparasitic causes, others possibly to bacteria or fungi not present in the United States.
Helicobasidium tanakae Miy. See *Morus*.
Hypoderma roseum (Pers.) Fr. See *Populus*.
Marsonia manshurica Naum. Numerous sub-circular ashen-white leaf spots on *J. manshurica* in Russia.
Mycosphaerella saccardoana Jaap. On old leaves of *J. regia* in Dalmatia.
Phyllosticta juglandina Sacc. Indefinite whitish leaf spots with dull-brown borders on *J. regia* in Italy and Portugal.
Phyllosticta juglandis (DC.) Sacc. Indefinite whitish leaf spots on *J. regia* in Russia, Italy, France, and Austria.
Septoria epicarpii Thuem. On epicarp of nuts of *J. regia* in Europe.
Septoria letendreana Sacc. On leaves of *J. regia* in France.
Septoria nigromaculans Thuem. On epicarp of nuts of *J. regia* in Austria.
- JUNCOIDES.** See *Luzula*.
- JUNIPERUS.** JUNIPER. Redcedar. Coniferous ornamentals and timber trees.
Asteridium juniperinum Cke. On needles of *Juniperus* sp. in Great Britain.
Clasterosporium glomerulosum Sacc. On leaves of *J. communis* in Germany.
Gymnosporangium amelanclieris (DC.) Ed. Fisch. The acial stage of this rust forms small galls on the lower leaf surfaces of *Amelanchier ovalis*. The telia stage occurs in indefinite swollen areas along the branches of *J. communis* and *J. nana* in Europe.
Gymnosporangium koreense (P. Henn.) Jacks. The acial stage occurs on leaves of *Cydonia japonica*, *C. oblonga*, and *Pyrus sinensis*, the telia on *J. chinensis* in Japan and Chosen. This rust has been introduced with nursery stock into Oregon and Connecticut.
Gymnosporangium mespilii (DC.) Kern. Aecia on leaves and stems of *Cotoneaster tomentosa*, *C. vulgaris*, *Crataegus grandiflora*, *C. laciniata*, *C. monogyna*, *C. orientalis*, *C. oxyacantha*, *C. pinatifida*, *C. tanacetifolia*, *Cydonia oblonga*, *Mespilus germanica*, and *Pyrus communis*. Telia dark chestnut-brown on fusiform enlargements on branches of *J. oxycedrus*, *J. phoenicea*, *J. sabina*, and *J. virginiana* in Europe and Asia.

JUNIPERUS—Continued.

- Gymnosporangium photiniae** (P. Henn.) Kern. The acacia occur on thickened discolored spots on the lower leaf surfaces of *Pourthinea villosa*. The telia occur on fusiform enlargements on the branches of *J. chinensis* in Japan and Chosen. Reported from Connecticut on imported stock.
- Gymnosporangium sabinae** (Dicks.) Wint. This rust produces swellings on the twigs, deformed fruit, and leaf spots on which the acacia occur. The acical hosts are *Pyrus betulaeifolia*, *P. communis*, *P. elaeagnifolia*, *P. michauxii*, *P. nivalis*, *P. salicifolia*, and *P. tomentosa*. The dark-brown telia occur on swellings on twigs and branches of *Juniperus chinensis*, *J. japonica*, *J. oxycedrus*, *J. phoenicea*, *J. sabina*, *J. sphaerica*, *J. tripartita*, and *J. virginiana* in Europe.
- Gymnosporangium terminali-juniperinum** Ed. Fisch. Acacia in small groups on brown leaf spots on *Sorbus latifolia* and telia on *J. communis* in Europe.
- Gymnosporangium yamadai** Miy. Acacia on leaves of *Malus* sp. (apple), *M. spectabilis* and *M. toringa* and the telia on *J. chinensis* in China and Japan.
- Herpotrichia mucilaginosa** Start. and Grev. On needles of *J. communis* in Sweden.
- Sarcosoma thwaitesii** (B. and Br.) Petch. Produces witches'-brooms on the stems and main branches of *J. bermudiana* in Ceylon.
- Stemphylium juniperinum** Karst. Kills the needles of *Juniperus* sp. in Finland.
- Venturia furcata** Fautr. On leaves of *J. sabina* in France.
- JUSSIAEA**. PRIMROSE WILLOW. Herbs.
- Puccinia jussiaeae** Speg. Cinnamon-brown rust pustules on leaves of *J. lanceolata* and *J. longifolia* in Argentina and Uruguay.
- Puccinia sphaeroidea** P. Henn. Brown rust pustules on stems and leaves of *Jussiaea* sp. in Japan.
- Septoria obsidionis** Speg. Small circular, often confluent, white leaf spots with broad purple borders on *J. longifolia* in Argentina.
- Septoria platensis** Speg. Small definite white leaf spots with purple margins on *J. longifolia* in Argentina.
- Septoria thomasi** Sacc. Subcircular light-buff leaf spots with brown-purple margins on *J. acuminata* in the Portuguese Island of St. Thomas.
- JUSTICIA**. Herbs cultivated for their showy flowers.
- Aecidium acanthacearum** Cke. Rust on leaves of *J. procumbens* and *J. uncinulata* in Ceylon and Uganda.
- Peronospora wildemaniana** P. Henn. and var. **macrospora** K. Saw. Downy mildew attacking the leaves of *J. procumbens* in Japan and India, causing irregular light yellowish-green to white spots, which finally occupy the entire leaf area.
- Puccinia fuhrmanni** Mayor. Rust on leaves of *J. secunda* in Colombia.
- Puccinia peraffinis** Syd. Brown rust pustules on yellow-brown circular leaf spots on *J. diffusa* in India.
- Puccinia shiraiana** Syd. Dark-brown rust pustules on leaves and stems of *J. procumbens* in Japan and Ceylon.
- Puccinia thwaitesii** Berk. Dark-brown rust pustules on large sunken brown leaf spots on *J. gendarussa* in Ceylon, Indo-China, Java, Sumatra, Celebes, Malaya, New Guinea, and the Philippines.
- Uromyces kwangensis** P. Henn. Brown rust pustules on leaves of *Justicia* sp. in the Congo.
- Ustilago thwaitesii** Berk. A doubtful smut species reported on *J. gendarussa* in Malaya.
- KADSUEA**. Tropical Asian woody climbers.
- Septoria kadsurae** Tassi. On leaves of *K. japonica* in Italy.
- KAGENECKIA**. Small evergreen trees from South America.
- Phyllachora negeriana** P. Henn. and Lind. Shiny black stromata on leaves of *K. oblonga* in Chile.
- KENNEDIA**. Woody trailers or twiners. See also *Hardenbergia*.
- Cercospora kennedyae** Cke. and Mass. Irregular and confluent cinnamon-brown leaf spots on *K. prostrata* in Australia.
- Phyllosticta kennedyae** Wint. Irregular, often confluent, dull-brown, then whitish, leaf spots on *K. ovata* in Portugal.
- Septoria kennedyae** Trav. and Spessa. On leaves of *Kennedia* sp. in Portugal.
- KENTIA**. See *Palmae*.
- KERRIA**. Shrubs.
- Helicobasidium tanakae** Miy. See *Morus*.
- Phomopsis striataeformis** Grove. On stems of *K. japonica* in Algeria.
- Septoria kerriae** Syd. Circular, pale-brown leaf spots on *K. japonica* in Japan.
- KIGELIA**. **KIGELKEIA** Ag. Ornamental African trees.
- Leptosphaeria dryadea** Sacc. On *K. pinnata* in the Philippines.
- Phyllosticta kigelia** Died. On leaves of *K. pinnata* in the Philippines.
- KIGELKEIA**. See *Kigelia*.
- KOCHIA**. SUMMER CYPRESS. MOCK CYPRESS. Herbs.
- Puccinia kockiae** Mass. Golden-brown to dark-brown rust pustules on leaves of *K. sedifolia*, *K. villosa*, and *Enchylaena tomentosa* in Australia.
- Uromyces kochiae** Syd. Rust on leaves of *K. prostrata* in Russia.
- KOELREUTERIA**. GOLDEN-RAIN TREE. Ornamental trees grown for their panicles of yellow flowers and handsome foliage.
- Phyllosticta koelreuteriae** Höll. Oblong ashen leaf spots with brown borders on *K. paniculata* in Hungary.
- Triphragmium koelreuteriae** Syd. Leaf rust on *K. paniculata* in Japan.
- Uncinula koelreuteriae** Miy. Powdery mildew on leaves of *K. bipinnata* in China.
- KBAUNHIA**. See *Wistaria*.
- LABURNUM**. GOLDEN CHAIN. Ornamental trees or shrubs, grown for their showy racemes of yellow flowers. See also *Cytisus*.
- Microsphaera guarinonii** Br. and Cav. Powdery mildew on leaves of *L. anagyroides* (*L. vulgare*) in Italy.
- Peronospora cytisi** Magn. See *Cytisus*.
- Septoria cytisi** Desm. Numerous subcircular whitish to brown leaf spots on *L. anagyroides* in Italy, France, and Hungary.
- Stereum purpureum** Pers. See *Prunus*.
- Uromyces genistae-tinctoriae** (Pers.) Fekl. See *Caragana*.
- LACTUCA**. LETTUCE. Hardy annual or perennial herbs, several species grown as vegetables; others weedy.
- Ascochyta lactucae** Rostr. On leaves and stems of *L. sativa* in Denmark.
- Cercospora lactucae** P. Henn. Dark areas on leaves of *L. raddeana* in Japan.
- Cercospora longissima** Trav. (*C. lactucae* Stevenson) (*C. lactucae* Welles). Circular to angular, often confluent, whitish leaf spots with rufous or brown margins on *L. sativa* in Italy, Brazil, Porto Rico, Cuba, China, and the Philippines.

LACTUCA—Continued.

- Puccinia lactucae** Diet. Brown rust pustules on leaves of *L. albiflora*, *L. brevisstris*, *L. denticulata*, *L. laciniata*, *L. raddeana*, *L. squarrosa*, and *L. thunbergiana* in Japan.
- Puccinia lactucae-debilis** Diet. Dull-brown powdery rust pustule on leaves of *L. debilis* and *L. stolonifera* in Japan.
- Puccinia lactucae-denticulatae** Diet. Powdery cinnamon-brown to chestnut-brown rust pustules on leaves of *L. denticulata* and *L. stolonifera* in Japan.
- Puccinia lactucae-repentis** Miy. Rust on leaves of *L. repens* in Japan.
- Puccinia lactucina** Syd. Yellow-brown to brown rust pustules on leaves of *L. scarioloides* in Persia.
- Puccinia lactucarum** Syd. Brown rust pustules on leaves and stems of *L. altissima*, *L. perennis*, *L. quercina*, *L. sagittata*, and *L. scariola* in Europe.
- Puccinia prenanthis** (Pers.) Lindr. Orange aecia and brown uredinia and telia on leaf blades and petioles of *L. muralis*, *L. sativa*, *L. stricta*, *L. virosa*, and *Prenanthes* spp. in Alaska, Europe, and Australia.
- Ramularia lactucae** Jaap. On leaves of *L. muralis* in Yugoslavia.
- Ramularia lactucosa** Lamb. and Fautr. Numerous small circular leaf spots on *L. sativa* in France.
- Ramularia lampsanae** (Desm.) Sacc. var. **lactucae-muralis** Jaap. Leaf spots on *L. muralis* in Germany.
- Septoria fernandezii** Unam. On leaves of *L. virosa* in Spain.
- Uredo proximella** Arth. Brown rust pustules on leaves of *L. intybacea* in Porto Rico, Cuba, and Dominican Republic.
- LAELIA**. Epiphytic orchids. See Orchidaceae.
- LAGENARIA**. GOURD. Cucurbitaceous vines.
- Phyllosticta lagenariae** Pass. On leaves of *L. leucantha* (*L. vulgaris*) in Italy.
- Phyllostictella cucurbitacearum** Tass. On leaves of *L. leucantha* (*L. vulgaris*) in Italy.
- LAGENOPHORA**. Small perennial scapose herbs.
- Puccinia lagenophorae** Cke. Brown rust pustules on leaves of *L. billardieri* in Australia.
- Septoria lagenophorae** McAlp. Circular to irregular ashen to dull-brown leaf spots on *L. billardieri* in Australia.
- Thecaphora lagenophorae** McAlp. Smut sori developing in the ovaries, converting entire flower heads into powdery, brown spore masses. On *L. emphysopus* in Australia.
- LAGERSTROEMIA**. CRAPE MYRTLE. Showy-flowered shrubs and trees.
- Cercospora lagerstroemiae** Syd. Circular to irregular gray-brown to white leaf spots on *L. speciosa* in the Philippines.
- Mollisia ravida** Syd. Effuse, pale, indefinite leaf spots on *L. speciosa* in the Philippines.
- Rhytisma lagerstroemiae** Rabh. Irregular black stromata on leaves of *L. indica*, *L. lanceolata*, and *L. speciosa* in India and the Philippines.
- Septoria lagerstroemiae** Sacc. and Scalia. Brown leaf spots on *L. indica* in Portugal.
- Uncinula australiana** McAlp. Powdery mildew on leaves and inflorescences of *L. indica* and *L. ovalifolia* in Japan and Australia.
- LAGURUS**. RABBIT TAIL GRASS. Hardy annual grasses.
- Puccinia laguri** Jasp. Rust on leaves of *L. ovatus* in Dalmatia.
- Septoria caruianiana** Sacc. Oblong pale ochraceous spots on leaves of *L. ovatus* in Malta.
- LAMIUM**. DEAD NETTLE. Annual and perennial herbs, some species cultivated as hardy border plants.
- Ascochyta lamiorum** Sacc. Ochraceous leaf spots on *L. album* in Italy.
- Ovularia lamii** (Fekl.) Sacc. On leaves of *L. album*, *L. amplexicaule*, and *L. purpureum* in Russia and Denmark.
- Peronospora lamii** A. Braun. Broadly effused dingy lilac, downy mildew patches on under leaf surfaces of *L. album*, *L. amplexicaule*, *L. maculatum*, *L. purpureum*, and *L. rubrum* in Europe and Tunis. Reported from New York.
- Phyllosticta albanica** Bub. Circular to irregular gray leaf spots with brown margins on *L. galeobdolon* in Yugoslavia.
- Phyllosticta lamii** Sacc. Whitish leaf spots with broad ochraceous surrounding zones on *L. album* and *L. ovata* in Siberia and Italy.
- Phyllosticta venziana** Mart. Small circular gray leaf spots on *Lamium* sp. in Italy.
- Ramularia exilis** Syd. Indefinite yellowish leaf spots on *L. galeobdolon* in Denmark and Germany.
- Ramularia lamicola** C. Mass. Angular, then confluent, leaf spots on *L. album* in Europe.
- Septoria didickel** Sacc. Circular to irregular, then confluent, white leaf spots with brown or dark-purple margins on *L. galeobdolon* in Denmark and Germany.
- Septoria lamicola** Sacc. and var. **intermedia** C. Mass. Whitish leaf spots with red margins on *L. album*, *L. amplexicaule*, *L. bithynicum*, *L. maculatum*, *L. ovata*, and *L. purpureum* in Europe.
- LANDOLPHIA**. PACURIA. Ag. Tropical shrubs mostly climbers.
- Colletotrichum paucipilum** Delacr. Brown anthracnose spots, often marginal, on leaves of *L. kleinii* in France and the Congo.
- Phyllosticta landolphiae** P. Henn. Marginal brown leaf spots on *L. kirkii* in Germany.
- Phyllosticta owariensis** Maubl. Small irregular whitish leaf spots with brown borders on *L. owariensis* in central Africa.
- LANSIUM**. LANSIA. Oriental trees, one species cultivated for its edible fruit.
- Bacterium citri** Haase. See Citrus.
- Corticium salmonicolor** B. and Br. See Citrus.
- LANTANA**. Shrubs, some species cultivated for the verberna-like flowers.
- Acanthostigma lantanae** Theiss. On leaves of *Lantana* sp. in Brazil.
- Aecidium lantanae** Mayor. Rust on leaves of *L. hispida* in Colombia.
- Phyllachora sororecula** Speg. Shiny black stromata on leaves of *Lantana* sp. in Paraguay and Brazil.
- Phyllosticta lantanae** Pass. Small irregular whitish leaf spots on *L. camara* and *Viburnum lantana* in Porto Rico and Italy.
- Prospodium tuberculatum** (Speg.) Arth. Powdery cinnamon-brown to black rust pustules on leaves of *L. camara*, *L. horrida*, *L. involucreta*, *L. lilacina*, *L. robusta*, *L. salvifolia*, and *L. tiliaefolia* in Mexico, South and Central America, and Cuba.
- Puccinia natalensis** Diet. and Syd. Dark-brown rust pustules on leaves of *L. salvifolia* and *L. trifolia* in Uganda and the Union of South Africa.
- Puccinia schimperiana** Syd. Rust on leaves of *L. citrifolia* in Abyssinia.
- Septoria lantanae** Gar. Small, irregular leaf spots on *L. camara* in Porto Rico.
- Uromyces dubiosus** P. Henn. Dark-brown powdery rust pustules on leaves of *Lantana* sp. in Brazil.
- LAPAGERIA**. CHILE BELLS. Half-hardy evergreen twiner.
- Phyllosticta jaffuelii** Speg. On leaves of *L. rosea* in Chile.

LAPEYROUSIA. African summer-flowering bulbs.

Uromyces anomathecae Cke. Brown elliptical rust pustules on both leaf surfaces of *L. cruenta* in Natal.

Uromyces delagoënsis Bub. Round to oblong golden-yellow to black rust pustules on both surfaces of leaves of *L. delagoënsis* in Mozambique and Portuguese East Africa.

LAPORTEA. Perennial herbs, shrubs or trees.

Aecidium laporteei P. Henn. Rust pustules on circular yellow-brown leaf spots on *L. bulbifera* in Japan.

LARDIZABALA. Evergreen climbers of Chile.

Physalospora lardizabala Speg. On leaves of *L. biternata* in Chile.

Septoria lardizabala P. Henn. Effuse brown leaf spots on *L. biternata* in Germany.

LARIX. LARCH. TAMARACK. Timber and ornamental trees.

Cercospora exosporioides Bub. On needles of *L. decidua* (*L. europaea*) in Bohemia.

Dasycephala calycina (Schum.) Fekl. (*D. willkommii* Hartig.) This fungus causes the most serious disease of the larch known and is widespread throughout Europe. It has also been reported from Newfoundland, and there is grave danger of the introduction of the parasitic strain of the fungus into the United States. The fungus occurs commonly on dead branches, the fruiting bodies appearing as small orange or yellow cups. It attacks living parts of the tree through wounds, and develops large, perennial cankers. The bark becomes black, resin oozes out and a flattened cavity forms, which increases year by year. Young trees may be girdled and killed, older trees persist often for years, but their merchantable value is lowered or destroyed, particularly when several or more cankers occur along the trunk. The hosts are *L. decidua*, *L. occidentalis*, *Abies pectinata*, *Pinus austriaca*, *P. laricina*, *P. pumilo*, and *P. sylvestris*.

Exostriola laricis Oud. Very injurious to leaves of *Larix* sp. in Holland.

Hartigiella laricis (Hart.) Syd. (*Meria laricis* Vuill.) Brown spots on needles, particularly those on young shoots of *Larix* sp. in Russia, France, and Germany. Leaf fall results, and often the death of young trees.

Heterosporium laricis Cke. and Mass. On leaves of *L. decidua* in Great Britain.

Hypodermella laricis Tub. On needles of *L. decidua* in Switzerland and Germany, causing a dwarfing of shoots and dying of foliage.

Melampsora larici-capraearum Kleb. See *Salix*.

Melampsora larici-epitea Kleb. See *Salix*.

Melampsora larici-pentandrae Kleb. See *Salix*.

Melampsora larici-populina Kleb. See *Populus*.

Melampsora laricis R. Hart. See *Populus*.

Melampsora larici-urbaniana Mat. See *Salix*.

Melampsoridium betulae (Schum.) Arth. This rust attacks the needles of *L. europaea* in Europe, Siberia, and Japan. The alternate stages occur on *Betula* spp. and have been reported from the United States.

Mycosphaerella laricina R. Hartig. Brown patches on needles of *L. decidua* in Germany, causing premature leaf fall.

Myxosporium abietinum Rostr. On bark of *L. decidua*, *Picea sitchensis*, *Pinus strobus*, and *Pseudotsuga taxifolia* in Denmark.

Phoma laricis Oud. On leaves of *L. decidua* in Denmark.

Phoma lineolata Desm. On branches and cone scales of *L. decidua* in Europe.

Phomopsis pseudotsugae Wils. See *Pseudotsuga*.

Phyllosticta lineolata Desm. On *L. decidua* in Germany.

Phytophthora fagi R. Hart. See *Fagus*.

LASIA. Perennial herbs, woody at the base.

Mycosphaerella lasiana Sacc. On leaves of *L. heterophylla* in Malaya.

LATANIA. Palm. See *Palmae*.

LATHYRUS. (Including *Orobus*). PEA. Sweet pea. Annual and perennial climbing or upright herbs and shrubs.

Ascochyta lathyri Trail. and var. **lathyri-odorati** Bub. and Kab. Large brown areas on leaves of *L. annuus*, *L. nissolia*, *L. odoratus*, and *L. sylvestris* in Scotland, Denmark, Bohemia, Italy, and Spain.

Ascochyta orobi Sacc. Irregular whitish leaf spots with dull-brown margins on *L. niger*, *L. (Orobus) vernus*, *Orobus lathyroides*, and *O. vernus* in Siberia, Russia, Italy, and Austria.

Cladosporium album Dowson. Irregular sunken faint-buff spots on both leaf surfaces and wings of stems of *L. odoratus* in Great Britain.

Cylindrosporium lathyri Bub. and Kab. Irregularly circular to angular ochraceous leaf spots with brown-purple margins on *L. vernus* in Italy and Austria.

Cylindrosporium orobicolum (Sacc.) Bub. Large indefinite brown-margined leaf spots on *L. variegatus*, *Orobus venetus*, and *O. vernus* in Russia, Bohemia, and Italy.

Entyoloma feurichii Krieg. Small sori in confluent brown leaf spots on *L. sylvestris* in Germany.

Gleosporium orobi Karst. Irregular dull-brown leaf spots on *Orobus vernus* in Finland.

Isariopsis carnea Oud. Leaf spot on *L. pratensis* in Italy and Holland.

Mycosphaerella lathyri A. Pot. Indefinite irregular ochraceous leaf spots on *L. pisiformis* in Siberia and Russia.

Ovularia deusta (Fekl.) Sacc. Large dark-brown areas on leaves of *L. pratensis*, *L. tuberosus*, and *Tetragonolobus siliquosus* in Siberia, Sweden, Italy, France, and Germany.

Peronospora fulva Syd. Downy mildew on leaves of *L. pratensis* and *L. vernus* in central and northern Europe.

Peronospora lathyri palustris Gäum. Downy mildew on leaves of *L. heterophyllus*, *L. palustris*, and *L. sativus* in Sweden, Switzerland, and India.

Peronospora orobi Gäum. Downy mildew on leaves of *L. montanus* and *L. tuberosus* in central and northern Europe.

Peronospora senneniana Sacc. Downy mildew on leaves of *L. nigrus* in Spain, Switzerland, Bohemia, Denmark, Norway, Austria, Poland, and Germany.

Phyllachora lathyri (Lév.) Theiss. and Syd. Black, crust-like stromata on both leaf surfaces of *L. grandiflorus*, *L. latifolius*, *L. pratensis*, *L. tuberosus*, *Onobrychis sativa*, *O. viciaefolia*, and *Vicia tenuifolia* in Siberia, Russia, Asia Minor, and Austria.

Phyllosticta lathyricola Bub. and Krieg. Small circular to irregular dull purple-brown leaf spots on *L. sylvestris* in Germany.

Phyllosticta lathyrina Sacc. and Wint. Pale ochraceous leaf spots with brown margins on leaves of *L. sylvestris* in Russia, Italy, and Switzerland.

Phyllosticta orobella Sacc. On leaves of *L. vernus* in Russia and Italy.

Physalospora lathyri (D. and M.) Sacc. On leaves of *L. amphicarpos* in Algeria.

Placosphaeria onobrychidis Sacc. See *Lupinus*.

Ramularia lathyri Höll. Oblong yellowish leaf spots on *L. hirsutus* in Hungary.

LATHYRUS—Continued.

- Rhizoctonia napi** West. See Brassica.
Rhytisma onobrychis Sacc. See Lupinus.
Septogloeum lathyri Lind. Large confluent pale-reddish leaf spots on leaves of *L. sylvestris* in Denmark.
Septoria fatreyana Sacc. On leaves of *L. sylvestris* in Italy.
Septoria fulvescens Sacc. Brown confluent leaf spots on *L. maritimus*, *L. pisiformis* and *L. sylvestris* in Russia, Bohemia, Italy, and Germany.
Septoria podgoricensis Bub. Circular to elliptical yellowish spots on leaves and stems of *L. angulatus* and *L. annuus* in Spain and Yugoslavia.
Septoria silvestris Pass. Leaf spots on *L. pratensis*, *L. sylvestris*, *L. tuberosus* and *Vicia sepium* in Sweden, Russia, Denmark, and Italy.
Septoria stipularis Pass. See Lotus.
Uromyces clavatus Diet. Yellow and cinnamon-brown rust pustules on leaves of *L. magellanicus*, *L. multiceps*, and *Vicia tenuifolia* in Chile, Argentina, and Brazil.
Uromyces lathyrinus Speg. Yellow and dark-brown rust pustules on leaves and stems of *L. clymenum* in Argentina.
Uromyces orobi (Pers.) Lév. Brown rust pustules on *L. cicera*, *L. latifolius*, *L. montanus*, *Orobis tuberosus*, and *Vicia unijuga* in Siberia and Europe.
Uromyces pisi (Pers.) De By. See Pisum.

LAURUS. LAUREL. SWEET BAY. Trees.

- Ascochyta laurina** Tass. Marginal white leaf spots with rufous margins on *L. mollis* in Italy.
Cercospora unicolor Sacc. and Penz. On leaves of *L. nobilis* in France.
Exobasidium lauri Geyl. Long sulcate excrescences on trunks of *L. canariensis* and *L. nobilis* in Portugal, Italy, Brazil, and the Canary Islands.
Gloeosporium nobile Sacc. Circular dull-yellowish anthracnose spots on leaves of *L. nobilis* in Italy and Austria.
Metasphaeria nobilis Sacc. Whitish leaf spots with dull-brown margins on *L. nobilis* in Italy and Portugal.
Phragmonaevia lauri Pat. Circular, then confluent, rufous to whitish leaf spots with brown margins on *L. nobilis* in Tunis.
Phyllosticta alantella Sacc. On leaves of *L. nobilis* in Spain and France.
Phyllosticta laurina d'Alm. Irregular dull-brown leaf spots with darker borders on *L. nobilis* in Portugal.

LAVANDULA. LAVENDER. Perennial herbs, subshrubs, and shrubs.

- Phoma lavandulae** Brierley. Diseased buds and shoots turn brown, the leaves wither and fall and the epidermis of the twigs splits away in shreds. On *L. spica* (*L. officinalis*) in Great Britain.
Septoria lavandulae Desm. Circular to irregular white leaf spots with raised purple margins on *L. spica*, and *L. stoechas* in Great Britain, France, Italy, Denmark, and Madeira.

LAVATERA. TREEMALLOW. Annual and perennial flower-garden herbs and some ornamental shrubs.

- Bamularia daniloi** Bule. Angular yellowish leaf spots becoming white on *L. rotundata* and *L. thuringiaca* in Spain and Montenegro.
LEDUM. LABRADOR TEA. Shrubs cultivated for their white flowers and evergreen foliage.

- Ascochyta ledicola** Oud. On leaves of *L. palustre* in Holland.
Coccomyces ledi Rehm. On living branches of *L. palustre* in Sweden.
Phyllosticta ledi Rostr. White to gray leaf spots with purple borders on *L. groenlandicum* in Greenland.
Sclerotinia heteroica Wor. A heteroecious species, one stage attacking the flowers and ovaries of *L. palustre*, the other shoots of *Vaccinium uliginosum* in Finland, Sweden, Russia, and Germany.

LENS. LENTIL. Herbs.

- Peronospora lentis** Gäum. Downy mildew on leaves of *L. esculenta* in Germany.
Rhizoctonia napi West. See Brassica.
Uredo lentis Lagerh. Dark brown, elongate rust sori on leaves and stems of *L. esculenta* in Ecuador.
Uromyces viciae-cracca Const. See Vicia.

LEONOTIS. LION'S-EAR. Shrubs cultivated for their scarlet or orange flowers.

- Aecidium leonotidis** P. Henn. Rust on leaves of *L. velutina* in central Africa.
Puccinia leonotidicola P. Henn. Brown rust pustules on leaves of *L. nepetifolia* in Porto Rico, Cuba, Bahamas, Jamaica, central and southern Africa.
Uredo cancerina P. Henn. A rust producing large cancerous swellings on the stems, which break irregularly and finally become woody. On *L. velutina* in Abyssinia.
Uredo leonoticola P. Henn. Rust on leaves of *Leonotis* sp. in Brazil.

LEONTODON. See Taraxacum.**LEONTOPODIUM**. EDELWEISS. Perennial herbs.

- Puccinia leontopodii** Vogl. Rust on leaves of *L. alpinum* in Switzerland.

LEPIDAGATHIS. Tropical herbs and shrubs.

- Aecidium lepidadathis** Syd. Rust on brown leaf spots on *L. hyalina* in India.

LEPTODERMIS. Low shrubs with small white or purple flowers.

- Coleosporium leptodermidis** (Barcl.) Syd. Golden rust pustules on leaves of *L. lanceolata* in India.

LEPTOSPERMUM. AGONIS Ag. Ornamental woody plants grown for their flowers and myrtlelike foliage.

- Phyllochora callistemoni** Rodw. Tar spot on leaves of *P. scoparium* in Tasmania.

LESCHENALTIA. Herbs and heathlike Australian plants.

- Puccinia gilgiana** P. Henn. Yellow and brown to black rust pustules on petioles, stems, and calyces of *L. linarioides* in Australia.

LESPEDEZA. BUSH CLOVER. Forage plants.

- Uromyces rugulosus** Pat. Black rust pustules on leaves of *L. yunnanensis* in China.
Woroninella vulcanica Wor. On *L. cytioides* in Java.

LEUCADENDRON. SILVER TREE. Shrubs or trees, the leaves densely covered with white silky hairs.

- Cercospora protearum** Cke. Circular dark-brown leaf spots on *L. argenteum* in the Union of South Africa.

LEUCAENA. Shrubs with Acacia-like foliage.

- Exosporium leucaenae** Stev. and Dal. Circular dull-yellow leaf spots on *L. glauca* in Porto Rico.
Ravenelia leucaenae Long. Brown powdery rust sori on leaves of *L. diversifolia* and *L. esculenta* in Mexico.
Ravenelia leucaenae-microphyllae Diet. Brown rust pustules on leaves of *L. angustissima* and *L. microphylla* in Mexico and Guatemala.
Ravenelia verrucosa Cke. and Ell. Yellow-brown to brown rust pustules on leaves of *L. lanceolata* in Mexico.

LEUCANTHEMUM. See Chrysanthemum.

LEUCOJUM. SNOWFLAKE. Hardy bulbous plants.

Puccinia schmidtiana Diet. See Phalaris.

Ramularia ucrainica Petr. Brown to gray-brown leaf spots on *L. vernum* in southeastern Europe.

Septoria malisora Bub. Elongate (up to 3 centimeters) gray-purple leaf spots with brown margins on *L. aestivum* in Yugoslavia.

LEUCOTHOE. Ornamental shrubs grown for their handsome evergreen leaves and white flowers.

Coniothyrium leucothoes P. Henn. Circular, then confluent, whitish leaf spots with brown margins on *Leucothoe* sp. in Brazil.

Exobasidium leucothoes P. Henn. Deforms leaves and branches of *Leucothoe* sp. in Brazil.

Rhytisma leucothoes P. Henn. Black stromata on leaves of *Leucothoe* sp. in Brazil.

LEUZEA. See Centaurea.

LEVISTICUM. LOVAGE. Perennials grown for their aromatic seed.

Bacterium levistici Osterw. Small, somewhat sunken, brown spots which are confluent, forming large irregular areas with yellow margins. On leaf blades and petioles and stems of *L. officinale* in Germany.

Puccinia bornmülleri P. Magn. Powdery dark-brown rust pustules on leaf blades and petioles and stems of *L. persicum* in Persia.

Ramularia levistici Oud. Leaf spots on *L. officinale* in Holland.

Ramularia schroeteri P. Syd. Subcircular ochraceous leaf spots with whitish margins on *L. officinale* in Yugoslavia and Austria.

Ramularia vestergreniana Allesch. Large brown marginal leaf spots on *L. officinale* in Russia and Finland.

Septoria levistici West. Circular to irregular broad brown, then whitish, leaf spots on *L. officinale* in Belgium and Spain.

LEYCESTERIA. HIMALAYA HONEYSUCKLE. Small Chinese shrubs with small flowers in whorls.

Septoria leycesteriae Sacc. and Speg. Subcircular to irregular ochraceous leaf spots with brown borders on *L. formosa* in Italy and Germany.

LICUALA. Dwarf fan palms. See Palmae.

LIGHTFOOTIA. African shrubs or herbs with small white, blue, or pink flowers.

Uredo campanularum Cke. and Mass. Rust on leaves of *Lightfootia* sp. in Madagascar.

LIGULARIA. Perennial herbs.

Aecidium nikkense P. Henn. and Shir. Yellow rust pustules on leaves of *L. stenocephala* in Japan.

Coleosporium ligulariae Thuem. Golden rust pustules on leaves of *L. sibirica* in Russia, Finland, Siberia, and Denmark.

Puccinia paulsenii Syd. Brown leaf rust on *L. altaica* in central Asia.

LIGUSTICUM. Hardy herbs used for naturalizing.

Puccinia aphanicandra Lindr. Powdery dark-brown rust pustules on *L. alatum* in Armenia.

Puccinia ligusticola S. Miy. Brown to black rust pustules on leaf blades and petioles of *L. scoticum*, *L. ibukienensis*, and *L. scoticum* in Japan.

LIGSTRUM. PRIVET. Ornamental woody plants cultivated for their handsome foliage and as hedge plants.

Aecidium klugkistanum Diet. Rust on leaves of *L. japonicum* in Japan.

Ascochyta ligustri Sacc. and Speg. (*Ascochyta ligustrina* [Pass.] Tass.) Dull-yellow leaf spots on *L. vulgare* in Italy and France.

Cercospora ligustri Roum. Numerous circular to irregular yellow leaf spots on *L. vulgare* in France.

Phyllosticta ligustri Sacc. Leaf spots on *L. vulgare* in Denmark, Italy, Portugal, and Austria.

Phyllosticta ligustrina Sacc. and Speg. Leaf spot on *L. vulgare* in France and Italy.

Phytophthora syringae Kleb. See Syringa.

Puccinia obtusata (Oth.) Ed. Fisch. See Phragmites.

Ramularia ligustrina Maubl. Circular to oval dull-yellow leaf spots with dark-brown margins on *Ligustrum* sp. in France.

Septoria japonica Thuem. On leaves of *L. japonicum* and *L. vulgare* in Italy and Germany.

Septoria ligustri (Desm.) Kicks. Pale ochraceous irregular leaf spots with reddish margins on *L. vulgare* in Belgium and France.

Septoria ligustrina Sacc. On leaves of *L. vulgare* in Bohemia.

LILIUM. Lily. Scaly bulbous plants with showy flowers.

Aecidium lili-cordifolii Diet. Circular to irregular spots are produced on both leaf surfaces up to 1 centimeter in diameter, in which yellow rust pustules appear. On *L. cordifolium* in Japan.

Aecidium safranoffianum Thuem. Yellow rust pustules appear on leaves of *L. martagon* in Siberia.

Bacillus lili Uyd. This bacterial disease attacks the leaves and bulb scales. The spots are deep brown in color, oval to elongate or more or less irregular on the bulb scales. The entire leaf blade soon becomes involved, turns brown, and droops. When young bulbs are attacked flowering shoots fail to develop. On *L. auratum*, *L. cordifolium*, *L. elegans*, *L. hansonii*, *L. longiflorum*, *L. meaeoloides*, *L. rowlettii*, *L. speciosum*, *L. tigrinum*, and *L. venustum* in Japan.

Botrytis liliorum Fujikuro. This fungus attacks mostly the leaves, first appearing as small spots about 1 millimeter in diameter, rapidly spreading over the entire surface, causing total decay of the host plant. "As bad as any three other lily diseases in Japan." On *L. longiflorum* in Japan.

Cercospora liliicola (Rich.) Sacc. On stems of *L. candidum* in France.

Cylindrosporium inconspicuum Wint. (*Cercospora lili inconspicua* [Wint.] v. Hoeh.) (*Cercospora hungarica* Baume.) Causes irregular, often confluent, spots on the leaves which are dark-brown, becoming paler at the center. On *L. martagon* and other species in France, Switzerland, and Austria.

Mycosphaerella cinxia Sacc. (*Sphaeria matorina* Sacc.) Causes a leaf blight of *L. candidum* in Italy.

Phyllosticta liliicola Sacc. A leaf-spot disease of *L. candidum* in Italy.

Puccinia sp. Intercepted on *Lilium* sp. from Ecuador.

Rhizopus necans Mass. A wound parasite, causing a soft rot of the bulbs, particularly during shipment. On *L. auratum*, *L. lancifolium*, *L. speciosum*, and other species in Japan, Great Britain, and Denmark.

Uromyces acidiformis (Strauss.) Rees. (*Uromyces lili Kunze*.) This rust produces yellow-brown to chestnut-brown rust pustules on dead spots on leaves, petioles, and stems. On *L. bulbiferum*, *L. candidum*, *L. carnaticum*, and *L. croceum* throughout central Europe and on *Fritillaria involucreta* in France.

Uromyces holwayi Lagerh. Leaf rust on *L. dauricum* and *L. maximowiczii* in Japan.

Uromyces japonicus Syd. Cinnamon-colored powdery rust pustules, later becoming darker on the upper surfaces of leaves of *L. maximowiczii* and *L. tigrinum* in Japan.

Uromyces lili (Lk.) Eckl. Brown rust pustules are produced in sunken spots on both surfaces of the leaves of *Fritillaria delphinensis*, *F. graeca*, *F. meleagris*, *F. messanensis*, *F. minor*, *F. ruthenica*, *Lilium auratum*, *L. bulbiferum*, *L. canadense*, *L. candidum*, *L. carnaticum*, and *L. croceum* in France, Switzerland, Italy, Yugoslavia, Russia, Montenegro, Spain, Japan, Austria, Hungary, Dalmatia, and Germany. Reported from Washington.

LINARIA. TOADFLAX. Low herbs.

Cintractia cingens (Berk.) De T. Smut sori on stems and leaves of *L. genistifolia* and *L. vulgaris* in Austria.

Peronospora flava Güm. Downy mildew on leaves of *L. vulgaris* in central and north Europe.

Peronospora linariae Fekl. Downy mildew on leaves of *L. arvensis* and *L. minor* in central and north Europe.

Phyllosticta linariae Sacc. See Antirrhinum.

Puccinia trabuti Roum. and Sacc. See Phragmites.

Septoria cymbalariae Sacc. and Speg. Circular dull-brown leaf spots on *L. cymbalaria* and *L. nivea* in Italy, France, and Spain.

Septoria gandulphi Sacc. and D. Sacc. Small orbicular to angular leaf spots on *L. commutata* in Italy.

Septoria lapadensis Jaap. Yellow-brown leaf spots on *L. commutata* in Dalmatia.

Ustilago hypogaea Tul. Doubtful smut species on *L. spuria* in the Isle of Wight.

LINUM. FLAX. Fiber plant. Several species cultivated as ornamentals.

Ascochyta lini Rostr. On stems of *L. catharticum* in Denmark.

Cercospora lini Wor. On leaves of *L. nervosum* in Russia.

Colletotrichum lini (Westerd.) Toehnal. (*C. linicolum* Pethybr. and Laff.) Irregular, shrunken, discolored cankers occur on seedlings of *L. usitatissimum*, involving the cotyledons and causing damping off. Similar dark rusty-brown areas occur on stems, leaves, calyces, buds, and capsules of older plants. The disease is carried in the seed. It has been reported from Ireland, Great Britain, Japan, and Formosa. It is not clear whether the flax anthracnose reported from the United States is identical.

Cylindrosporium melitense Sacc. On leaves and flowers of *L. strictum* in Italy.

Entyoma lini Oud. Smut sori on yellowish circular leaf spots on *L. usitatissimum* in Holland.

Fusicladium lini Sor. On leaves of *L. usitatissimum* in Belgium and Germany.

Oidiaster radialis (Willd.) Pascher. (*Asterocystis radialis* Willd.) This fungus attacks the roots of *L. usitatissimum*, *Brassica* spp., *Plantago* spp., and *Veronica* spp. in Europe, Argentina, and Morocco, causing a yellowing and stunting of the plants.

Peronospora lini Schroet. Downy mildew on leaves of *L. catharticum* in Scandinavia, Finland, Russia, Austria, Switzerland, and Germany.

Phyctaena linicola Speg. Yellowish areas on leaves and stems of *L. usitatissimum* in Argentina. Recently reported from several localities in the United States as a serious disease.

Phoma sp. On stems of *L. usitatissimum* in Ireland.

Pleosphaerulina lini Lebed. On leaves and stems of *L. perenne* in Russia.

Polyspora lini Pethybr. The brown or stem-break disease of flax (*L. usitatissimum*) is serious in Ireland and is a seed-borne disease. The cotyledons are first attacked, the disease spreading successively to leaves, stems, branches, and seed bolls. Stem lesions may cause half-grown plants to break over and die. On older plants numerous brown spots cover the various organs, diseased areas in a field taking on a brown appearance rather than the golden-yellow color produced by healthy plants. Stems develop a brittleness, indicating injury to the fiber. The disease also occurs in Great Britain.

Ramularia lini Lebed. On leaves and stems of *L. perenne* in Russia.

Rhizoctonia napi West. See Brassica.

Septogloeum linicola Speg. Pale effuse areas on leaves and stems of *L. usitatissimum* in Argentina.

Tylenchus dipsaci Kuehn. See Narcissus.

La brûlure or viasbrand disease of flax is a widely distributed disease of flax (*L. usitatissimum*) in Belgium, France, and Ireland, due to an unknown cause.

LIPARIS. TWAYBLADE. Terrestrial and epiphytic orchids. See Orchidaceae.**LIPPIA.** LEMON VERBENA. Herbs and shrubs grown for their ornamental flowers. One species used for ground cover or lawns.

Aecidium evansi P. Henn. Rust on leaves of *L. asperifolia* in the Union of South Africa.

Phyllosticta aloysiae Speg. Subcircular dull-brown leaf spots with white centers on *L. citriodora* in Argentina.

Prospodium lippiae (Speg.) Arth. Powdery golden-yellow rust pustules on *L. asperifolia*, *L. calli-carpaefolia*, *L. dulcis*, *L. lycioides*, *L. myriocephala*, *L. strigosa*, and *L. umbellata* in Argentina, Mexico, Guatemala, Cuba, and Costa Rica.

Puccinia accedens Syd. Powdery brown rust pustules on leaves of *L. aristata* and *L. arvensis* in Brazil and Uganda.

Puccinia conjuncta Diet. and Holw. Brown rust pustules on leaf blades and petioles and on stems of *L. pringlei* in Mexico.

Puccinia elatipes Arth. and Holw. Rust on leaves of *L. myriocephala* in Costa Rica and Guatemala.

Puccinia lippicola Pat. and Har. Brown rust pustules on circular brown spots on *L. adoensis* in central Africa.

Puccinia lippilvora Syd. Rust on leaves of *Lippia* sp. in the Congo.

Puccinia paraguayensis Speg. Powdery cinnamon-brown rust pustules on *L. lycioides* in Brazil.

Puccinia permagma Arth. and Holw. Brown leaf rust on *L. myriocephala* in Costa Rica.

Puccinia peruviana Syd. Powdery dark-brown rust pustules on leaves of *L. urticoides* in Peru.

Puccinia senilis Arth. Yellow to brown powdery rust pustules on leaves of *L. myriocephala* in Guatemala.

Puccinia vunguenti Mayor. Rust on leaves of *L. americana* in Colombia.

Uredo lippiae Diet. and Holw. Brown, powdery rust pustules on leaves of *L. pringlei* in Mexico.

LIRIODENDRON. TULIP TREE. YELLOW-POPLAR. Timber and ornamental trees.

Diaporthe delitescens Bomm., Rouss., and Sacc. On *L. tulipifera* in Belgium.

Mycosphaerella elatior Sacc. and Speg. See Magnolia.

Myxosporium tulipiferae Died. On twigs of *L. tulipifera* in Germany.

LISIANTHUS. Herbs or shrubs cultivated for their flowers.

Phyllosticta lisianthi Syd. Circular to subcircular, brown leaf spots with raised margins on *L. eszerius* in Jamaica.

Uredo lisianthi Pat. Rust on leaves of *L. elegans* in Brazil.

LISTERA. Terrestrial orchids. See Orchidaceae.**LITHOCARPUS.** See Pasania.**LITHOSPERMUM.** GROWWELL. Herbaceous, hardy perennials, used in rock gardens.

Aecidium lithospermi Thuem. Rust on leaves of *L. arvense* and *L. purpureocaruleum* in Austria and Macedonia.

Peronospora lithospermi Güm. Downy mildew on leaves of *L. arvense* in central and north Europe.

Puccinia arnaldi Har. and Diet. Rust on leaves, stems, and sepals of *L. fruticosum* in France.

Ramularia lithospermi Lebed. On leaves of *L. officinale* in Russia.

LITSEA. Trees or shrubs.

Asterocaulium saccardoi Syd. Forms yellow galls on leaves of *L. glaucescens* in Mexico.

Cercospora litseae P. Henn. Circular dull-brown leaf spots on *L. glauca* in Japan.

Cercospora litseae-glutinosa Syd. On leaves of *L. glutinosa* in the Philippines.

Endothella litseae Racib. Black shiny irregular to circular stromata on leaves of *L. javanica* in Java.

Gloeosporium litseae Petch. Anthracnose on *Litsea* sp. in Ceylon.

Helminthosporium maculosum Sacc. On *L. perrottetii* in the Philippines.

Laestadia litseae (B. and Br.) Cke. Small pale-brown leaf spots on *Litsea* sp. in Australia.

Oligostroma apiculatum (Sacc. and Berl.) Theiss. and Syd. Shiny black somewhat irregular stromata on leaves of *L. dealbata* in Australia.

Phyllachora cantanensis Syd. Black stromata on leaves of *L. glutinosa* in China.

Phyllachora laurinearum Rac. Circular to elliptical black stromata on yellow-green subcircular leaf spots on *L. chrysocoma* and *Tetranthera* sp. in Java.

Phyllachora lepidia Syd. Black stromata on leaves of *L. tayabensis* in the Philippines.

Phyllachora litseae Koord. Black stromata on indefinite yellowish spots on *L. polyantha* in Java.

Puccinia litseae (Pat.) Diet. and P. Henn. Chestnut-brown rust pustules on leaves of *L. glauca* in Japan.

Teichosporella negeriana Sacc. and Syd. Black fungus layers on leaves of *L. caustica* in Chile.

Venturia litseae Syd. On leaves of *L. glutinosa* in the Philippines.

LIVISTONA. Fan palms. See Palmae.**LLOYDIA.** Small bulbous plants.

Asteroma lloydiae Cruchet. Pale-brown, then darker, areas on stems and leaves of *L. serotina* in Switzerland.

Puccinia bessei Cruchet. Dull-brown rust pustules on leaves and stems of *L. serotina* in Switzerland.

LOASA. Flower-garden annuals.

Aecidium rehderianum Magn. Rust on leaves of *L. aurantiaca* and *L. papaverifolia* in Germany.

Septoria loasae F. Tassl. On leaves of *L. vulcanica* in Italy. Spegazzini has given the same name to a species on *Loasa* sp. in Argentina.

LOBELIA. Annual and perennial flower-garden and border plants.

Ascochyta lobeliae Petch. On *L. nicotianaeifolia* in Ceylon.

Cercospora ochracea Sacc. and Malbr. On leaves of *L. urens* in France.

Cercospora (?) tupae Speg. Pale, indefinite spots on leaves of *L. tupa* in Chile.

Gloeosporium hawaiiense Thuem. Anthracnose on leaves of *L. macrostachys* in Hawaii.

Heterosporium lobeliae Speg. Indefinite leaf spots on *L. salicifolia* in Chile.

Heterosporium tupae Speg. Indefinite leaf spots on *L. bridgesii* in Chile.

Mycosphaerella lobeliae Petch. On leaves of *L. nicotianaeifolia* in Ceylon.

Phoma devastatrix Berk. On *Lobelia* (cult.) in Great Britain.

Phyllosticta bridgesii Speg. Irregular whitish leaf spots on *L. bridgesii* in Chile.

Puccinia aucta Berk. and Muehl. Yellow to brown rust pustules on leaves of *L. anceps*, *L. pedunculata*, and *L. platycalyx* in Australia.

Puccinia vacua Diet. and Holw. Brown rust pustules on leaves of *Lobelia* sp. in Mexico.

Rhizoctonia destruens Tass. See Solanum.

LOESELIA. Greenhouse plants related to *Gilia*.

Puccinia fumosa Holw. Brown to black rust pustules on leaves of *L. ciliata* and *L. coccinea* in Costa Rica and Guatemala.

LOLIUM. DARNEL, RYE GRASS. Pasture grasses.

Cladochytrium caespitis Griff. and Maubl. Attacks young plants of *Lolium* sp. in France, causing stems and leaf sheaths to turn brown, accompanied by a rotting of the roots.

Fusarium loliae Duc. Attacks the leaves of *L. italicum* in France.

Fusarium nivale (Fr.) Sor. See Secale.

Hypochnus fuciformis (Berk.) McAlp. Produces bright-pink glutinous fungus layers on spikelets of *Agropyron scabrum*, *Agrostis alba*, *Bromus mollis*, *B. sterilis*, *Danthonia pilosa*, *Festuca bromoides*, *Lolium perenne*, *Medicago denticulata*, and *Silybum marianum* in Great Britain and Australia.

Ovularia lolii Volk. Subcircular red-brown leaf spots on *L. italicum* and *L. perenne* in France and Switzerland.

Ovularia pulchella (Ces.) Sacc. and var. *lolii-italici* Ferr. See Dactylis.

Puccinia brachypus Speg. See Bromus.

Sclerospora macrospora Sacc. See Triticum.

Septoria lolii (Cast.) Sacc. Small subcircular blackish spots on glumes of *L. perenne* in France.

Septoria passerinii Sacc. See Hordeum.

Sorosporium lolii Thuem. A smut forming black spore masses in ovaries of *L. perenne* in Austria.

Thecaphora westendorpii Fisch. Dark-brown, irregular smut sori in spikelets of *L. perenne* in Belgium and Germany.

Tilletia lolii Awd. Stinking smut in ovaries of *L. arvense*, *F. lincolnum*, *L. multiflorum*, *L. perenne*, *L. remotum*, and *L. temulentum* in Europe.

Typhula graminum Karst. Forms small black sclerotia in yellowed leaves, resulting at times in destruction of seedlings of *Hordeum* spp., *L. multiflorum*, *L. perenne*, *L. temulentum*, and *Triticum* sp. in Great Britain, Sweden, and Germany.

Ustilago lolii Magn. A smut deforming the inflorescences of *L. temulentum* in Egypt.

Urocystis bolivari Bub. and Frag. A smut of *L. perenne* in Spain, producing systemic infections and preventing the development of normal heads.

LONGCHOCARPUS. LANCEPOD. Tropical trees or climbing shrubs.

Aecidium menyharthi P. Henn. Rust on leaves of *Lonchocarpus* sp. in South Africa.

Diorchidium manaosense P. Henn. Dull-brown rust pustules on leaves of *L. rariflorus* in Brazil.

Endodothella lonchocarpicola (P. Henn.) Theiss. and Syd. Dull-black stromata on lower leaf surfaces of *Lonchocarpus* sp. in Brazil.

Ophiodothella atomaculans (P. Henn.) v. Hoehn. Black stromata on both leaf surfaces of *Lonchocarpus* sp. in Brazil.

Phyllachora lonchocarpi Pat. and Har. Tar spot on leaves of *Lonchocarpus* sp. in the Congo.

Phyllachora lungusaensis P. Henn. Dull-black stromata on leaves of *Lonchocarpus* sp. in Tanganyika.

Ravenella bakeriana Syd. Brown rust pustules on leaves of *Lonchocarpus* sp. in Brazil.

Ravenella lonchocarpi Lagh. and Diet. Cinnamon-brown rust pustules on leaves of *L. campestris* and *L. latifolius* in Brazil, Cuba, and Guatemala.

LONGICERA. HONEYSUCKLE. Ornamental shrubs and climbers.

Actinonema loniceræ-alpigenae Allesch. Small circular to irregular dark-brown leaf spots on *L. alpigena* in Germany.

Loniceræ—Continued.

- Ascochyta periclymeni** Thuem. Circular dull-yellow to gray leaf spots on *L. periclymenum* in Portugal.
- Ascochyta sarmenticia** Sacc. Subcircular to angular whitish leaf spots on *L. caprifolium* and *L. xylosteum* in France.
- Ascochyta tatarica** Allesch. On *L. tatarica* in Germany.
- Ascochyta vulgaris** Kab. and Bub. Circular to angular yellow-brown leaf spots with purple-brown margins on *L. xylosteum* in Austria.
- Cercospora periclymeni** Wint. On leaves of *L. caprifolium* and *L. periclymenum* in Russia, Denmark, and Germany.
- Criella loniceræ** P. Henn. and E. Nym. Brown concave spots on leaves of *Loniceræ* sp. in Java.
- Doidhella periclymeni** (Fekl.) Theiss. and Syd. On *L. periclymenum* in Germany.
- Kabatia latemarensis** Bub. Dull-yellow to white leaf spots with broad purple margins on *L. caerulea* and *L. xylosteum* in Italy and Austria.
- Kabatia mirabilis** Bub. Angular dull-yellow to whitish leaf spots with black margins on *L. alpigena* and *L. nigra* in Austria.
- Lasiobotrys loniceræ** Kunz. and Schm. Circular black spots on leaves and branches of *Loniceræ* spp. in India, Siberia, and Europe. What has been referred to this species in California is distinct.
- Leptosphaeria periclymeni** Oud. On branches of *L. periclymenum* in Russia.
- Marsonia staritzii** Bres. Small, then confluent, black leaf spots on *L. tatarica* in Germany.
- Melasma loniceræ** Jacz. Black stromata on leaves of *L. maackii* and *L. mazimowiczii* in Austria.
- Mycosphaerella elymenia** Sacc. Subcircular ochraceous to gray leaf spots on *L. caprifolium* and *L. periclymenum* in Denmark, Italy, and France.
- Mycosphaerella implexicola** (R. Maire.) Jaap. Gray leaf spots with brown margins on *L. impleza* in Dalmatia and Spain.
- Ophiobolus minor** Bub. On branches of *L. xylosteum* in Bohemia.
- Phyllosticta alpigena** Sacc. Black leaf spots on *L. nigra* in Italy.
- Phyllosticta caprifolii** (Op.) Sacc. Subcircular gray leaf spots on *L. caprifolium*, *L. impleza*, and *L. pallasi* in Siberia, Russia, Dalmatia, France, and Italy.
- Phyllosticta nitidula** Dur. and Mont. Large brownish leaf spots on *L. impleza* in Algeria.
- Phyllosticta sydowii** Bres. On leaves of *L. latifolia* in Germany.
- Puccinia longirostris** Kom. Brown rust pustules on small sunken yellow spots on *L. altmanni* and *L. turkestanica* in Turkestan.
- Ramularia loniceræ** Vogl. and f. *hispanica* Gz. Frag. Circular to irregular chestnut-brown leaf spots on *L. alpina* and *L. hispanica* in Italy and Spain.
- Rhabdospora xylostet** Lamb. and Fautr. On branches of *L. xylosteum* in France.
- Rhytisma lonicericola** P. Henn. Carbonous black stromata on leaves of *L. japonica*, *L. maackii*, and *L. tschonoskii* in Japan and Russia.
- Rhytisma xylostet** Naumoff. On leaves of *L. xylosteum* in Russia.
- Septoria obscurata** Thuem. Circular ashen-colored leaf spots with dark purple margins on *L. periclymenum* and *L. tatarica* in Russia and Portugal.
- Septoria xylostet** Sacc. and Wint. Small whitish leaf spots with black margins on *L. nigra*, *L. tatarica*, and *L. xylosteum* in Russia and Switzerland.
- Trabutia loniceræ** Rac. Circular black stromata on leaves of *L. japonica* in Java.
- LOPEZIA.** Greenhouse plants, cultivated for their attractive flowers. Grown outdoors in the South.
- Puccinia fuchsiae** Syd. and Holw. See Fuchsia.
- LOPHANTHUS.** Erect plants with clusters of blue flowers.
- Aecidium lophanthi** P. Henn. Rust on leaves of *L. rugosus* in Japan.
- LOTUS.** Herbs or subshrubs, grown for their various-colored flowers.
- Cercospora loti** Höll. Circular red-brown leaf spots on *L. siliguosus* in Hungary.
- Mitralia sclerotiorum** Rostr. See Medicago.
- Ovularia sphaeroidea** Sacc. On leaves of *L. corniculatus*, *L. uliginosus*, *Vicia sativa*, and *V. villosa* in France, Denmark, Sweden, Italy, and Austria.
- Peronospora lotorum** Syd. Downy mildew on leaves of *L. corniculatus* and *L. uliginosus* in central and northern Europe.
- Ramularia loticola** Massal. On leaves of *L. corniculatus* in Italy.
- Ramularia schulzeri** Bäuml. See Anthyllis.
- Septoria stipularis** Pass. On leaves, stems, and stipules of *Lathyrus aphaca* and *Lotus corniculatus* in Denmark and Italy.
- Uromyces anthyllidis** (Grev.) Schroet. See Anthyllis.
- Uromyces libycus** Trott. Rust on leaves of *L. pusillus* in Tripoli.
- Uromyces loti** Blytt. Brown rust pustules on leaves of *Euphorbia cyparissia*, *Lotus angustissimus*, *L. corniculatus*, *L. palustris*, and *L. uliginosus* in Japan and Europe.
- LUCUMA.** SAPOTE. CANISTEL. MAMEY COLORADO. Tropical trees grown for fruit.
- Phyllosticta lucumae** Syd. Circular to subcircular whitish leaf spots with raised purple-brown margins on *L. nerifolia* in Uruguay.
- Uredo lucumae** Arth. and Johnst. Rust on leaves of *L. nervosa* in Cuba.
- Uromyces lucumae** Diet. Powdery brown leaf spots on *Lucuma* sp. in Brazil.
- LUFFA.** TOWEL GOURD. DISHCLOTH GOURD.
- Corticium koleroga** (Cke.) v. Hoehn. See Coffea.
- LUNARIA.** HONESTY. Herbaceous annuals and perennials.
- Helminthosporium lunariae** Pöhl. Circular leaf spots on *L. annua* (*L. biennis*) in Italy.
- Peronospora lunariae** Gäum. Downy mildew on leaves of *L. annua* and *L. rediviva* in central Europe.
- LUPINUS.** LUPINE. Herbaceous legumes.
- Ascochyta lupinicola** Petr. On leaves of *Lupinus* sp. in Bohemia and Austria.
- Bacillus elegans** Hegyi. Yellow, then brown, spots on leaves of *L. albus*, *L. angustifolius*, and *L. luteus* in Hungary.
- Ceratophorum setosum** Kirch. See Cytisus.
- Chalaropsis thielavioides** Peyr. Said to cause a root rot of *L. albus* in Italy.
- Chrysocelis lupini** Lagerh. and Diet. Rust on leaves of *L. clarkei*, *L. paniculatus*, and *Lupinus* sp. in Colombia, Costa Rica, and Peru.
- Gloeosporium lupinus** Bondar. Anthracnose on *Lupinus* sp. in Brazil.
- Rhytisma onobrychidis** Sacc. Black stromata on leaves and stems of *Lathyrus latifolius*, *Lupinus* sp., and *Hedysarum coronarium* in Italy, France, and Germany.
- Uromyces anthyllidis** Grev. See Anthyllis.
- Uromyces elatus** Syd. Yellow and brown rust pustules on leaves of *L. ramosissimus*, *L. sativus*, and *L. tomentosus* in Bolivia and Peru.
- Uromyces lupinicolus** Bub. Powdery brown rust pustules on leaves of *L. albus*, *L. angustifolius*, and *Lupinus* sp. in Spain and Bohemia.

LUPINUS—Continued.

Uromyces montanus Arth. Cinnamon-brown rust pustules on leaves of *L. mexicanus* and *L. montanus* in Mexico and Guatemala.

Uromyces renovatus Syd. Powdery brown rust pustules on leaves of *L. albus*, *L. angustifolius*, *L. hispanicus*, *L. luteus*, and *L. termis* in Europe, northern Africa, and Madeira.

Uromyces rugosus Arth. Powdery cinnamon-brown to dark-brown rust pustules on leaves of *Lupinus* sp. in Mexico.

LUZULA. JUNCOIDES Ag. WOOD-RUSH. Grass or rushlike perennial herbs.

Ascochyta teretiuscula Sacc. and Roum. On leaves of *L. campestris* and *L. vernalis* in France and Yugoslavia.

Phyllachora luzulae (Rabh.) Cke. Dull-black stromata on both leaf surfaces of *Luzula* sp. in Germany.

Puccinia luzulae-maximae Diet. Leaf rust on *L. maxima* in Europe.

Puccinia luzulina Syd. Cinnamon-brown rust pustules on leaves of *L. alopecurus* in South America.

Puccinia oblongata (Lk.) Wint. Dull-brown to black rust pustules on red-brown irregular leaf spots on *L. campestris*, *L. forsteri*, *L. maxima*, *L. multiflora*, *L. nivea*, *L. pilosa*, and *L. sylvatica* in Europe.

Puccinia tenuispora McAlp. Yellow-brown to dark-brown rust pustules on leaves and stems of *L. campestris* and *L. oldfieldii* in Australia and Tasmania.

Septoria chanosiana Ferr. Purple-brown leaf spots on *L. lutea* in Italy.

Septoria luzulae Schroet. On leaves of *L. forsteri* in Yugoslavia.

Uredo antarctica Berk. Rust on leaves of *L. crinita* in the Campbell Islands.

Urocystis luzulae (Schroet.) Wt. Black smut pustules on leaves of *L. multiflora* and *L. pilosa* in Denmark and Germany.

Ustilago hyperborea Blytt. Smut in fruit of *L. hyperborea* in Norway.

Ustilago liebmanni P. Henn. A smut transforming the ovaries of *Luzula* sp. in Mexico into brown spore masses.

Ustilago viuyckii Oud. and Beyer. A smut forming black spore masses in ovaries of *L. campestris* in Holland.

LYCASTE. Terrestrial and epiphytic orchids. See Orchidaceae.**LYCHNIS.** CAMPION. CORN COCKLE. MALTESE CROSS. Flower-garden herbs.

Ascochyta cookei Mass. Blotches on leaves of *L. alba* (*L. vespertina*) in Great Britain.

Ascochyta dianthi (A. and S.) Berk. See Dianthus.

Fabraea inolexa Bres. and Cap. On leaves of *L. flos-jovis* in Italy.

Gloeosporium lychnidis Oud. Anthracnose on leaves of *L. dioica* (*L. diurna*) in Holland.

Marsonia delastrei (Delacr.) Sacc. Dull-yellow indefinite areas on leaves and stems of *L. chalcadonica*, *L. dioica*, *Silene latifolia*, *S. noctiflora*, and *Agrostemma githago* in Europe and Siberia. Reported from Wisconsin.

Peronospora lychnidis Güm. Downy mildew on leaves of *L. coronaria* in Germany.

Phyllosticta punctiformis (Desm.) Allesch. On leaves of *L. dioica* in Scotland.

Puccinia lychnidis-miqueliana Diet. Rust on leaves of *L. miqueliana* in Japan.

Ramularia chalcadonica Allesch. Ashen leaf spots on *L. chalcadonica* in Germany.

Ramularia lychnicola (Cke.) On leaves of *L. chalcadonica*, *L. dioica*, and *Silene latifolia* in Great Britain, Austria, and Germany.

Septoria lychnidicola P. Brun. Circular to oblong ochraceous leaf spots with brown margins on *L. flos-cuculi* in France.

Septoria lychnidis Desm. Irregular red-brown leaf spots on *L. chalcadonica*, *L. dioica*, *L. flos-jovis* and *Gypsophila altissima* in Siberia, Italy, France, and Great Britain.

Septoria melandrii Pass. Subcircular reddish leaf spots with dull-brown margins on *L. diurna* and *L. alba* (*L. vespertina*) in Italy. Reported from Wisconsin.

Uromyces behenii (DC.) Ung. Rust on leaves of *L. indica* in India.

Uromyces crassivertex Diet. Cinnamon-brown to black rust pustules on leaves of *L. miqueliana* in Japan.

Uromyces lychnidicola Speg. Rust on leaves of *L. magellanica* and *L. patagonica* in Argentina.

LYCII M. MATRIMONY VINE. Ornamental shrubs and vines.

Ascochyta destructiva Kab. and Bub. Subcircular, often confluent, ochraceous leaf spots on *L. barbarum* in Bohemia.

Ascochyta lycii Rostr. Ashen leaf spots with dull-brown margins on *L. barbarum* and *L. halimifolium* in Denmark.

Microsphaera mougeotii Lévl. Powdery mildew on leaves of *L. barbarum*, *L. europaeum*, *L. ovatum*, and *L. ruthenicum* in Europe.

Phyllosticta destructiva Desm. See Althaea.

Puccinia afra Wint. Powdery brown rust pustules on leaves, peduncles, and sepals of *L. afrum* in Spain and the Union of South Africa.

Puccinia lycii Kalchbr. Brown rust pustules on leaves of *L. tubulosum* in the Union of South Africa. Reported from Ohio.

Puccinia lycicola Speg. Brown rust pustules on leaves of *L. argenteum* and *L. patagonicum* in Argentina.

Puccinia turgida Syd. Powdery brown to dark-brown rust pustules on leaves of *L. europaeum* in Palestine.

Septoria lycii Paul. and Gz. Frag. On leaves of *L. helimifolium* (*L. vulgare*) in Spain.

Septoria lycicola Speg. Circular definite leaf spots on *L. cestroides* in Argentina.

Uredo megalospora Speg. Rust on leaves of *Lycium* sp. in Argentina.

LYCOPERSICON. (*Lycopersicon*.) TOMATO. Herbaceous plants, cultivated for their fruit.

Bacillus caulivorus Prill. and Delacr. See Solanum.

Bacterium briosii Pavar. Bacteriosis of leaves, stalks, and fruits of *L. esculentum* in Italy and France. The leaves wither up, new buds are deformed, and long brown spots appear on the stems.

Bacterium lycopersici Burgwitz. This bacterium is reported as the cause of "blossom-end rot" of fruit of *L. esculentum* in Russia.

Chrysophyctis endobiotica Schlib. See Solanum.

Diplodina lycopersicola (v. Bond.) Mont. Large black spots on tops or sides of both green and ripe fruit of *L. esculentum* in Russia. The spots are somewhat depressed, warty, black at the centers, and brown at the margins, with surrounding dirty-green areas. Probably not distinct from *Phoma destructiva* Plowr.

Fusarium erubescens A. and v. Ov. Small black sunken spots on fruit of *L. esculentum* in Germany.

Hainesia lycopersici Speg. On leaves of *L. esculentum* in Argentina.

Phoma ferraristi Ciferri. Causes a fruit rot of *L. esculentum* in Italy. *Ramularia ferraristi* Cif. is described as a stage of the same fungus. Probably not distinct from *P. destructiva* Plowr.

LYCOPERSICUM—Continued.

Phytophthora cryptogea Pethyb. and Laffer. This fungus has been found as one of the causes of damping off of seedlings of *L. esculentum* in Great Britain. It also attacks *Cineraria* seedlings.

Diseased plants show brown areas at the soil level and collapse, breaking over at the point of infection.

Phytophthora melongenae K. Saw. See Solanum.

Phytophthora mexicana Hots. and Hartge. This downy mildew is said to cause a rot of fruit of *L. esculentum* in Mexico and to be capable of attacking other plant parts as well. Infected fruit have been imported into the United States.

Phytophthora parasitica Dastur. See Ricinus.

Puccinia pitteriana Arth. See Solanum.

Rhizoctonia sp. See Vigna.

Sclerotium setosum Bewley and Shearn. Affected plants stop growing; the lower leaves turn yellow and die prematurely, and the terminal growths are pale and weak. The stems become hollow and yellow, and a dark-brown discoloration spreads up the stems from the ground level. Finally all the leaves wither and the plants die. The roots decay and shrivel. Numerous minute black globular sclerotia occur in the rotted tissues of the roots and the bases of the stems. On *L. esculentum* in England. This Sclerotium has been connected with *Colletotrichum tabificum* (Hallier) Pethybr.

Vermicularia varians Duc. is also considered synonymous.

Spongospora subterranea (Walls.) Lang. See Solanum.

Vermicularia capsici Syd. See Capsicum.

Vermicularia varians Duc. See Solanum.

Verticillium pulverulentum Gouven. Circular, sharply delimited, dark spots on fruit of *L. esculentum*, causing premature fall in Holland.

LYGODIUM. CLIMBING FERN.

Aphelenchus olesistis Ritz. Boz. See Begonia.

Helminthosporium filicicola P. Henn. Effuse leaf spots on *Lygodium* sp. and *Selaginella* sp. in Peru and Brazil.

Milesia blechni Syd. See Blechnum.

Mycosphaerella ditissima Syd. On fronds of *L. flexuosum* in the Philippines.

Puccinia lygodii (Har.) Arth. (*Uredo lygodii* Har.) Yellow-brown rust pustules on fronds of *Lygodium* sp. in Brazil.

LYONIA. Ornamental shrubs grown for their white flowers and dense foliage.

Exobasidium fawcettii Mass. On leaves of *L. jamaicensis* in Jamaica. Probably not distinct from *E. vaccinii* Fckl.

LYSILOMA. Trees and shrubs of the American Tropics.

Ravenelia lysilomae Arth. Yellow-brown to chestnut-brown rust pustules on leaves of *L. tergeminia* in Mexico.

Ravenelia sololensis Arth. and Holw. Rust on leaves of *L. acapulcensis* and *L. bahamensis* in Mexico, Cuba, and Guatemala.

LYSIMACHIA. LOOSESTRIPE. Erect or creeping leafy herbs.

Aecidium lysimachiae-japonicae Diet. Leaves of *L. japonica* covered with rust pustules in Japan.

Phyllosticta lysimachiae Allesch. Irregular, then confluent, greenish-ashen, then pale ochraceous, leaf spots on *L. vulgaris* in Russia and Germany.

Puccinia dicteliana Syd. Yellow and dark-brown rust pustules on leaves of *L. clethroides* and *L. fortunei* in Japan.

Ramularia lysimachiae Thuem. On leaves of *L. nummularia*, *L. thyrsiflora*, and *L. vulgaris* in Siberia, Yugoslavia, Denmark, Austria, Bohemia, and Germany. Reported from Wisconsin.

Ramularia lysimachiarum Lindr. Brown to dark-green, then yellow, leaf spots on *L. nummularia* in Sweden, Finland, Denmark, and Great Britain.

Septoria bresadolana Krieger. Reddish leaf spots, becoming grayish white at the centers, on *L. vulgaris* in Germany.

Septoria lysimachiae West. Brown leaf spots on *L. nummularia*, *L. thyrsiflora*, and *L. vulgaris* in Siberia and Europe.

Septoria nambuana P. Henn. Red-brown leaf spots on *L. brachystachys* in Japan.

LYTHRUM. Spiked or purple loosestrife. Herbs or subshrubs.

Aecidium lythri Diet. and Neg. Rust on leaves of *L. hyssopifolium* in Chile.

Cercospora lythri (West.) Niessl. On leaves of *L. salicaria* in Spain, Belgium, Italy, and Germany.

Doassansia winteriana (Wint.) Magn. A smut producing small globose punctiform galls on both leaf surfaces of *L. hyssopifolium* in Australia.

Septoria brissacana Sacc. and Let. Small dull-yellow leaf spots with dark margins on *L. salicaria* in Denmark, France, Russia, and Germany.

MAACKIA. Trees grown for their handsome foliage and panicles of white flowers.

Uromyces amurensis Kom. Leaf rust on *M. amurensis* in Japan.

MABA. Tropical trees and shrubs with hard ebonylike wood.

Aecidium bicolor Sacc. Rust on leaves of *M. buxifolia* and *M. natalensis* in India and the Union of South Africa.

Aecidium melaleucum Syd. Leaf rust on *M. buxifolia* in India.

Aecidium rhytismoideum B. and Br. var. **mabae** P. Henn. See Diospyros.

Ascochyta mabiana Sacc. Dull-brown leaf spots with darker margins on *M. abyssinica* in Abyssinia.

Exobasidium symploci-japonicae Kusano. See Symplecos.

Phyllachora mabae (P. Henn.) Theiss. and Syd. Circular shiny-black stromata on leaves of *M. inconstans* in Brazil.

MACARANGA. Trees or shrubs.

Cercospora macarangae Syd. Circular ochraceous leaf spots on *M. tanaria* in the Philippines.

Marsonia pavonina Syd. Leaf spots on *M. bicolor* in the Philippines.

Phyllachora macarangae P. Henn. Black stromata on circular brown leaf spots on *Macaranga* sp. in the Philippines.

MACFADYENA. Tall climbing tropical shrubs.

Leptothyrium heterospermum Speg. Shield-shaped black fruiting bodies on yellow leaf spots on *M. cynanchoides* in Argentina.

MACLURA. TOXYLON Ag. OSAGE ORANGE. BOW WOOD. Cultivated as a hedge plant and for its

evergreen foliage and orange-like fruit.

Ectostroma maclurae Thuem. On leaves of *M. aurantiaca* (*Toxylon pomiferum*) in Italy and Portugal.

Septoria maclurae P. Brun. Subcircular to irregular olive or dull-yellow leaf spots with brown margins on *M. aurantiaca* (*T. pomiferum*) in France.

Uredo maclurae Speg. Rust on leaves of *M. aurantiaca* (*T. pomiferum*) and *M. ucora* in Argentina.

MACROZAMIA. Cycaslike plants.

Cryptosporella macrozamia P. Henn. On leaf blades and petioles of *M. fraseri* in Australia.

MAESA. Tropical and subtropical shrubs.

Amazonia peregrina Syd. On *M. indica* and *M. laza* in Ceylon and the Philippines.

Fusidium maesa P. Henn. Pale or ashen leaf spots on *M. lanceolata* in Tanganyika.

Physalospora chaenostoma Sacc. On leaves of *Maesa* sp. in Natal.

MAGNOLIA. Woody plants cultivated for their handsome foliage and showy flowers.

Ascochyta magnoliae Thum. On leaves of *M. grandiflora* in Italy.

Gloeosporium haynaldianum Sacc. and Roum. Anthracnose on leaves of *M. grandiflora* and *M. soulangeana* in Denmark, Italy, and France.

Hendersonia magnoliae Sacc. Irregular whitish leaf spots with brown margins on *M. grandiflora* in France and Italy.

Mycosphaerella elatior Sacc. and Speg. and var. *apula* Sacc. On leaves of *M. grandiflora* and *Liriodendron tulipifera* in Spain and Italy.

Phyllosticta kobus P. Henn. Irregular dull-brown leaf spots on *M. kobus* in Japan.

Phyllosticta yulan Tassi. Large pale-brown leaf spots on *M. denudata* (*M. yulan*) in Italy.

Septoria caerulea Tassi. Fuscous leaf spots on *M. denudata* in Italy.

Sphaeropsis magnoliae Magnag. On *Magnolia* sp. in Italy.

MAGYDARIS. Perennial herbs.

Puccinia magydaridis Pat. and Trab. Cinnamon-brown to black rust pustules on leaves of *M. panacina* and *M. tomentosa* in Algeria, Tunis, and Spain.

MAHONIA. See Berberis.**MAIANthemum.** UNIFOLIUM Ag. Small spring-blooming woody perennials.

Cercospora majanthemi Fekl. On leaves of *M. bifolium* in Denmark, Belgium, Sweden, Russia, Siberia, and Austria. Reported from Ohio.

Mycosphaerella asteroma (Fr.) Lind. See *Convallaria*.

Puccinia majanthemi Diet. Brown rust pustules on both leaf surfaces and on petioles of *M. bifolium* in Japan.

Ramularia rubicunda Bres. On leaves of *M. bifolium* in Russia, Germany, and Austria.

Septoria majanthemi West. Dull-gray indefinite leaf spots on *M. bifolium* in Belgium and Germany.

MALLOTUS. MONKEY-FACE TREE. Tropical trees or shrubs.

Cercospora laguensis (Sacc.) Yates. On leaves of *M. moluccanus* in the Philippines.

Cronartium malloeti Raeb. Golden-yellow to red-brown rust pustules on leaves of *M. moluccanus* in Java.

Helminthosporium insigne Sacc. On *M. philippinensis* in the Philippines.

Phyllosticta marmorta Cke. Confluent leaf spots on *M. philippinensis* in India and the Philippines.

Uromyces malloeti P. Henn. Brown rust sori on leaves of *M. moluccanus* in New Guinea and the Philippines.

MALUS. APPLE. CRAB.

Asteroma mali Desm. Circular brown leaf spots on *Malus* sp. in Germany.

Bacterium mali Brez. Reported as the cause of a canker disease of trunks and limbs of *Malus* (apple) in central Europe.

Bacterium sp. Blossom blight. See *Pyrus*.

Botryosphaeria mali Putt. Causes a cankering of branches and trunks of *Malus* (apple) in the Union of South Africa. Death of infected branches, and even of entire trees, may be brought about through girdling.

Cercospora porriga Speg. Causes a drying of young fruits of *Malus* sp. in Argentina.

Colletotrichum piri Noack. See *Pyrus*.

Coniothecium chromatoporum Cda. This fungus is reported as the cause of the blister and fruit-cracking disease of *Pyrus* and *Malus* in the Union of South Africa, Australia, Tasmania, New Zealand, and India. Dark specks appear in groups on the twigs and branches, finally becoming reddish-brown blisters. Diseased twigs die back. On the fruit the fungus produces a scurfy condition or russetting. Young fruit when attacked crack on expanding.

Dendrodochium pulchrum Marchal. On *Malus* (apple) in Belgium.

Diaporthe mali Miura. Pale, discolored areas, from 1 to 2 centimeters in diameter, on the leaves of *Malus* (apple) in Japan, producing curling and defoliation. Young shoots and twigs also show irregular brown areas which cause die-back. Spots on the fruit are circular to irregular, coalescing and involving the entire fruit in a brown rot. A serious disease throughout Japan. *Diaporthe mali* Bres., reported from Australia, is apparently distinct.

Diaporthe perniciosus Marchal. See *Prunus*.

Diplodia griffoni Sacc. and Trav. This fungus has been confused with *Physalospora cydoniae* but is considered distinct. It causes a serious canker disease of twigs of *Malus* (apple) and *Pyrus* (pear) in Europe and New Zealand. The cankers are slightly depressed, reddish brown, zonate, and with the surfaces studded with the pycnidia. Old cankers are dull brown.

Dothiorella vinosa Marchal. See *Prunus*.

Fusarium sp. A blossom blight of *Malus* sp. has been attributed to this fungus in Great Britain.

Gymnosporangium sabiniae (DC.) Wint. See *Juniperus*.

Gymnosporangium yamadai Miy. See *Juniperus*.

Helicobasidium tanakae Miy. See *Morus*.

Hydnum schiedermayeri Heufler. A wound parasite of *Malus* sp. in Europe, causing a rot of bark and wood of trunk and limbs. Reported from Pennsylvania.

Labrella piricola Bres. and Sacc. See *Pyrus*.

Laestadia mali (Fekl.) Sacc. On leaves of *Malus* sp. in Belgium and Germany.

Leptosphaeria vagabunda Sacc. Causes a fruit spot and rot of *Malus* sp. in Great Britain.

Marsonia mali P. Henn. Subcircular, then confluent, dull-brown leaf spots with purple margins on *Malus* in Japan.

Melanobasidium mali Maubl. Oval or elongate whitish leaf spots with brown margins on *Malus* in Spain.

Melasma mali Oud. On leaves of *Malus* in Holland.

Mycosphaerella pomacearum Sacc. On leaves of *Malus* sp. and *Cydonia* sp. in China and France.

Mycosphaerella pomi (Pass.) Lind. On leaves of *Malus* in Italy.

Myxosporium mali Bres. Brown spotting and rot of fruit of *Malus* in Great Britain and on dead twigs in Germany.

Ochropsora pallida Rostr. See *Sorbus*.

Phacidiella discolor (Mont. and Sacc.) A. Poteb. Causes a fruit rot and trunk canker of *Malus* (apple) and *Pyrus* (pear) in Europe.

Phyllosticta mali Prill. and Delacr. Small elongate or circular ochraceous, then dirty-gray, leaf spots with fuscous margins on *Malus* sp. in Denmark, Russia, Bulgaria, and France. Reported from West Virginia.

Phyllosticta briardi Sacc. Brown or ochraceous leaf spots with darker borders on *Malus* and *Pyrus* sp. in Russia, Great Britain, France, Malta, and Hungary, often causing defoliation.

MALUS—Continued.

- Phytophthora syringae** Kleb. Produces a dark-brown rather firm and elastic rot of fruit of *Malus* in Europe. See, also, *Syringia*.
- Plectodiscella piri** Wor. See *Pyrus*.
- Pleospora pomorum** Horne. Causes a brown spotting and rot of fruit of *Malus* in Great Britain.
- Polypetus purpureus** Horne. Associated with fruit spotting and rotting of *Malus* in Great Britain. *P. aureus* Horne, *P. pomi* Horne, and *P. recurvatus* Horne have been found in similar situations.
- Polystigma ochraceum** (Wahlb.) DC. See *Prunus*.
- Sclerotinia mali** Tak. Produces brown areas on leaf blades and petioles, flower stalks, and branches and a brown rot of young fruit of *Malus* in Japan.
- Sclerotium stellatum** Horne. Fruit spot and rot of *Malus* in Great Britain.
- Spilocaea pomi** Fr. Round sunken brown spots on ripe fruit of *Malus*, especially during storage, in Russia, Great Britain, and Germany. Called the "apple-speck" disease.
- Sporodesmium cerebriforme** McAlp. "Eye-scale" disease of *Malus* in Australia.
- Stagonospora mali** Delacr. See *Pyrus*.
- Stereum purpureum** Pers. See *Prunus*.
- Trichoseptoria fructigena** Maubl. See *Cydonia*.
- Trichothecium candidum** Walbr. "White-apple blotch" is marked by small round spots with brown-black borders, causing a brownish rot and mummification of fruit of *Malus* in Europe. Probably not distinct from *T. roseum*.
- Valsa mali** Miy. and Yan. Oblong to irregular slightly elevated brown spots on branches of *Malus* in Japan. These areas dry out, become sunken, more or less darkened, and the surface cracks.
- A trunk canker** of apple (*Malus*), with which bacteria and a *Cytospora*-like fungus are associated, has been reported from Great Britain. Large, rapidly extending, sunken cankers form on the lower part of the trunk, soon girdling and killing infected trees.

MALVA. MALLOW. Annual or perennial herbs.

- Ascochyta malvae** H. Zimm. Subcircular ochraceous, then white, leaf spots on *M. neglecta* and *M. vulgaris* in Austria and Germany.
- Ascochyta malvicola** Sacc. Subcircular or sinuous whitish leaf spots on *M. rotundifolia* and *M. sylvestris* in Italy and Austria.
- Ascochyta montenegrina** Bub. Subcircular, often confluent, yellow leaf spots on *M. sylvestris* in Yugoslavia.
- Cercospora malvarum** Sacc. Large olive-colored leaf spots on *M. moschata* in Spain and Denmark. Reported from Kansas.
- Cercospora polymorpha** Bub. Subcircular, oblong, or angular yellowish leaf spots with narrow dark-brown borders on *M. sylvestris* and *M. verticillata* in China and Yugoslavia.
- Phyllosticta destructiva** Desm. See *Althaea*.
- Puccinia heterogena** Lagerh. Chestnut-colored rust pustules on leaves and stems of *Althaea rosea*, *M. crispa*, *M. nicaeensis*, and *M. peruviana* in Ecuador.
- Ramularia malvae** Fckl. Circular to elongate white leaf spots on *M. moschata*, *M. rotundifolia*, and *M. sylvestris* in Denmark and Germany.
- Septoria heterochroa** Desm. Small subcircular gray-brown, then whitish leaf spots on *M. neglecta*, *Althaea officinalis*, and *Sida* sp. in Costa Rica, Russia, Denmark, Italy, Belgium, and Great Britain.
- MALVASTRUM**. FALSE MALLOW. Mallowlike herbs.
- Aecidium malvastriicola** P. Henn. Rust on leaves of *M. spicatum* in Argentina.
- MANDEVILLA**. Tall tropical climbers.
- Phyllosticta gelsenii** E. and E. var. *mandevillae* Sacc. and Seal. Subcircular gray leaf spots on *M. suaveolens* in Portugal.
- Uredo mandevillae** Mayor. Rust on leaves of *M.* (*Echites*) *mollissima* and *M. tomentosa* in Colombia and Trinidad.

MANETTIA. Twinning plants grown for their flowers.

- Puccinia niedertleitii** P. Henn. Cinnamon-brown rust pustules on leaves of *M. bicolor*, *M. ignita*, and *M. leianthiflora* in Argentina and Paraguay.
- Uromyces tener** Schroet. Powdery brown rust pustules on leaves of *M. gracilis* in Brazil.
- MANGIFERA**. MANGO. Tropical fruit and shade trees.

- Bacillus mangiferae** Doidge. This bacterial blight first appears on mango (*M. indica*) fruits as small water-soaked areas which rapidly enlarge, turn brown, and crack, corky tissue developing. Gumming generally occurs. Similar irregular brown spots occur on leaf blades and petioles, young twigs, peduncles, and pedicels. Diseased fruits fall. The disease is serious in Portuguese east Africa and the Union of South Africa.
- Cercospora mangiferae** Koord. Small circular numerous brown leaf spots on *M. indica* in Java, Philippines, and Haiti.
- Coccomyces vilis** Syd. and Butl. Irregular spots with purple margins on leaves of *M. indica* in India.
- Corticium salmonicolor** B. and Br. See *Citrus*.
- Cornartium kemangae** Racib. Brown rust pustules on irregular leaf spots on *M. kemanga* in Java.
- Diplodia cacaicola** P. Henn. See *Theobroma*.
- Endothia havenensis** Brun. See *Eucalyptus*.
- Eutype erumpens** Mass. See *Theobroma*.
- Gloeosporium mangiferae** P. Henn. (*Gloeosporium mangae* Noack.). These anthracnose fungi, reported as causing bloom blight, fruit rot, and leaf spotting of the mango (*M. indica*) in Trinidad, Brazil, and elsewhere, are certainly synonymous with *Colletotrichum gloeosporioides*, widespread in the United States.
- Leptothyrium circumscissum** Syd. Large irregular or angular leaf spots on *M. indica* in the Philippines. Especially destructive to seedlings.
- Macrophoma** sp. Attacks the fruit of *M. indica* in Brazil, causing them to blacken and fall to the ground, rotting or mummifying.
- Meliola mangiferae** Earle. Black circular to irregular superficial fungus patches on leaves of *M. indica* in the West Indies, the Philippines, and Malaya.
- Oidium** sp. A powdery mildew attacking the leaves of *M. indica* has been reported from India and Cuba.
- Phyllachora** sp. Tar spot on leaves of *M. indica* in the Philippines.
- Phyllosticta mortoni** Fairm. Numerous small deep-brown angular leaf spots on *M. indica* in the Bahamas, Mexico, and Florida.
- Rosellinia bunodes** B. and Br. See *Citrus*.
- Sphaerostilbe repens** B. and Br. See *Hevea*.
- Stilbella flavida** (Cke.) Lind. See *Coffea*.
- Zimmermaniella trispora** P. Henn. Black stromata on leaves of *M. indica* in Java.
- Eucalyptus disease**. See *Eucalyptus*.

MANIHOT. CASSAVA. Sometimes called mandioca. Ceara rubber tree. Tropical trees and shrubs with many economic uses.

Asochyta manihotis P. Henn. Circular pale-yellow-brown leaf spots on *M. utilisima* in Tanganyika.

Basillus manihotus Arthaud. and Ber. Young shoots rot internally, the vascular bundles decompose, wither, and die. On older shoots elongate gum pockets form and a yellow gum exudes. On *M. palmata* and *M. utilisima* in Brazil.

Cercospora cearae Petch. Irregular, then confluent, red-brown, then gray, spots on *M. glaziovii* and *M. piakhyensis* in India, Ceylon, Malaya, and Uganda, causing the leaves to crumple up and fall.

Cercospora henningsii Allesch. Subcircular to irregular pale-ochraceous leaf spots on *M. palmata* and *M. utilisima* in Porto Rico, Dominican Republic, Cuba, Ceylon, Siam, Indo-China, the Philippines, and tropical Africa.

Cercospora manihotis P. Henn. Circular or effuse dark-fuscescent leaf spots on *M. utilisima* in Brazil, China, the Philippines, and central Africa.

Cercospora pseudoidium Speg. Subcircular pale-brown leaf spots on *M. utilisima* in Paraguay and Brazil.

Colletotrichum lussoniense Sacc. Anthracnose on *M. utilisima* in the Philippines.

Colletotrichum manihotis P. Henn. Circular whitish leaf spots with reddish-brown surrounding zones on *M. utilisima* in Brazil.

Diplodia cacaoicola P. Henn. See Theobroma.

Fomes lamaensis Murr. See Hevea.

Fomes lignosus Klotzsch. See Hevea.

Haplographium manihotica Vinc. Circular brown, zoned leaf spots on *M. glaziovii* in Brazil.

Macrophoma manihotis P. Henn. On leaves of *M. utilisima* in Tanganyika.

Mycosphaerella manihotis Syd. Circular brown leaf spots on *M. utilisima* in Brazil.

Phyllosticta manihotica Syd. On leaves of *M. dichotoma*, *M. glaziovii*, and *M. heterophylla* in the Philippines.

Rosellinia arcuata Petch. See Thea.

Septogloeum manihotis Zimm. Brown leaf spots on *M. utilisima* in Java, India, Ceylon, and east Africa. Probably the same as *Cercospora henningsii* Allesch.

Stagonospora cassavae Wolk. A wound parasite attacking rooted cuttings of *M. utilisima* in Java.

Uromyces carthagenensis Speg. Yellow and dark-brown rust pustules on leaves and stems of *M. carthagenensis* and *M. tuediana* in Argentina.

Uromyces janiphae (Wint.) Arth. Rust on leaves of *M. utilisima* in Porto Rico, Cuba, Dominican Republic, Trinidad, and South America.

Uromyces manihotica P. Henn. Powdery dark-brown rust pustules on leaves of *Manihot* sp. in Brazil.

Uromyces manihotis P. Henn. Brown to black rust pustules on leaf blades and petioles, stems, and flowers of *M. glaziovii* and *M. utilisima* in Brazil. 'Produces a witches'-broom effect, as well as galls, on twigs and branches.

Uromyces manihotis-catingae P. Henn. Powdery brown rust pustules on leaves of *M. catingae* and *M. glaziovii* in Brazil.

MAPANIA. Tropical plants of the sedge family.

Puccinia mapaniae Racib. Powdery dark-brown rust pustules on *Mapania* sp. in Java.

MAPROUNEA. Tropical trees and shrubs.

Aecidium maprouneae P. Henn. Rust on irregular, fuscous leaf spots on *Maprounea* sp. in Peru and Brazil.

Phyllachora maprouneae P. Henn. Black stromata on yellowish to violet indefinite leaf spots on *M. surinamensis* in Peru.

Uredo maprouneae P. Henn. Rust on leaves of *M. guianensis* in Peru.

MARANTA. ARROWROOT. Foliage plants.

Puccinia cannae (Wint.) P. Henn. See Canna.

Rosellinia bunodes B. and Br. See Citrus.

Sclerotium sp. See Saccharum.

Sphaerostilbe repens B. and Br. Causes a soft rot of the underground stems of *M. arundinacea*. See also Hevea.

MARATTIA. Tropical ferns.

Leptosphaeria cafra Thuem. Large, more or less circular, leaf spots on *M. salicifolia* in the Union of South Africa.

MARIANA. See Silybum.

MARKHAMIA. Tropical evergreen trees and shrubs.

Phragmidium markhamiae P. Henn. Brown rust pustules on leaves of *M. zansibarensis* in Tanganyika.

MARRUBIUM. HOARHOUND. Aromatic herbs.

Phyllosticta marrubii McAlp. Pale-brown irregular leaf spots on *M. vulgare* in Australia.

Ramularia marrubii Mass. Small Rufous angular leaf spots on *M. album*, *M. creticum*, and *M. vulgare* in Italy, Estonia, and Yugoslavia.

Uredo marrubii Rabb. Rust on leaves of *Marrubium* sp. in Italy.

MARSDENIA. Twinning shrubs.

Puccinia aquatoriensis Syd. Gray-brown rust pustules on leaves of *Marsdenia* sp. in Ecuador.

Puccinia marsdeniae Diet. and Holw. Yellow-brown to black rust pustules on leaves of *M. mexicana* in Mexico and Guatemala.

Septoria asclepiadeae Sacc. See Cynanchum.

MAETINEZIA. See Palmae.

MARTYNIA. UNICORN PLANT. Annual and perennial plants sometimes cultivated for the peculiar seed pods.

Cercospora decolor Pass. Circular gray leaf spots on *M. lutea* in Argentina and Italy.

Phyllosticta martyniae Thuem. More or less circular leaf spots on *M. lutea* in Portugal.

MASDEVALLIA. See Orchidaceae.

MATRICARIA. FALSE CAMOMILE. Annual or perennial weedy herbs.

Cylindrosporium matricariae Diet. On leaves of *M. chamomilla* in Germany.

Entyloma matricariae Rost. Smut sori on leaves of *M. discoidea*, *M. inodora*, and *Tripleurospermum disciformis* in Denmark and Sweden.

Entyloma tralii Mass. Smut sori in small subcircular to ovate spots on stems and leaves of *M. inodora* in Great Britain.

Peronospora leptosperma (De By.) Gäum. See Chrysanthemum.

Septoria matricariae Höll. On leaves of *M. chamomilla* and *M. discoidea* in Hungary and Germany.

MATTHIOLA. STOCK. Sometimes called gilliflower. Flower-garden plants.

Ascochyta matthiolae Oud. On pods of *M. incana* in Holland.

Bacterium matthiolae Br. and Pav. Pale-green irregular spots first appear, becoming brown and causing deforming and curling of the leaves. Inflorescences are checked in their development. On *M. annua* in Italy.

Heminthosporium roridum Tode. On leaves of *M. incana* in Europe.

Peronospora matthiolae Gäum. Downy mildew on leaves of *M. incana* in central and north Europe.

Puccinia oudemansii Tranzsch. Powdery brown rust pustules on leaf blades and petioles of *M. nudicaulis* and *Parrya pinnatifida* in Turkestan.

Sclerotinia matthiolae Lendner. Causes a stem rot and blight of *M. vallesiaca* in Switzerland. Long black sclerotia form in the rotted stems.

Septoria henriquesii Thum. and f. *santonensis* P. Brun. On leaves of *M. incana* in France and Portugal.

Urocystis coralloides Rostr. Irregular gray coral-shaped smut masses on roots of *M. sinuata* and *Turritis glabra* (*Arabis perfoliata*) in France and Denmark.

MAURITIA. Fan palms. See *Palmae*.

MAXILLARIA. Epiphytic orchids. See *Orchidaceae*.

MAYTENUS. Trees and shrubs.

Aecidium mayteni Pazsch. Rust on leaves of *M. brasiliensis* and *Maytenus* sp. in Brazil.

Heterobotrys paradoxa Sacc. and subsp. *chilensis* Sacc. and Syd. On leaves of *M. magellanica* in Chile.

Phyllosticta mayteni Speg. Leaf spots on *M. magellanica* in Chile.

Septoria mayteni Wint. On leaves of *M. magellanica* in Tierra del Fuego.

Syncarpella missionum (Speg.) Theiss. and Syd. Black carbonous stromata on living twigs of *M. ligustrina* in Argentina.

MAZUS. Low annual or perennial rock-garden plants.

Aecidium foetidum Diet. Rust on leaves of *M. rugosus* in Japan.

MEDICAGO. MEDICK. Alfalfa. Herbaceous perennials cultivated as forage plants.

Ascochyta affinis Jaap. Circular to elliptical white leaf spots with brown borders on *M. arabica* in Dalmatia.

Hypochnus fuciformis (Berk.) McAlp. See *Lolium*.

Laestadia insidiosa Mass. On leaves of *M. sativa* in Tasmania.

Leptosphaeria circinans (Fckl.) Sacc. See *Asparagus*.

Marsonia medicaginis Voss. Circular to elliptical dull-yellow leaf spots on *M. lupulina* in Europe.

Mitrella sclerotium Rostr. Forms black sclerotia in the roots, infected plants dying. On *M. sativa*, *Lotus corniculatus*, *L. uliginosus*, and *Lotus* sp. in Europe.

Mycosphaerella destructiva B. and Br. On leaves of *M. sativa* in Australia.

Ovularia medicaginis Br. and Cav. On leaves of *M. sativa* in Italy.

Peronospora aestivalis Syd. Downy mildew on leaves of *M. denticulata*, *M. falcata*, *M. lupulina*, *M. media*, *M. minima*, and *M. sativa* in Europe and Australia. Occurs sparingly in the United States.

Phyllosticta bonanseaana Sacc. Indefinite yellow areas on leaves of *M. arborea* in Mexico.

Rhizoctonia destruens Tassi. See *Solanum*.

Rhizoctonia napl West. See *Brassica*.

Septoria medicaginis Rob. and Desm. Subcircular to irregular whitish leaf spots on *M. lupulina* and *M. sativa* in Argentina, Russia, Sweden, Denmark, Italy, and France.

Tylenchus dipsaci Kuehn. This nematode is a serious pest of alfalfa (*M. sativa*), greatly reducing yields by stunting the plants. It has been reported from several localities in the western United States. See *Narcissus*.

Typhula trifolii Rostr. See *Trifolium*.

Uredo medicaginicola Speg. Rust on stems and petioles of *M. sativa* in Argentina.

Uredo medicaginis Speg. Rust on leaves of *M. denticulata* in Argentina.

Uromyces anthyllidis (Grev.) Schroet. See *Anthyllis*.

Uromyctis alfalfae (Lagerh.) Magn. The crown-wart of alfalfa (*M. sativa*) is a serious disease in Europe, India, and South America. The disease has been found in limited areas in the United States. It is characterized by warts or galls at the base of the stems. These bodies are irregular in shape, varying in size from a fraction of an inch to 3 or 4 inches, and of a corallike appearance. Affected plants show dead and dying stems with yellow leaves.

MEDINILLA. Evergreen shrubs with showy flowers.

Colletotrichum medinillae Rangel. Anthracnose on leaves of *M. magnifica* in Brazil.

Laestadia medinillae Rangel. Large concentrically zoned leaf spots on *M. magnifica* in Brazil.

Phyllosticta lageniformis Rangel. On leaves of *M. magnifica* in Brazil.

Phyllosticta medinillae Rangel. On leaves of *M. magnifica* in Brazil.

MEIBOMIA. TICK CLOVER. Mostly weedy annual and perennial herbs, partly woody. Some species cultivated for forage.

Aecidium amagensis Mayor. Rust on leaves of *M. tortuosa* in Colombia.

Cercospora atropunctata Rac. Circular to irregular, often confluent, brown spots on leaves of *M. umbellata* in Java.

Phyllachora desmodii P. Henn. Black stromata on leaves of *M. scalpe* and *Meibomia* sp. in India and central Africa.

Physopella meibomiae Arth. Rust on leaves of *M. supina* in Trinidad.

Uredo amagensis Mayor. Rust on *M. tortuosa* in Colombia.

Uredo desmodii-heterocarpi Petch. Rust on leaves of *M. heterocarpa* in Ceylon.

Uredo desmodii-parvifolii Petch. Rust on leaves of *M. parvifolia* in Ceylon.

Uredo desmodii-pulchellii Syd. Powdery yellow-brown rust pustules on leaves of *M. pulchella* in Hongkong.

Uredo desmodii-triquetri Petch. Rust on leaves of *M. triquetra* in Ceylon.

Uromyces capitatus Syd. Rust on leaves of *M. yunnanensis* in China.

Uromyces castaneus Syd. Brown rust pustules on leaves of *M. incana* in Brazil.

Uromyces huallagensis P. Henn. Brown rust pustules on leaves of *Meibomia* sp. in Peru.

Uromyces mexicanus Diet. and Holw. Yellow-brown to black rust pustules on leaf blades and petioles and stems of *M. uncinata* and *Meibomia* sp. in Mexico and Costa Rica.

Uromyces orbicularis Diet. Chestnut-brown rust pustules on leaves of *Meibomia* sp. in Brazil.

Uromyces tenuistipes Diet. and Holw. Brown rust pustules on leaves of *M. strobilacea* and *Meibomia* sp. in Mexico.

MELALEUCA. Ornamental shrubs and trees.

Laestadia melaleuca (Berk.) Sacc. Circular dull-brown leaf spots on *Melaleuca* sp. in Australia.

Phyllachora melaleucæ Syd. Shiny black circular to irregular stromata on leaves of *M. spinosa* in Australia.

Phyalospora melaleucæ (Lév.) Sacc. On leaves of *M. leucadendron* in France.

Phyllosticta leucadendri P. Henn. Circular brown leaf spots on *M. leucadendron* in Australia.

MELASPHAERULA. DIASIA Ag. TWIN SPUR. Small bulbous plants.

Uromyces melasphaerulæ Syd. Brown to black rust pustules on orbicular to irregular spots on both surfaces of leaves of *M. graminea* in the Union of South Africa.

MELIA. CHINABERRY. Ornamental trees, also used for windbreaks.

Cercospora congoensis Syd. Small olivaceous leaf spots on *Melia* sp. in the Congo.

Cercospora mellicola Speg. Small subcircular whitish leaf spots on *M. azedarach* in Argentina.

Cercospora subseissilis Syd. Leaf spot on *M. azedarach* in Ceylon, India, China, and the Philippines.

Gloeosporium mellicola Speg. On leaves of *M. azedarach* in Argentina.

Peronosplasmopora portoricensis Lamk. Irregular, diffuse yellow areas on leaves of *M. azedarach* in Porto Rico, causing premature leaf fall.

Phyllachora meliæ Pat. Small circular shiny black stromata on leaves of *M. azedarach* in Indo-China.

Septobasidium acaciæ Saw. See *Acacia*.

Septoria meliæ Syd. Small circular white leaf spots on *M. azedarach* in the Union of South Africa.

MELICA. MELIC GRASS. Perennial grasses.

Aseochyta phyllachoroides Sacc. and Malbr. On leaves of *M. altissima* in France.

Cintractia melicæ (Sorok.) Det. Smut in ovaries of *M. ciliata* in Central Asia.

Phyllachora melicicola Speg. Tar spot on *M. violacea* in Argentina.

Phyalospora montana Sacc. On leaves and stems of *Melica* sp. and *Sesleria caerulea* in Italy, Spain, and Switzerland.

Puccinia helmerllana Bub. Yellow-brown to black rust pustules on leaves of *M. ciliata* and *M. cupani* in Russia, Italy, and Switzerland.

Puccinia rangiferina S. Ito. Black rust on leaves and sheaths of *Melica* sp. and *Calamagrostis arundinacea* in China and Japan.

Puccinia trebouxii Syd. Rust on leaves of *M. ciliata* in Russia.

Septoria melicæ Pass. Rufous leaf spots on *M. picta* and *M. uniflora* in Europe.

Septoria simplex (Schroet.) Sacc. On *M. nutans* in Lapland and Sweden.

Uredo Jozankensis Ito. Golden rust sori on dark-brown leaf spots on *M. nutans* in Japan.

Urocystis bornmulleri Magn. Smut on leaves and inflorescences of *M. cupani* in Syria.

Uromyces graminis (Niessl.) Diet. See *Seseli*.

Uromyces seseli-graminis Ed. Fisch. See *Arrhenatherum*.

Ustilago trebouxii Syd. Dark-brown linear smut sori on leaves of *M. ciliata* and *Triticum* (*Agropyron*) *crispatum* in Russia.

MELILOTUS. SWEET CLOVER. Forage plants.

Cercospora melloti Oud. Circular to oblong white leaf spots on *M. officinalis* in Russia, Italy, and Holland.

Entyloma melloti McAlp. Small circular slightly raised pustules producing large, often confluent, patches on leaves of *M. indica* in Australia.

Peronospora melloti Syd. Downy mildew on leaves of *M. alba*, *M. altissima*, *M. indica*, *M. officinalis*, and *M. parviflora* in India and Europe.

Septoria melloti (Lasch.) Sacc. Small subcircular pale leaf spots on *M. alba*, *M. officinalis*, and *M. vulgaris* in Italy, Russia, and Germany.

Stagonospora carpathica Baeuml. See *Vicia*.

Uromyces haeumerianus Bub. Brown rust pustules on leaves and stems of *M. neapolitana* and *M. officinalis* in Spain and Russia.

MELIOSMA. Trees and shrubs grown for their handsome foliage and large panicles of white flowers.

Aecidium meliosmae-myrianthæ P. Henn. and Shir. Rust on leaves of *M. myriantha* and *M. simplicifolia* in Japan and India.

Aecidium meliosma-pungentis P. Henn. and Shir. Rust deforming leaf blades and petioles of *M. pungens* in Japan.

Gophana mirabilis Rac. Rust on leaves of *Meliosma* sp. in Java.

Phakospora meliosmae Kusano. Yellow to brown rust pustules on leaves of *M. myriantha* in Japan and the Philippines.

Phyllachora meliosmae Racib. Circular black stromata on leaves of *Meliosma* sp. in Java.

Uredo meliosmae Petch. Leaf rust on *M. wightii* in Ceylon.

MELISSA. BALM. Hardy perennial herbs.

Cercospora sennensis Gz. Frag. Leaf spot on *M. officinalis* in Spain.

Cylindrosporium melissæ C. Mass. Irregular leaf spots on *M. officinalis* in Spain and Italy.

Phleospora melissæ (Desm.) Parisi. Numerous small angular to irregular brown spots causing leaves to curl up and fall. On *M. officinalis* in Italy, France, and Spain.

Phyllosticta latemarensis Kab. and Bub. On leaves of *M. officinalis* in Hungary.

Phyllosticta melissæ Bub. Small circular brown leaf spots on *M. officinalis* in Hungary.

MELITTIS. Perennial herbs, used for border plantings.

Phyllosticta montellæ Sacc. Circular or angular dark-red leaf spots, becoming whitish, on *M. melissophyllum* in Italy.

Septoria melittidis Sacc. and Speg. Whitish indefinite leaf spots on *M. grandiflora* and *M. melissophyllum* in France and Italy.

MELOTHRIA. Slender herbaceous annual or perennial vines.

Puccinia melothricola Syd. Leaf rust on *M. mucronata* in the Philippines.

Uredo cantoniensis Yates. Rust on leaves of *M. indica* in China.

Uredo melothriæ P. Henn. Rust on leaves of *M. tomentosa* in Abyssinia.

Uromyces hellerianus Arth. Brown rust sori on *M. guadalupensis* and *M. scabra* in Porto Rico, Cuba, and Guatemala.

MENISPERMUM. MOONSEED. Hardy, attractive semiwoody climbers.

Ascochyta jenissensis Sacc. Subcircular pale-brown leaf spots on *M. dahuricum* in Siberia.

Gloeosporium menispermii Ailesch. Anthracnose on leaves of *M. canadense* in Germany and Monaco.

Guignardia menispermii (Speg.) Theiss. and Syd. On leaves of *Menispermum* sp. in Brazil and Paraguay.

Phyllosticta maculiformis var. *menispermii* Sacc. On leaves of *M. canadense* in France.

Septoria menispermii Thuem. Ochraceous leaf spots with gray-brown margins on *M. dahuricum* in Siberia and Russia.

MENTHA. MINT. Strong-scented perennial herbs.

Peronospora stigmatica Raunk. Downy mildew on flowers of *M. aquatica* and *M. arvensis* in Denmark, Sweden, and Russia.

Phyllosticta menthae Bres. On leaves of *M. arvensis* in Germany.

Physoderma menthae Schroet. On stems and occasionally leaves of *M. aquatica* in Germany.

MENTZELIA. BLAZING STAR. Showy flowered plants, sometimes cultivated.

Coleosporium mentzeliae (Diet. and Holw.) Arth. Powdery golden rust sori on leaves of *M. hispida* in Mexico.

MENZIESIA. Low deciduous shrubs allied to Rhododendron.

Chrysomyxa menziesiae Diet. Numerous brown rust pustules on leaves of *M. pentandra* in Japan.

MERIANA. See Watsonia.

MESEMBRYANTHEMUM. FIG MARIGOLD. Low-growing succulents.

Peronospora mesembryanthemi Verwoerd. Downy mildew on leaves, stems, flower stalks, and calyces of *Mesembryanthemum* spp. in the Union of South Africa.

Puccinia mesembryanthemi McOwan. Golden and black rust pustules on leaves of *M. micranthum* in the Union of South Africa.

Septoria confuens McAlp. Gray-white leaf spots on *M. aequilaterale* in Australia.

MESPILOS. MEDLAR. Woody plants cultivated for fruit.

Ascochyta mespili Pass. Irregular brown, then gray or gray-brown, leaf spots on *M. germanica* in France.

Entomosporium mespili (DC.) Sacc. See Cotoneaster.

Gymnosporangium mespili (DC.) Kern. See Juniperus.

Laetadia mespili Fautr. On leaves of *M. germanica* in France.

Ovularia necans (Pass.) Sacc. See Cydonia.

Phyllosticta mespilicola Rota-Rossi. Circular to irregular large dull-brown leaf spots on *M. germanica* in Italy.

Phyllostictiella mespili (Sacc.) Tass. Subcircular, ochraceous leaf spots with rufous margins on *M. germanica* in Denmark, Malta, and Italy.

Sclerotinia mespili Schell. This fungus causes dark-brown leaf blotches from 1 centimeter in diameter up to, as well as brown rot and mummification of the fruit of *M. germanica* and *Cydonia* sp. in Switzerland and Great Britain.

Septoria mespili Sacc. Ochraceous leaf spots with rufous margins on *M. germanica* in France, Malta, and Italy.

METROSIDEOS. IRON TREE. Trees and shrubs planted for their showy flowers.

Phyllosticta metrosideri P. Brun. Circular pale gray-brown leaf spots with red-brown margins on *M. villosa* in Tahiti.

Phyllosticta salicifolia P. Brun. Round to irregular gray leaf spots with brown borders on *M. salicifolia* in France.

A serious disease, due to an unknown cause, destroyed large areas of *M. polymorpha* in Hawaii.

MEUM. Perennial herbs with delicate ornamental foliage.

Aecidium mei Schroet. Rust on leaf blades and petioles of *M. athamanticum* and *M. mutellinum* in central Europe.

Protomyces macrosporus Ung. See Coriandrum.

MEZONEURUM. Large, usually armed, climbing shrubs.

Ravenelia aculeifera Berk. Rust on leaves of *M. enneaphyllum* in Ceylon.

MICHELIA. Trees and shrubs with magnoliolike foliage and fragrant flowers.

Phyllachora micheleae Syd. Black circular stromata on leaves of *M. nilagirica* in India.

Phyllostictiella micheleae (P. Henn.) Tassi. On leaves of *M. fuscata* in Germany.

MICONIA. Tropical trees and shrubs with large, showy, strongly-veined leaves.

Catacauma pululahuensis (Pat.) Theiss. and Syd. Tar spot on leaves of *Miconia* sp. in Brazil.

Catacaumella miconiae (P. Henn.) Theiss. and Syd. Shiny black stromata on leaves of *Miconia* sp. in Brazil.

Cronartium egenulum Syd. Yellowish-red rust sori on leaves of *M. theezans* in Japan and Brazil.

Cyclothea miconiae (Syd.) Theiss. Fruiting bodies form on red-brown, then black, spots on lower leaf surfaces of *M. rigidiuscula* in Brazil.

Dothidina fiebrigii (P. Henn.) Theiss. and Syd. Black fruiting bodies on small brown spots on *Miconia* sp. in Paraguay.

Dothidina miconiae (P. Henn.) Theiss. and Syd. Black stromata on leaves of *M. laevigata*, *M. prasina*, and *M. sintenisii* in Porto Rico and Brazil.

Dothidina peribueyensis (Speg.) Chardon. Small hemispherical dark-brown fruiting bodies on brown spots on leaves of *M. calvescens*, *M. laevigata*, *M. lepidota*, *M. prasina*, *M. sintenisii*, *Lasiandra obscura*, and *Tibouchina multiceps* in Porto Rico, Colombia, and Brazil.

Hysterostromina miconiae (P. Henn.) Theiss. and Syd. Thick dull-black stromata on leaves of *Miconia* sp. in Brazil.

Phaeofabraea miconiae Rehm. On leaves of *Miconia* sp. in Brazil.

Phyllachora miconicola Speg. On leaves of *Miconia* sp. in Costa Rica.

Physalospora miconiae (Duby.) Sacc. On leaves of *M. calvescens* in Brazil.

Rosellinia bunodes B. and Br. See Citrus.

Septoria miconiae Garm. Brown leaf spots on *M. impatiolaris* and *M. laevigata* in Porto Rico.

MICROCITRUS. Small spiny trees allied to Citrus.

Bacterium citri Hasse. See Citrus.

MICROGLOSSA. Shrubs and climbers.

Uredo microglossae Petch. Leaf rust on *M. Zeylanica* in Ceylon.

MICROSTYLIS. See Orchidaceae.

MIKANIA. Climbing hempweed. Shrubs or twining herbs.

Aecidium mikaniae P. Henn. Round dull-brown leaf spots on *M. confertissima* in Brazil.

Apospora pachyspora Rehm. On leaves of *Mikania* sp. in Brazil.

Ascomycetella punctoidea Rehm. Sessile black stromata on leaves of *M. vismiaeifolia* in Brazil.

Cercospora sp. Leaf spot on *M. scandens* in Fiji.

Cercospora mikaniaeacola Stevens. Small circular white leaf spots with dark-brown margins on *Mikania* sp. in Porto Rico.

Chrysospora mikaniae Arth. Yellow rust pustules on leaves of *M. buddleiaefolia* and *Mikania* sp. in Brazil and Bolivia.

Endophylloides portoricensis Whetzel and Olive. Yellow rust sori on leaves of *M. cordifolia*, *M. odoratissima*, and *M. scandens* in Porto Rico, Guatemala, Trinidad, and Brazil.

Gibbera mikaniae (P. H.) Rick. and Theiss. Black stromata on leaves of *Mikania* sp. in Brazil.

Mollisia mikaniae Rehm. Indefinite yellow leaf spots on *M. discolor* in Brazil.

Septoria mikaniae Wint. Circular to irregular leaf spots on *Mikania* sp. in Porto Rico and Brazil. Reported from Texas.

MILUM. MILLET GRASS. Perennial grass.*Laetadia microspora* (Awd.) Sacc. On leaves of *M. effusum* in Germany.*Leptosphaeria nigrans* (Desm.) Ces. and De N. See Aira.*Puccinia millii* Erikss. Rust on leaves of *M. effusum* and *M. paradoxum* in Sweden, Norway, Denmark and Spain.*Puccinia piptatheri* Laegerh. Yellow-brown to dark-brown rust pustules on leaves of *M. multiflorum* in Portugal.**MILLETTIA.** Tropical trees producing hard, durable timber.*Bacillus milletiae* Kawakami. Irregular, hard, woody knots, 5 to 10 millimeters in diameter and up to 10 centimeters long, for the most part on old stems of *M. floribunda* var. *brachybotrys* in Japan.*Dierchidium woodii* (Kalehr. and Cke.) Syd. Black powdery rust sori on leaves of *M. caffra* in Natal.*Microcyclus angolensis* Sacc. and Syd. Irregular black stromata on leaves of *M. thonningii* in Angola.*Phyllachora luzonensis* P. Henn. Tar spot on leaves of *M. cavitensis*, *M. merrillii*, and *Derris philippinensis* in the Philippines.*Phyllachora milletiae* P. Henn. Black stromata on red-brown leaf spots on *Millettia* spp. in the Congo and Tanganyika.*Phyllosticta milletiae* P. Henn. Circular to angular and confluent pale-brown leaf spots on *Millettia* sp. in the Congo.*Physalospora congensis* P. Henn. Circular effuse pale-brown leaf spots on *Millettia* sp. in the Congo.**MILTONIA.** Orchid. See Orchidaceae.**MIMOSA.** SENSITIVE PLANT. Woody or herbaceous tropical plants grown for their showy flowers and feathery foliage.*Cercospora mimosae* Pat. Circular dull-brown leaf spots on *M. floribunda* in Ecuador.*Cercospora sensitivae* Speg. Rufous spots on leaves of *M. sensitiva* in Argentina.*Mycosphaerella mimosicola* P. Henn. On leaves of *M. asperata* in Brazil.*Phyllachora michelii* Speg. Black stromata on indefinite brown leaf spots on *M. procurrens* in Brazil.*Phyllachora rhytismoides* (Cda.) Sacc. Tar spot on leaves of *Mimosa* sp. in India.*Puccinia mimosae* (Speg.) Syd. Dark cinnamon rust sori on young twigs and inflorescences of *Mimosa* sp. in Argentina.*Ramularia mimosae* Stev. and Dal. Indefinite spots, often involving entire area of leaflets of *M. pudica* in Porto Rico.*Ravenelia bahiensis* P. Henn. A rust deforming the branches of *M. remansoana* in Brazil.*Ravenelia mainsiana* Arth. and Holw. Brown to black rust pustules on leaves of *M. albida* in Guatemala.*Ravenelia mimosae-albidae* Diet. Rust on leaves of *M. albida* in Mexico, Costa Rica, and Guatemala.*Ravenelia mimosae-caeruleae* Diet. Cinnamon to dark-brown rust pustules on leaves of *M. caerulea* in Mexico.*Ravenelia mimosae-sensitivae* P. Henn. Brown rust pustules on leaves of *M. sensitiva* in Colombia and Argentina.*Ravenelia mimosicola* Arth. Brown rust pustules on leaves of *M. polyanthoides* and *M. stipitata* in Mexico.*Uromyadium cubense* Arth. and Johnst. Leaf rust on *M. asperata* in Cuba.**MIMUSOPS.** Tropical trees with milky juice, some producing edible fruit.*Phyllosticta mimusopidis* P. Henn. Red-brown leaf spots on *M. schimperii* in Abyssinia.*Phyllosticta mimusopis* Cufino. Circular to irregular white leaf spots on *M. crassifolia* in Italy.*Physalospora placida* Syd. Circular gray-brown leaf spots with raised purple-brown margins on *Mimusops* in Portuguese East Africa.*Uromyces mimusops* Cke. Brown rust pustules on leaves of *M. caffra* and *M. elengi* in India, Portuguese East Africa, and the Union of South Africa.**MIRABILIS.** FOUR-O'CLOCK. Erect herbs.*Aecidium mirabilis* Diet. and Holw. Rust sori on irregular yellow leaf spots on *Mirabilis* sp. in Mexico.*Gloeosporium mirabilis* Bres. Anthracnose on stems of *M. jalapa* in Portugal.**MISCANTHUS.** Tall perennial grasses with large terminal, feathery, fan-shaped panicles.*Didymosphaeria striatospora* Syd. On stems and leaf sheaths of *M. sinensis* in Japan.*Ephelis japonica* P. Henn. Black stromata forming between the compressed spikelets, deforming them. On *M. tinctorius* and *Paspalum thunbergii* in Japan.*Phyllachora miscanthi* Syd. Black stromata on leaves of *M. sinensis* and *Miscanthus* sp. in China and the Philippines.*Puccinia erythropus* Diet. See Calamagrostis.*Puccinia eulaliae* Barcel. Purple-brown rust pustules on leaves of *M. condensatus*, *M. sacchariflorus*, *M. sinensis*, and *Pollinia japonica* in Japan and India.*Puccinia miyoshiana* Diet. Brown to black rust pustules on leaves of *M. cotulifer* and *M. sibiricus* in Japan and Siberia.*Sclerospora* sp. See Saccharum.*Sclerospora philippinensis* Weston. See Zea.*Sphaelia miscanthi* P. Henn. Black elongate sclerotia formed in ovaries of *M. sinensis* in Japan.*Ustilago kusanoi* Syd. Dark-olive smut sori destroying the panicles and rachides of *M. sacchariflora* and *M. sinensis* in Japan and the Philippines.**MITELLA.** BISHOP'S CAP. Low perennial herbs with creeping rootstocks.*Puccinia asiatica* (Kom.) Syd. Rust on leaves of *M. nuda* and *Tiarella polyphylla* in Asia.**MOLDAVICA.** See Draccephalum.**MOMORDICA.** BALSAM APPLE. Annual or perennial tropical climbing herbs, cultivated for ornament and fruit.*Aecidium momordicae* Juel. Rust on leaves of *Momordica* sp. in Brazil.*Ascochyta elaterii* Sacc. Indefinite yellow spots on leaves of *Ectellium elaterium* (*M. elaterium*) in Italy.*Cercospora elaterii* Sacc. On leaves of *M. (Ecballium) elaterium* in Italy.*Leandria momordicae* Rangel. Circular, then confluent, gray leaf spots with narrow pale-brown margins on *M. charantia* in Brazil.*Phyllosticta momordicae* F. Tassi. Effuse grayish-white leaf spots on *M. involucreta* in Italy.*Puccinia momordicae* Kalehr. and Cke. Powdery brown rust pustules on leaves of *M. cordifolia* in the Union of South Africa.*Uredo momordicae* Petch. Rust on leaves of *M. charantia* in Ceylon.**MONADENIA.** Orchid. See Orchidaceae.

MONTANOA. Shrubs, used for subtropical bedding.

Aecidium montanoae Diet. and Holw. Rust on petioles and branches of *Montanoa* sp. in Mexico.

Uromyces montanoae Arth. and Holw. Cinnamon-brown rust pustules on leave of *M. dumicola*, and *Montanoa* sp. in Costa Rica.

MONSTERA. Ceriman. Root climbing evergreen tropical aroids with edible fruit.

Neohenningsia brasiliensis P. Henn. On leaves of *Monstera* sp. in Brazil.

MONTIA. Small annual and perennial herbs, one species used as a salad plant.

Cercospora montiae Rostr. Blackened areas, often over entire surface of leaves of *M. minor* in Denmark.

Sorosporium montiae Rostr. Black smut sori in leaves and stems of *M. minor* in Denmark.

Tolyposporium junci (Schroet.) Wor. Smut in ovaries, peduncles, and culms of *M. minor* and *Juncus* spp. in Denmark and Germany.

MORAEA. MOREA. Bulbous plants resembling *Iris*.

Phyllosticta macrotheca (Thuem.) Gz. Frag. A leaf spot on *M. grandiflora* in Spain.

Puccinia moraeae P. Henn. Small brown rust pustules on both surfaces of leaves of *M. edulis* in the Union of South Africa.

Uromyces geissorhizae P. Henn. See Geissorhiza.

Uromyces moraeae Syd. Rust on leaves of *M. spathacea* in the Union of South Africa.

MORINDA. INDIA MULBERRY. Woody plants.

Cercospora morindae Syd. Circular dull-yellow leaf spots on *M. tinctoria* in India.

Phylospora morindae Koord. Circular or oblong yellow leaf spots on *M. citrifolia* in Java.

MORINGA. HYPERANTHERA Ag. HORSE-RADISH TREE. Tropical ornamental trees.

Puccinia moringae Koord. Rust on pale-yellow leaf spots on *M. oleifera* (*M. pterygosperma*) in Java.

MORUS. MULBERRY. Ornamental trees cultivated for their fruit and the leaves used for silkworm rearing.

Aecidium mori Syd. and Butl. Yellow rust sori on leaves of *M. alba* in the Philippines, India, Java, Japan, Formosa, and China.

Aecidium moricola P. Henn. Rust on leaves of *M. indica* in Java.

Ascochyta miyabei Tan. On branches of *M. alba* in Japan.

Ascochyta mori Maire. On leaves of *M. alba* in France.

Cercospora snelliana Reich. Attacks leaves of *M. alba* in Egypt.

Clasterosporium mori Syd. Brown leaf spots on *M. alba* in China and Japan.

Clasterosporium putrefaciens Frank. On leaves of *M. alba* and *Vitis vinifera* in the Caucasus Mountains.

Coryneum mori Mon. Cankers are produced on trunks and branches, causing a die-back and stunting of trees of *Morus* spp. in Japan and India. It is particularly serious on small trees in nurseries.

Cryptosporium mori Fekl. On leaves of *M. alba* in Germany.

Cylindrosporium maculans Bereng. On leaves of *M. alba* and *M. nigra* in Russia, Poland, Yugoslavia, Albania, and Denmark.

Cytodiplospora mori Miy. Parasitic on twigs of *M. alba* in Japan.

Didymella mori K. Hara. On twigs of *M. alba* in Japan.

Gibberella moricola (Ces. and De N.) Sacc. On branches of *M. alba*, *M. nigra*, and *Broussonetia* sp. in China, Italy, Austria, Hungary, and Germany.

Helicobasidium mompa Tan. and **forma macrosporum** Hara. The fungus attacks the roots which become covered with a network of purple-brown mycelial strands. Growth of infected plants is stopped, twigs and branches die back, and any new leaves formed are small. Death of the infected plants finally results. The hosts are *Beta vulgaris*, *Chamaecyparis obtusa*, *C. pisifera*, *Cryptomeria japonica*, *Cydonia japonica*, *Daucus carota*, *Diospyros kaki*, *Edgeworthia chrysantha*, *Hebe brasiliensis*, *Ipomoea batatas*, *Morus* spp., *Photinia glabra*, *Prunus domestica*, *P. pseudocerasus*, *Pyrus japonica*, *Quercus acuta*, *Q. glandulifera*, *Q. serrata*, *Rhus vernicifera*, *Salix* spp., *Sambucus racemosa*, and *Thuja* sp. in Japan, Uganda, and Straits Settlements.

Helicobasidium tanakae Miy. Circular, then large and irregular, velvety, thick brown fungus layers on trunks and twigs of *Firmiana platanifolia*, *Grossularia*, *Juglans*, *Kerria japonica*, *Malus*, *Morus*, *Paulownia tomentosa*, *Pittosporum undulatum*, *Prunus armeniaca*, *P. mume*, *P. salicina*, *Pyrus sinensis*, *Salix*, *Thea sinensis*, *Vitis*, and *Zanthoxylum* in Japan. Said to be parasitic.

Hypochnus cucumeris Frank. See Cucumis.

Massaria phorcioides I. Miy. On twigs of *M. alba* in Japan.

Mycosphaerella maculiformis (Pers.) Schroet. See Acer.

Mycosphaerella mori (Pass.) Fekl. Irregular pale-brown leaf blotches, with red-brown margins and fruit spots on *M. alba*, *M. nigra*, and *M. rubra* in Europe, South America, Japan, India, Uganda, and Canada. Infected leaves turn yellow and fall.

Mycosphaerella morifolia Pass. On leaves of *M. alba* in China and Italy. Reported from Indiana.

Ovulariopsis moricola Delacr. Effuse whitish leaf spots on *M. alba* in Madagascar and China.

Phoma mororum Sacc. Attacks twigs of *M. alba* in Italy, "often causing great injury."

Phyllosticta berlesiana Allesch. and var. **socialis** Ferr. Brown leaf spots on *M. alba* in Italy.

Phyllosticta consors Sacc. Brown leaf spots on *M. alba* in Mexico.

Phyllosticta kuwaicola K. Hara. The "circle blotch of mulberry" (*M. alba*) in Japan is characterized by minute brown leaf spots, circular to irregular, increasing in size and becoming rufous, then ashen, and concentrically zoned. The leaves are finally destroyed. Diseased twigs become brown and die. On older twigs reddish spots occur, which become black and sunken.

Phyllosticta langarum Ferr. Round to irregular, whitish leaf spots with brown margins on *M. alba* in Italy.

Phyllosticta morifolia Pass. Dirty-gray leaf spots with dull chestnut-brown margins on *M. alba* and *M. nigra* in Russia and Italy.

Phyllosticta osteospora Sacc. Rufous leaf spots on *M. alba*, *M. nigra*, *Frazinus* sp., *Populus nigra*, and *Rhamnus* sp. in Italy and France.

Phylospora minuta Miy. Slightly raised spots on living twigs of *M. alba* in Japan.

Sclerotinia shiraiana P. Henn. Forms sclerotia in fruit of *M. alba* and *M. rubra* in Japan and Formosa.

Sphaeropsis mori Berl. Attacks *Morus* in Italy.

Stagonospora mori I. Miy. This fungus attacks twigs of *M. alba* in Japan, causing them to disintegrate.

Thyrococcum sirakoffi Bub. Causes a die-back of twigs and cankers on trunk and limbs of *Morus* spp. in central Europe.

Uncinula mori Miy. Powdery mildew on lower leaf surfaces of *M. alba* in Japan and China.

Ustilago haesendockii West. A very doubtful species of smut reported on roots of *M. alba*, *M. nigra*, and *M. rubra* in Belgium, Italy, and Argentina.

A disease due to an unknown cause appears yearly on *Morus* in Indo-China, causing the curling, drying up, and premature fall of the leaves, and forming swellings on the trunk.

MOSCHOSMA. Showy-flowered small shrubs or tall herbs.

Aecidium moschomatis P. Henn. Rust on petioles and stems of *M. multiflora* in east Africa.

MUCUNA. Annual beanlike vines, grown as forage and green manure crops.

Cercospora mucunae Syd. Brown circular to irregular leaf spots on *M. urens* and *Mucuna* sp. in Brazil and the Philippines. Reported from Alabama.

Sphaerophragmium mucunae Rac. Brown rust pustules on leaves of *Mucuna* sp. in Java.

Uromyces illotus Arth. and Holw. Dark-brown powdery rust pustules on leaves of *M. andreana* in Guatemala.

Uromyces mucunae Rabh. (*Uredo mucunicola* P. Henn.). Brown to black powdery rust pustules on lower leaf surfaces of *M. capitata*, *M. pruriens*, *Stizolobium deringianum*, *S. lyoni*, and *S. niveum* in the Congo, India, Ceylon, Japan, Formosa, and the Philippines.

MUEHLENBECKIA. CALACINUM Ag. Climbing or erect plants.

Cercospora platensis Speg. On leaves of *M. sagittata* in Argentina.

Gloeosporium muehlenbeckiae Briost and Cav. Anthracnose on pseudo-leaves of *M. platyclada* in Italy.

Puccinia muehlenbeckiae (Cke.) Syd. Brown rust pustules on leaves of *M. adpressa* and *M. gracillima* in Australia.

Septoria muehlenbeckiae F. Tassi. Leaf spot on *M. complexa* in Italy.

Uromyces politus (B. and Br.) McAlp. See Jacksonia.

MULGEDIUM. Annual, biennial, or perennial herbs, allied to *Lactuca*.

Aecidium minusensis Thuem. Rust on leaves of *Mulgedium* sp. in Siberia.

Ectostroma mulgedii Thuem. Black, often confluent, spots on stems of *M. sibiricum* in Siberia.

Ovularia mulgedii Bub. Indistinct, irregular yellow leaf spots with brown centers on *M. alpinum* in Yugoslavia.

Puccinia minusensis Thuem. Brown rust pustules on leaves of *M. sibiricum* and *M. tataricum* in Siberia, Russia, Finland, and Sweden.

Puccinia mulgedii Syd. Yellow and brown rust pustules on leaves of *Mulgedium* spp. in Europe and Siberia.

Puccinia tatarica Tranzsch. Rust on stems and leaves of *M. tataricum* in Turkestan and Tibet.

Septoria mulgedii Thuem. Large irregular gray leaf spots with gray-brown margins on *M. sibiricum* in Siberia.

MUSA. BANANA. ABACA. PLANTAIN. Tropical fruit and fiber plants.

Bacillus sp. (?). A bacterial stem and bud-rot of banana and abaca has been reported from the Philippines. It is characterized by a soft black rot of the terminal bud, which extends down the central cylinder and has a disagreeable odor. Entire fields are reported to have been destroyed by this disease.

Bacillus musae Rorer. Causes a serious bacterial disease of the banana and plantain in Trinidad and probably in British Guiana. Lower leaves are first attacked, the petioles giving way, causing the blades to droop and turn yellow. The disease progresses upward, finally involving the terminal bud, after which the plants rot down to the ground. Infected plants show discolored vascular bundles at all stages. Even plants not severely diseased fail to produce fruit.

Bacillus musarum Zeman. Another bacterial disease reported as serious on varieties of bananas in Argentina, Paraguay, and probably Brazil. Leaves of infected plants turn yellow and droop, the petioles lose their turgidity and turn black. In severe cases further terminal growth ceases, and the stem dries out and dies from the apex downward. Suckers may start, but soon succumb. Vascular bundles are discolored and exude a yellow liquid when cut, which turns to red and finally black upon exposure. This disease may be identical with the preceding.

Bacterium celebensis Gäum. This bacterial disease, called the blood disease, is reported as a very serious enemy of the banana (*Musa*) in the Celebes of the Dutch East Indies. Its progress is at first slow, showing as a discoloration of one of the younger leaves, with yellow-brown stripes spreading out from the midrib. The remainder of the plant is attacked suddenly at fruiting time, and within a few days the leaves turn yellow, droop, and the bunch sags down and decays. A reddish slime exudes from the bundles when they are cut across. The fruits turn yellow and become filled with a slimy brownish-red fluid.

Bacterium musae Gäum. A second bacterial disease of the banana (*Musa*) reported from the Dutch East Indies, in which the heart leaf stops growing and the outer leaf sheaths split longitudinally, followed by a premature breaking down and wilting of the leaves. The vascular bundles are discolored.

Bunchy-top. A disease of the banana, plantain, and abaca due to an unknown cause, possibly of the group of infectious mosaics, reported as serious in Australia, Fiji, Bonin Islands, Ceylon, and Egypt. Other hosts are *Canna* and *Saccharum officinarum*. The disease checks the growth of infected plants, the leaves bunching together at the top, due to the stunting of the leaf stalks. Diseased plants never produce fruit, remaining stunted. The leaves are smaller and lighter in color than normal and very brittle, becoming ridged or corrugated with age. The bulbs show small brown or yellow flecks internally. The roots are for the most part dead, except immediately adjoining the base of the plant.

Cercospora musae A. Zimm. Indefinite brown leaf spots on *Musa* in Fiji, Java, and Ceylon. Reported as a "dangerous disease."

Cercospora musarum Ashby. Found on leaves of *Musa* in Jamaica in connection with a black spotting disease.

Coccomyces musae (Lév.) Sacc. On leaves of *Musa* in Central America.

Fusarium cubense E. F. Sm. The "Panama" or wilt disease of *Musa* sp. is the most serious disease known of this important tropical fruit. While only a few varieties, including the Gros Michel, are subject to attack, they are unfortunately the most important ones economically. The disease occurs in Central and South America, Mexico, the West Indies, Hawaii, the Philippines, and probably in India, Java, and Australia. Thousands of acres of banana land have been abandoned, due to the ravages of this disease in Central America. The entire export banana industry of Surinam was destroyed.

A diseased plant first shows a yellowing of the lower or outer leaves and petioles. The transition from normal green to yellow is sudden and proceeds from the margin inward. Diseased leaves wilt rapidly and break over a few inches from the stem and hang downward. Finally, all the leaves are involved, the innermost folded leaf resisting longest. The plant remains with dry brown, hanging leaves for a time until blown over. In severe cases young plants are attacked and remain stunted, finally dying, as do older plants. A longitudinal splitting of the outer leaf sheaths accompanies this stage. Fruit is rarely produced and is always abnormal, small, yellow, and bitter tasting. The vascular bundles are discolored, yellow at first, gradually darkening, and in advanced stages black, the blackening involving all tissues. The disease is commonly carried in offshoots.

Guignardia musae Reieb. Irregular reddish-brown leaf spots on *M. paradisiaca* in Java.

Halesia tallingii Koord. Purple, irregular, sunken leaf spots on *Musa* sp. in Java.

MUSA—Continued.

Heart rot. The abaca plant (*M. textilis*) is seriously attacked by a heart-rot disease which threatens the Manila hemp industry in parts of the Philippines. This disease has doubtfully been attributed to *Fusarium cubense* Sm. The terminal bud of infected plants develops a black rot with a disagreeable saline odor. Various varieties of edible bananas are also subject to the disease.

Laestadia musae Syd. On leaves of *Musa* sp. in the Congo.

Leptosphaeria musarum Sacc. and Berl. On leaves of *M. paradisiaca* in Ceylon and Portuguese St. Thomas.

Marasmius semustus B. and C. This gill fungus has been reported from the West Indies, Trinidad, Hawaii, and Mauritius as the cause of a root disease of the banana, and an extensive literature has developed. Its status is much the same as *M. sacchari* on *Saccharum*. (q. v.).

Mycosphaerella musae (Speg.) Syd. Leaf spot on *M. cavendishii*, *M. sapientum* and *M. textilis* in Guam and the Philippines.

Phoma musae Carpenter. The "freckle" disease is serious on *M. cavendishii* in Hawaii and has been reported on *Musa* from the Philippines. The fruit spots are at first minute, circular, gray, with indefinite borders, becoming larger and slightly erumpent. Similar spots occur on the leaves, at times causing defoliation. Diseased fruit are undersized, discolored, and ripen unevenly, and even when only slightly attacked are rendered unfit for export.

Phoma musarum Cke. On *Musa* in Bermuda and Spain.

Phyllachora musae (Kl.) Sacc. On leaves of *Musa* sp. in India and the Congo. Referred by Theissen to *Macrophoma musae* (Sacc.) Berl. and Vogl.

Phyllostictia gastoni (Roum.) Tassi. On leaves of *M. sapientum* in Haiti.

Sphaerostilbe musarum Ashby. The so-called "bonnygate" disease of Jamaican bananas is attributed to this fungus. Leaves of diseased plants show a narrow zone of brown along the margins, bordered by a yellow band. Inner leaves usually bear broad yellow bands. Suckers grow slowly and seldom form fruit. Old stems may break over, a black-rotted area having formed at the base upon which the yellow fruiting bodies of the fungus appear.

Thielaviopsis paradoxa (De Sey.) v. Roeh. See *Saccharum*.

Tylenchus musicola Cobb. Plants of *Musa* of all ages are attacked in the British West Indies, the outer leaves dying and the fruit, if developed, drying up. The roots die, becoming black in color. The outer part of the bulb from which the roots develop likewise becomes black and rotted.

Tylenchus similis Cobb. This nematode attacks the roots of *Musa* spp. and *Saccharum officinarum* in Jamaica, Hawaii, and Fiji. The roots die back, diseased tissue becoming cinnabar-red and finally dark-purple.

Uromyces musae P. Henn. Powdery brown to black elongate rust sori on under leaf surfaces of *Musa* sp. in the Congo.

MUSCARI. GRAPE HYACINTH. Hardy spring flowering bulbs.

Aecidium muscari Link. Forms yellow rust pustules on leaves of *M. comosum* and *M. racemosum* in England, Italy, France, Switzerland, Dalmatia, and Hungary. Said to be distinct from *Uromyces scillarum*.

Heterosporium ornithogali Klotzsch. See *Ornithogalum*.

Physoderma muscari Poir. On leaves of *M. comosum* in France.

Puccinia lilacearum Duby. See *Ornithogalum*.

Puccinia lojkaiana Thuem. See *Ornithogalum*.

Sclerotium sp. A storage rot due to a *Sclerotium* (*Botrytis*) commonly infects imported Holland bulbs of *Muscari* spp. Infected bulbs are matted together by a web of white mycelium in which small brown to black sclerotia appear. Other related bulbous plants may also be attacked.

Septoria muscari P. Brun. Large irregular spots with brown margins on leaves of *M. comosum* in France. This species is probably identical with *S. scillae* West.

Septoria muscari-neglecti Bub. On leaves of *M. neglectum* in Yugoslavia.

Septoria scillae West. Causes a leaf spot of *Muscari* sp., *Scilla bifolia*, and *S. nutans* in Belgium, Italy, Portugal, Spain, and Germany.

Urocystis colchici (Schlecht.) Rab. See *Colchicum*.

Uromyces ornithogali Lévl. See *Gagea*.

Uromyces scillarum (Grev.) Wint. See *Hyacinthus*.

Ustilago vaillantii Tul. See *Hyacinthus*.

MUTISIA. Showy flowered shrubs.

Phyllachora mutisiae Speg. A doubtful species on *Mutisia* sp. in Argentina.

Puccinia crassicutis Syd. Yellow and black rust pustules on *M. clematis* in Bolivia.

Puccinia mutisiae Lagerh. Brown rust pustules on leaves and stems of *Mutisia* sp. in Ecuador

Puccinia mutisiicola Speg. Brown rust pustules on leaves of *M. subspinoso* in Argentina.

Septoria jaffueli Speg. On leaves of *M. latifolia* in Chile.

Septoria mutisiae Speg. Ashen-colored leaf spots on *M. subspinoso* in Argentina.

MYOPORUM. Heathlike shrubs, grown for their small white or purplish flowers.

Bagnisiella endopyria Sacc. On leaves of *M. platycarpum* in Australia.

Septoria myopori Cke. and Mass. Circular whitish leaf spots with dull-brown margins on *M. insulare* in Australia.

MYOSOTIS. FORGET-ME-NOT. Low perennial or annual herbs.

Aecidium kabatianum Buh. Rust on leaves of *M. sparsiflora* and *M. stricta* in central Europe.

Cylindrosporium myosotidis Sacc. and var. **boraginis** Sacc. Leaf spot on *M. palustris*, *M. scorpioides*, *Borago officinalis* and *Symphitum* sp. in Poland, France, and Spain.

Entyloma canescens Schroet. Smut sori in gray circular leaf spots on *M. caespitosa*, *M. hispida*, *M. intermedia*, and *M. sylvatica* in France, Denmark, Italy, and Germany.

Peronospora myosotidis De By. Downy mildew on leaves of *M. arenaria*, *M. arvensis*, *M. hispida*, *M. intermedia*, *M. palustris*, *M. sylvatica*, *M. sparsiflora*, *M. stricta*, and *M. versicolor* in Europe.

Puccinia myosotidis Transsch. Black rust sori on leaves and stems of *M. sylvatica* in Turkish.

Puccinia novo-zelandica Bub. Dark-brown rust pustules on leaves of *M. capitata* in New Zealand.

Synchytrium globosum Schroet. See *Potentilla*.

MYRCIA. Tropical shrubs.

Catacauma myrciae (Lévl.) Theiss. and Syd. Dark-black stromata on leaves of *Myrcia* sp. in Brazil.

Puccinia sanguinolenta P. Henn. Dark-brown rust pustules on leaves of *Myrcia* sp. in Brazil.

Uredo myrciae Mayor. Rust on leaves of *M. acuminata* in Colombia.

MYRCIARIA. Jaboticaba. Brazilian fruit trees and shrubs.

Laestadia cambuciae Rangel. On leaves of *M. plicati-costata* in Brazil.

Uredo roebiae Pult. Rust on fruit and leaves of *M. jaboticaba* and *M. plicati-costata* in Brazil.

MYRICA. BAYBERRY. Ornamental woody plants with attractive fruits, some edible.

Dothidella koordersii (P. Henn.) Theiss. and Syd. Black stromata on leaves of *M. germanica* in Java.

Dothidina disciformis (Wint.) Theiss. and Syd. Black stromata on leaves of *Myrica* sp. in the Union of South Africa.

MYRICA—Continued.

- Microcyclus koordersii** P. Henn. Black discoid stromata on lower leaf surfaces of *M. javanica* in Java.
- Peronospora rubribasis** B. and Br. Downy mildew on brown leaf spots on *M. gale* in Great Britain.
- Rhytisma nitidum** Lév. Black tar spots on *M. aethiopica* in Abyssinia.
- MYRICARIA.** FALSE TAMARIX. Shrubs allied to Tamarix.
- Marsonia myricariae** Rostr. On leaves of *M. germanica* in Norway.
- Phyllosticta germanicae** Speg. On leaves of *M. germanica* in Italy.
- Puccinia involvens** (Voss.) Syd. Yellow-brown to black elongate rust pustules on *M. germanica* in France, Belgium, Switzerland, and Austria.
- Pyrenopeziza tamaricis** (Roum.) Sacc. On branches of *M. germanica*, *Tamarix africana*, and *T. narbonensis* in Argentina, Great Britain, France, and Germany.
- MYRISTICIA.** NUTMEG. Tropical trees with aromatic fruit.
- Corticium salmonicolor** B. and Br. See Citrus.
- Eutype erumpens** Mass. See Theobroma.
- Fomes lamaensis** Murr. See Hevea.
- Marasmius** spp. See Theobroma.
- Perisporium myristicae** P. Henn. Black superficial globular perithecia on leaves of *Myristica* sp. in Java.
- Phomopsis** sp. Causes a light-brown rot of fruit of *M. fragrans*, preventing seed germination, in Grenada.
- Phytophthora** sp. An undetermined species of this genus has been reported from Java, attacking the leaves and twigs of *Myristica* (nutmeg).
- MYRRHIS.** MYRRH. Perennial herbs.
- Puccinia chaerophylli** Purl. See Chaerophyllum.
- MYRSINE.** Tropical shrubs and trees with coriaceous leaves and white or yellow flowers.
- Chaetothyrilum punctiforme** Rick. On leaves of *Myrsine* sp. in Brazil.
- Corynelia fruticula** (Pat.) v. Hoehn. Black carbonous superficial fruiting bodies, more or less completely enveloping fruit of *M. africana* and *Rapanea melanophleos* in China, India, and the Union of South Africa.
- Napieladium piriforme** Speg. Circular brown leaf spots on *M. floribunda* in Argentina.
- Phyllachora myrsincola** Doidge. Black stromata on leaves of *M. (Rapanea) melanophleos* in the Union of South Africa.
- Phyllachora pittieri** Theiss. and Syd. Tar spot on leaves of *Myrsine* sp. and *Xylosma salzmanni* in Costa Rica.
- Phyllachora sinik-lagaraik** Speg. Circular shiny black stromata on leaves of *M. floribunda* in Brazil.
- Phyllohendersonia nitida** (E. and E.) Tass. On leaves of *Myrsine* sp. in Hawaii.
- Scolecodothis circularis** (Bres.) Theiss. and Syd. Shiny black stromata on brown leaf spots on *Myrsine* sp. in Brazil.
- Uromyces myrsines** Diet. See Ardisia.
- MYRTUS.** MYRTLE. Shrubs grown for their aromatic qualities and attractive foliage, flowers and fruit.
- Ascochyta apiospora** Cke. and Mass. See Backhousia.
- Ascochyta myrticola** Maire. and Sacc. Subcircular reddish leaf spots with dark-purple margins on *M. communis* in Italy and Corsica.
- Cercospora myrti** Erikss. Subcircular purplish leaf spots on *M. communis* and *M. italica* in Scotland, Sweden, Denmark, Spain, and Austria.
- Cercospora saecardiana** Scalia. Indefinite dark-purple leaf spots on *M. communis* in Sicily.
- Depazea myrticola** Rabenh. On leaves of *M. communis* in Italy.
- Macrophoma nuptialis** Bub. Large yellow-brown leaf spots on *M. communis* in Yugoslavia.
- Mycosphaerella myrticola** Speg. On leaves of *M. chequen* in Chile.
- Napieladium fumago** Speg. See Eugenia.
- Phyllosticta costalis** Speg. On leaves of *M. chequen* in Chile.
- Phyllosticta nuptialis** Thuen. Subcircular dull-yellow to white leaf spots with violet margins on *M. communis* and *M. italica* in Albania, Dalmatia, Yugoslavia, Madeira, and Portugal.
- Rehmiodothis myrtincola** (Rehm.) Theiss. and Syd. Shiny black stromata on leaves of *Myrtus* sp. in Ecuador.
- NAGEIA.** See Podocarpus.
- NANDINA.** Small shrubs with red or white berries.
- Phyllosticta nandinae** Tassi. Irregular dull-yellow leaf spots on *N. domestica* in Italy.
- NAPOLEONA.** Shrubs of tropical Africa.
- Phyllosticta napoleonae** Thuem. Irregular whitish leaf spots on *N. imperialis* in Portugal.
- NARCISSUS.** Daffodil. Hardy spring flowering bulbous plants.
- Bacillus croci** Miuza. See Crocus.
- Botrytis parasitica** Cav. The black sclerotia of this fungus are common on imported *Narcissus* bulbs, and it has doubtless been introduced into all parts of the United States. See under Tulipa.
- Cercospora narcissi** Bond. Causes spotting on both surfaces of the leaves of *N. poeticus* in France.
- Coleosporium narcissi** Grove. Brown rust pustules on both surfaces of leaves of *N. poeticus* in England.
- Fusarium bulbigenum** Cke. and Mass. Reported by Massee as a very serious disease of *Narcissus* in Great Britain. The disease has also been reported from Holland. It is characterized by small yellowish to brown leaf spots which finally become covered with pale salmon-colored spore masses. The bulbs also ultimately become infected. Work by recent investigators, however, has quite clearly shown that *Fusarium bulbigenum* is a secondary organism following nematode attack and not the cause of a separate disease.
- Phyllosticta narcissi** Aderh. Large brown spots without definite margins on leaves of *N. poeticus* in Germany.
- Phyllosticta odemensis** Sacc. and Syd. On leaves of *Narcissus* (cultivated varieties) in Holland.
- Puccinia schroeteri** Pass. Large oblong spots are produced on both surfaces of the leaves, with dull-violet margins. Brown to black rust pustules appear on these spots. The disease occurs in Great Britain, Italy, Austria, Belgium, and Germany on *N. pseudonarcissus* and *N. radiiflorus*.
- Ramularia narcissi** Chittenden. This disease is characterized by yellow stripes down the foliage and a dwarfing of the entire plant, involving also the flowerstalks. Yellow-brown spots up to 1 centimeter in length also occur on leaves, which later become covered with a fine white powder. The host is *N. poeticus* in variety in England. The fungus may be synonymous with the following species as well as with *Cercospora narcissi* Bond.
- Ramularia vallambrosae** Br. and Cav. A brown spotting of leaves and flower stalks in Italy of *N. biflorus*, *N. odoratus*, *N. poeticus*, and *N. pseudonarcissus*.
- Septoria narcissi** Pass. Attacks tips of living leaves of *Narcissus* (cultivated varieties) in Italy, Belgium, and Holland.

NARCISSUS—Continued.

- Stagonospora narcissi** Höll. Produces large irregular brown spots on living leaves of *N. poeticus* in Hungary.
- Tylenchus dipsaci** Kuehn. (*T. devastatrix* Kuehn.) The eelworm or nematode disease of bulbs (*Narcissus* and *Hyacinthus*) is beyond any doubt the most serious bulb disease known at present. On growing plants the disease is characterized by twisted leaves, with swollen yellow spots and stripes. Affected leaves as a rule remain stunted and may even die. The minute worms causing the disease can be found in these diseased areas. In the bulbs the presence of the nematode is noted by dark rings. The disease occurs in Holland, Great Britain, and France. It has been introduced into the United States with Dutch bulbs and it is known from a few localities. It has proved a serious pest of *Allium* spp., *Avena sativa*, *Fraxia* sp., *Linum usitatissimum*, *Medicago sativa*, *Pisum sativum*, *Secale cereale*, *Solanum tuberosum*, *Trifolium pratense*, and other economic plants in Europe and Japan.
- Urocystis colchici** (Schlecht.) Rab. See Colchicum.
- NARTHECIUM.** BOG ASPHODEL. Perennial rhizomatous herbs.
- Entyloma ossifragi** Rostr. Smut sori on gray, angular leaf spots on *N. ossifragum* in Denmark.
- Heterosporium ossifragi** Rostr. Produces spots on leaves of *N. ossifragum* in Denmark.
- NASTURTIIUM.** WATER-CRESS. Semiaquatic herbs. Now referred to *Radicula*. (q. v.)
- Aecidium nasturtii** Hazsl. Rust on leaves of *Radicula* (*Nasturtium*) sp. in Hungary.
- Peronospora nasturtii-montani** Gäum. Downy mildew on leaves of *N. montanum* in Japan.
- Phyllosticta anceps** Sacc. Small subcircular ochraceous-yellow leaf spots on *N. amphibium* and *N. anceps* in England and France.
- Phyodermma magnusiana** Krieg. Small oblong spots on leaf blades and petioles of *N. palustre* in Germany.
- Septoria uliginosa** Speg. Definite subcircular whitish leaf spots on *N. bonaërense* in Argentina.
- NELUMBO** (*Nelumbium*). LOTUS. Water lilies.
- Cercosporina nelumbi** Hori. On leaves of *N. nucifera* (*N. speciosum*) in Japan and China.
- NEMESIA.** Flower-garden annuals.
- Albugo evansii** Syd. White rust on leaves of *Nemesia* sp. in the Union of South Africa.
- Cronartium flaccidum** (Alb. and Schw.) Wint. See Paonia.
- NEPENTHES.** PITCHER PLANT. Aquatic pitcher plants.
- Phyllachora nepenthidis** Rac. Circular to elliptical shining black stromata on leaves of *N. melamphora* in Java.
- Phyllosticta nepenthaearum** F. Tassi. Subcircular dull-brown leaf spots on *N. sanguinea* in Italy.
- NEPETA.** CATNIP. GROUND IVY. Hardy perennial herbs.
- Erysiphe taurica** Lév. See Althaea.
- Septoria bornmuelleri** Syd. Irregular dirty yellow-brown leaf spots, often destroying entire leaf, on *N. teucrifolia* in Persia.
- Septoria catariae** Bub. Yellow leaf spots, often with greenish-gray surrounding zones, on *N. beltranii* and *N. cataria* in Spain and Hungary.
- NEPHELIUM.** LYCHEE. LITCHI. Tropical fruit trees.
- Ekoascus** sp. Leaf curl of *N. litchi* (*Litchi chinensis*) in India.
- Phyllosticta nephelei** Delacr. See Durio.
- NEPHROLEPIS.** SWORD FERN.
- Entyloma nephrolepidis** Rac. Smut sori in fronds of *N. biserrata* (*N. acuta*) in Java.
- Milesina columbiensis** (Diet.) Arth. Rust on fronds of *N. pendula* and *N. rivularis* in Colombia and Porto Rico.
- NERIUM.** OLEANDER. Glaucons, evergreen-leaved shrubs.
- Ascochyta oleandri** Sacc. and Speg. White leaf spots with brown margins on *N. oleander* in Denmark, France, Italy, and Portugal.
- Cercospora neriella** Sacc. On leaves of *N. oleander* in Italy. Reported from Louisiana.
- Phyllosticta glaucispora** Delacr. White leaf spots with dull-brown margins on *N. oleander* in Spain.
- Phyllosticta nerilicola** Brun. Pale-ochraceous leaf spots on *N. oleander* in France.
- Phyllosticta nerii-oleandri** Stem. Leaf spot on *N. oleander* in the Caucasus.
- Septoria nerilicola** Pass. On leaves of *N. oleander* in Italy.
- Septoria oleandrina** Sacc. On *N. oleander* in Bermuda, France, and Italy.
- NICOTIANA.** TOBACCO. Erect, rank-growing herbs with narcotic or poisonous properties.
- Ascochyta nicotianae** Pass. Irregular dull-brown leaf spots on *N. affinis* and *N. tabacum* in Italy, Sweden, Russia, and possibly some of the West Indies.
- Bacillus caulivorus** Prill. and Delacr. See Solanum.
- Bacillus maculicola** Delacr. Bacterial leaf spot of *N. tabacum* in France.
- Bacterium pseudozoogloeae** Honing. Said to cause black rust disease of leaves of *N. tabacum* in the Dutch East Indies.
- Cercospora raciborskii** Sacc. and Syd. Circular brown to whitish zoned leaf spots on *N. tabacum* in Java, India, Ceylon, Fiji, Australia, and Uganda.
- Colletotrichum nicotianae** Avena. Irregular confluent dark-brown anthracnose spots on stems and branches of *N. tabacum* in Brazil.
- “Coreova.” This disease of tobacco (*N. tabacum*), due to an unknown cause, is reported from Argentina under the native name. Numerous short (1 to 3 millimeters) dark lines, resembling insect burrows, appear on the leaves, uniting to form much longer ones. The leaf tissue between dies and the leaves wilt. Similar lesions from 2 to 6 centimeters long occur on the stems. Unaffected portions of leaves continue to expand, causing swelling and distortion of the leaf blades. The pith of the stem darkens and dries up.
- Cytospora nicotianae** Avena. Irregular yellow areas on stems of *N. tabacum* in Brazil.
- Diplodia cacaocola** P. Henn. See Theobroma.
- Macrophoma tabaci** Avena. Irregular gray spots on the stems of *N. tabacum* in Brazil.
- Oidium tabaci** Thuem. A powdery mildew forming circular white powdery spots on leaves of *N. tabacum* in the Union of South Africa, Italy, and Portugal.
- Peronospora hyoscyami** De By. (*Peronospora nicotianae* Speg.) A downy mildew attacking cotyledons, leaves, and roots of *N. glauca*, *N. tabacum*, and *Hyoscyamus niger* in Europe, Australia, Mexico, and in limited areas in the United States. Causes the serious blue mold disease in Australia.
- Phoma solanicola** Prill. and Delacr. Irregular yellow, then brown, areas on stems and branches of *N. tabacum* and *Solanum tuberosum* in Brazil, Spain, Italy, and France. The leaves wither when the areas are extensive. Avena-Sacca reports *Plowrightia solanicola* Avena as a perfect stage.
- Phyllosticta capsulicola** Sacc. and Speg. On capsules of *N. dilatata* in Italy.
- Phyllosticta nicotianicola** Speg. Ashen leaf spots on *N. acutiflora* in Argentina.
- Phyllosticta tabaci** Pass. Irregular, often confluent, ashen-colored leaf spots on *N. tabacum* in Italy and Yugoslavia.
- Phytophthora melongenae** K. Saw. See Solanum.

NICOTIANA—Continued.

Phytophthora nicotianae v. B. de Haan. (*Peronospora nicotianae* Speg.) This downy mildew causes a damping-off of tobacco seedlings. In addition it attacks leaves, roots, and stems of older plants and is destructive in Ceylon, Sumatra, Borneo, Java, Brazil, and Argentina.

Placophæria nicotianae Avena. Irregular dark-brown or black patches at bases of stems of *N. tabacum* in Brazil.

Rhizoctonia sp. See Vigna.

Sclerotinia nicotianae Oud. and Konig. Rotten areas, covered with gray mold on leaves and stems, in which black sclerotia form on *N. tabacum* in Holland.

Septoria diversa Sacc. and Syd. Rufous leaf spots with ashen-colored centers on *N. longiflora* in Argentina.

Septoria nicotianae Pat. Brown to white zoned leaf spots on *N. affinis*, *N. rustica*, and *N. tabacum* in Russia, Ecuador, and Jamaica. The frog-eye disease.

Septoria tabacina McAlp. Leaf spot on *N. tabacum* in Australia.

Uredo nicotianae Anast. and Spenk. Powdery brown rust sori on leaves of *N. quadrivalvis*, *N. sylvestris*, and *N. tabacum* in Brazil and Italy.

NIDULARIUM. Epiphytic bromeliads related to the pineapple.

Uredo nidularii P. Henn. Linear brown rust pustules on leaves of *N. longiflorum* in Brazil.

NIEREMBERGIA. CUPFLOWER. Tender perennial herbs.

Entyloma nierembergiae Lagh. Smut sori on pale circular leaf spots on *N. spathulata* in Ecuador.

Puccinia nierembergiae Lév. Powdery black rust pustules on leaves of *Nierembergia* sp. in Peru.

NIGELLA. LOVE-IN-A-MIST. Hardy annuals grown for their interesting flowers and seed pods.

Cercospora nigellae Höll. On leaves of *N. arvensis* in Hungary.

NIOBE. See Hosta.

NIPA. Oriental thatch palm with edible fruit. See Palmae.

NITRARIA. NITER BUSH. Low, rigid shrubs.

Aecidium nitrariae Pat. Rust on leaves of *N. tridentata* in Algeria.

NOTHOFAGUS. Ornamental South American trees.

Cytaria darwini Berk. Causes a rot of the wood of branches of *Nothofagus* sp. in Chile.

Exoascus entomosporus (Thaxt.) Sacc. and Trot. Fruiting layer on lower leaf surfaces of *N. antarctica* var. *bicrenata* and var. *uliginosa* in Patagonia. The leaves are deformed.

Uncinula magellanica Thaxt. Powdery mildew on leaves of *N. antarctica bicrenata* in Patagonia.

Uncinula nothofagi Thaxt. Similar to above species.

NYCTANTHES. NIGHT JASMINE. Trees or shrubs with numerous white fragrant flowers.

Cercospora puttemansii P. Henn. Dull-brown leaf spots on *N. arbor-tristis* in Brazil.

Ramularia sp. Areolate mildew on leaves of *N. arbor-tristis* in India.

NYMPHAEA. WATER LILY (including *Nuphar*). SPATTER-DOCK. Herbaceous aquatics.

Aecidium nymphaeoides DC. Rust on leaves of *N. alba*, *N. luteum*, and *Limnanthemum nymphaeoides* in France, Italy, Switzerland, Belgium, and Germany.

Ovulariella nymphaearum (Allesch.) Kab. and Bub. (*Gloeosporium nymphaearum* Allesch.) Subcircular to irregular, often confluent, reddish to gray-brown leaf spots, pale at the centers, on *N. alba*, *N. douglasiana*, *N. copensis*, *N. flava*, *N. luteum*, and *N. ortgiesiana* in Europe.

Phyllosticta hydrophila Speg. Subcircular sunken dull-brown leaf spots on *N. alba* in Italy.

Phyllosticta nymphaeae Pass. On leaves of *N. alba* in Italy.

Physoderma tenue (Nowak.) Karst. On petioles of *Nymphaea* sp., *Acorus calamus*, and *Iris pseudacorus* in Denmark, France, Silesia, Finland, and Germany.

Pythium undulatum H. E. Peters. Downy mildew on leaf blades and petioles of *N. luteum* and *N. alba* in Denmark.

Ramularia nymphaeae Bres. Black circular leaf spots on *N. alba* in England and Germany.

Rhamphospora nymphaeae Cunn. Smut on leaves of *N. lotus*, *N. rubra*, and *N. stellata* in India.

Septoria nupharis Ranof. On leaves of *N. luteum* in Yugoslavia.

OCHLANDRA. See Bambuseae.

OCHNA. Woody plants.

Cocconia capensis Doidge. On *O. arborea* in the Union of South Africa.

Phyllachora ochnae Pat. and Har. Tar spot on leaves of *Ochna* sp. in Madagascar and the Philippines.

OCIMUM. BASIL. Annual and perennial herbs.

Aecidium leiocarpum Syd. Leaf rust on *O. canum* in India.

Aecidium ocimi P. Henn. Golden-yellow rust sori on circular brown leaf spots on *O. canum*, *O. graveolens*, and *O. suave* in India and Abyssinia.

Coleosporium plectranthi Barcl. See *Plectranthus*.

Phyllosticta basilici Brun. Gray-brown leaf spots on *O. basilicum* in France.

OCOTEA. Tropical trees.

Cephalosporium tumefaciens Wint. On *O. tristis* in Brazil.

Oligostroma archavaletae (Speg.) Theiss. and Syd. Black tar spot on leaves of *O. acutifolia* in Uruguay.

Phyllachora ocoteae P. Henn. Irregular black stromata on leaves of *Ocotea* sp. in Brazil.

Phyllachora ocoticola Stev. and Dalby. Angular dull-black stromata on leaves of *O. leucorylon* in Porto Rico.

ODONTOGLOSSUM. Epiphytic orchids. See Orchidaceae.

OENANTHE. Aquatic perennial herbs.

Entyloma debonianum Sacc. Smut sori on stems of *O. globulosa* in Malta.

Entyloma oenanthae R. Maire. White smut pustules on leaves of *O. apifolia* in Corsica.

Physoderma vagans Schroet. See *Ranunculus*.

Puccinia oenanthae (Diet.) T. Miy. Dark-brown rust pustules on leaves of *O. stolonifera* in Japan.

OENOTHERA. EVENING PRIMROSE. Flower-garden plants with bright-colored flowers.

Cercospora oenotherae Speg. Circular dull-brown leaf spots on *O. mollissima* in Argentina.

Puccinia luxurians Diet. and Neg. Brown rust pustules on leaves of *O. multifida* in Chile.

Uredo oenothericola Speg. Powdery, dark ferruginous rust pustules on leaves of *O. mollissima* in Argentina.

OLEA. OLIVE. Fruit and ornamental trees.

Ascochyta oleae Scal. On leaves of *O. europaea* in Sicily.

Bacterium olivae Cif. Found in connection with dieback of twigs of *O. europaea* in Italy.

Cercospora cladosporioides Sacc. On leaves of *O. europaea* in Italy and Algeria.

Ceuthospora oleae Kalcbr. and Cke. Circular dull-brown leaf spots on *O. capensis* in the Union of South Africa.

Ceuthospora phacidioides Grev. and var. *oleae* Scalia. On leaves of *O. europaea* in Sicily and Austria.

Coniothyrium oleae Pol. On leaves of *O. europaea* in Italy.

OLEA—Continued.

- Cylindrosporium olivae** Petri. Large sunken yellowish-purple spots with dark purple margins on fruit of *O. europaea* in Italy.
- Cystospora oleae** Butl. Rust on pale-brown leaf spots on *O. dioica* in India.
- Gloeosporium olivarum** d'Alm. Anthracnose on fruit of *O. europaea* in France, Italy, Spain, Portugal, and Algeria.
- Macrophoma dalmatica** (Thuem.) Berl. and Vogl. Subcircular dull-brown spots on fruit of *O. europaea* in Spain, Italy, and Austria.
- Perischizon oleifolium** (Kalch. and Cke.) Syd. On leaves of *O. capensis* in the Union of South Africa.
- Phyllosticta insulana** Mont. Small pale leaf spots with narrow brown margins on *O. europaea* in Spain, Italy, and France.
- Phyllosticta oleae** Petri. Leaf spots on *O. europaea* in Italy.
- Septoria oleae-chrysophyllae** Pass. On leaves of *O. chrysophylla* in Abyssinia.
- Septoria oleaginea** Thuem. Irregular pale-ochraceous leaf spots on *O. europaea* in Austria.
- Septoria olivae** Pass. and Thuem. On leaves of *O. europaea* in Portugal.
- Stictis panizzeae** De N. Causes the leaf disease of *O. europaea* known as Brusca in Spain and Italy.
- OLEARIA. SHAWIA** Ag. DAISY TREE. Australasian trees and shrubs.
- Acidium oleariae** McAlp. Rust on leaves and stems of *O. axillaris* in Australia.
- Puccinia oleariae** McAlp. Bright orange then brown rust pustules on leaves of *O. argophylla* in Tasmania.
- Uredo brownii** Syd. Rust on leaves of *O. angustifolia* in New Zealand.
- ONCIDIUM**. Epiphytic orchids. See Orchidaceae.
- ONCOSPERMA**. See Palmae.
- ONOBRYCHIS. HOLY CLOVER**. Perennial forage plants.
- Anthostomella** sp. Black leaf spots, often confluent, with yellow surrounding zones, on *O. sativa* in Italy.
- Ascochyta orobi** Sacc. var. **onobrychidis** Prill. and Délacr. On leaves of *O. sativa* in France and Yugoslavia.
- Cylindrosporium onobrychidis** (Syd.) Died. On *O. sativa* in Europe.
- Oularia bornmülleriana** P. Magn. Yellow leaf spots on *O. tournefortii* in Asia Minor.
- Peronospora ruegeriae** Gäm. Downy mildew on leaves of *O. sativa* in Switzerland.
- Phylachora lathyri** (Lév.) Theiss. and Syd. See Lathyrus.
- Physalosporina ducllerii** Maire. On leaves of *O. caputgallii* in French North Africa.
- Ramularia onobrychidis** Allesch. Clear light-brown leaf spots with darker margins on *O. sativa* and *O. viciifolia* in Russia, Germany, Sweden, Yugoslavia, France, and Denmark.
- Rhizisma onobrychidis** DC. Irregular warty dull-black patches, on leaves of *O. sativa* and *Vicia cracca* in Siberia and France.
- Uromyces onobrychidis** (Desm.) Lév. Powdery brown rust pustules on *O. arenaria*, *O. gracilis*, *O. montana*, *O. sativa*, and *O. viciifolia* in Europe.
- ONOCLEA**. Sensitive fern.
- Taphrina hiratsuka** Nish. Fruiting areas on leaves of *O. sensibilis* in Japan.
- ONONIS. REST-HARROW**. Annual, biennial, and perennial border and rock-garden plants.
- Napcladadium ononidis** (Auerw.) Sacc. On leaves of *O. repens* in France.
- Peronospora ononidis** Wils. Downy mildew on leaves of *O. hircina*, *O. repens*, and *O. spinosa* in Russia, Denmark, Switzerland, Austria, and Germany.
- Ramularia winteri** Thuem. On leaves of *O. arvensis*, *O. hircina*, *O. repens*, and *O. spinosa* in Switzerland, Denmark, and Germany.
- Uromyces ononidis** Pass. Cinnamon to dark brown rust pustules on leaves of *O. alopecuroides*, *O. antiquorum*, *O. campestris*, *O. columna*, *O. hircina*, *O. procurrens*, *O. repens*, *O. rotundifolia*, and *O. spinosa* in Europe and Asia Minor.
- Uromyces sphaeropleus** Cke. Dark-brown rust pustules on *Ononis* sp. in India.
- ONOPORDON. COTTON THISTLE**. Coarse, woolly, old-world herbs.
- Puccinia acanthii** Syd. Powdery brown rust pustules on leaves of *O. acanthium* in Spain and Germany.
- Puccinia onopordi** Syd. Brown rust pustules on leaves of *O. cardunculum* in Syria.
- Ramularia onopordi** Massal. Subcircular yellow leaf spots on *O. acanthium* and *O. ceratonicum* in Spain, Italy, Yugoslavia, and Germany.
- Septoria onopordonis** P. Nag. Leaf spot on *O. acanthium* in Russia.
- ONOSMA**. Herbs grown for their golden-colored flowers.
- Acidium onosmatis** Thuem. Rust on leaves of *O. gmelini* and *O. simplicissima* in Siberia.
- OPHIOGLOSSUM. ADDER'S-TONGUE**.
- Helminthosporium crepini** West. On leaves of *O. vulgatum* in France, Belgium, and Denmark.
- Helminthosporium diedickei** P. Magn. On *O. vulgatum* in Germany.
- OPHIPOGON. SNAKE'S-BEARD**. Herbs grown for their small bluish or white flowers.
- Fusicladium transversum** Sacc. Greenish to black areas on leaves of *O. japonicus* in Italy.
- Septoria aemula** Tassi. Brown leaf spots on *O. spicatum* in Italy.
- Septoria ophiopogonis** Pass. Brown areas at the tips of leaves of *O. japonicus* in Italy.
- Uredo ophiopogonis** Syd. Cinnamon-brown rust sori on leaves of *O. japonicus* in Italy.
- OPHRYS**. Terrestrial orchids. See Orchidaceae.
- OPULASTER**. See Physocarpus.
- OPUNTIA. PRICKLY PEAR**. Sometimes cultivated for ornament or forage.
- Acidium opuntiae** P. Magn. On *Opuntia* sp. in Bolivia.
- Ascochyta opuntiae** Sealia. On *O. ficus-indica* in Italy.
- Didymosphaeria opulenta** Sacc. Causes yellow-brown blotches on joints of *O. ficus-indica* in Italy and Malta.
- Phyllosticta opuntiae** Sacc. and Speg. and var. **microscopicae** Cifer. Whitish spots on *O. ficus-indica* and other species in Italy and Bermuda.
- Sclerotium opuntiarum** Speg. On *Opuntia* spp. in Argentina.
- Septoria ficus-indicae** Vogl. On *O. ficus-indica* in Italy.

ORCHIDACEAE.

The diseases attacking the various genera of the *Orchidaceae* are considered under the latter head for reasons similar to those given under *Bambuseae*.

- Acidium alaskanum** Trel. This rust forms small, pale to light brown spots on both surfaces of the leaves of *Habenaria* sp. and other orchids in Alaska.
- Amerosporium rhodospermum** McAlp. On leaves of *Diuris pedunculata* in Australia.
- Amerosporium vanillae** P. Henn. On leaves of *Vanilla aromatica* in Java.
- Ascochyta orchidis** Rabenh. On leaves of *Orchis* sp. in Germany.
- Ascospora** sp. On living leaves of *Evelyna discolor* in Peru.
- Asterina liparidis** Rac. On leaves of *Liparis* sp. in Java.

ORCHIDACEAE—Continued.

Asterinella epidendri (Rehm.) Theiss. The fungus produces small, black, shield-shaped fruiting bodies on the upper surfaces of leaves of *Epidendrum* sp. in Brazil.

Atichia vanillae (Pat.) V. Hoeh. On leaves of *Vanilla planifolia* in Tahiti.

Bacillus cyripedii S. Hori. This bacterium causes the brown spot disease of orchids which is described as follows: Dirty-cinnamon or light-umber colored spots appear on the leaf-blades. The spots enlarge with great rapidity, so that within a few days entire green leaves can become discolored. After some days the diseased parts become brownish and finally deep chestnut-brown; their surfaces become more or less wrinkled, with loss of luster; their margins diffused, faint in color, and depressed. The lower surfaces of the leaves beneath the spots assume more or less rapidly a pale color and only gradually develop the same color as the upper surfaces. When the spots develop on the lower portions of a leaf, the upper part soon becomes yellowish and dies off without being directly attacked by the bacteria. The rotting also spreads downward into the stem, and if the diseased leaf is not cut off in time, the entire stock will soon be destroyed.

A similar orchid disease called the "brown spot" produces circular brown sunken spots on the upper leaf surfaces. As the spots enlarge the shape becomes somewhat irregular and the periphery more or less undulated, but it is well defined by the deeply sunken area from the surrounding healthy part. The color of the spots differs somewhat according to the different hosts; some spots assume a yellowish-brown, others a deep chestnut-brown color. Inoculation experiments demonstrated the two diseases to be identical, differences being due to varying texture of the leaves of the hosts. Brown rot occurred on succulent leafed species and brown spot on firm, tough-leaved species.

Hori states that the disease is greatly feared by gardeners, since it occurs on the most valuable orchids and spreads very rapidly during the summer season. The host species are: *Cypripedium godefroiae*, *C. raynaldianum*, *C. philippinense*, *Phalaenopsis amabilis*, *P. aphrodite*, and *P. schilleriana*. The disease has been found in Japan and Formosa on plants in hothouses as well as on plants collected in the wild.

Bacillus farnetianus Pavar. Pavarino has reported this disease as attacking the leaves and pseudobulbs of *Oncidium ornithorhynchum* and *Cattleya crispa* in the botanical garden, Pavia, Italy. The disease is characterized by small, indefinite areas showing as transparent spots which increase in size, becoming visible on both sides of the leaves as brown to black dry, depressed areas.

Bacillus kramerianii Pavar. This disease attacks *Oncidium kramerianum* in Italy. At first the diseased areas are transparent, later becoming irregular and finally brown and dry. Leaves and pseudobulbs are attacked.

Bacillus polacii Pavar. This disease causes irregular, black, depressed spots on leaves of *Odontoglossum citrosum* in Italy.

Bacterium briosianum Pavar. The leaves of *Vanilla planifolia* are said to be seriously attacked by this bacterium in the botanical garden at Pavia, Italy. Small, irregular, pitch-black spots are first seen which soon become lighter in color. The disease progresses rapidly, drying out and killing the leaves and in time involving the entire plant. This disease and the three preceding ones have all been reproduced by artificial inoculations, according to Pavarino.

Bacterium cattleyae Pavar. This disease is said to produce brown spots on the leaves and pseudobulbs of *Cattleya warneri* and *C. loddigesii* in the botanical garden of Pavia, Italy.

Bacterium dendrobii Pavar. Said to cause a withering of leaves and shoots of *Dendrobium nobile* in Italy.

Bacterium oncidii Peglion. This disease produces violet-brown spots on the leaves of *Oncidium* spp. in Italy. The entire leaf area is soon involved, the leaf losing its turgor and falling. Hori states that the causative organism should be known as *Bacillus oncidii* (Peg.) Hori, and that it is closely related to his *B. cyripedii*.

Black rot. A black rot of *Cattleya* leaves and bulbs has been reported from England. It is stated to be "A dreadful disease which carries off plants in a short time." Cause unknown.

Botryodiplodia digitata Maubl. On pseudobulbs of *Cattleya mossiae* in hothouses in France.

Botryospheria egenula Syd. and Burt. On leaves of *Cymbidium* sp., in Assam.

Calonectria copelandii P. Henn. On leaves of *Orchidaceae* in the Philippines.

Calospora vanillae Mass. Anthracnose on leaves of *Dendrobium* spp., *Oncidium* spp., and *Vanilla planifolia* in Madagascar, Tahiti, Colombia, West Indies, Seychelles, Reunion, and Mauritius. Said to be the same as *Gnomoniopsis vanillae* Stonem. on *Vanilla* in greenhouses in the United States.

Cercospora angrei Gaill. and Feuill. On leaves of *Angraecum fragrans*, *Cattleya* sp., and *Odontoglossum crispum* in Mauritius.

Cercospora epipactidis C. Mass. On leaves of *Epipactis palustris* in Italy.

Cercospora odontoglossi Prill. and Delacr. On the under sides of leaves of *Odontoglossum* spp., including *Odontoglossum crispum*, in greenhouses in France. This species is said to cause much damage.

Ceuthospora cattleyae Sacc. and Syd. (*C. minima* Delacr.) This fungus produces dark-brown spots on the leaves of *Cattleya intermedia* in Holland. The leaves ultimately die as a result of the fungus attack.

Chaetodiplodia vanillae A. Zimm. On rotting leaves of *Vanilla* sp. in Java. Probably not distinct from *Diplodia cacaoicola* P. Henn. See Theobroma.

Chytridiaceae (?). A leaf-spot disease of *Cattleya*, *Dendrobium*, *Odontoglossum*, and other cultivated orchids has been briefly described by Brierley. It is characterized by a series of concentric zones of green to pale-straw color and brown through purple to black. These vary in diameter from $\frac{1}{8}$ to $1\frac{1}{2}$ inches, occurring either as complete circles or segments of circles. The fungus has not yet been identified, but inoculations with diseased material readily reproduce the disease. It is thought that the fungus concerned belongs in the *Chytridiaceae*.

Ciliella epidendri (Rehm) Sacc. and Syd. On leaves of *Epidendrum* sp. in Brazil.

Cladochytrium olivieri Har. Dark leaf spots on *Orchis incarnata* and *O. laxiflora* in France.

Cladosporium sp. (*C. orchidis* Cke. and Mass. [?]). Forms olive-green blotches, which turn to a brownish-purple on living leaves of *Dendrobium* spp., *Coelogyne* spp., *Oncidium* spp., *Phalaenopsis* sp., and other cultivated orchids in England.

Coleosporium arundinae Syd. This rust produces orange-yellow pustules on the under sides of leaves of *Arundina chinensis* and *A. speciosa* in Java and Formosa.

Coleosporium bletiae Diet. Characterized by brown to golden-yellow rust pustules which fade out to white on the under sides of leaves of *Bletia hyacinthina* in Japan. The fungus has also been reported from one locality in central California on *Phajus wallichii* (*P. grandifolius*), it doubtless having been introduced from Japan.

Coleosporium merrillii P. Henn. Bright-yellow rust pustules on the under sides of leaves of *Spathoglottis chrysantha* and other undetermined orchids in the Philippines.

Colletotrichum dichaeae P. Henn. On leaves of cultivated *Dichaea vaginata* in the botanical Garden at Berlin.

ORCHIDACEAE—Continued.

Colletotrichum effiguratum Syd. Irregular, yellow-brown leaf spots with distinct purple-brown margins on *Paphiopedilum roezlii* in Germany.

Colletotrichum macrosporum Sacc. On *Orchidaceae* in Brazil.

Colletotrichum orchidearum Allesch. On leaves of *Bulbophyllum lobbiti*, *B. longiflorum*, *Cattleya eldorado*, *Cymbidium* sp., *Coelogyne mayeriana*, *Epidendrum macrostachyum*, *Eria ornata*, *E. stellata*, *Eulophia saundersiana*, *Laelia crispata*, *Oncidium pulvinatum*, *Physisiphon loddigesii*, *Pholidota imbricata*, *Pleurothallis tribuloides*, *Spathoglottis plicata*, and *Surcanthus puyoniiformis* in the botanical garden at Munich, as well as in Colombia, Brazil, and the Philippines on native plants.

Colletotrichum orthianum A. Kostlan. On leaves and leaf sheaths of *Cattleya* sp., *Coelogyne cristata*, *Cypripedium insigne*, *Dendrobium* sp., *Dendrobium nobile*, and *Vanda coerulesa* in the botanical garden at Berlin.

Colletotrichum roseolum P. Henn. On pseudobulbs of *Stanhopea oculata* in the botanical garden at Berlin.

Colletotrichum vanillae Scalia. Forms brown spots on leaves of *Vanilla odorata* and *V. planifolia* in Sicily and Ceylon.

Colletotrichum vinosum P. Henn. On pseudobulbs of *Stanhopea oculata* in the botanical garden at Berlin.

Cylindrina delavayi Pat. On leaves of *Liparis liliifolia* in China.

Diplodia dendrobii Cooke and Mass. On leaves of *Dendrobium speciosum* in Australia.

Diplodia henriquesiana Trav. and Spessa. On pseudobulbs of *Cattleya labiata* in Italy.

Diplodia sobraliae (P. Henn.) Taub. On leaves of *Sobralia sessilis* in the botanical garden at Berlin.

Fusarium sp. Attacks the bases of leaves of *Vanilla planifolia* in Brazil, extending into the stems and forming small, clear, irregular brown spots, which finally become confluent.

Fusicadum vanillae A. Zimm. Appears as a brown fungus layer growing over living leaves of *Vanilla* sp. in Java.

Gloeosporium affine Sacc. This fungus produces white spots on both surfaces of leaves and pseudobulbs of *Bulbophyllum lobbiti*, *Cattleya mendelii*, *Coelogyne cristata*, *Maxdevalia psittacina*, *Odontoglossum crispum*, *Pleurothallis lansbergii*, *Vanilla* sp., and other cultivated orchids in Italy and Germany.

Gloeosporium beyrodtii Klitz. Produces dark-brown blotches which enlarge rapidly, soon causing the death of leaves of *Vanda coerulesa* in Germany.

Gloeosporium bidgoodii Cke. On *Odontoglossum* spp. and *Oncidium* sp. in hothouses in England.

Gloeosporium bussei P. Henn. On leaves of *Vanilla* sp. in the botanical garden at Berlin.

Gloeosporium cattleyae Sacc. On leaves of *Cattleya mossiae* in France.

Gloeosporium coelogyne Syd. Forms pale spots on leaves of *Coelogyne viscosa* in the botanical garden at Berlin.

Gloeosporium dendrobii Maubl. Characterized by large, pale, ochre-yellow spots on leaves of *Dendrobium farmerii* in France.

Gloeosporium epidendri P. Henn. Large black spots on flower stems of *Epidendrum* sp. in the botanical garden at Berlin and on *Epidendrum ciliaris* in Spain.

Gloeosporium intermedium Sacc. var **epidendri** Sacc. On leaves of *Orchidaceae* including *Epidendrum* spp., in France.

Gloeosporium laeliae P. Henn. Yellow-brown spots covering entire leaf surfaces of leaves of *Laelia* sp. in the botanical garden at Berlin.

Gloeosporium maxillariae Allesch. On leaves of *Maxillaria rufescens* in the botanical garden at Berlin.

Gloeosporium noackianum Allesch. On leaves of *Orchidaceae* in Brazil.

Gloeosporium oncidii Oud. Produces brown spots on the leaves and leaf-sheaths of *Maxillaria infestans*, *M. rufescens*, and *Oncidium ansiferum* in Holland and Germany, and on *Oncidium sphacelatum* in Mexico.

Gloeosporium ornithidii Allesch. Produces large, irregular spots which are ochre-colored and often confluent, on leaves of *Ornithidium densum* in greenhouses in Germany.

Gloeosporium pallidum (Karst.) Hariot. f. **physosiphonis-loddigesii** Allesch. On leaves of *Physisiphon loddigesii* and *Liparis longipes* in greenhouses in Germany.

Gloeosporium phaji Maubl. Produces round spots, which are white at first, becoming black and finally whitish at the centers, on living leaves of *Phajus* spp. and *Phajus wallichii* in greenhouses in France.

Gloeosporium sobraliae Maubl. Forms spots near the tips of the leaves, with a brown-black line of demarcation between healthy and diseased tissue, on *Sobralia* sp. in greenhouses in France.

Gloeosporium stanhopaeae Allesch. On leaves of *Stanhopea* sp. in greenhouses in Germany.

Gloeosporium stanhopicolum P. Henn. On leaves of *Stanhopea* sp. in the botanical garden at Berlin.

Gloeosporium vandopsisidis Keissler. On leaves of *Vandopsis* sp. in the Solomon Islands.

Gloeosporium vanillae Cooke and Mass. On leaves of *Vanilla planifolia* in Colombia, Mauritius, Ceylon, and other parts of the world and on *Vanda sanderiana* in the Philippines.

Guignardia traversii Sacc. On *Cattleya* sp. in France.

Guignardia majori (Car.) Lind. On leaves of *Vanilla planifolia* in Italy.

Hemileia americana Mass. This rust has been reported on the under sides of leaves of *Cattleya dowiana* and *Oncidium cavendishianum* in greenhouses in England, the plants having come originally from Central America. Powdery golden-yellow pustules are produced. The form on *O. cavendishianum* should be referred to *Hemileia oncidii* Griff. and Maubl.

Hemileia oncidii Griff. and Maubl. (*Uredo behnickiana* P. Henn.) This species produces bright-yellow pustules on the lower leaf surfaces of *Epidendrum vitellinum*, *Lycaste skinneri*, *Oncidium crispum*, *O. dasystachy*, *O. forbesii*, *O. marshallianum*, and *O. varicosum* in greenhouses in Brazil, France, Ireland, and Germany.

Hemileia phaji Syd. Characterized by yellow rust pustules on the lower surfaces of leaves of *Phajus* spp. including *P. blumei* and *P. wallichii* in Java. Grove states that these fungi (*Hemileia* spp.) might easily become dangerous parasites if allowed to spread.

Hendersonia epidendri Keissler. On leaves of *Epidendrum bifidum* in the island of St. Croix.

Hypodermium orchidearum Cke. and Mass. This fungus produces spots in groups which are often 1 to 2 inches long, causing the leaves of *Cymbidium eburneum* in Great Britain to turn yellow and die.

Leptostroma orchidearum Mntg. On living leaves of *Cypripedium macranthum* in Siberia.

Leucothyridium (?) **vanillae** Avena. Irregular black fungus crusts on leaves and stems of *Vanilla planifolia* in Brazil.

Macrophoma cattleyicola P. Henn. Characterized by numerous large, brown-black spots on pseudobulbs of *Cattleya labiata* and *C. peruviana* in the botanical garden at Berlin.

Macrophoma cattleyicola P. Henn. var. **brassavolae** Keissler. Forms spots on living leaves of *Brassavola* sp. at the Kew gardens, England.

ORCHIDACEAE—Continued.

Macrophoma miltoniae DaCam. On leaves of *Miltonia candida* in the botanical garden, Coimbra, Portugal.

Macrophoma oncidii P. Henn. Said to cause the death of leaves of *Oncidium pulvinatum* in the botanical garden at Berlin. Also reported on leaves of *O. sphecelatum* and *Cattleya* sp. from Mexico.

Macrophoma vanillae Avena. Large irregular, then confluent, ashen leaf spots with dark margins on *Vanilla planifolia* in Brazil.

Melampsora repentis Plowr. (*Caeoma orchidis* Wint.) This rust produces pale yellow spots 1 to 2 centimeters long and generally somewhat elongated, on the under sides of the leaves of *Gymnadenia conopsea*, *G. alida*, *Habenaria bifolia*, *H. chlorantha*, *H. conopsea*, *H. solstitialis*, *Listera ovata*, *Ophrys arandifera*, *O. muscifera*, *Orchis ambustifolia*, *O. bifolia*, *O. incarnata*, *O. latifolia*, *O. maculata*, *O. mascula*, *O. militaris*, *O. palustris*, *O. sambucina*, *O. undulata*, and *Phalaenopsis esmeralda* in British Guiana, Soain, Austria, Denmark, Great Britain, Hungary, Russia, Sweden, Switzerland, and Germany. The urdo and telial stages occur on *Salix repens*, *S. aurita* and *S. triandra*.

Me anospora coemansii West. On leaves of *Oncidium* sp. in Belgium.

Meioida sp. On living leaves of *Pleurothallis ruscifolia* in Porto Rico and *Laelia alba* in Mexico.

Myriocopron (?) **vanillae** Avena. Black fruiting bodies on extensive confluent ashen leaf spots on *Vanilla planifolia* in Brazil.

Nectria bulbicola P. Henn. On bulbs of *Cyrtopodium* sp., *Gomezia planifolia*, *Gongora galeata*, *Maxillaria rufescens*, *M. variabilis*, *Oncidium pulvinatum*, *Polystachya* sp., and *Sophonitis cernua* in the botanical garden at Berlin. The species has been referred to *N. ochroleuca* (Schw.) Bark., a common tropical species.

Nectria bulbophylli P. Henn. On *Bulbophyllum lobtii* in the botanical garden at Berlin.

Nectria goshankiniana Wahrlich. On roots of *Vanda tricolor* in orchid houses in Russia.

Nectria vandae Wahrlich. On roots of *Vanda suavis* in orchid houses in Russia.

Nectria vanillae A. Zimm. Brown spots on stems and leaves of cultivated *Vanilla* in Java.

Nectria vanillicola P. Henn. On leaves of *Vanilla aromatica* in Java.

Pestalozzia vanillae Avena. Irregular, often marginal, brown leaf spots on *Vanilla planifolia* in Brazil.

Phoma obtrudens F. Tassi. On leaves of *Phalaenopsis longifolia* in Italy.

Phoma oncidii Speg. (*Phoma corrientina* Speg.) Forms indefinite white spots on leaves of *Oncidium* sp. in Argentina and *O. papilio* in hothouses in Denmark.

Phoma oncidii-sphacelati F. Tassi. On living leaves of *Oncidium sphacelatum* in Mexico.

Phyllosticta bletiae A. Zimm. On living leaves of cultivated *Bletia hyacinthina* in Austria.

Phyllosticta decidua Ferraris. Characterized by pale ochre-colored spots, circular in shape, with brown margins, on leaves of *Goodyeria repens* in Italy.

Phyllosticta donckelaarii Westendorp. Forms large grayish spots with red-brown margins on the under sides of leaves of *Oncidium* sp. in hothouses in Belgium.

Phyllosticta epipactidis Diet. Produces dark-brown spots, becoming lighter at the centers, 1 to 1½ centimeters in diameter, on leaves of *Epipactis violacea* in Germany.

Phyllosticta laeliae Keissler. The fungus produces light-colored spots on the leaves and leaf sheaths of *Laelia alba* and *L. furfuracea* in Mexico.

Phyllosticta masdevalliae P. Henn. On leaves of *Masdevallia chimaera* in the botanical garden at Berlin.

Phyllosticta nigramaculans Sacc. On leaves of *Orchidaceae* in Brazil.

Phyllosticta pleurothallidis Keissler. On living leaves of *Pleurothallis longissima* in Costa Rica.

Phyllosticta renantherae Keissler. Forms indistinct, light-colored, subcircular spots on living leaves of *Renanthera storied* in the Philippine Islands.

Phyllosticta stanhopeae Allesch. The fungus does not form definite spots but attacks the whole leaf, which soon dies. The hosts are *Stanhopea* spp. in Germany.

Phyllosticta valparadisiaca Speg. Forms circular brown spots 1 to 3 millimeters in diameter, on both sides of leaves of undetermined orchids from Chile.

Phyllosticta vanillae P. Henn. Produces large pale-colored spots on leaves of *Vanilla aromatica* in Java.

Physalospora campyospora Sacc. On leaves of *Orchidaceae* in Brazil.

Physalospora cattleyae Maubl. and Lasn. On *Cattleya mossiae* in greenhouses in France.

Physalospora orchidearum P. Henn. On stems of *Laelia schilleriana* and *Tainia stellata* in the botanical gardens at Berlin.

Physalospora vanillae A. Zimm. On leaves of *Vanilla planifolia* in Java.

Physalospora wildemaniana Sacc. On leaves of *Orchidaceae* in Brazil.

Phytophthora sp. A very serious disease of the cultivated vanilla (*Vanilla planifolia*) due to a *Phytophthora* species or related fungus is known to occur in Seychelles and various South Sea Islands. There are no published records of the disease and no studies have been made of it.

Phytophthora faberi Maubl. This fungus, which causes one of the most serious diseases known, the black rot, and pod canker of rubber (*Hevea brasiliensis*), cacao (*Theobroma cacao*) and other economic plants, has been reported as attacking *Dendrobium naccarthiae* in Ceylon.

Puccinia aurea Wint. Produces yellow to brown rust pustules on both leaf surfaces of *Monadenia rufescens* in south Africa.

Puccinia cinnamomea Diet. and Holw. Produces scattered cinnamon-colored rust pustules on the lower leaf surfaces of unknown species of *Orchidaceae* in Mexico.

Puccinia habenariae P. Henn. Produces brown rust pustules on both sides of the leaves and on stems of *Habenaria* spp. in east Africa.

Puccinia orchidearum-phalaridis Kleb. Characterized by yellow, then brown, rust pustules on the lower surfaces of leaves of *Gymnadenia conopsea*, *Habenaria bifolia*, *H. chlorantha*, *H. montana*, *Listera* sp., *L. ovata*, *Orchis incarnata*, *O. latifolia*, *O. maculata*, *O. majalis*, *O. mascula*, *O. militaris*, *O. morio*, *O. purpurea*, and *Phalaris arundinacea* in Denmark, France, Great Britain, Holland, Switzerland, Austria, and Germany.

Puccinia satyrii Syd. This rust forms yellow-brown rust pustules on the under sides of the leaves of *Satyrion carneum* in south Africa.

Sclerotium orchidearum P. Henn. Forms globose sclerotia on the stems 0.8 to 1.2 millimeter in diameter, white at first, then yellow, and finally brown-black, smooth, white within. Found on *Dichaea vaginata* and *Vanda tricolor* in the botanical gardens at Berlin.

Septoria codonorchidis P. Henn. On leaves of *Codonorchis poeppigii* in Chile.

Septoria epipactidis Sacc. Characterized by long light-colored spots on the leaves of *Epipactis violacea* and *Epipactis* sp. in Italy and Germany.

Septoria gymnadeniae Thuem. On leaves of *Gymnadenia conopsea* in Siberia.

Septoria listerae Allesch. On leaves of *Listera ovata* in Germany.

Septoria orchidearum West. Produces brown spots on leaves and leaf sheaths of *Cocloglossum viride*, *Habenaria bifolia*, *Listera ovata*, *Orchis latifolia*, *O. maculata*, *O. mascula*, and *O. morio* in Belgium, France, and Italy.

ORCHIDACEAE—Continued.

- Septoria posekensis** Sacc. On leaves of Orchidaceae in Siberia.
- Septoria thelymitrae** McAlp. On leaves of *Thelymitra aristata* in Australia.
- Stigmatodopsis palawanensis** Syd. On living leaves of *Dendrobium* sp. in the Philippines.
- Stibella bulbicola** P. Henn. On leaf bases of *Epidendrum* sp., *Gomeza planifolia*, *Oncidium pulvinatum*, *Sorcanthus pugioniformis*, and *Stanhopea* sp. in the botanical gardens at Berlin.
- Tubercularia** sp. Chlorotic or livid irregular areas on stems of *Vanilla planifolia* in Brazil.
- Tylenchus olesistus** Ritz-Bos. See *Begonia*.
- Uredo aurantiaca** Mortem. Yellow rust pustules on the leaves of *Oncidium cavendishianum* in greenhouses in France and Italy.
- Uredo carnosa** Speg. Rust on *Catasetum fimbriatum* in Brazil.
- Uredo cyrtopodii** Syd. Brown rust pustules on the under surfaces of the leaves of *Cyrtopodium* sp. and *Bletia* sp. in Brazil, Cuba, and Germany; in the latter case on imported plants.
- Uredo epidendri** P. Henn. Brown rust pustules on leaves of *Epidendrum* sp. in Brazil.
- Uredo guacae** Mayor. On *Epidendrum rigidum* and *E. umbellatum* in Ecuador.
- Uredo gynandrearum** Cda. Rust on leaves of *Habenaria maculosa* and other Orchidaceae in Cuba, Porto Rico, Trinidad, and Central and South America.
- Uredo lynckii** Berk. Rust on tropical species of *Spiranthes* growing in greenhouses in England.
- Uredo nigropunctata** P. Henn. Forms yellow rust pustules on the under sides of leaves of *Bletia patula* and *B. purpurea* in Cuba, Porto Rico, Haiti, Bahamas, South America, and Florida and on the leaves of *Stanhopea* spp. in Brazil.
- Uredo oncidii** P. Henn. Rust on leaves of *Oncidium lanceanum* in Brazil.
- Uredo phaji** Racib. Forms golden-yellow rust pustules on the lower sides of leaves of *Phaius blumei* in Java. This species is probably identical with *Hemileia phaji* Rac.
- Uredo pleurothallidis** Keissl. Rust found on plants of *Pleurothallis dinotherii* in greenhouses in Germany.
- Uredo pustulata** P. Henn. Characterized by brown rust pustules on both surfaces of the leaves of *Stenorrhynchus* spp., including *S. lanceolatus* in Brazil.
- Uredo satyrii** Mass. Rust on *Satyrium corifolium* in the Union of South Africa.
- Uredo scabies** Cooke. Rust on small, irregular spots on both sides of the leaves of *Vanilla planifolia* in Colombia.
- Uredo wittmackiana** P. Henn. and Klitz. Rust on round yellow or brownish spots on the upper surfaces of leaves of *Epidendrum* sp. in Mexico.
- Uromyces citrifolius** Bab. (*U. thelymitrae* McAlp. [?]) Rust on leaves of *Microstylis* sp. or *Thelymitra* sp. in New Zealand.
- Uromyces joffrini** Delacr. Reported on the fruits of *Vanilla planifolia* from Tahiti. A doubtful species.
- Uromyces microtidis** Cooke. Produces irregular groups of yellowish-brown rust pustules on living leaves of *Microstylis perfoliata* and *M. pulchella* in New South Wales, Tasmania, and Chatham Island.
- Uromyces orchidearum** Cke. and Mass. This rust is characterized by golden-brown rust pustules, mostly on the upper surfaces of the leaves (æcia hypophyllous), of *Chiloglottis diphylla* and *C. gunnii* in New South Wales, Victoria, and Tasmania.
- Uromyces stenorrhynchi** P. Henn. Brown rust pustules on the lower surfaces of the leaves of *Stenorrhynchus* sp. in Peru.
- Uromyces thelymitrae** McAlp. The golden-yellow to chestnut-brown pustules are produced on the leaves and leaf sheaths, either in scattered groups or gregarious, on *Thelymitra aristata*, *T. antennifera*, *T. crinita*, and *T. flexuosa* in Victoria, Australia, and *T. javanica* in Java.
- ORCHIS**. Terrestrial orchids. See Orchidaceae.
- ORIGANUM**. MARJORAM. Sweet herbs cultivated for their ornamental flowers and foliage.
- Phyllosticta origani** P. Tassi. Small circular to irregular leaf spots on *O. heracleoticum* in Italy.
- Puccinia caulicola** Schneid. See *Thymus*.
- Puccinia rubsaameni** P. Magn. Rust on leaves of *O. vulgare* in Denmark, Switzerland, Germany, and Austria.
- Septoria origanicola** Allesch. and var *majoranae* Bres. Large, irregular, often confluent, brown leaf spots on *O. majorana* and *O. vulgare* in Germany.
- ERMOSIA**. NECKLACE TREE. Tropical trees.
- Puccinia ormosiae** Arth. Rust on leaves of *O. krugii* in Porto Rico.
- ORNITHIDIUM**. Epiphytic orchids. See Orchidaceae.
- ORNITHOGALUM**. STAR-OF-BETHELEHEM. Dwarf, hardy bulbous plants.
- Aecidium ornithogalum** Bub. Yellow rust pustules on the leaves of *O. tenuifolium* in Moravia. Possibly the same as *Puccinia lilacearum*.
- Heterosporium ornithogali** Klotzsch. On leaves of *Muscari comosum*, *M. neglectum*, *Ornithogalum* sp., and *Gagea* sp. in Great Britain, Yugoslavia, and Germany. Reported from Washington.
- Puccinia kalebrenneriana** De Toni. Yellow rust pustules turning to brown on both leaf surfaces of *O. altissimum* in South Africa.
- Puccinia lilacearum** Duby. Rufous-brown rust pustules on the leaves of *Bellevalia romana*, *Endymion nutans*, *Hyacinthus ramosus*, *Muscari comosum*, *M. racemosum*, *Ornithogalum narbonneense*, *O. nutans*, *O. pyrenaicum*, *O. tenuifolium*, *O. umbellatum*, and *Scilla bifolia* in Great Britain, France, Russia, Italy, Belgium, Holland, Hungary, Austria, and Germany.
- Puccinia iohkaiana** Thiem. Black linear rust pustules on both leaf surfaces of *O. boucheanum*, *O. chloranthum*, *O. nutans*, *O. pyrenaicum*, *O. refractum*, *O. prasadum*, *O. umbellatum*, *Muscari botryoides*, and *M. racemosum* in France, Italy, Hungary, Austria, India, and Germany.
- Puccinia ornithogali-thyrsoideis** Diet. Brown rust pustules on stems and leaves of *O. thyrsoideis* in the Union of South Africa.
- Septoria ornithogalea** Oud. Pale linear spots on stems and capsules of *O. pyrenaicum* and *O. umbellatum* in Holland.
- Septoria ornithogali** Pass. On leaves of *O. umbellatum* in Italy.
- Septoria ornithogalicola** Höll. Large yellowish spots with indefinite margins on both surfaces of the leaves of *O. boucheanum* in Hungary.
- Synchytrium niessli** Bub. Small galls, which are dirty-white in color with a brown band, are produced on the leaves of *O. umbellatum* in Austria.
- Urocystis ornithogali** Koern. Black powdery smut sori on the leaves of *O. umbellatum* in Germany. This species is said not to be distinct from *U. colchici*, which has been reported sparingly on other hosts in the United States.
- Uromyces maireanus** Syd. This rust forms yellow-brown pustules, which later become dark brown, on irregular pale spots on both leaf surfaces of *O. sessiliflorum* in Algeria.
- Uromyces ornithogali** Lév. See *Gagea*.
- Ustilago peglerae** Bub. A smut destroying the anthers of *O. lacteum* in the Union of South Africa.
- Uromyces scillarum** (Grev.) Wint. See *Hyacinthus*.

ORNITHOPUS. Leguminous plants, one species cultivated for forage.

Peronospora ornithopi Gäum. Downy mildew on leaves of *O. perpusillum* in Denmark and Germany.

Uromyces ornithopodioides Gz. Frag. Rust on leaves of *O. ichomocarpus* in north Africa.

OROBUS. See Lathyrus.

OROXYLUM. EAST INDIAN TRUMPET FLOWER. Ornamental Asiatic trees.

Phyllosticta oroxylonis P. Henn. On leaves of *O. indicum* in Brazil.

ORTHOSIPHON. Annual or perennial herbs.

Puccinia pallida Syd. Rust on leaves of *Orthosiphon* sp. in the Union of South Africa.

ORYZA. RICE. Annual grasses cultivated for grain.

Achlya prolifer (Nees) De By. Seeds of *O. sativa* in nursery beds are covered by a cottony mycelium and rotted. The disease may also attack young shoots and roots of seedlings; in Formosa.

Asecochya oryzae Catt. On leaves of *O. sativa* in Italy, Japan, and Brazil.

Cercospora oryzae Miy. Small brown linear spots on leaves and glumes of *O. sativa* in Japan, China, Brazil, Porto Rico, and Cuba. It is not certain that the western form is the same as the oriental. *Cercospora* sp. common throughout Louisiana may be the same.

Chaetophoma glumarum Miy. Whitish areas on glumes of *O. sativa*, hindering formation of grain in Japan.

Chaetophoma oryzae Cav. On sheaths, peduncles, and glumes of *O. sativa* in Italy.

Coniothyrium anomele Miy. White spots with dark-brown borders, mostly at tips and along margins of leaves of *O. sativa* in Japan.

Coniothyrium brevisporum Miy. Irregular white spots finally involving entire leaf area of *O. sativa* in Japan and China.

Coniothyrium japonicum Miy. Brown spots, becoming whitish, commonly at tips and along margins of leaves of *O. sativa* in Japan.

Diplodia oryzae Miy. On leaves and glumes of *O. sativa* in Japan.

Diplodella oryzae Miy. On leaves and glumes of *O. sativa* in Japan.

Entyloma oryzae Syd. Leaf smut on *O. sativa* in the Philippines.

Gnomonia oryzae Miy. On glumes of *O. sativa* in Japan.

Helminthosporium oryzae v. B. de Haan. The "sesame" spot disease or seedling blight is one of the most serious diseases reported on rice, 10 to 60 per cent of the seedlings often being destroyed. It is known to occur in Japan, Formosa, Java, India, and the Philippines. A similar, if not identical, disease is known from Portuguese East India, Federated Malay States, Italy, and restricted areas in Louisiana. The disease is most destructive to seedlings, but also attacks leaves, culms, and panicles of older plants. In the case of seedlings, the tips of the cotyledons become dark brown and the fungus spreads to new leaves. Numerous small brown spots edged with yellow occur on the leaves. These spots coalesce, becoming irregular, and as a result the leaves die. Diseased culms are yellow or pale brown and the heads either can not push out at all or are bent and distorted. Brown or grayish spots may also occur on the various parts of the head. The disease has been transferred by inoculation to a number of other economic grasses. It has been reported from a restricted area in Louisiana, but it will be desirable to prevent further introductions, especially since other and more virulent strains may gain entrance.

Helminthosporium sigmoideum Cav. Effuse black areas on sheaths, leaves, and stems of *O. sativa* in Italy.

Hendersonia oryzae Miy. Brown spots on glumes and leaves of *O. sativa* in Japan. This disease is said to prevent the development of the grain.

Leptosphaeria iwamotoi Miy. On leaves of *O. sativa* in Japan, Chosen, and Uganda.

Leptosphaeria salvinii Catt. On leaf sheaths of *O. sativa* in Italy.

Melanomma glumarum Miy. Brown, then paler, spots on glumes and culms of *O. sativa* in Japan, China, India, and the Philippines. Causes abortion of the grain.

Metasphaeria albescens Thuem. Small brown-black spots on the leaves of *O. sativa*, increasing rapidly in size and becoming yellowish-brown, in Japan.

Metasphaeria oryzae (Catt.) Sacc. On leaves of *O. sativa* in Italy.

Mosaic. The mosaic disease of sugar-cane has been transmitted to rice experimentally, and might become of importance under favorable conditions.

Mycosphaerella hondai Miy. White spots on leaves of *O. sativa* in Japan and Chosen.

Mycosphaerella malinverniana Catt. On leaves of *O. sativa* in Italy and India.

Mycosphaerella oryzae (Catt.) Sacc. This fungus attacks the leaves of *O. sativa*, checking the development of the grain. It occurs in Japan, Italy, Portugal, and Austria.

Mycosphaerella shiraiana Miy. Gray to black spots on both leaf surfaces and occasionally on glumes of *O. sativa* in Japan and China.

Napcladium jenseanum Rac. Linear red-brown spots on leaves of *O. sativa* in Java.

Nigrospora panici Zimm. See Triticum.

Oospora oryzetorum Sacc. This disease attacks the heads when they are near maturity, a whitish bloom spreading over the panicles. Diseased heads are largely blasted or destroyed. On *O. sativa* in the Philippines.

Ophiobolus cariceti (B. and Br.) Sacc. (*Ophiochaeta graminis* Hara). The take-all disease which attacks wheat and other cereals so seriously in many parts of the world has been reported on rice (*O. sativa*) in Japan and Italy. See Triticum.

Ophiobolus oryzae Miy. On leaves and glumes of *O. sativa* in Japan.

Phaeosphaeria cattanii (Thuem.) Miy. On leaves and stems of *O. sativa* in Italy.

Phaeosphaeria oryzae Miy. Brown areas beginning at the margins and tips of leaves and finally involving entire leaf area. Also on glumes of *O. sativa* in Japan.

Phyllosticta japonica Miy. Dark-brown, then whitish, spots on both sides of leaves and on glumes of *O. sativa* in Japan. This disease finally destroys infected leaves.

Phyllosticta niurai Miy. Large areas on leaves become white and the leaves die. On *O. sativa* in Japan and the Philippines.

Phyllosticta necatrix Thuem. On culms, leaves, and leaf sheaths of *O. sativa* in Italy.

Phyllosticta oryzae (Cke. and Mass.) Miy. Dark-brown irregular spots on leaf sheaths in Formosa, Japan, and India.

Pleosphaerulina oryzae Miy. Large areas on leaves of *O. sativa* whitened, in Japan.

Pseudomonas oryzae Ishiyama. This bacterium attacks the leaves, causing the "white speck" disease of *O. sativa* in Japan. The disease also attacks the stems and causes losses up to 20 per cent.

Puccinia graminis Pers. forma **oryzae** Küss. A rust on rice (*O. sativa*) has been reported from Spain and other parts of Europe and named as above. A rust referred to as *Puccinia oryzae* has also been reported, but is doubtless the same. The rust involved here is a strain of the common and widespread black stem rust of cereals. This strain is unknown in the United States.

Pyrenochaeta oryzae Shir. Diseased areas occur on the leaves and glumes of *O. sativa* in Japan. The leaves are greatly weakened and the plants stunted, since root development is poor.

Rhizoctonia destruens Tass. See Solanum.

ORYZA—Continued.

Sclerospora sp. An undetermined species of *Sclerospora* has been reported on *O. sativa* in Japan. The species involved may possibly be *S. macrospora* which occurs to a limited extent in the United States, but this is probably not the case and this downy mildew should be as carefully guarded against as the similar oriental diseases on corn and sugar cane.

Sclerospora macrospora Sacc. See *Triticum*.

Sclerotium sp. A serious stem-rot disease due to an undetermined species of *Sclerotium* is reported from Java, Japan, China, and the Philippines. From India a similar disease is reported, attacking the leaf-sheaths and causing light-brown areas which ultimately spread over and kill infected plants. These diseases may be referable to one of the following species.

Sclerotium glumale Ces. Gray to black spots on glumes of *O. sativa* in Borneo.

Sclerotium irregulare Miy. Irregular to globular or oval rust-brown sclerotia produced on greenish-gray spots with dark brown margins on leaf sheaths of *O. sativa* in Japan.

Sclerotium oryzae Catt. The bases of stems of diseased plants are discolored at the lowest internodes, with dark-gray mycelium present in the hollow stems, in which appear numerous small, round, black, shining sclerotia. The disease may begin at a node some distance up the stem, involving the leaf sheaths at the same time. Diseased plants are stunted, the culms lodge and little or no grain is formed. This is said to be the most serious of known rice diseases. It occurs in Japan, Italy, India, Ceylon, Portugal, British Guiana, the Philippines, and to a very limited extent in Louisiana and Arkansas. The disease is carried for the most part on seed grain and gained entrance to the United States in this manner. Prohibition of further importations should be effective in preventing continued spread of the disease, provided proper steps are taken to eliminate present centers of infection.

Septoria curvula Miy. On leaves of *O. sativa* in Japan.

Septoria longispora Miy. Gray areas on leaves of *O. sativa* in Japan. Often sufficiently serious to check development of grain.

Septoria miyake Sacc. and Trav. On glumes of *O. sativa* in Japan and the Philippines.

Septoria oryzae Catt. On leaves, leaf sheaths, and glumes of *O. sativa* in Japan, China, Brazil, and Italy.

Septoria poae Catt. On leaves of *O. sativa* in Brazil.

Sphaeronema oryzae Miy. Produces white spots on glumes of *O. sativa* in Japan.

Sphaeropsis japonicum Miy. On glumes of *O. sativa* in Japan.

Sphaeropsis oryzae (Catt.) Sacc. On leaves and leaf sheaths of *O. sativa* in Italy and Brazil.

Tilletia horrida Tak. This smut forms black spore masses in the ovaries, generally concealed by the glumes, but at times forcing them apart. The smut is usually not visible until the grain is threshed. The disease occurs in Burma, India, British Guiana, Indo-China, Siam, Japan, China, and the Dutch Indies. It has been introduced into the United States on seed rice and was at one time serious in the Carolinas, but is said to have been eradicated. It also occurs sparingly in other Southern States, but not in California.

Tylenchus angustus Butl. The "ufra" disease of rice, known only from India, is caused by a nematode and is considered one of the most serious diseases of this crop. The worm attacks the aboveground parts of the plant. The ends of the leaves turn brown and this browning continues until the entire plant is involved. Plants are often stunted, with either withered or normal appearing leaves. Brown stains or lesions appear at one or more of the upper nodes just below the base of the peduncles, the stalks of which become deep brown or black, flaccid, and shrunken. The upper part of the stalk may be swollen into a spindle-shaped swelling composed of the ear and inclosing sheaths or the ear may push its way out. Normal grains as a rule are not produced. The nematodes may be carried on seed grain, or on empty glumes and other diseased plant parts.

Ustilagioides vires (Cke.) Tak. The so-called green smut or false smut disease of rice attacks the individual grains which are transformed into hard green masses, more or less twice the size of normal grains. These sclerotia are orange-yellow near the surface and white at the center. The disease occurs in India, Ceylon, Japan, Formosa, China, Sumatra, Malacca, Indo-China, Portuguese East India, Philippines, and Straits Settlements. It has been sparingly introduced into the southern United States, but is not known in California.

ORYZOPSIS. Tufted perennial grasses.

Puccinia oryzopidis Syd. and Butl. Yellow and dark-brown rust pustules on leaves of *O. molinoides* in India.

Septoria trapezuntica Bub. On leaves of *O. miliacea* in Asia Minor.

OSBECKIA. Tropical herbs and shrubs.

Rehmiodothis osbeckiae (B. and Br.) Theiss. and Syd. Shining black convex stromata on leaves of *O. virgata* in Ceylon.

OSMANTHUS. Ornamental woody plants grown for their handsome foliage and white, fragrant flowers.

Accidium osmanthi Syd. and Butl. Rust on leaves of *Osmanthus* sp. in India.

Phyllosticta osmanthi Tass. White leaf spots on *O. aquifolius* in Italy.

Phyllosticta osmanthicola Trinch. On leaves of *O. fragrans* in Italy.

Septoria osmanthi Tass. Irregular white spots on leaves of *O. aquifolius* in Italy.

OSMORHIZA. Aromatic herbs with edible roots.

Puccinia philippii Diet. and Neg. Yellow and brown rust sori on leaf blades and petioles of *O. berterii* and *O. glabrata* in Chile and Argentina.

OSMUNDA. Royal fern.

Mycosphaerella collista Syd. On fronds of *O. regalis* in Germany.

OSTEOSPERMUM. Yellow-flowered shrubs.

Ascochyia atro-punctata Wint. Circular, often confluent, yellow leaf spots on *O. moniliferum* in the Union of South Africa.

OSTRYA. HOP HORNBEEAM. Ornamental and timber trees.

Exoascus ostryae Mass. Brown spots on leaves of *O. carpinifolia* in Italy.

Gloeosporium ostryae Thuem. Irregular dark-brown anthracnose spots on leaves of *O. virginiana* in Portugal.

Gnomonia ostryae DeN. On leaves of *O. carpinifolia* in Italy.

Mamiania ostryae (P. Henn.) Theiss. and Syd. On *O. japonica* in Japan.

OURISIA. Small herbs used in border plantings.

Puccinia apus Diet. and Neg. Brown rust sori on leaves of *O. racemosa* in Chile.

Puccinia ourisiae Diet. and Neg. Rust on leaves of *O. alpina*, *O. coccinea*, and *O. pygmaea* in Chile.

Puccinia quillensis Diet. and Neg. Powdery rust pustules on leaves of *Ourisia* sp. in Argentina.

OXALIS. Bulbous or tuberous herbs with cloverlike leaves; a few species cultivated for their edible roots or tubers.

Cercospora oxalidis Grove. Small circular leaf spots on *O. acetosella* in Great Britain.

Cylindrosporium oxalidis Traill. Circular brown leaf spots on *O. acetosella* in Scotland.

OXALIS—Continued.

Mycosphaerella depazeaeformis (Ces. and DeN.) Wakef. Circular, rufous leaf spots on *O. acetosella*, *O. corniculata*, and *O. stricta* in Uganda and Germany.

Ovularia oxalidis Oud. On leaves of *O. tropaeoloides* in Holland.

Phyllosticta oxalidicola Speg. On leaves of *Oxalis* sp. in Argentina. A doubtful species.

Phyllosticta oxalidicola P. Henn. Yellow-brown leaf spots on *Oxalis* sp. in Jamaica.

Puccinia oxalidis Diet. and E. Brown rust pustules on leaves of *O. newwiedii* and *Oxalis* sp. in Mexico and Brazil.

Septoria oxalidis Lind. Circular ashen leaf spots with dull-brown margins on *O. acetosella* in Denmark.

Septoria oxalidis-japonicae Pat. Sunken rufous to dark-brown leaf spots on *O. japonica* in Tibet.

Stagonospora hygrophila Sacc. On leaves of *O. acetosella* in Great Britain.

Stigmatea depazeaeformis (Auersw.) Schroet. On leaves of *O. acetosella* in Austria and Germany.

Possibly a form of *Mycosphaerella depazeaeformis*.

Uredo oxalidis Lév. Leaf rust on *O. corniculata*, *O. corymbosa*, *O. martiana* and *O. stricta* in Ecuador, Argentina, France, Switzerland, and Belgium.

Uromyces oxalidis Pazschke. Rust on petioles and scapes of *Oxalis* sp. in Brazil.

XYCOCCUS. Cranberry. See also *Vaccinium*.

Ascochyta oxycocci P. Henn. On leaves of *O. macrocarpus* in Germany.

Sclerotinia oxycocci Woron. Gray mold on branches and leaves and soft rot of fruit of *O. palustre* and *O. quadrupetalus* in Denmark and Finland.

XYPETALUM. Herbs and subshrubs.

Puccinia lagoensis P. Henn. Black rust pustules on leaves of *Oxypetalum* sp. in Brazil.

Puccinia oxypetali P. Henn. Stems shortened, swollen, and completely deformed, dark-brown rust pustules developing on the swollen areas and on leaves and flower buds on *O. banksii* in Brazil.

Septoria oxypetali F. Tassi. Small white leaf spots with black margins on *O. caeruleum* in Italy.

Uromyces cabo-friensis P. Henn. Black rust pustules on leaves of *Oxypetalum* sp. in Brazil.

XYTROPIS. Perennial herbs and shrubs.

Ovularia oxytropidis Jacq. On leaves of *O. pilosa* in Russia.

Peronospora oxytropidis Gaum. Downy mildew on leaves of *O. campestris* in Switzerland.

Phyllosticta oxytropidis Höll. Circular ochraceous leaf spots on *O. pilosa* in Hungary.

Septoria oxytropidis Died. On leaves of *O. pilosa* in Germany.

Septoria oxytropidis E. Rost. Large ashen leaf spots on *O. lapponica* in Norway.

Uromyces astragali (Opiz.) Sacc. See *Astragalus*.

PACHYRHIZUS. CACARA. Ag. Yam bean. Jicama. Climbing herbs with large tuberous roots used for food and starch production.

Phakopsora pachyrhizi Syd. Dull-brown rust sori on leaves of *P. angulatus* and *P. erosus* in Japan, Formosa, and the Philippines.

PACOURIA. See *Landolphia*.**PAEONIA**. PEONY. Showy spring and autumn blooming herbaceous perennials.

Aecidium paeoniae Kom. Leaf rust on *P. albiflora* in Russia.

Botrytis paeoniae Oud. This fungus causes a blighting of leaves and buds and a rotting of stems.

Black sclerotia are formed on old stems which enable the fungus to live over from season to season.

The disease is reported as serious in Great Britain, Holland, Germany, and other parts of Europe.

All varieties of cultivated paeonies are subject to attack as well as *Conrallaria*. The connection of

this disease with the American *Botrytis* blight is in doubt.

Cronartium flaccidum (Alb. and Schw.) Wint. (*C. paeoniae* Cast.) A rust producing yellow-brown to red-brown pustules on the leaves. Infected leaves present the appearance of being studded with long hairs on the under surfaces. One stage of the rust occurs on *Pinus silvestris*. It also occurs on *Asclepias pneumonanthis*, *A. speciosa*, *Cynanchum fuscatum*, *C. nigrum*, *C. purpurascens*, *Crammatocarpus volubilis*, *Impatiens balsamina*, *Nemesia versicolor*, *Paonia albiflora*, *P. angustifolia*, *P. anomala*, *P. arborea*, *P. arietina*, *P. broterii*, *P. corallina*, *P. daurica*, *P. decora*, *P. edulis*, *P. fimbriata*, *P. formosa*, *P. grandiflora*, *P. herbacea*, *P. hybrida*, *P. moultan*, *P. officinalis*, *P. peregrina*, *P. rosea*, *P. sessiliflora*, *P. sinensis*, *P. splendens*, *P. tenuifolia*, *Pedicularis palustris*, *Schizanthus grahami*, *Tropaeolum canariensis*, *T. lobbianum*, *T. major*, *T. minor*, *Verbena erimoides*, *V. leucoroides*, *V. officinalis*, *Vincetoxicum lazum*, and *V. officinalis* in Europe, Siberia, and Japan.

Leptothyrella paeoniae P. Henn. On leaves of *P. obovata* in Japan.

Leptothyrium paeoniae Vogl. Circular to oblong pale-yellow spots on leaves of *P. arborea* in Italy.

Mycosphaerella moultan Siem. Leaf spot on *P. moultan* in Russia.

Phyllosticta baldensis Massal. A leaf spot producing fungus on *P. decora*, *P. officinalis*, and *P. peregrina* in Denmark, Switzerland, Bulgaria, and Italy.

Phyllosticta moultan Pass. Dark-brown spots on leaves of *P. moultan* in Italy.

Phyllosticta paeoniae S. and P. Produces indefinite spots on leaves of cultivated *Paonia* varieties, including *P. corallina* in Great Britain.

Puccinia martianoffiana Thuem. Small sunken spots in which appear brown-black rust pustules on both surfaces of the leaves of *P. anomala* in Siberia.

Ramularia paeoniae Vogl. On leaves of *P. peregrina* in Italy.

Ramularia sardoa Sacc. and Trav. On leaves of *P. corallina triternata* in Italy.

Septoria macrospora Sacc. On leaves of *P. officinalis* in Italy and Russia.

Septoria martianoffiana Thuem. Produces a leaf spot on *P. anomala* in Siberia.

Septoria servica Syd. Suborbicular to irregular brown spots with black margins on both surfaces of leaves of *P. decora* in Yugoslavia.

PALA. See *Astonia*.**PALIURUS**. Woody plants, sometimes grown for their attractive foliage and curiously-shaped fruits.

Ascochyta paliuri Sacc. Yellow-brown leaf spots on *P. aculeatus* in Italy.

Phyllosticta camusiana Sacc. Leaf spots on *P. aculeatus* and *P. australis* in France and Italy.

Phyllosticta paliuri (Lév.) Cke. Black leaf spots on *P. australis* in Russia.

Septoria ascochytae Sacc. Olivaceous or dull-yellow leaf spots on *P. aculeatus* in Italy.

PALMAE. PALMS. The diseases of palms are considered in one group for reasons similar to those explained under *Bambuseae*.

Aphelethus cocophila Cobb. The red-ring disease of the coconut (*Cocos nucifera*) is serious in British Guiana, Trinidad, Tobago, Grenada, and Panama. It doubtless occurs in other parts of the West Indies and Central America. Externally it is marked by a yellowing and browning of leaves which works from below upwards. This phase takes place fairly rapidly, involving even the unfolded leaves. Green nuts of all sizes and the flowers are shed, the branches of the inflorescences withering. The bud rots. Internally a red zone appears in the stem, extending upward several feet, then breaking into longitudinal streaks, and finally into scattered dots. Similar red spots or streaks occur in the leaf petioles. The roots are discolored and finally die. The reddened tissues are filled with the nematodes which cause the disease.

PALMAE—Continued.

- Ascochyta depazeoides** (Dur. and Mont.) Tass. On leaves of *Phoenix dactylifera* and *Chamaerops humilis* in Portugal and Algeria.
- Ascochyta kentiae** Maubl. Elongate pale-brown leaf spots with dark-brown borders on *Kentia* sp. in Algeria.
- Ascochyella passeriniana** (Thuen.) Tass. On leaves of *Chamaerops* sp. and *Phoenix* sp. in Italy.
- Ascochyella pinnarum** (Pass.) Tass. On leaves of *Phoenix dactylifera* in Italy.
- Auerswaldia decipiens** Rehm. A doubtful species of tar-spot fungus on *Arenga mindorensis* and *A. tremula* in the Philippines.
- Bacillus coli** Esch. See Bud-rot.
- Bagnisiopsis bactridis** (Rehm.) Theiss. and Syd. Black stromata on leaves of *Bactris* sp. in Brazil.
- Bagnisiopsis diplothemii** (Rehm.) Theiss. and Syd. Dull-black stromata on leaves of *Diplothemium littorale* in Paraguay.
- Botryosphaeria palmigena** (B. and C.) Bomm. and Rouss. On leaves of *Chamaedorea* sp. in Costa Rica.
- Brachysporium arecae** (B. and Br.) Sacc. Leaf spot on *Areca catechu* in India and Ceylon.
- Bud rot.** A disease of coconuts and other palms, characterized by a soft rot of the bud which destroys the tree, has been reported from nearly all tropical palm-growing regions and constitutes the most serious menace to these valuable plants. Workers on the problem do not agree as to the causative organism, and in fact several are undoubtedly concerned. The form of the disease on *Cocos nucifera* and *Elais guineensis* in Cuba, Jamaica, Santo Domingo, Cayman Islands, Trinidad, British Guiana, Surinam, British Honduras, and possibly also in the eastern tropics has been attributed to a strain of *Bacillus coli* Esch.
- The disease is first noted through the falling of immature nuts, staining of opening flower spikes to a rich chocolate brown and the dying and bending over of center leaves. A dark wet rot develops at the base of affected spikes and spreads around the leaf sheaths. Finally the entire bud is destroyed by a foul-smelling soft rot, the leaves are brown and dead, hanging downward and all nuts have fallen. The entire top of a diseased tree ultimately falls away, leaving the bare dead trunk. Losses have been very large. For further discussion of bud-rot diseases see *Phytophthora*.
- Camarotella astrocaryae** (Rehm.) Theiss. and Syd. Black stromata on both leaf surfaces of *Astrocaryum* sp. in Brazil.
- Catacauma mucosum** (Speg.) Theiss. and Syd. (*Phyllachora cocoicola* P. Henn.) Tar spot on leaves of *Cocos australis* in Brazil.
- Cercospora acrocomiae** Stevenson. Oval red-brown, then gray, leaf spots with dark-brown margins on *Acrocomia media* in Porto Rico.
- Cercospora calamicola** P. Henn. Subcircular, black leaf spots on *Calamus caryotoides* in Queensland.
- Cercospora licualse** Syd. On leaves of *Licualla spinosa* in the Philippine Islands.
- Cercospora palmicola** Speg. Elliptical, often confluent, leaf spots on *Cocos australis* in Brazil.
- Cercospora prelsii** Bub. On leaves of *Phoenix* sp. in Russia.
- Cercospora virens** Sacc. On leaves of *Licualla* sp. in Malaya.
- Colletotrichum arecae** Syd. Anthracnose on leaves of *Areca catechu* in the Philippines.
- Colletotrichum paucisetum** Petch. Anthracnose on leaves of *Cocos nucifera* in Ceylon.
- Coniothyrium cocois** P. Henn. Dull-brown leaf spots on *Cocos nucifera* in east Africa.
- Coryneum cocoëis** P. Henn. Dull-brown leaf spots with darker brown margins on *Cocos nucifera* in east Africa.
- Didymosphaeria astrocaryi** v. Hoeh. On leaves of *Astrocaryum vulgaris* in Guiana.
- Diplodia cacaicola** P. Henn. See Theobroma.
- Dothidina palmicola** (Speg.) Theiss. and Syd. (*Auerswaldia palmicola* Speg.) Irregular to linear brown-black stromata on leaves of *Acrista monticola*, *Cocos yatai*, *Diplothemium littorale*, *Phoenix dactylifera*, and *Trithrinax brasiliensis* in Porto Rico, Brazil, Argentina, Paraguay, and Uruguay.
- Exosporium durum** Sacc. The leaves of *Cocos nucifera* in the Philippines turn yellow and become covered with small spots on which occur numerous dark-colored raised fruiting bodies.
- Exosporium pulchellum** Sacc. Numerous small, dark-colored, raised fruiting bodies on leaves of *Areca catechu* in the Philippines.
- Fomes lamaensis** Murr. See Hevea.
- Fomes lignosus** Klotzsch. See Hevea.
- Fusicladium livistonae** Karst. Black fungus layers on petioles of *Livistona sinensis* in Denmark.
- Gloeosporium catechu** Syd. Anthracnose on leaves of *Areca catechu* in the Philippines.
- Gloeosporium coccophilum** Wakef. Anthracnose on leaves of *Cocos nucifera* in Ceylon.
- Gloeosporium ptychospermatis** P. Henn. Large white spots on leaves of *Ptychosperma* sp. in Java.
- Graphiola appianata** Syd. and Butl. Small black, hard, raised fruiting bodies on leaves of *Phoenix sylvestris* in India.
- Graphiola arengae** Rac. Hard raised fruiting bodies on circular yellow leaf spots on *Arenga ambong* and *A. saccharifera* in Java and the Philippines.
- Graphiola borassi** Syd. and Butl. On leaves of *Borassus flabelliferus* in India.
- Graphiola cocoïna** Pat. Small black, raised, hard fruiting bodies on small brown-leaf spots on *Cocos nucifera* in Fiji and Gambier Islands.
- Graphiola cylindrospora** Syd. On leaves of *Livistona chinensis* in the Philippines.
- Graphiola macrospora** Penz. and Sacc. Globular black, hard fruiting bodies on leaves of *Calamus* sp. and *Plectocomia* sp. in Java.
- Haplosporella astrocaryi** P. Henn. On leaves of *Astrocaryum* sp. in Brazil.
- Helminthosporium coryphae** Syd. On leaves of *Corypha elata* in the Philippines.
- Helminthosporium incurvatum** Ch. Bern. On leaves of *Cocos nucifera* in Ceylon and Java.
- Helminthosporium philippinum** Sacc. On *Arenga tremula* in the Philippines.
- Heterosporium chamaeropsis** Oud. On leaves of *Chamaerops excelsa* in Holland.
- Heterosporium minutulum** C. and M. On leaves of *Chamaerops humilis* in Great Britain.
- Hysterostomella elaeicola** Maubl. Irregular black carbonous fruiting bodies in pale sunken leaf spots on *Elais guineensis* in Dahomey.
- Laestadia ecophilla** (Cke.) Sacc. On leaves of *Cocos nucifera* in British Guiana.
- Lembosia diplothemii** P. Henn. On leaves of *Diplothemium maritimum* in Brazil.
- Leptosphaeria briosiana** Pöhl. On leaves of *Jubaea spectabilis* in Italy.
- Leptosphaeria molleriana** d'Alm. and da Cam. On leaves of *Cocos romanzoffiana* in Portugal.
- Leptostroma phoenicis** Oud. and Fautr. On leaves of *Phoenix dactylifera* in France.
- Meliola furcata** Lévl. Black superficial fungus patches on leaves of *Acrista monticola* and *Coccothrinax alta* in Porto Rico.
- Melittosporiopsis gigantospora** Rehm. On leaves of *Rhaphidophyllum* sp. in the Philippines.
- Melittosporiopsis pachycarpa** Rehm. On leaves of *Rhaphidophyllum* sp. in Chile.
- Merrilopeltis hoehneltii** Rehm. On petioles of *Arenga mindorensis* and *A. saccharifera* in the Philippines.

PALMAE—Continued.

- Metasphaeria cocogena** (Cke.) Sacc. On leaves of *Cocos nucifera* in Demerara.
- Mollisia copelandi** Rehm. On leaves of *Caryota* sp. in the Philippines.
- Mycosphaerella** sp. Leaf spot on *Roystonea regia* in Cuba.
- Mycosphaerella chamaeropsis** Trav. Irregular ochraceous-gray leaf spots with broad purple margins on *Chamaerops humilis* in Italy.
- Mycosphaerella claudis** Beeli. Large irregular gray leaf spots on *Elaeis* sp. in Belgian Congo.
- Mycosphaerella gastonis** (Sacc.) Lind. On leaves of *Cocos nucifera* in Samoa, Tahiti, Ceylon, and the Philippines.
- Mycosphaerella ptychospermatis** Rehm. On leaves of *Ptychosperma macarthurii* in the Philippines.
- Ovularia palmicola** Pantanelli. On *Chamaerops* sp. and *Prithardia* sp. in Italy and Tripoli.
- Palawania grandis** (Niessl.) Syd. Black stromata on leaves of *Calamus* sp. and *Oncosperma horrida* in India, Java, and the Philippines.
- Parmulinea rehmi** (Maubl.) Theiss. and Syd. Dull-black stromata on yellow indefinite leaf spots on *Bactris* sp. in Brazil.
- Pestalozzia brevipes** Prill. and Delacr. Irregular yellow leaf spots on *Corypha australis* and *Kentia chamaerops* in France.
- Phaeochora acrocomiae** (Mont.) Theiss. and Syd. Black stromata on leaves of *Acrocomia sclerocarpa* and *A. yatay* in French Guiana, Brazil, and Paraguay.
- Phaeochora chamaeropsis** (Cke.) v. Hoeh. Shining black stromata on brown leaf spots on *Chamaerops humilis* in Morocco and Spain.
- Phaeotrabiella perisporioides** (Sacc.) Theiss. and Syd. Black stromata on circular to irregular leaf spots on *Phoenix dactylifera* in Oceania.
- Phomatospira migrans** Rehm. Large irregular, concentrically zoned areas on leaves of *Arenga saccharifera* in the Philippines.
- Phomopsis palmicola** (Wint.) Sacc. and var. *arecae* Sacc. Extensive irregular dead areas on leaves of *Areca catechu* in the Philippines.
- Phyllachora calamigena** (B. and Br.) Sacc. Brown stromata on brown leaf spots on *Calamus rudentum* in Ceylon.
- Phyllachora palmicola** Speg. Black stromata on long yellowish leaf spots on *Copernicia cerifera* and *Trithrinax brasiliensis* in Brazil and Argentina.
- Phyllachora roystoneae** Johnst. and Brun. Conspicuous black carbonaceous stromata on midribs of leaves of *Roystonea regia* in Cuba.
- Phyllosticta** sp. Leaf spot on *Caryota urens* in India.
- Phyllosticta chamaeropsis** Pöhl. Oblong brown leaf spots on *Chamaerops* sp. in Italy.
- Phyllosticta coccis** Allesch. Small oblong whitish leaf spots on *Cocos* sp. in Brazil.
- Phyllosticta cocoina** Sacc. Subcircular leaf spots on *Cocos nucifera* in Tahiti.
- Phyllosticta cocophila** Pass. Oblong to irregular whitish leaf spots with dull-brown margins on *Cocos flexuosa* and *C. nucifera* in Italy and the Philippines.
- Phyllosticta cocos** Cke. Irregular pale-brown leaf spots on *Cocos nucifera* in India and Italy.
- Phyllosticta daemonoropsis** Sacc. On leaves of *Daemonorops* sp. in Malaya.
- Phyllosticta palmigena** Sacc. On leaves of *Dictyosperma alba* in Malaya.
- Phyllosticta sabalicola** Szabo. On petioles of *Sabal blackburniana* in Hungary.
- Physalospora calani** Syd. Oblong, white leaf spots with brown margins on *Calamus tenuis* in India.
- Phytophthora arecae** (Colem.) Pethyb. The "koleroga" disease of the betel palm (*Areca catechu*) is serious in India. The fungus has also by inoculation been found capable of causing a rot of potato (*Solanum tuberosum*) tubers. The leaves of diseased palms turn yellow and droop, the flowers and immature fruit fall, and the flower stalks blacken and rot. This rot often spreads to the leaf sheaths and apical bud. The disease is one of the bud-rot type.
- Phytophthora faberi** Maubl. This downy mildew fungus has been found in Porto Rico, Jamaica, Ceylon, and the Philippines, causing typical bud rot of *Cocos nucifera* and other palms. The symptoms given for bud rot under that heading above apply here. The fungus has also been recently reported from limited areas in Florida in connection with the death of coconut palms. See also *Theobroma* for a further account of the fungus.
- Phytophthora melongenae** K. Saw. See *Solanum*.
- Phytophthora palmivora** Butler. The bud-rot disease of coconut (*Cocos nucifera*) and Palmyra palm (*Borassus flabellifer*) in India, Ceylon, Fiji, and Jamaica is attributed to this species. With this disease the inner leaves turn brown and break over and the heart leaf can usually be drawn out, having been rotted off at the base. All the young leaves turn yellow and the center column falls out, leaving a ring of the older leaves which may persist for some time. Young nuts cease growing. During this early stage brown sunken spots occur at the bases of young leaves and extend through to the bud, which soon becomes a pasty mass with an offensive odor. The inner and outer faces of the spots on leaf bases may show a white mildew. Development of the disease continues, the leaves all falling off and the tree dying.
- Phytophthora parasitica** Dastur. See *Ricinus*.
- Piaostroma diplothemii** Syd. Black shining stromata on leaves of *Diplothemium maritimum* in Brazil.
- Pleospora kentiae** Maubl. On leaves of *Kentia* sp. in Algeria.
- Pucciniopsis guaranitica** Speg. Small black fruiting pastules on brown leaf spots on *Cocos yatay* in Brazil.
- Ramularia thrincae** Rth. On leaves of *Thrinax nudicaulis* in Madeira.
- Rhipidocarpon javanicum** (Pat.) Theiss. and Syd. On leaves of *Nipa fruticans* in Java and the Philippines.
- Septoria calami** P. Henn. Pale indefinite leaf spots on *Calamus caryotoides* in Queensland.
- Septoria cocos** Patch. On leaves of *Cocos nucifera* in Ceylon.
- Septoria martiniae** Thuem. Irregular dull-brown leaf spots on *Martinia caryotefolia* in Portugal.
- Septoria palmorum** Sacc. On leaves of *Corypha elata*, *C. umbraculifera*, and *Latania borbonica* in Italy and the Philippines.
- Sphaerodothis arenziae** (Racib.) Shear. Black, shiny, irregular stromata on leaves of *Arenga obtusifolia*, *A. saccharifera*, *Caryota propinqua*, and *Caryota* sp. in Java and the Philippines.
- Stagonospora kentiae** Maubl. Whitish leaf spots with brown margins on *Kentia* sp. in Algeria.
- Thielaviopsis paradoxa** (DesSeyn.) v. Hoeh. Causes the "stem-bleeding" disease of *Cocos nucifera* in Ceylon and other parts of the oriental Tropics. See *Saccharum* for further discussion.
- Trabutia atroquinans** (Wint.) Theiss. and Syd. Black circular to elliptical stromata on leaves of *Mauritia flexuosa* in Brazil.

PALMAE—Continued.

Ustilago dactylifera Speg. In fruit of *Phoenix dactylifera* in central Africa. Beyond doubt only *Aspergillus niger* again. *Ustilago phoenicis* Cda. reported from Egypt on the same substratum is doubtless also the same.

Zukalla stuhlmanniana P. Henn. Black superficial fungus layers (sooty mold) on leaves of *Cocos nucifera*, *Phoenix dactylifera*, and *Ravenala madagascariensis* in Zanzibar and Tanganyika.

The "baloud" or "white" disease attacks *Phoenix dactylifera* in Algeria, often destroying the trees. The cause is unknown. Other diseases of the trunks, leaves, and fruit of this host are also reported from Algeria, the causes likewise being unknown.

PANAX. GINSENG. Hardy perennials with aromatic roots.

Bacterium arallavorum Uyeda. Causes a red rot of stems and roots of *P. quinquefolium* in Korea.

Bacterium panaxi Nak. and Tak. Produces an amber-colored rot in roots and a brown rot of stems of *P. quinquefolium* in Japan and Korea.

Colletotrichum panacicola Nak. and Tak. Leaf blight of *P. quinquefolium* in Korea.

Phoma panacicola Nak. and Tak. Black dry rot of roots of *P. quinquefolium* in Korea.

Phoma panacis Nak. and Tak. Produces brown to silvery-gray spots and cankers on stems of *P. quinquefolium* in Korea.

Phyllosticta panax Nak. and Tak. Brown leaf spots on *P. quinquefolium* in Korea; called the "snake-eye" disease.

Portia hypobrunnea Petch. See Hevea.

Rosellinia arcuata Petch. See Thea.

Rosellinia bunodes B. and Br. See Citrus.

Uredo panacis Syd. Dull-yellow rust pustules on leaves of *P. pseudoginseng* in India.

PANCRATIUM. Summer and winter flowering bulbs.

Aecidium amaryllidis Syd. and Buitl. Leaf rust on *Pancratium* sp. in India.

PANDANUS. SCREW PINE. Tropical woody plants and small trees remarkable for their aerial roots and the spiral arrangement of the long leaves.

Anthostomella pandani (Rabh.) Sacc. On leaves of *P. furcatus* in India.

Ellisiodothis pandani Syd. Black opaque stromata on leaves of *P. luzonensis* in the Philippines.

Leptosphaeria pandani Tass. On leaves and sheaths of *P. utilis* in Italy.

Phyllosticta pandanicola Young. Grayish-white to light-brown spots on leaves of *Pandanus* sp. in Porto Rico.

Pirostoma farnetianum Pöhl. On leaves of *P. utilis* in Italy, causing much damage.

Plenodomus inaequalis Sacc. and Trott. On leaves of *P. utilis* in Italy.

Stigmata pandani Pat. Circular dull-brown leaf spots on *P. odoratissimus* in Gambier Islands.

PANICULARIA. See Glyceria.**PANICUM**. GUINEA GRASS. PARA GRASS. BROOMCORN MILLET. A very large genus of annual and perennial grasses. Many are weeds, but others of importance as grain and forage plants.

Acrothecium lunatum Wak. Small brown leaf spots on *P. frumentaceum*, *Eleusine coracana*, and *Setaria italica* in India.

Balanis claviceps Speg. Forms sclerotia in spikelets of *P. carinatum* and *P. nodosum* in the Philippines and Argentina.

Balanis paspali P. Henn. Forms sclerotia in spikelets of *P. amplexicaule* in the Philippines.

Balanis trinitensis Cke. and Mass. Hard black globose stromata in place of ovaries on *P. palmifolium* in Trinidad.

Balanisella pulvinula (B. and Br.) Petch. Forms white to honey-colored stromata in young shoots of *Panicum* sp. in Ceylon, arresting their development.

Beniowskia graminis Rac. On leaves of *P. nepalense* in Java.

Chaetodiplodina graminicola Speg. On leaves of *P. laxum* in Argentina.

Chlorocytrium graminis Buesgen. See Festuca.

Cerebella antidotala Subra. Black corrugated fungus masses replacing ovaries of *P. antidotala* in India.

Cerebella burmanensis Subra. Black corrugated fungus masses in ovaries of *P. setigerum* in India.

Cerebella cynodontis Syd. See Cynodon.

Cerebella iniquans (B. and Br.) Petch. On *P. javanicum* and *Paspalum longiflorum* in India, *P. royleanum*, *P. (Syntherisma) sanguinalis*, and *P. scrobiculatum* in India and Ceylon.

Mycosphaerella panicola P. Henn. On leaves of *Panicum* sp. in Brazil.

Phaeodonthis stenostoma (E. and E.) Theiss. and Syd. Black stromata on leaves of *P. brizanthum* in central Africa.

Phyllachora acutispora Speg. Dull-black stromata on upper leaf surfaces of *Panicum* sp. in Brazil.

Phyllachora apiculata Speg. Small linear black stromata on leaves of *P. grumosum* in Argentina.

Phyllachora bokensis P. Henn. Dull-black stromata on oblong dull-brown leaf spots on *Panicum* sp. in the Congo.

Phyllachora bonariensis Speg. Small, subcircular, smooth black stromata on leaves of *P. bambusoides* in Venezuela and Argentina.

Phyllachora congruens Rehm. Black stromata on leaves of *P. carinatum* in the Philippines.

Phyllachora heterospora P. Henn. Shiny black stromata on large yellow leaf areas on *P. maximum* and *Panicum* sp. in Ecuador, the Congo, and the Union of South Africa.

Phyllachora panicl (Rehm.) Theiss. and Syd. Dull-black stromata on brown leaf spots on *Panicum* sp. in Brazil.

Phyllachora panicl-sulcati (P. Henn.) Theiss. and Syd. Black stromata on leaves of *P. sulcatum* in Costa Rica.

Phyllachora pazschkeana Syd. Circular or elliptical black stromata on leaves of *P. pilosum* and *P. sciurotes* in Brazil and Colombia.

Phyllachora raciborskii Theiss. and Syd. Black stromata on leaves of *P. nepalense* in Java.

Phyllachora sanguinolenta Theiss. and Syd. Shiny black circular stromata on leaves of *P. sanguinolentum* in the Congo.

Phyllachora seriata Theiss. and Syd. Black shiny oblong stromata on yellow leaf spots on *P. palmarum* and *Panicum* sp. in the Philippines.

Phyllachora stenospora (B. and Br.) Sacc. Tar spot on leaves of *P. patens* and *P. trigonum* in Ceylon and the Philippines.

Phyllachora vanderystii Theiss. and Syd. Black elliptical stromata on leaves of *P. plicatum* in Ceylon and the Congo.

Phyllachora urvilleana Speg. Tar spot on leaves of *P. urvilleana* in Argentina.

Phyllosticta panicl Young. Indefinite brown leaf spots on *P. maximum* in Porto Rico.

Physalospora panicl Rehm. On *Panicum* sp. in Brazil.

Puccinia goyazensis (P. Henn.) Syd. Rust on leaves of *Panicum* sp. in Brazil.

Puccinia huberi P. Henn. Brown to black rust pustules on leaves of *P. fasciculatum*, *P. ovalifolium*, *P. trichoides*, *P. paniculatum*, *P. tenellum*, *P. utowanum*, and *Paspalum virgatum* in Porto Rico, Cuba, Mexico, and Brazil.

PANICUM—Continued.

- Puccinia insolita** Syd. Powdery brown elongate rust sori on leaves of *P. maximum* in the Congo.
- Puccinia negrensis** P. Henn. Brown rust pustules on dull-brown leaf spots on *P. negrensis* in Brazil.
- Puccinia oahuensis** E. and E. Powdery yellow-brown rust pustules on leaves of *Panicum* sp. and *Syntherisma sanguinale* in Hawaii and Japan.
- Puccinia orientale** Syd. and Butl. Yellow-brown rust sori on leaves of *P. prostratum*, *P. trigonum*, and *Panicum* sp. in India and the Philippines.
- Puccinia panicophila** Speg. Brown rust sori on leaves of *P. insulare* in Argentina.
- Puccinia puttemansii** P. Henn. Leaf rust on *Panicum* sp. in Brazil.
- Septoria tandilensis** Speg. On leaves and sheaths of *P. nitidum* in Argentina.
- Sorosporium africanum** Syd. Black powdery smut masses destroying the ovaries of *P. trichops* in Portuguese East Africa.
- Sorosporium cryptum** McAlp. Smut sori concealed by the glumes and converting the ovaries into black spore masses. On *P. bicolor* and *P. effusum* in Australia.
- Sorosporium ovarium** Grifl. Gray-brown smut sori in ovaries of *P. caespitosum* in Mexico, rupturing to expose black powdery spore masses.
- Sorosporium panicl** MacKin. and var. **kinshasaensis** Beeli. Smut in ovaries of *P. flavidum*, *P. kinshasaensis*, and *Panicum* sp. in New South Wales and the Congo.
- Sphaedotheca chudaei** Har. and Pat. Long ovoid smut sori in inflorescences of *P. turgidum* in Morocco.
- Telimena graminis** (v. Hoeh.) Theiss. and Syd. Circular to oblong black stromata on leaves of *Panicum* sp. in Samoa.
- Telimena panicl** Theiss. and Syd. Dull-black stromata on leaves of *P. patens* in the Philippines.
- Tilletia ayresii** Berk. Buff spore masses produced in the ovaries of *P. maximum* in Mauritius.
- Tilletia courtetiana** Har. and Pat. Smut sori consisting of olive-brown spore masses, deforming ovaries of *P. proliferum* in the Congo.
- Tilletia magnusiana** F. de W. Black smut masses in ovaries of *P. geniculatum* in the West Indies.
- Tilletia tumefaciens** Syd. Powdery smut sori deforming leaves and culms of *P. antidotale* in India.
- Tilletia verrucosa** Cke. and Mass. Smut in ovaries of *P. coloratum* and *P. miliare* in South Africa.
- Tolyposporium bogoriense** Racib. Smut sori in flowers of *Panicum* sp. in Java.
- Tolyposporium minus** Schroet. Ashen smut sori in ovaries of *Panicum* sp. in Brazil, rupturing to expose black spore masses.
- Uredo panicl-montani** Petch. Leaf rust on *P. montanum* in Ceylon.
- Uredo panicl-prostrati** Syd. Powdery brown rust pustules on leaves of *P. prostratum* in India.
- Uredo panicl-villosi** Petch. Leaf rust on *P. villosum* in Ceylon.
- Uromyces leptodermus** Syd. Powdery brown rust pustules on leaves of *P. barbinode*, *P. isachne*, *P. javanicum*, and *P. prostratum* in Porto Rico, Cuba, Peru, Guatemala, and India. Also known from southern Florida.
- Uromyces linearis** B. and Br. Brown, then black, linear rust sori on leaves of *P. flavidum*, *P. miliare*, *P. proliferum*, and *P. repens* in India, Ceylon, China, Japan, and the Philippines.
- Uromyces puttemansii** Rangel. Leaf rust on *P. mellinum* in Brazil.
- Uromyces superfluus** Syd. Leaf rust on *P. antidotale* in India.
- Ustilaginoida congenisii** P. Henn. Globose to subtorbinate olivaceous sclerotia in ovaries of *P. brizanthum* and *P. maximum* in the Congo.
- Ustilaginoida ochraceae** P. Henn. Sclerotia in ovaries of *P. auritum* in the Philippines.
- Ustilaginoida usambarensis** P. Henn. Ashen-olivaceous sclerotia deforming the ovaries of *P. laxum* in Porto Rico and Tanganyika.
- Ustilago balansae** Speg. Powdery dark-brown smut masses in panicles of *Panicum* sp. in Argentina.
- Ustilago bonariensis** Speg. Powdery dark-olive smut sori covered by a whitish pseudomembrane in spikelets of *P. elephantipeds*, *P. spectabile*, and *Pennisetum tristachyon* in Argentina.
- Ustilago cacheutensis** Speg. Black powdery smut masses destroying the panicles of *P. leucophaeum* in Argentina.
- Ustilago confusa** Mass. Smut in ovaries of *P. paradoxum* in Australia.
- Ustilago digitaliae** (Kze.) Rbh. Flowers, rachides, and culms destroyed and replaced by black smut sori on *P. glabrum*, *P. repens*, *P. teneriffae*, and *Syntherisma sanguinale* in Ceylon, India, Formosa, Portuguese East Africa, Sudan, Italy, and Germany.
- Ustilago formosana** K. Saw. Linear smut sori, up to 14 centimeters long, inclosed at first by a grayish-white membrane, which ruptures, exposing the black spore masses; in inflorescences and culms of *P. proliferum* in Formosa.
- Ustilago globigena** Speg. Smut sori in spikelets of *P. spectabile* in Argentina.
- Ustilago heterospora** P. Henn. Powdery olivaceous spore masses deforming the ovaries of *P. maximum*, *Setaria aenea*, and *S. glauca* in tropical Africa.
- Ustilago manilensis** Syd. Smut sori in the spikes of *P. indicum* in the Philippines.
- Ustilago negeriana** Diet. Black spore masses in ovaries of *P. urvilleanum* in Chile.
- Ustilago operta** Syd. and Butl. Smut in ovaries of *P. prostratum*, *P. ramosum*, and *P. villosum* in India.
- Ustilago panicl-frumentacei** Bref. Black smut sori covered with a persistent gray membrane involving inflorescences, stems, nodes, and young shoots and causing abortion of fruiting heads. On *P. (Echinochloa) frumentaceum* in India.
- Ustilago panicl-gracilis** MacKin. Smut in ovaries of *P. gracile* in New South Wales.
- Ustilago panicl-latifolii** P. Henn. Black powdery smut sori in ovaries of *P. latifolium* in Brazil.
- Ustilago panicl-leucophaei** Bref. Spikelets of *P. leucophaeum* in Brazil and Guatemala transformed into black powdery smut masses.
- Ustilago panicl-petrosi** Syd. Dark-brown powdery smut sori destroying the inflorescences of *P. petrosum* in Brazil.
- Ustilago panicl-proliferi** P. Henn. Smut sori in inner folded leaves, aborting the inflorescences and covered by thin transparent membranes which soon rupture, exposing dusty olive-black spore masses, on *P. paspaloides* and *P. proliferum* in Mexico.
- Ustilago paradoxa** Syd. and Butl. Smut sori attacking the individual ovaries of *P. frumentaceum* in India. The sori are round, gray, and covered by persistent membranes, which are broken only by handling, exposing the black spore masses.
- Ustilago pretoriensis** Pole-Evans. Long powdery dark olivaceous smut sori destroying the ovaries of *P. helopodis* in the Union of South Africa.
- Ustilago rickerii** Clint. Olive-black dusty smut masses in ovaries of *P. paspaloides* in Cuba.
- Ustilago trichophora** (Link.) Kze. Smut sori in ovaries of *P. colonum* and *Tricholaena (Panicum) teneriffae* in India and north Africa.
- Ustilago ugandensis** P. Henn. and var. **macrospora** Beeli. Smut sori consisting of olivaceous powdery spore masses at apex of culms, in rachides and flowers of *Panicum* sp. in tropical Africa.
- Ustilago vesiculosa** P. Henn. Smut sori in panicles of *Panicum* sp. in Brazil.

- PAPAYER.** POPPY. Biennial or perennial herbs with milky juice. Economic and flower-garden plants. Oil produced from the seed, opium and its derivatives from the juice.
- Ascochyta papaveris** Oud. On leaves of *P. nudicaule* in Nowaja Semla.
- Entyloma fuscum** Schroet. See Glaucium.
- Gloeosporium** sp. Anthracnose on leaves of *P. somniferum* in Japan.
- Helminthosporium papaveris** K. Saw. Large irregular brown leaf spots on *P. somniferum* in Formosa. Lesions also occur on the stems and circular yellow to brown spots on the capsules. Infected plants are often destroyed. "The most dangerous disease of the cultivated poppy in Formosa."
- Helminthosporium papaveris** P. Henn. On *Papaver* sp. in Tanganyika.
- Macrosporium papaveris** Parisi. On green capsules and leaves of *P. somniferum* in Italy.
- Peronospora arborescens** (Berk.) De By. A downy mildew forming pale-brown leaf spots, more especially near the tips and margins. These spots dry up and fall out and entire leaves may wither. On the lower surfaces a gray-violet fungus layer forms. On *P. dubium*, *P. rhoeas*, *P. somniferum* and *Meconopsis cambrica* in Japan, India, Siberia, Algeria, Egypt, and Europe.
- Peronospora argemone** Gäum. Downy mildew on leaves of *P. argemone* in Scotland, Scandinavia, Austria, and Germany.
- Peronospora cristata** Tranz. As above on *P. hybridum* in Asia Minor.
- Septoria rhoeadis** F. Tassi. Violet-brown circular and confluent leaf spots on *P. rhoeas* in Italy.
- PAPHIOPELIDUM.** Terrestrial or epiphytic orchids. See Orchidaceae.
- PAPYRIUS.** See Broussonetia.
- PARAMIGNYA.** Evergreen climbing shrubs.
- Acidium paramignyae** Racib. Rust on circular leaf spots on *P. blumei* and *P. monophylla* in Java, Ceylon, and India.
- Bacterium citri** Haase. See Citrus.
- Helminthosporium makiingense** Syd. On branches of *P. monophylla* in the Philippines.
- PARIS.** Small, hardy, rhizomatous plants.
- Cercospora paridis** Erikss. Dark-brown leaf spots on *P. quadrifolia* in Russia, Sweden, Switzerland, Bohemia, Austria, and Denmark.
- Septoria paridis** Passer. Small irregular whitish leaf spots with dull-brown margins on *P. quadrifolia* in Russia and Italy.
- Tubercinia paridis** (Ung.) Vesterg. Smut on leaves and stems of *P. quadrifolia* in Russia, Sweden, Germany, and Austria.
- Urocystis colchici** (Schlecht.) Rab. See Colchicum.
- PARKIA.** CUPANG. Tall, unarmed leguminous trees.
- Endodothella parkiae** (P. Henn.) Theiss. and Syd. Black stromata on brown leaf spots on *Parkia* sp. in Brazil.
- Leptothyrella manaoensis** P. Henn. Black fruiting bodies on leaves of *P. auriculata* in Brazil.
- Phyllachora parkiae** P. Henn. Circular shiny black stromata on indefinite yellowish leaf spots on *P. roxburghii* and *P. timoriana* in the Philippines.
- PARNASSIA.** Sometimes called grass of Parnassus. Low growing, hardy perennial herbs.
- Puccinia uliginosa** Juel. Leaf rust on *P. palustris* and *Carex vulgaris* in Europe.
- PAROSELA.** See Dalea.
- PARRYA.** Low perennial herbs with thick caudices and scapelite peduncles.
- Puccinia oudemansii** Tranzsch. See Matthiola.
- PARSONSIA.** **HELYGIA** Ag. Twining apocynaceous shrubs.
- Acidium parsoniae** Fetch. Leaf rust on *P. spiralis* in Ceylon.
- PARSONSIA.** See Cuphea.
- PARTHENIUM.** Annual or perennial herbs or shrubs, mostly heavily pubescent.
- Entyloma calendulae** (Oud.) De By. See Calendula.
- Uredo parthenii** Speg. Leaf rust on *P. hysterophorus* in Argentina.
- PASANIA.** **LITHOCARPUS** Ag. Ornamental trees grown for their handsome foliage.
- Taphrina kusanoi** Ikeno. Fruiting areas on lower leaf surfaces of *P. cuspidata* in Japan.
- PASCALIA.** See Wedelia.
- PASPALUM.** Perennial grasses, including valuable forage grasses as well as weeds.
- Balanisa gigas** Racib. Globose yellow-brown stromata, white within, on *Paspalum* sp. in Java.
- Balanisa paspali** P. Henn. Black rugulose stromata in spikes of *Paspalum* sp. in Oceania and the Philippines.
- Cerebella inquinans** (B. and Br.) Petch. See Panicum.
- Claviceps deliquescens** (Speg.) Haum. Sclerotia in ovaries of *P. dilatatum*, *P. distichum*, *P. hieronymi*, *P. larranagai*, *P. plicatulum*, and *P. quadrifarium* in Argentina. Probably not distinct from *C. paspali*, which is widespread in the United States.
- Claviceps lutea** A. Möll. Yellow oblong curved sclerotia in spikelets of *Paspalum* sp. in Brazil.
- Endodothella platensis** (Speg.) Theiss. and Syd. Elliptic to linear black stromata on leaves of *Paspalum* sp. in Argentina.
- Ephedra japonica** P. Henn. Inflorescences deformed by blackish stromata which form between the compressed spikelets of *P. scrobiculatum*, *P. thunbergium*, and *Miscanthus* sp. in Japan.
- Helminthosporium mayaguezense** Miles. Oval yellow spots with narrow dark-brown margins on leaves and culms of *P. conjugatum* in Porto Rico.
- Phyllachora acuminata** Starb. Tar-spot on leaves of *Paspalum* sp. in Argentina.
- Phyllachora infusans** Wint. Irregular linear black stromata on *Paspalum* sp. in Brazil.
- Phyllachora paspaleola** P. Henn. Dull-black stromata on leaves of *P. compressum* and *P. conjugatum* in Brazil and Colombia.
- Phyllachora winkleri** Syd. Tar spot on leaves of *P. scrobiculatum* in Ceylon and Tanganyika.
- Puccinia huberi** P. Henn. See Panicum.
- Puccinia macra** Arth. and Holw. Leaf rust on *P. candidum* in Guatemala.
- Puccinia maublancii** Rangel. Leaf rust on *P. densum* in Brazil.
- Puccinia pilgeriana** P. Henn. Dark-brown rust sori on leaves of *Paspalum* sp. in Brazil.
- Puccinia tubulosum** (Pat. and Gaill.) Arth. Aecia in circular yellow areas 1 to 2 centimeters in diameter on leaves of *Solanum tomentosum* and *S. torum* in Mexico, Porto Rico, Cuba, Jamaica, and Central America. Brown powdery uredo and telial sori on leaves of *P. compressum*, *P. conjugatum*, *P. humboldtianum*, *P. paniculatum*, *P. plicatulum*, *Syntherisma digitata*, *S. sanguinale*, and *Valota insularis* in Mexico, Central America, the West Indies, and Trinidad. Also known from Texas.
- Sorosporium paspali** McAlp. Smut sori in inflorescences of *P. scrobiculatum*, destroying them and leaving the elongated remains surrounded by a black spore mass. Occurs in Ceylon, India, Australia, and the Philippines.
- Telimena graminella** Syd. Opaque black stromata on leaves of *Paspalum* sp. in the Philippines.
- Telephora inquinans** B. and Br. Black smut sori in seed of *P. scrobiculatum* in Ceylon.
- Tilletia ulei** Schroet. Black smut sori destroying the ovaries of *P. scrobiculatum* in Brazil.
- Uredo paspali-longiflorae** Petch. Leaf rust on *P. longiflorum* in Ceylon.
- Uredo paspalina** Syd. Brown leaf rust on *P. scrobiculatum* in the Philippines.

PASPALUM—Continued.

- Uredo paspali-perrottetii** Petch. Leaf rust on *P. perrottetii* in Ceylon.
Uredo paspali-scribiculari Syd. Brown rust sori on upper leaf surfaces of *P. scribicularum* in India and Ceylon.
Ustilaginoides ochracea P. Henn. On *Paspalum* sp. in Polynesia.
Ustilago microspora Schroet. and P. Henn. Smut sori reducing panicles of *P. dilatatum* and *Paspalum* sp. in Brazil and Uruguay to oliveaceous powdery spore masses.
Ustilago paspali Speg. Black powdery spore masses destroying ovaries and glumes of *Paspalum* sp. in Uruguay and Argentina.
Ustilago paspali-dilatati P. Henn. Black smut masses replacing inflorescences of *P. dilatatum* in Brazil.
Ustilago paspali-thunbergii P. Henn. Black powdery smut masses in the inflorescences of *P. thunbergium* in Japan.
Ustilago rabenhorstiana J. Kuehn. Smut sori in inflorescences of *P. (Syntherisma) sanguinale* in India.
Ustilago roylei Syd. and Butl. Black smut sori in spikes of *P. royleanum* in India.
Ustilago schroeteriana P. Henn. Black powdery smut sori in spikes of *Paspalum* sp. in Brazil.
Ustilago subnitens Schroet. and P. Henn. Smut on *Paspalum* in Brazil.
Ustilago venezuelana Syd. Smut on *Paspalum* sp. in Brazil.
Ustilago verrucosa Schroet. Black smut in ovaries of *P. distichum* in Brazil.

PASSERINA. Heathlike shrubs.

Marsonia anduricensis (Ces.) P. Magn. On stems of *P. annua* in Europe.

PASSIFLORA. PASSION FLOWER. Herbs, shrubs, and trees, many species climbers cultivated for fruit and as ornaments.

- Aecidium passiflorae** P. Henn. Leaf rust on *Passiflora* sp. in Tanganyika.
Ascochyta passiflorae Penz. and Sacc. On peduncles of *P. hybrida* and *P. incarnata* in Italy.
Asterina megalospora B. and C. Black superficial fungus patches on *Passiflora* spp. in Porto Rico, Cuba, Ecuador, Peru, and Brazil.
Colletotrichum passiflorae Slem. Anthracnose on *P. edulis* in Russia.
Didymella passiflorae v. Hoch. On leaves of *Passiflora* sp. in Samoa.
Didymosphaeria innumerabilis Wint. Circular to irregular whitish leaf spots on *Passiflora* sp. in Brazil.
Helminthosporium stahlil Stevens. Small irregular to large and diffuse brown leaf spots on *P. foetida* in Porto Rico.
Melampsora passiflorae Har. Rust on leaves of *P. lutea* in France.
Mycosphaerella passiflorae Rehm. Leaf spot on *Passiflora* sp. in South America.
Ovulariopsis sp. On leaves of *Passiflora* sp. in India.
Phyllosticta passiflorae McAlp. Large irregular leaf spots on *P. edulis* in Australia.
Phyllosticta superficialis Stevens. Irregular, brown leaf spots on *P. scrofula* in Porto Rico.
Puccinia scleriae (Paz.) Arth. The aecial stage of this rust occurs on circular, yellow-brown leaf spots on *P. rubra* and *P. tuberosa* in Porto Rico, Cuba, Trinidad, Jamaica, and Peru.
Uromyces appeianus Gassn. Leaf rust on *P. foetida* in Brazil and Uruguay.

PASTINACA. PARSNIP. Fleshly edible roots.

- Cereospora pastinacae** Karst. Small circular brown spots with white centers on leaf blades and petioles of *P. sativa* and *Daucus carota* in Great Britain, Finland, Russia, and Germany. Reported from Indiana.
Cylindrosporium pimpinellae C. Mass. var. *pastinacae* Sacc. Small dry spots on leaves of *P. sativa* and *P. urens* in Bulgaria, Italy, and Germany.
Entyloma pastinacae Jaap. Smut sori in grayish-white leaf spots on *P. sativa* in Dalmatia.
Gloeosporium achaenitcola Rost. See *Petroselinum*.
Phyllachora pastinacae Rost. Indefinite, finally confluent, pale-green, then brown, leaf spots on which stromata occur on *P. sativa* in France, Belgium, Sweden, Denmark, Russia, and Italy.
Protomyces macrosporus Ung. See *Coriandrum*.
Ramularia pastinacae Bub. Circular to irregular dark grayish-green, then brown, leaf spots with white centers on *P. sativa* and *Daucus carota* in Yugo-slavia, Dalmatia, Great Britain, Finland, and Austria. Reported from North Dakota.

PATRINIA. FEDIA. Ag. Hardy herbaceous perennials with yellow or white Valerian-like flowers.

- Aecidium patriniae** P. Henn. Leaf rust on *P. scabiosaefolia* and *P. villosa* in Japan.
Puccinia melanoplaea Syd. Black rust sori on leaves of *P. palmata* in Japan.
Puccinia patriniae P. Henn. Chestnut-brown rust sori on leaves of *P. villosa* in Japan.
Puccinia patriniae-gibbosae Mura. Leaf rust on *P. gibbosa* in Japan.

PAULLINIA. Twining tropical shrubs.

Phaeoplopora neetroides (Rehm.) Sacc. and Syd. Rust-brown galls on leaves of *Paullinia* sp. in Brazil.

PAULOWNIA. Ornamental trees grown for their showy flower panicles and large leaves.

- Gloeosporium kawakamii** Miy. Causes witches'-brooms on *P. tomentosa* (*P. imperialis*) in Japan. The leaves on these brooms remain small, pale green in color, and curled, with red-brown spots on blades and petioles. Seedlings often die.
Helicobasidium tanakae Miy. See *Morus*.
Mycosphaerella paulowniae Shir. and Hara. Leaf spots on *P. tomentosa* in Japan.
Phyllosticta paulowniae Sacc. Sinuous and confluent ochraceous leaf spots on *P. tomentosa* in France, Italy, and Russia.
Septoria paulowniae Thuem. More or less circular gray leaf spots on *P. tomentosa* in Italy and France.

Valsa paulowniae Miy. and Hemmi. Attacks the twigs of *Paulownia* sp. and *P. tomentosa* in Japan, spreading into the branches and finally destroying the entire tree. Considered serious, being compared with chestnut blight in virulence.

PAVETTA. Tropical shrubs and small trees allied to *Ixora*.

- Aecidium pavettae** Berk. Leaf rust on *P. hispidula* and *P. indica* in Ceylon, India, and Uganda.
Aecidium transvaaliae P. Henn. and Evans. Leaf rust on *Pavetta* sp. in the Union of South Africa.
Cocconia placenta (B. and Br.) Sacc. See *Symplocos*.
Endophyllum griffithiae Rac. See *Randia*.
Hemileia mildbraedii Syd. Powdery yellow rust pustules on leaves of *P. oliveriana* in central Africa.
Phoma pavettae F. Tassi. On branches of *P. indica* in Italy.
Woodiella natalensis Sacc. and Syd. Small leaf spots on *P. obovata* in Natal.
PAVONIA. Malvaceous herbs or shrubs.
Caecoma pavoniae Diet. Golden rust sori on leaves of *Pavonia* sp. in Brazil.
Phyllachora pusilla Syd. Tar spot on leaves of *P. schrankii* in Brazil.
Puccinia exilis Syd. Yellow-brown rust pustules on leaves of *P. leucantha* and *P. rosea* in Guatemala and Brazil.

PAVONIA—Continued.

Puccinia pavoniae P. Henn. Ochraceous-brown rust pustules on leaves of *Pavonia* sp. in Tanganyika.

Puccinosira pallidula (Speg.) Lagh. Yellow rust sori on leaves of *P. paniculata* and *Triumfetta* spp. in Central and South America and the West Indies.

Ramularia aguierei Speg. Indefinite areas on leaves of *P. sepium* in Argentina.

Rostrupia praelonga Speg. Linear yellow-brown rust pustules on leaves and branches of *P. polymorpha* in Argentina.

Uromyces pavoniae Arth. Leaf rust on *P. racemosa* in Porto Rico.

PEDICULARIS. WOOD BETONY. Lousewort. Mostly perennial herbs with finely cut foliage.

Conartium flaccidum (Alb. and Schw.) Wint. See *Paeonia*.

Melampsora pedicularis Vogl. Leaf rust on *P. verticillata* in Switzerland.

Peronospora pedicularis Palm. Downy mildew on leaves of *P. lapponica* in Sweden and Norway.

Plasmopara densa Rabb. Schroet. Downy mildew on leaves of *P. palustris*, *P. sibirica*, *Alectorolophus alpinus*, *Barisia alpina*, *Euphrasia officinalis*, *E. pratensis*, and *Odonites rubra* in Europe.

Puccinia oederi Blytt. Brown rust sori on leaves of *P. oederi* in Norway.

Puccinia paludosa Plowr. The yellow aecia of this rust occur on *P. mexicana* and *P. palustris*, the other stages on Carex spp. in Europe.

Puccinia pedicularis Thuem. Leaf rust on *Pedicularis* sp. in Siberia.

Ramularia filiformis Lindr. Leaf spots on *P. sylvatica* in Sweden.

Ramularia obtusens Thuem. On leaves of *P. foliosa* and *P. palustris* in Austria and Switzerland. Reported from California.

Rhytisma pedicularis (DC.) Rehm. Black confluent stromata on leaves of *P. incarnata* and *P. alpina* in Italy.

Septoria pedicularium Fr. On stems of *Pedicularis* sp. in Europe.

PELARGONIUM. GERANIUM. Herbaceous plants used as potted plants for indoor planting and for bedding.

Cercospora sp. On leaves of *Pelargonium* sp. in China.

Coniothyrium trabuti Riza. Indefinite yellow patches on leaves of *P. peltatum* in Morocco.

Fusarium pelargonii Crouan. Causes a black rot of stems, the leaves turning yellow, on *Pelargonium* sp. in Great Britain, France, and Denmark.

Gloeosporium pelargonii Cke. and Mass. Anthracnose, causing leaves of *Pelargonium* sp. in Great Britain to wither and droop.

Leptosphaeria pelargonii Rehm. Irregular whitish spots on leaves of cultivated *Pelargonium* in Brazil.

Macrosporium macalpineanum Sacc. and Syd. Large irregular and confluent yellow-brown leaf spots with purple margins on *P. zonale* in Australia and Ceylon.

Puccinia granularis Kaich. and Cke. Brown rust sori in circles on lower leaf surfaces of *P. aconitophyllum*, *P. alchemilloides*, and other species, as well as *Begonia* sp., in the Union of South Africa. Serious in greenhouses.

Puccinia morrisoni McAlp. Orange and brown to black rust pustules on leaves of *P. australe* in Australia.

Puccinia pelargonii (Thuem.) Syd. Yellow and brown rust sori on leaves of *P. alchemilloides* in the Union of South Africa.

Septoria geranii Rob. and Desm. See *Geranium*.

Septoria pelargonii Syd. Circular to irregular yellow-brown to red-brown leaf spots on *P. cuculatum* in the Union of South Africa.

PELLAEA. CLIFF BRAKE. Small rock-loving ferns.

Uredo pellaeae Diet. and Neg. Rust on leaves of *P. ternifolia* in Chile.

PELTOPHORUM. Tropical leguminous trees.

Phyllachora peltophori Syd. Shiny black stromata on leaves of *P. africanum* in the Union of South Africa.

PENNISETUM. PEARL MILLET. NAPIER GRASS. Stout grasses with bristly spike-like inflorescences.

Acrothecium penniseti Mitra. The fungus attacks the leaves, leafsheaths, and ears of *P. typhoidum* in India. The leaf spots are small, yellow brown, and oval to oblong. The lower leaves are destroyed and infected heads fail to set grain.

Balanisia claviceps Speg. Black stromata in spikes of *Pennisetum* sp. and *Setaria* sp. in Argentina.

Beniowskia penniseti Wakel. Small pale-red leaf spots on *P. purpureum* and *Holcus sorghum* in Uganda.

Cerebella cenchroidis Subra. Black corrugated fungus masses in ovaries of *P. cenchroides* in India.

Cintractia cotumellifera (Tul.) McAlp. See *Andropogon*.

Didymella penniseti Syd. On leaves of *P. tristachyon* in Columbia.

Leptostroma penniseti P. Henn. Effuse dull-brown leaf spots on *P. japonicum* in Japan.

Neovossia bardayana Bref. Smut sori in ovaries of *P. triflorum* in India.

Phyllachora minutissima (Welw. and Curr.) Sm. Small elliptical stromata on leaves of *Pennisetum* sp. in Angola.

Phyllachora penniseti Syd. Black stromata on leaves of *P. benthami* and *P. sphacellatum* in Congo and the Union of South Africa.

Phyllachora pennisetina Syd. Tar spot on leaves of *P. alopecuroides* in China.

Puccinia apoda Har. and Pat. Brown to black rust sori on rufous leaf spots on *P. setosum* in the Congo.

Puccinia arthuri Syd. Leaf rust on *P. crinitum* and *P. mexicanum* in Mexico.

Puccinia burmeisteri Speg. Leaf rust on *P. tristachyon* in Argentina.

Puccinia penniseti Zimm. Brownish-yellow to black rust sori on brown leaf spots on *P. spicatum* and *P. typhoidum* in India and east Africa.

Tolyposporium penicillariae Bref. Smut sori bright green to black, projecting beyond the glumes and finally breaking to expose deep-brown spore masses. Attacking individual grains or small groups of grain of *P. spicatum* and *P. typhoidum* in India, Egypt, and other parts of Africa.

Ustilaginoides penniseti Miy. Sclerotia in ovaries of *P. compressum* in China.

Ustilago bonariensis Speg. See *Panicum*.

Ustilago kamerunensis Syd. Smut sori destroying inflorescences of *Pennisetum* sp. in Congo.

Ustilago pappiana Bacc. Black smut sori in ovaries of *P. orientalis* and *P. ruppelii* in Abyssinia.

Ustilago penniseti Rabenh. Black smut sori in ovaries of *P. asperifolium*, *P. cenchroides*, *P. dichotomum*, *P. fasciculatum*, *P. ruppelii*, *P. vulpinum*, and *P. villosum* in Madeira, China, Arabia, Abyssinia, Egypt, Algeria, and Tripoli.

Ustilago penniseti-japonici P. Henn. Smut sori destroying flowers of *P. compressum*, *P. japonicum*, and *P. purpurascens* in China and Japan.

Ustilago scheffleri Syd. Smut sori up to 3 centimeters long, covered by ashen membranes which rupture, exposing black powdery spore masses in inflorescences of *P. inclusum* in tropical Africa.

PENTAS. Tender herbs and subshrubs.

Puccinia pentadictyla Grove. Leaf rust on *P. verticillata* in Uganda.

Puccinia pentadis P. Henn. Brown rust sori on leaves of *P. mombassana* in tropical Africa.

Puccinia pentadis-carneae Wakef. Leaf rust on *P. carnea* in Uganda.

PENTSTEMON. Sometimes called beardtongue. Tubular-flowered bedding and border plants.

Phyllosticta pentastemonis Cke. and f. *pentastemonis-azurei* Allesch. Irregular brown leaf spots on *P. azureus*, *P. grandiflorus*, and *P. hybridus* in Great Britain and Germany.

Puccinia mexicana Diet. and Holw. Brown rust pustules on sunken, circular leaf spots on *P. campanulatus* in Mexico.

Puccinia pentastemonum Lév. Brown rust sori on leaves of *Pentstemon* sp. in France.

PEPEROMIA. Tropical and subtropical herbs, a few species cultivated as foliage plants.

Albugo tropica Lagh. White rustlike pustules on leaves of *P. pellucida* in the West Indies and Ecuador.

Uredo peperomiae P. Henn. Irregular, dull yellow rust sori on leaves of *Peperomia* sp. in Brazil.

Uredo piperis P. Henn. See *Piper*.

PERDICUM. See *Gerbera*.**PERESKIA.** Shrubby or clambering vines with edible fruits.

Uromyces pereskiae Diet. Light-brown rust pustules on *Pereskia* sp. in Brazil.

PERILLA. Herbs, some species with colored foliage.

Coleosporium perillae Syd. Golden-yellow rust pustules on leaves of *P. arguta*, *P. crispata*, *P. nan-kinensis*, *P. ocimoides*, *Elsholtzia cristata*, *Keiskea japonica*, *Mosla formosana*, and *M. punctata* in Japan, Formosa, Manchuria, and India.

PERIPLLOCA. SILK VINE. Ornamental vines grown for their handsome glossy foliage.

Ascochyta periplocae Kab. and Bub. Large round to irregular dark-brown, then ashen, leaf spots with brown-purple margins on *P. graeca* in Bohemia and Hungary.

Melampsora periplocae Miy. Leaf rust on *Periploca* sp. in China.

Phyllosticta periplocae F. Tassi. Circular, ashen leaf spots on *P. graeca* in Italy.

Septoria periplocae Höll. Subcircular, brown leaf spots on *P. graeca* in Hungary.

Uredo periplocae P. Henn. Ochraceous rust sori on irregular yellow leaf spots on *P. linearis* in tropical Africa.

PERISTROPHE. Greenhouse plants grown for their flowers.

Accidium peristrophes Syd. Leaf rust on *Peristrophe* sp. in India.

Synchytrium ryzii Syd. Small leaf galls on *Peristrophe* sp. in India.

PERNETTYA. Plants grown for their attractive berries, evergreen foliage, and white or pinkish flowers.

Hypocrella pernettyae Pat. On leaves of *P. repens* in Java.

Stereum purpureum Pers. See *Prunus*.

PERSEA. Avocado. Sometimes called alligator pear. Tropical and subtropical fruit trees.

Cercospora (?) lingue Speg. On leaves of *P. lingue* in Chile.

Endothia havanensis Bruner. See *Eucalyptus*.

Fomes lamaensis Murr. See *Hevea*.

Mycosphaerella perseae Miles. Large irregular leaf spots, ashen above, brown beneath on *P. americana* in Porto Rico.

Phyllachora gratissima Rehm. Black stromata on irregular red-brown leaf spots on *P. americana* in Porto Rico, Costa Rica, and Ecuador.

Physalospora perseae Doidge. Produces a canker disease of *P. americana* in the Union of South Africa. Sunken discolored areas appear, gradually girdling twigs and branches. The dead bark dries out, cracks, and finally falls away. Brown, deeply cracking corky growths on the lower ends of the fruit are attributed to the same fungus.

Piononthe capillacea Sacc. Causes a die back of *P. americana* in the Philippines.

Rhytisma perseae Gandara. Black stromata on large, irregular brown leaf spots on *P. americana* in Mexico. Probably not distinct from *Phyllachora gratissima* Rehm.

Rosellinia spp. (*R. pepo* Pat.). See *Citrus*.

Eucalyptus disease. See *Eucalyptus*.

PERTYA. Oriental shrubs.

Accidium pertyae P. Henn. Rust pustules on circular dull-brown leaf spots on *P. ovata* and *P. scandens* in Japan.

PETASITES. BUTTERBUR. Hardy perennial herbs.

Accidium petasitidis Syd. Leaf rust on *P. albus*, *P. japonicus*, *P. nivus*, *P. officinalis*, and *P. tomentosus* in Japan and central Europe.

Coleosporium petasitis Lév. See *Pinus*.

Bamularia cervina Speg. and var. *petasitis* Bauml. On leaves of *P. albus* in Hungary.

Venturia petasitidis (Fekl.) Sacc. On leaves of *P. officinalis* in Europe.

PETIVERIA. Shrubby herbs.

Rosellinia bunodes B. and Br. See *Citrus*.

PETROSELINUM. PARSLEY. Biennial herbs cultivated for ornament and edible foliage.

Bacillus nelliæ Welles. Bacterial wilt of *P. sativum* (*P. hortense*) in the Philippines.

Bacillus petroselini Pot. Small yellow leaf spots with brown margins on *P. sativum* in Russia.

Fusicladium depressum (B. and Br.) Sacc. and var. *petroselini* Sacc. On leaves of *P. sativum* in France and Spain.

Gloeosporium achaenicolæ Rostr. Ashen-colored areas on fruit and peduncles of *P. sativum* and *Pastinaca sativa* in Denmark.

Macrosporium cheiranthi (Lib.) Fr. Circular, gray leaf spots on *P. sativum* in France.

Puccinia petroselini (DC.) Lindr. Cinnamon to dark-brown rust pustules on leaf blades and petioles and on stems of *P. sativum*, *Aethusa cynapioides*, and *Anethum graveolens* in Argentina and Europe.

Septoria petroselini Desm. and var. *segetum* P. Brun. Leaf spots on *P. hortense*, *P. sativum*, and *P. segetum* in Australia, Barbados, Argentina, Union of South Africa, and Europe.

PETUNIA. Small herbs cultivated for their showy flowers.

Entyloma petuniae Speg. Smut sori in pale indefinite spots, often occupying entire area of leaf blades of *P. nyctaginiiflora* in Argentina.

Phyllosticta petuniae Speg. Circular, then angular, dull-brown leaf spots on *Petunia* sp. in Great Britain.

Bamularia petuniae Cke. Large, somewhat circular, dull-yellow leaf spots on *Petunia* sp. in Great Britain.

PHAJUS. Epiphytic and terrestrial orchids. See *Orchidaceae*.**PHALAENOPSIS.** See *Orchidaceae*.

PHALARIS. REED CANARY GRASS. Stout, hardy, ornamental grasses.

- Entyloma brefdi* Krieger. Smut sori in leaf spots on *P. arundinacea* in Germany.
Heterosporium phragmites (Opiz.) Sacc. and var. *inflorescentiae* Bub. On leaves of *P. arundinacea* and *Phragmites communis* in France and Bohemia.
Leptosphaeria culmicola (Fr.) Karst. See *Bromus*.
Mycosphaerella chlorina (Cke.) Lind. On leaves of *P. arundinacea* in Great Britain.
Ophiobolus cariceti (B. and Br.) Sacc. See *Triticum*.
Phyllosticta crastophila Sacc. Dull-yellow areas on leaves of *P. arundinacea* and *Setaria verticillata* in Italy.
Physalospora phalaridis Zem. On *P. bulbosa* in Argentina.
Physoderma gerhardtii Schroet. See *Glyceria*.
Puccinia brevicornis Ito. Leaf rust on *P. arundinacea* in Japan.
Puccinia orchidearum-phalaridis Kleb. See *Orchidaceae*.
Puccinia schmidiana Diet. Yellow and powdery brown to black rust pustules on leaves of *P. arundinacea*, *Leucojum aestivum*, and *L. vernum* in Italy, Austria, and Germany.
Puccinia winteriana P. Magn. Yellow and brown to black rust sori on leaves of *P. arundinacea* and *Allium ursinum* in Spain, France, Switzerland, Great Britain, Denmark, Holland, and Germany.
Sclerospora kriegeria Magn. Downy mildew on leaves of *P. arundinacea* in Germany.
Sclerospora macrospora Sacc. See *Triticum*.
Septoria phalaridis Cocc. and Mor. Pale-yellow spots on leaves and sheaths of *P. brachystachys* in Italy.
Tilletia menieri Har. and Pat. Smut sori turning the ovaries of *P. arundinacea* into brown spore masses. Reported from France.

PHASEOLUS. BEAN. Lima-bean. Herbs cultivated as vegetables and forage plants.

- Acanthostigma heterochaeta* Syd. and Buhl. Brown or yellow-brown leaf spots on *Dumassia villosa*, *Dumassia* sp., *P. mungo*, and *P. radiatus* in India.
Ascochyta boltschauseri Sacc. Brown spots concentrically zoned and causing premature death of the leaves of *P. lunatus*, *P. vulgaris*, and *Vicia faba* in New Zealand, Russia, Bohemia, Switzerland, Holland, Norway, and Denmark.
Ascochyta bormullerii Syd. On leaves of *P. acutifolius* in Mexico.
Ascochyta phaseolorum Sacc. Indefinite ochraceous leaf spots on *P. mungo*, *P. multiflorus*, *P. nanus*, *P. radiatus*, *P. vulgaris*, *Vigna catjang*, and *Vicia faba* in Japan, Uganda, and Europe.
Asteroma phaseoli Brun. Felty-black spots on pods of *P. vulgaris* in the Philippines.
Cercospora lussoniensis Sacc. Leaf spot on *P. lunatus* in the Philippines.
Cercospora stuhlmanni P. Henn. Irregular dull-brown leaf spots on *P. vulgaris* in Tanganyika.
Cercospora zonata Wint. See *Vicia*.
Cercosporina caracallae Speg. Angular ashen leaf spots on *P. caracalla* in Argentina.
Cylindrosporium phaseoli Rabh. On leaves of *Phaseolus* sp. in Germany.
Diplodia phaseolina Sacc. Small black spots on ripening pods of *P. lunatus* and *P. vulgaris* in the Philippines.
Hypochnus cucumeris Frank. See *Cucumis*.
Leptosphaeria circinans (Fekl.) Sacc. See *Asparagus*.
Macrophoma phaseolina F. Tassi. Irregular pale-brown leaf spots on *P. ornithopus* in Italy.
Mosaic. A mosaic disease, said to resemble a similar disease on soy (*Sofa*), is reported from Japan and may be the same as the mosaic of legumes occurring in the United States.
Mycosphaerella phaseolicola Desm. Pale-reddish leaf spots on *P. vulgaris* in Japan and France.
Mycosphaerella phaseolorum Siem. On leaves of *P. mungo* and *Sofa maz* in Russia.
Phyllachora phaseoli (P. Henn.) Theiss. and Syd. Shiny black stromata on leaves of *P. adenanthus* and *Phaseolus* sp. in Porto Rico and Brazil.
Phyllachora phaseolina Syd. Circular black stromata on rust-brown leaf spots on *P. calcaratus* and *Vigna* sp. in the Philippines and Amboina.
Phyllosticta noackiana Allesch. Subcircular ochraceous leaf spots on *Phaseolus* sp. in Brazil.
Phyllosticta phaseolorum Sacc. and Speg. See *Vigna*.
Physopella concors Arth. Brown powdery rust pustules on leaves of *P. lunatus*, *Dolichos lablab*, and *Teramnus uncinatus* in Porto Rico.
Pleosphaerulina phaseoli Syd. On leaves of *P. semierectus* in the Philippines.
Rhizoctonia sp. See *Vigna*.
Rhizoctonia dimorpha Matz. Attacks leaves and stems of *Phaseolus* sp. and *Vigna catjang* in Porto Rico.
Rhizoctonia macrosclerotia Matz. Reported on petioles and stems of *Phaseolus* sp. in Porto Rico.
Sclerophoma phaseoli Karak. On pods of *P. multiflorus* in Russia.
Septoria phaseoli Maubl. Irregular greenish leaf spots on *P. mungo* and *Phaseolus* sp. in Russia and Brazil.
Uredo vignae Bres. See *Vigna*.
Uromyces phaseolicola Speg. Leaf rust on *P. prostratus* in Argentina.
- PHYGOPTERIS.** WOOD FERN.
Exoascus filicinus (Rostr.) Sacc. See *Polystichum*.
Uredinopsis filicina P. Magn. Yellow-brown rust pustules on leaves of *P. polypodioides* (*Dryopteris polypodioides*) in Europe.
- PHYLLODENDRON.** CORK TREE. Ornamental trees.
Ascochyta phelodendri Kab. and Bub. Round to irregular brown leaf spots with white centers on *P. amurense* in Bohemia.
Coleosporium phelodendri Kom. Fg. Golden rust sori on leaves of *P. amurense* in Chosen, Manchuria, and Japan.
- PHILADELPHUS.** MOCK ORANGE. *Syringa*. Ornamental shrubs.
Aecidium philadelphii Diet. Leaf rust on *P. coronarius* in Japan.
Ascochyta fuscescens Kab. and Bub. Indefinite dull-brown to ashen areas on leaves of *P. coronarius* in Bohemia.
Ascochyta philadelphii Sacc. and Speg. Subcircular ochraceous leaf spots on *P. coronarius* in Europe.
Phragmidium philadelphii Pass. Leaf rust on *P. coronarius* in Yugoslavia.
Phyllosticta coronaria Pass. Large brown leaf spots on *P. coronarius* in Russia, Italy, Belgium, and Denmark.
Stereum purpureum Pers. See *Prunus*.
- PHILIBERTIA.** Climbing shrubs.
Aecidium philibertiae P. Henn. Yellow rust sori on leaf blades and petioles of *P. flava* in Argentina.
Cercospora gillesii Speg. On leaves of *P. gillesii* in Argentina.
Napcladium asclepiadinum Speg. Pale indefinite leaf spots on *P. rotata* in Argentina.
Phyllosticta gillesii Speg. Circular definite whitish leaf spots on *P. gillesii* in Argentina.

- PHILLYREA.** Ornamental woody plants grown for their handsome evergreen foliage.
- Aecidium phillyreae** DC. Leaf rust on *P. latifolia* and *P. media* in Dalmatia and Portugal.
- Ascochyta bacillifera** Wint. Irregular dull-brown to gray leaf spots on *P. angustifolia* in Portugal.
- Fusicladium radiosum** (Lib.) Lind. var. **microsporum** Sacc. On leaves of *P. media* in Spain.
- Phyllosticta goritensis** Sacc. Circular dull-brown leaf spots on *P. angustifolia* and *P. latifolia* in Italy.
- Phyllosticta phillyreae** Sacc. Gray-ochraceous leaf spots with rufous margins on *P. latifolia* and *P. media* in Italy and France.
- Phyllosticta phyllicola** Rabh. On leaves of *Phillyrea* sp. in Germany.
- Phyllosticta romana** D. Sacc. Subcircular to oblong ashen leaf spots on *P. media* in Italy.
- Septoria crateriformis** (Dur. and Mont.) Sacc. On leaves of *P. latifolia* and *P. media* in France and Algeria.
- Septoria erythrostoma** (Dur. and Mont.) Sacc. On leaves of *Phillyrea* sp. in Algeria.
- Septoria phillyreae** Thum. Small gray leaf spots on *P. latifolia* in Portugal.
- Zaghouania phillyreae** Pat. Golden-yellow rust sori on leaf blades and petioles and on stems of *P. angustifolia*, *P. intermedia*, *P. latifolia*, *P. media*, and *P. vilmoriniana* in Great Britain, France, Corsica, Italy, Sicily, Dalmatia, Spain, Portugal, Austria, Tunis, and Algiers.
- PHILODENDRON.** Shrubby or tree-like ornamental aroids, usually climbing.
- Colletotrichum philodendri** P. Henn. Circular whitish leaf spots with dark brown margins on *P. krebsii* and *Philodendron* sp. in Brazil and Porto Rico.
- Gloeosporium aracearum** P. Henn. See Caladium.
- Mycosphaerella philodendri** (Pat.) Lind. Sunken white leaf spots with brown margins on *P. pertusum* in Venezuela.
- Phyllachora engleri** Speg. See Anthurium.
- Phyllosticta philodendri** Allesch. On leaves of *P. pertusum* in Brazil.
- Septoria aracearum** Sacc. Pale-brown leaf spots on *P. pertusum* in Italy.
- PHLEUM.** TIMOTHY. Hay and pasture grass.
- Dilophia graminis** (Fckl.) Sacc. See Triticum.
- Entyoma camusianum** P. Har. Smut sori in numerous circular to irregular dull-brown leaf spots on *P. arenarium* in France.
- Septoria culmifida** Lind. Small gray-white leaf spots with broad purple margins on *P. pratense*, *Arrhenatherum avenaceum*, *Dactylis glomerata*, and *Poa trivialis* in Denmark and Great Britain.
- PHLOMIS.** JERUSALEM SAGE. Stout herbs grown for their dense axillary whorls of large flowers.
- Ascochyta phlomidis** Jaap. On stems of *P. fruticosa* and *P. tuberosa* in Dalmatia and Russia.
- Erysiphe taurica** Lév. See Althaea.
- Ovularia phlomidis** Bub. and Wrobl. On leaves of *P. tuberosa* in Galicia and Russia.
- Phyllosticta cuestasae** Gz. Frag. On leaves of *P. herba-venti* in Spain.
- Puccinia excelsa** Barel. Dark-brown rust pustules on leaves of *P. lamiifolia* in India.
- Puccinia phlomidis** Thum. Powdery dull-brown rust sori on leaves of *P. armeniaca*, *P. brevilabris*, *P. canescens*, *P. herba-venti*, *P. nissoli*, and *P. tuberosa* in Siberia, Russia, Turkestan, Asia Minor, Italy, Spain, and Yugoslavia.
- Septoria barrasii** Gz. Frag. On leaves of *P. purpurea* in Spain.
- PHLOX.** Showy-flowered annual and perennial garden herbs.
- Phyllosticta decussata** Syd. Circular to irregular pale-yellow leaf spots with purple margins on *P. decussata* in Germany.
- Phyllosticta phlogis** Vesterg. Subcircular dirty-white leaf spots with brown margins on *P. drummondii* in Sweden.
- Pyrenochaeta phloxidis** Mass. The stems of cultivated *Phlox* are attacked just above the ground line, cankers resulting, with consequent death of the stems. Occurs in Great Britain.
- Septoria vogliniana** Sacc. and Trott. Irregular, often confluent, chestnut-brown rust sori on leaves of *P. drummondii* in Italy.
- PHOENIX.** DATE PALM. See Palmae.
- PHORMIUM.** FLAX LILY. NEW ZEALAND FLAX. Fiber plant.
- Heterosporium** sp. Large irregular sooty blotches on lower leaf surfaces of *P. tenax* in New Zealand rotting the fiber.
- Melampsora** sp. Leaf rust on *Phormium* sp. in New Zealand.
- Phaeoseptoria** sp. On leaves of *P. tenax* in New Zealand, destroying the fiber.
- Phyllosticta haematocycla** Berk. Broad pale-brown leaf spots with red borders on *P. tenax* in Germany and Portugal.
- Physalospora phormii** Schroet. On leaves of *P. tenax* in Germany.
- Ramularia phormii** Waters. The most serious disease of *P. tenax* in New Zealand is the so-called yellow-leaf disease attributed in part to the above fungus, although other causes have been assigned in times past. Diseased plants show a yellowing and dying of the leaves which soon involves and destroys the entire plant. The seat of the trouble is in the roots, which rot off.
- Septoria** sp. Longitudinal purple stripes on leaves of *P. tenax* in New Zealand, ruining the fiber.
- Septoria melanogramma** Tass. On *P. tenax* in Italy.
- PHOTINIA.** Ornamental woody plants grown for their attractive flowers and fruits.
- Corticium salmonicolum** B. and Br. See Citrus.
- Gloeosporium tremellinum** Gäum. Anthracnose on leaves of *P. notoniana* in Java.
- Guignardia photiniae** Alm. and Da Cam. On leaves of *Photinia* sp. in Portugal.
- Gymnosporangium photiniae** (P. Henn.) Syd. See Juniperus.
- Phyllosticta photiniae** Thum. Grayish-white leaf spots with red margins on *P. glabra* in Italy.
- Helicobasidium mompa** Tan. See Morus.
- PHRAGMITES.** Reed. Large grasses.
- Ascochyta arundinis** Fautr. and Lam. On leaves of *P. communis* in France.
- Ascochyta donacina** Sacc. On culms of *P. communis* and *Donax* sp. in Italy and France.
- Brachysporium phragmitis** Miy. On leaves of *P. communis* in China.
- Heterosporium phragmites** (Opiz.) Sacc. See Phalaris.
- Leptosphaeria eustoma** (Fr.) Sacc. See Scalea.
- Leptostromella phragmitis** Brun. On leaves of *P. communis* in Japan and France.
- Mycosphaerella lineolata** (Desm.) De N. See Calamagrostis.
- Napieladium arundinaceum** (Cda.) Sacc. On *P. communis* in Japan, Dalmatia, Switzerland, and Austria. Reported from Nebraska.
- Napieladium laxum** Bub. Long dark leaf spots with yellow surrounding areas on *P. communis* in Austria.
- Phyllosticta phragmitis** Nagorni. On leaves of *P. communis* in Russia.
- Physalospora festucae** (Lib.) Sacc. See Andropogon.
- Pirostoma circinans** Fr. On leaves, leaf sheaths, and stems of *P. communis* in Mongolia, France, Italy, Belgium, Sweden, and Germany.
- Puccinia invenusta** Syd. Yellow-brown to black rust sori on leaves of *P. karka* in India.
- Puccinia islaeae** (Thum.) Wint. See Arundo.

PHLAGMITES—Continued.

- Puccinia morioakaensis** S. Ito. Black rust pustules on leaves of *P. communis* and *P. longivalvis* in Japan.
- Puccinia obtusata** (Oth.) Ed. Fisch. Yellow and black rust pustules on leaves of *P. communis*, *Ligustrum vulgare*, and *L. thota* in Japan and Germany.
- Puccinia okatamaensis** S. Ito. Black linear rust sori on leaves of *P. communis* and *P. longivalvis* in Japan.
- Puccinia tepperi** Ludw. Leaf rust on *P. communis* in Australia.
- Puccinia trabutii** Roum. and Sacc. Leaf rust on *P. gigantea* and *Linaria reflexa* in Persia and French North Africa.
- Puccinia trallii** Plowr. Red-brown to black rust sori on leaves of *P. communis* and *Rumex acetosa* in Australia and Europe.
- Sclerospora macrospora** Sacc. See *Triticum*.
- Septoria arundinacea** Sacc. Oblong ochraceous leaf spots on *P. communis* in Italy and France.
- Septoria curva** Karst. On culms of *P. communis* in Finland.
- Septoria littoralis** Speg. and var. **culmicola** Gz. Frag. On leaves of *P. communis* in Spain and Italy.
- Septoria paludosa** Kab. and Bub. On leaves of *P. communis* in Bohemia.
- Septoria phragmites** Sacc. and var. **minor** Sacc. Subovate whitish leaf spots with dull-brown margins on *P. communis* in France, Spain, and Italy.
- Ustilago grandis** Fr. Smut sori at the nodes as black powdery spore masses, causing constrictions of the nodes and shortening of the internodes, on *P. communis* and *P. longivalvis* in Europe.
- Ustilago mirabilis** Sorok. Black powdery smut sori on *P. communis* in Russia.
- PHYLLANTHUS**. STAR GOOSEBERRY. OTABEITE GOOSEBERRY. Tropical trees and shrubs, some species cultivated for ornament and fruit.
- Accidium albicans** Arth. and Holw. Leaf rust on *P. acuminatus* in Guatemala and Costa Rica.
- Accidium luzoniense** P. Henn. Rust on circular yellow leaf spots on *Phyllanthus* sp. in the Philippines.
- Accidium phyllanthi** P. Henn. Leaf rust on *P. flexuosus* and *Phyllanthus* sp. in Indo-China, New Guinea, Japan, and Uganda.
- Accidium phyllanthinum** Syd. Leaf rust on *P. reticulatus* in India and the Philippines.
- Catacauma egregium** (Rehm.) Theiss. and Syd. Black stromata on leaves of *Phyllanthus* sp. in Brazil.
- Catacauma phyllanthopilum** P. Henn. Black stromata on leaves of *P. brasiliensis* and *Phyllanthus* sp. in Peru and the Philippines.
- Phakopsora phyllanthi** Diet. Brown leaf rust on *P. acidus* (*P. distichus*) and *Phyllanthus* sp. in India and the Philippines.
- Ravenelia appendiculata** Lagh. and Diet. Rust on leaves of *P. acuminatus*, *P. galeottianus*, and *Phyllanthus* sp. in Ecuador, Mexico, and Guatemala.
- Ravenelia emblicae** Syd. Brown rust pustules on leaves of *P. emblica* and *P. polyphyllus* in India, Burma, and Ceylon.
- Ravenelia pygmaea** Lagh. and Diet. Dark-brown rust pustules on deformed petioles and stems of *Phyllanthus* sp. in Ecuador.
- Schroeteriaster fenestrala** Arth. Brown rust pustules on leaves of *P. acidus* (*P. distichus*) in Porto Rico.
- Schroeteriaster ulei** Syd. Leaf rust on *Phyllanthus* sp. in Brazil.
- Uredo phyllanthi** P. Henn. Leaf rust on *P. conami* and *Phyllanthus* sp. in Trinidad and Brazil.
- Uredo phyllanthi-longifolii** Petch. Leaf rust on *P. longifolius* in Ceylon.
- Uredo phyllanthi-reticulati** Petch. Leaf rust on *P. reticulatus* in Ceylon.
- PHYLLAUREA**. See *Codiaeum*.
- PHYLLOSTACHYS**. Bamboo. See *Bambuseae*.
- PHYMOSIA**. See *Sphaeralcea*.
- PHYSALIS**. GROUND CHERRY. HUSK TOMATO.
- Ascochyta alkekengi** Massal. Subcircular to angular dull-brown concentrically zoned leaf spots on *P. alkekengi* in Italy.
- Ascochyta physalicola** Oud. On stems of *P. alkekengi* in Holland.
- Ascochyta physalina** Sacc. Dull-yellow leaf spots on *P. alkekengi* in Italy.
- Cercosporina physalidicola** Speg. Indefinite, pale-brown leaf spots on *P. viscosa* in Argentina.
- Phyllosticta physaleos** Sacc. and var. **calycicola** Speg. Subcircular, ochraceous leaf spots with brown margins on *P. alkekengi*, *P. peruviana*, and *P. viscosa* in Ceylon, Argentina, Russia, and Italy.
- Vermicularia varians** Duc. See *Solanum*.
- PHYSOCARPUS. OPULASTER**. Ag. NINEBARK. Ornamental shrubs with white flowers, attractive inflated pods, and bright green foliage.
- Phyllachora physocarp** Jacz. Black stromata on small red leaf spots on *P. amurensis* in Siberia.
- PHYSOSIPHON**. Epiphytic orchids. See *Orchidaceae*.
- PHYTEUMA**. HORNED RAMPION. Hardy herbaceous perennials used in borders and alpine gardens.
- Cercospora phytumatis** (Frank.) Sacc. Black leaf spots with white centers on *P. spicatum* in Denmark and Germany.
- Montagnellina stellaris** (Pers.) Theiss. and Syd. Black stromata on leaves of *P. scheuchzeri* and *P. spicatum* in Europe.
- Peronospora phytumatis** Fekl. Downy mildew on leaves of *P. betonicaefolium*, *P. nigrum*, *P. orbiculare*, and *P. spicatum* in Switzerland, Norway, Denmark, Austria, Russia, and Germany.
- Phyllosticta phytumatis** Gz. Frag. Leaf spot on *P. spicatum* in Spain.
- Pyrenopeziza phytumatis** Fekl. On leaves of *Phyteuma* sp. in Switzerland.
- Ramularia phytumatis** Sacc. and Wint. Ochraceous leaf spots with brown margins on *P. betonicaefolium*, *P. halleri*, *P. hedraeanthifolium*, *P. michelii*, *P. nigrum*, *P. orbiculare*, and *P. spicatum* in Switzerland, Italy, Spain, Bohemia, Austria, and Germany.
- Septoria phytumatis** Siegm. Pale leaf spots with dull-brown margins on *Phyteuma* sp. in Italy.
- Septoria phytumatum** Sacc. On leaves of *Phyteuma* sp. in Austria.
- Uromyces caricis-sempervirentis** Ed. Fisch. Aerial stage on *P. betonicaefolium* and *P. orbiculare* in Switzerland, Austria, and Germany. Other stages on *Carex* spp.
- Uromyces phytumatum** (DC.) Ung. Powdery brown rust pustules on leaves of *P. betonicaefolium*, *P. halleri*, *P. hemisphaericum*, *P. michelii*, *P. nigrum*, *P. orbiculare*, and *P. spicatum* in Europe.
- PHYTOLACCA**. POKEBERRY. POKEWEED. Herbs and woody plants, some species ornamental.
- Ascochyta phytolacca** Sacc. and Scalia. Marginal dull-brown leaf spots on *P. americana* (*P. decandra*) in Portugal.
- Heterosporium dalmaticum** Jaap. On stems of *P. americana* in Dalmatia.
- Phyllosticta phytolaccicola** Brun. Small circular brown leaf spots on *P. acinosa* in France.
- Ramularia harai** P. Henn. Effuse dull-brown leaf spots on *P. acinosa* in Japan.
- Septoria patouillardii** Sacc. and Syd. Circular dull-brown leaf spots on *Phytolacca* sp. in Ecuador.
- Septoria phytolacca** Cav. Yellow leaf spots on *P. americana* (*P. decandra*) in Russia, Yugoslavia, and Italy.

PICEA. SPRUCE. Ornamental and timber trees.

- Ascochyta piniperda** Lindau. Attacks leaves and twigs of *P. abies*, *P. sitchensis*, and *Tsuga* sp. in Belgium, Russia, Germany, and Austria. The fungus also causes a damping-off of seedlings. Reported from North Carolina on *Picea* spp.
- Barclayella deformans** (Barcl.) Diet. Elongate yellow rust pustules on needles and young twigs of *P. smithiana* (*P. morinda*) in India.
- Chrysomyxa abietis** (Walbr.) Ung. Small elongate yellow rust pustules on yellow or reddish spots on needles of *P. engelmannii*, *P. abies*, *P. hondoensis*, and *P. pungens* in Japan and Europe.
- Chrysomyxa rhododendri** DeBy. See *Rhododendron*.
- Coccomyces piceae** (Fekl.) Rehm. On needles of *P. abies* in Germany.
- Cucurbitaria piceae** Borthw. Thin black crustlike stromata enveloping buds of *P. pungens* in Scotland.
- Dasyscypha abietis** Sacc. On branches of *P. abies* in Finland.
- Dasyscypha calycina** (Schum.) Fekl. See *Larix*.
- Fabraea abietina** Sacc. On the upper surfaces of needles of *P. abies* in Italy.
- Gloeosporium conigenum** Sacc. and Roum. Anthracnose on cone scales of *P. abies* in France.
- Hypodermella macrosporum** (Hart.) Lagerh. Black linear fruiting bodies on needles of *P. abies* in Switzerland, Bohemia, and Germany.
- Lophodermium abietis** Rostr. Black areas on needles of *P. glauca* (*P. canadensis*), *P. abies*, *P. smithiana* (*P. morinda*), *P. sitchensis*, *Pseudotsuga taxifolia*, and *Taxus baccata* in Sweden and Denmark.
- Macrophoma excelsa** (Karst.) Berl. and Vogl. var. *infestans* Ohl. On needles of *P. abies* in Russia.
- Myxosporium abietinum** Rostr. See *Larix*.
- Oospora abietum** Oudem. See *Abies*.
- Peridermium piceae** Barcl. Rust on needles of *P. smithiana* in India.
- Peridermium piceae-hondoensis** Diet. Rust on needles of *P. hondoensis* in Japan.
- Peridermium thomsonii** Berk. Rust on needles of *P. smithiana* in India.
- Phlyctaena strobilina** Karst. and Har. On cone scales of *P. abies* in France.
- Phomopsis occulta** Trav. On twigs of *P. abies* in Denmark, France, and Great Britain.
- Physalospora abietina** Prill. and Delacr. See *Abies*.
- Phytophthora fagi** R. Hart. See *Fagus*.
- Sclerotopsis piceana** (Karst.) Died. On twigs and needles of *Picea* sp., *Abies alba* and *A. nordmanniana* in Europe.
- Thekopsora areolata** (Walbr.) P. Magn. See *Prunus*.
- PIERIS.** Ornamental shrubs grown for their handsome white flowers.
- Exobasidium pieridis** P. Henn. Forms concave galls on leaves of *P. ovalifolia* in Japan and India.
- Phyllosticta andromedae** West. On leaves of *P. floribunda* and *Zenobia cassinefolia* in Belgium.
- Rhytisma piceum** Berk. Circular black stromata on leaves of *P. ovalifolia* in Indo-China and India.
- Rhytisma pieridis** Pat. Black irregular fruiting areas on leaves of *P. ovalifolia* in Japan.
- PILEA.** Mostly weedy herbs, one or two species cultivated as curiosities.
- Uredo pileae** Barcl. Leaf rust on *P. trinervia* in India.
- PILOCARPUS.** Shrubs or small trees.
- Phyllosticta pilocarpi** Pat. Circular white leaf spots with red margins on *P. pinnatifolius* in Brazil.
- Puccinia pilocarpi** Cke. Black rust pustules on leaves of *P. pinnatus*, *P. pinnatifolius*, and *P. seloanus* in Brazil, Argentina, and Paraguay.
- PINGICULA.** BUTTERWORT. Small acaulescent herbs.
- Ustilago pingicululae** Rostr. Smut sori in anthers of *P. alpina* and *P. vulgaris* in Denmark, Estonia, and Switzerland.
- PINUS.** PINE. Ornamental and timber trees.
- Bacterium** sp. Bacteria have been found associated with a witches' broom disease of *P. pinaster* (*P. maritima*) in France.
- Bacillus pini** Vuill. Causes galls on branches of *P. halepensis* in France.
- Brachysporium pini-insularis** P. Henn. On needles of *P. insularis* in the Philippines.
- Cercospora pini-densiflorae** Hori. and Nambu. Yellow-brown spots, usually on the upper half of needles of *P. densiflora* in Japan. The disease starts in the upper part of a plant and works down, killing it in a short time. Serious in nursery beds.
- Coleosporium asterum** (Diet.) Syd. See *Aster*.
- Coleosporium caecaliae** Oth. See *Caecalia*.
- Coleosporium euphrasiae** (Schum.) Wint. Rust on needles of *P. montana* and *P. sylvestris* in Tibet, Siberia, and Europe. The alternate stage occurs on *Alectrolophus* spp. and *Euphrasia* spp.
- Coleosporium inulae** Rabh. See *Inula*.
- Coleosporium melampyri** Tul. Rust on needles of *P. montana* and *P. sylvestris* in Japan and Europe. Alternate stage on *Melampyrum* spp.
- Coleosporium petasites** Lév. Yellow rust pustules on needles of *P. sylvestris* in Japan, Siberia, and Europe. Alternate stage on leaves of *Petasites* spp.
- Coleosporium pulsatillae** Lév. See *Anemone*.
- Coleosporium senecionis** Fr. See *Senecio*.
- Coleosporium sonchi** Lév. See *Sonchus*.
- Coleosporium tussilaginis** (Pers.) Lév. Yellow rust pustules on needles of *P. sylvestris* in Europe. Alternate stage on *Tussilago farfara*.
- Cronartium flaccidum** (Alb. and Schw.) Wint. See *Paeonia*.
- Cronartium ribicola** Fisch. The blister rust of white pine (*P. strobus*) was introduced with nursery stock into the United States from 1900 to 1910 and now threatens the existence of this important timber tree in all parts of the country. All five-needled pines, including *P. albicaulis*, *P. armandi*, *P. aristata*, *P. balsamiana*, *P. cembra*, *P. griffithii* (*P. excelsa*), *P. flexilis*, *P. koraiensis*, *P. lambertiana*, *P. monticola*, and *P. peuce*, are susceptible. The uredo and telial stages occur on species of *Ribes* and *Grossularia*, practically all species being subject to attack. The disease occurs throughout Europe, in Japan, Siberia, and western Canada.
- On the pine the fungus causes swellings on the bark of trunk, branches, and twigs, and after several years breaks through to the surface as blisterlike orange spore masses. The areas of infection may vary from one to many hundred, often so numerous as to form long continuous cankers. Trees of all ages are susceptible and are destroyed by girdling when the infections become numerous. Small trees are killed in from 4 to 6 years. The long incubation period of the mycelium in the tissues without external symptoms, other than the swelling of the bark, makes difficult or impossible the elimination of diseased plants by inspection. The uredo and telial stages occur on the lower leaf surfaces of the *Ribes*, the former appearing as powdery yellow pustules, the latter consisting of numerous small brown, hairlike projections.

PINUS—Continued.

- Cucurbitaria pithyophila** (Schm. et Kunze) De N. Parasitic on branches of *P. sylvestris* and *Abies alba* (*A. pectinata*) in Great Britain and continental Europe. In Japan the fungus causes a serious disease of *A. firma* called millet-canker. Cankers form on trunks and larger branches and finally take on a gnarled appearance. The bark over these areas cracks and numerous small granules form beneath.
- Cylindrosporium aculeolum** Bres. On needles of *P. sylvestris* in Germany.
- Cytosporina septospora** Dor. Yellow blotches on needles of *P. montana* in Russia.
- Dasyascypha calycina** (Schuem.) Fckl. See Larix.
- Diplodia pinea** Kickx. A wound parasite attacking terminal shoots of *Pinus* spp. in Europe and the Union of South Africa.
- Dothichiza pini** Sacc. On bark of *P. sylvestris* in Germany.
- Fusarium blasticola** Rostr. Causes a damping-off of coniferous seedlings, including *P. montana* and *Cryptomeria japonica* in Formosa, Denmark, and Germany. Dark patches with a water-soaked appearance occur on the stems, followed by collapse of the infected plants.
- Hypoderma pinicola** Brunch. Brownish or yellow-gray areas on needles of *P. sylvestris* in Europe, often causing defoliation.
- Hypoderma brachysporium** (Rostr.) Tub. (*Hypoderma strobilicola* Brunch.) Black linear fruiting bodies causing yellowing and premature fall of needles of *P. strobus* in Europe. Reported from Pennsylvania.
- Hypodermella sulcigena** (Duby.) Tub. Causes browning and premature shedding of needles of *P. pinaster*, *P. montana* (*P. pumila*), and *P. sylvestris* in northern Europe.
- Lachnella pini** Brun. Injures twigs of *Pinus* sp. in Scandinavia.
- Leptostroma pinastri** Desm. Needles turn red and fall. The disease is reported capable of destroying whole plantations of *P. pinaster*, *P. montana*, and *P. sylvestris* in Italy, France, Belgium, Austria, and Germany.
- Leptothyrella mougeotiana** Sacc. and Roum. On needles of *Pinus* sp. in France.
- Lophodermium gilvum** Rostr. On needles of *P. nigra* (*P. austriaca*) in Denmark.
- Melampsora pinitorqua** Rostr. See Populus.
- Myosphaerella pinifolia** Duc. On needles of *P. pinaster* in France.
- Myxosporium abietinum** Rostr. See Larix.
- Peridermium brevius** Barcl. Needle rust on *P. griffithii* (*P. excelsa*) in India.
- Peridermium complanatum** Barcl. Two forms of this rust attacking *P. excelsa* and *P. longifolia* in India are recognized; f. *acicola* is a leaf form, while f. *corticola* occurs on the trunk and limbs causing cankers similar to those produced by *Peridermium strobii* on *P. strobus*. It is thought that the alternate host may be *Crataegus* or *Rosa*.
- Peridermium corticola** Link. Rust on branches of *P. halepensis* in Dalmatia.
- Peridermium guatemalense** Arth. and Kern. Rust on needles of *P. filifolia* in Guatemala.
- Peridermium japonicum** Syd. Needle rust on *P. thunbergii* in Japan.
- Peridermium kuriense** Diet. A rust producing fusoid swellings on branches of *P. pumila* in Japan.
- Peridermium orientale** Cke. Needle rust on *P. griffithii* and *P. longifolia* in India.
- Peridermium pini-thunbergii** Diet. Yellow masses of rust spores on needles of *P. massoniana* and *P. thunbergii* in Japan.
- Peridermium praelongum** Syd. Needle rust on *P. thunbergii* in Japan.
- Phacidium infestans** Karst. Black stromata on needles of *Pinus* spp. in northern Europe. Premature needle fall results.
- Phomopsis pitya** (Sacc.) Grove. See Pseudotsuga.
- Phyllosticta strobiligena** Desm. On cone scales of *P. strobus* in Germany.
- Phytophthora fagi** R. Htg. See Fagus.
- Phytophthora omnivora** DeBy. See Fagus.
- Septoria acuum** Oud. On needles of *P. sylvestris* in Russia and Holland.
- Sphaeropsis necatrix** Petri. and Adan. Destroys seeds in the cones of *P. pinea* in Italy.
- PIPEX.** PEPPER. CUBEB. Trees, shrubs, and a few herbs, erect or climbing, with aromatic or pungent properties.
- Actinodothis piperis** Syd. Black carbonaceous stromata on leaves of *P. retrofractum*, *P. sarmentosum*, and *Piper* sp. in the Philippines and Ambona.
- Cercospora pipericola** Sacc. and Syd. Subcircular spots on leaves of *P. hispidum* in Jamaica.
- Cercospora piperis** Pat. Circular brown leaf spots on *Piper* sp. in Ecuador.
- Cercospora portoricensis** Earle. Numerous dull-brown angular leaf spots on *P. aduncum*, *P. hispidum*, *P. peltatum*, and *P. umbellatum* in Salvador, Porto Rico, Cuba, and Haiti.
- Colletotrichum necator** Mass. Anthracnose on *Piper* sp. in Malaya.
- Colletotrichum piperis** Petch. Anthracnose on *P. betle* and *P. nigrum* in Ceylon.
- Corticium salmonicolor** B. and Br. See Citrus.
- Cyclodithis pulchella** Syd. Stromata in circles on brown leaf spots on *P. carylistachyum* and *P. medium* in Porto Rico and the Philippines.
- Oidium** sp. Powdery mildew on leaves of *P. betle* in Ceylon.
- Phyllosticta pipericola** Syd. Angular pale-brown leaf spots on *P. nigrum* in Java.
- Phyllosticta piperis** Tassi. Dull-brown leaf spots on *P. betle* and *P. longum* in Italy and Ceylon. The same name has been given to a species on *P. nigrum* in Java by Hennings.
- Physalospora piperina** Syd. On stems of *P. nigrum* in India.
- Ramularia piperis** P. Henn. Circular to angular brown to black leaf spots on *Piper* sp. in Congo.
- Rhizoctonia destruens** Tassi. See Solanum.
- Rosellinia bunodes** B. and Br. See Citrus.
- Septoriopsis piperis** Stev. and Dal. Irregular dirty-white leaf spots with purple-brown borders on *P. medium* in Porto Rico.
- Stigmatea piperis** Rehm. On leaves of *Piper* sp. in Brazil.
- Uredo piperis** P. Henn. Brown rust pustules on circular dark-brown leaf spots on *Piper* sp. and *Peperomia hernandifolia* in Porto Rico, Peru, and Brazil.
- PIPTADENIA.** Leguminous trees similar to Acacia.
- Endodothella piptadeniae** (P. Henn.) Theiss. and Syd. Black shiny stromata on leaves of *Piptadenia* sp. in Peru and Brazil.
- Phyllachora piptadenicola** Speg. Tar spot on leaves of *P. rigida* in Argentina.
- Puccinia papulifera** Syd. Leaf rust on *P. latifolia* in Brazil.
- Puccinia piptadeniae** P. Henn. Brown rust pustules on leaves of *Piptadenia* sp. in Brazil.
- Ravenelia cebil** Speg. Leaf rust on *P. macrocarpa* and *P. peregrina* in Porto Rico and Argentina.
- Ravenelia henningsiana** Diet. Yellow-brown to dark-brown rust pustules on leaves of *Piptadenia* sp. in Brazil.
- Ravenelia simplex** Diet. Leaf rust on *P. communis* in Brazil.
- Ravenelia vilis** Syd. Leaf rust on *Piptadenia* sp. in Brazil.

FISCIDIA. ICHTHYOMETHIA Ag. Fish poison tree.

Cercospora piscicidae P. Henn. Dull yellowish-brown spots on leaves of *P. erythrina* in Jamaica.

ISONIA. Erect, armed trees or shrubs.

Aecidium pisoniae Arth. and Johnst. Leaf rust on *P. aculeata* in Cuba.

ISTACIA. PISTACHE. Trees or shrubs.

Coryneum pistaciae Pat. Dull-brown leaf spots on *P. lentiscus* in Tunis.

Mycosphaerella pistaciae Cke. On leaves of *P. lentiscus* in Italy.

Phyllosticta lentisci (Pass.) Allesch. and var. **maculicola** Bub. Small, ashen leaf spots on *P. lentiscus* in Italy and Montenegro.

Phyllosticta terebinthi Pass. Large irregular dull-brown to gray leaf spots on *P. terebinthus* in Italy.

Pileolaria terebinthi (DC.) Cast. Leaf rust on *P. atlantica* and *P. terebinthus* in Dalmatia and French North Africa.

Septoria pistaciae Desm. (*Phleospora pistaciae* Petr.). Black irregular leaf spots on *P. lentiscus* and *P. vera* in Asia Minor, Russia, Albania, Yugoslavia, and Italy.

Uromyces terebinthi (DC.) Wint. Leaf rust on *P. falcata*, *P. lentiscus*, *P. mutica*, *P. terebinthus*, and *P. vera* in Europe, Asia Minor, Persia, and Kurdistan.

PISTIA. WATER LETTUCE. Tropical duckweed.

Phyllosticta stratiotes Tassi. Ochraceous indefinite leaf spots on *P. stratiotes* in Italy.

PISUM. PEA. Annual or perennial herbs cultivated as food and forage plants.

Aecidium pisi-formosii Syd. Leaves and stems of *P. formosum* in Persia covered with yellow rust sori.

Bacillus leguminiperdus V. Oven. Produces brown, sunken, irregular areas on pods of *P. sativum* in Germany. Has been found capable by inoculation of attacking pods of *Lupinus* spp. and other legumes.

Bacterium seminum Cayley. This bacterial disease of the pea (*P. sativum*) reported from Great Britain, is carried in the seed and is present in most parts of the plant without visible signs until flowering time. The stems then become brown with a water-soaked appearance, brown linear streaks appear at the bases of petioles, the leaves are spotted yellow, and the pods have an abnormal roughened appearance.

Brachysporium pisi Oudem. Black moldy patches occur on the leaves of *P. sativum*, which turn yellow and die, causing a seedling blight, in Holland.

Cercospora pisa-sativae Stevenson. Small circular yellow to brown leaf spots with darker margins on *P. sativum* in Porto Rico and Cuba.

Marsonia deformans Cke. and Mass. Discolored spots on leaf blades, stipules and petioles of *P. sativum* in Australia.

Peronospora pisi Syd. Downy mildew on leaves of *P. arvense* and *P. sativum* in India, France, Holland, Switzerland, Russia, Scandinavia, Bohemia, and Germany.

Phoma sp. Causes a leaf spot on *P. sativum* in China.

Phyllosticta pisi West. Subcircular to oblong dull-brown or yellow leaf spots with darker margins on *P. sativum* in Belgium.

Rhizoctonia napi West. See Brassica.

Tylenchus dipsaci Kuehn. See Narcissus.

Uromyces pisi (Pers.) DeBy. Cinnamon-brown to dark brown rust pustules on leaves of *P. arvense*, *P. sativum*, *Lathyrus angustifolius*, *L. heterophyllus*, *L. latifolius*, *L. nissolia*, *L. pisiiformis*, *L. platyphyllus*, *L. pratensis*, *L. sativus*, *L. setifolius*, and *L. sylvestris* in Europe, Japan, Siberia, India, and New Zealand. The acial stage occurs on species of *Euphorbia*. Occurs sparingly in the Middle West.

Ustilago entorrhiza Schroet. A doubtful species of root smut reported on *P. sativum* in Germany.

PITCAIRNIA. Bromeliads.

Puccinia pitcairniae Lagh. Rust on leaves of *P. palmerum* and *Pitcairnia* sp. in Ecuador and Brazil. Reported from New Mexico.

Septoria pitcairniae Syd. Yellow-brown leaf spots on *Pitcairnia* sp. in Brazil.

PITHECOCTENIUM. MONKEY COMB. Ornamental vines cultivated for their showy flowers.

Prospodium amphiphilii (Diet. and Holw.) Arth. Leaf rust on *P. echinatum*, *P. hexagonum*, *P. muricatum*, and *Amphilophium molle* in Mexico, Isle of Pines, and Costa Rica.

Puccinia cordobensis P. Henn. Rust on branches of *P. cynanchoides* in Argentina.

Puccinia depallens Arth. and Holw. Leaf rust on *P. muricatum* in Costa Rica and Guatemala.

Puccinia medusa Speg. Powdery rust pustules on leaves of *P. clematideum* in Argentina.

Puccinia phiyetopus Syd. Brown to black powdery rust pustules on leaves of *P. hexagonum* in Mexico.

Puccinia pithecoctenii Pazschke. Dull-brown to dark-brown rust pustules on leaves of *Pithecoctenium* sp. in Brazil.

PITHECOLOBIUM. Tropical trees and shrubs planted for ornament and shade.

Corticium sp. See Thea.

Corticium salmonicolor B. and Br. See Citrus.

Cylindrodadium pithecolobii Petch. On leaves of *P. saman* in Ceylon.

Diorehidium acanthostephium Syd. Leaf rust on *Pithecolobium* sp. in Brazil.

Gloeosporium pithecolobii Koord. Large irregular gray-brown spots on leaves of *P. lobatum* in Java.

Maravalia pallida Arth. and Thaxt. Leaf rust on *P. latifolium* in Trinidad.

Microstoma pithecolobii Lamk. White irregular spots on lower leaf surfaces of *P. saman* in Porto Rico.

Montagnellina pithecolobii (Racib.) v. Hoeh. Black stromata on yellowish leaf spots on *P. lobatum* in Java.

Phyllosticta pithecolobii Young. (*P. pithecolobii-monensis* Young.). More or less circular pale-yellow leaf spots with dark-brown slightly raised margins on *P. unguis-cati* in Porto Rico.

Ravenelia amazonica Syd. Leaf rust on *P. corymbosum* in Brazil.

Ravenelia minuta Syd. Leaf rust on *Pithecolobium* sp. in Brazil.

Ravenelia pileolarioides Syd. Leaf rust on *Pithecolobium* sp. in Brazil.

Septoria pithecolobii Roum. Leaf spot on *Pithecolobium* sp. in Paraguay.

Stigmachora controversa (Starb.) Theiss. and Syd. Shiny, black stromata on leaves of *P. scalare* and *Pithecolobium* sp. in Bolivia and Argentina.

Uredo bomfimensis P. Henn. Brown rust sori on yellow circular leaf spots on *Pithecolobium* sp. in Brazil.

Uredo pithecolobii P. Henn. Rust deforming twigs of *Pithecolobium* sp. in Brazil.

Uredo raciborskii Sacc. and Syd. Brown leaf rust on *P. lobatum* in Java.

Uromyces albescens Syd. Brown rust pustules on light-brown leaf spots on *P. glomeratum* in Brazil.

- PITTSOPORUM.** Hardy or half-hardy evergreen woody plants.
Helicobasidium tanakae Miy. See *Morus*.
Mycosphaerella rubiginosa (Cke.) Lind. On leaves of *P. rubiginosum* in Australia.
Phyllosticta pittospori P. Brun. Subcircular to angular light-brown leaf spots on *P. tobira* in France and Italy.
Physalospora pittospori d'Alm. and Cam. Brown apical leaf spots on *Pittosporum* sp. in Portugal.
Septoria pittospori P. Brun. Circular deep-brown leaf spots on *P. tobira* in France and Italy.
Trabutia pittospori (P. Henn.) Theiss. and Syd. Irregular black opaque stromata on leaves of *P. abyssinicum* in Abyssinia.
Uredo pittospori P. Henn. Rust on leaves of *P. abyssinicum* in Abyssinia.
- PLACEA.** Amaryllislike bulbous plants from Chile.
Puccinia placeae Diet. and Neg. Yellow to dark-brown rust pustules on leaves of *Placea* sp. in Chile.
- PLAGIANTHUS.** Trees or shrubs.
Puccinia plagianthis McAlp. Brown rust pustules on leaves and flower stalks of *P. sidoides* in Tasmania.
- PLATANUS.** PLANE TREE. Erroneously called sycamore. Shade trees.
Guignardia cylindrica (Sacc. and Speg.) Lind. On leaves of *P. orientalis* in Italy and Denmark.
Phyllosticta platani Sacc. and Speg. On leaves of *P. orientalis* in Russia and Italy.
Microstroma platani Eddel. and Eng. Small circular grayish-white leaf spots on *P. occidentalis* in Sweden and Germany.
- PLATYCODON.** BALLOON FLOWER. Perennial, glaucous, erect herbs used in border plantings.
Septoria platycodonis Syd. Circular to irregular, white leaf spots on *P. grandiflorum* in Japan.
- PLECTOCOMIA.** East Indian climbing palms. See *Palmae*.
- PLECTRANTHUS.** Herbs and subshrubs with blue to lilac flowers.
Aecidium plectranthi Barcl. Rust on leaves of *P. coetsa*, *P. glaucocalyx*, and *P. scrophularioides* in Japan and India.
Coleosporium plectranthi Barcl. Yellow to golden rust pustules on leaves of *P. excisus*, *P. gerardinus*, *P. glaucocalyx*, *P. inflexus*, *P. longitubus*, *P. trichocarpus*, and *Ocimum* sp. in Japan and India.
Puccinia luandensis Syd. Brown powdery rust pustules on leaves of *P. laxiflorus* and *P. nudiflorus* in the Union of South Africa.
Puccinia plectranthi Thuen. Brown leaf rust on *P. laxiflorus* in the Union of South Africa.
- PLECTRONIA.** Woody plants cultivated to their flowers.
Aecidium baumanum P. Henn. Leaf rust on *P. abbreviata* in the Union of South Africa.
Aecidium busseanum P. Henn. Rust on red-brown leaf spots on *P. hispida* in Tanganyika.
Aecidium incomparabile Syd. Rust forming galls on leaves of *P. arnoldiana* in the Congo.
Aecidium plectroniae Cke. Leaf rust on *P. barbata* and *P. guinezi* in Natal and Queensland.
Hemileia canthii B. and Br. Golden rust pustules on leaves of *P. campanulata*, *P. ciliata*, *P. huillensis*, *P. parviflora*, and *P. peduncularis* in Ceylon, India, Philippines, and the Union of South Africa.
Uredo detergibilis (Thuen.) P. Henn. Rust on leaves of *P. huillensis* in the Union of South Africa.
- PLEUROSPERMUM.** Biennial or perennial hardy herbs.
Phyllosticta pleurospermi Died. Angular gray leaf spots on *P. austriacum* in Germany.
- PLEUROTHALLIS.** See *Orchidaceae*.
- PLUMBAGO.** LEADWORT. Subshrubs or herbs cultivated for their phloxlike flowers.
Ascochyta plumbaginicola P. Henn. On stems of *P. europaea* in Italy and Germany.
Ascochyta plumbaginis Sacc. Ochraceous-brown leaf spots on *P. europaea* in Italy and Germany.
Cercospora plumbaginea Sacc. and D. Sacc. Subcircular olivaceous, then light-buff, leaf spots on *P. europaea* in Italy.
Phyllosticta larpentae Tassi. Subcircular ashen-white leaf spots on *P. larpentae* (*Cerastostigma plumbaginoides*) in Italy.
- PLUMERIA.** FRANGIPANI. Tropical trees with milky juice grown for their showy flowers.
Ascochyta plumeriae P. Henn. On *Plumeria* sp. in Brazil.
Coleosporium dominicensis (Berk.) Arth. Golden rust pustules on leaves of *P. lutea*, *P. rubra*, and *Plumeria* sp. in Guatemala, Peru, and the West Indies.
Coleosporium plumeriae Pat. Golden rust pustules on leaves of *P. alba*, *P. emarginata*, *P. krugii*, *P. obtusa*, and *P. rubra* in Cuba, Porto Rico, Mexico, Guatemala, and Guadeloupe.
Phyllosticta plumeriae Tassi. Small gray to white leaf spots on *P. alba* in Italy.
Septoria plumeriae Sacc. and Syd. Circular light-brown zoned leaf spots on *P. obtusa* in Central America and the Bahamas.
- POA.** BLUE GRASS. Perennial grasses, some species cultivated as lawn, forage, and ornamental grasses.
Ascochyta kerguelensis P. Henn. On leaves of *P. kerguelensis* (*Festuca scoparia*) in the Kerguelen Islands.
Cladochytrium graminis Buesgen. See *Festuca*.
Claviceps setulosa Sacc. Sclerotia in ovaries of *Poa* sp. in Switzerland and France.
Dilophia graminis (Fekl.) Sacc. See *Triticum*.
Entyloma crepinianum Sacc. and Roum. Doubtful smut species on *Poa* sp. in France.
Hadtrochium virescens Sacc. and Roum. var. *poae* Sacc. Oblong brown leaf spots on *Poa* sp. in France, Germany, and Austria.
Leptosphaeria sparsa (Fekl.) Sacc. On leaves and culms of *Poa* sp., *Agrostis* sp., and *Calamagrostis* sp. in Switzerland.
Phyllachora poae (Fekl.) Sacc. Black stromata on leaves of *P. alpina*, *P. nemoralis*, *P. pratensis*, and *P. sudetica* in central Europe.
Phyllosticta kerguelensis P. Henn. On leaves of *P. cookii* in the Kerguelen Islands.
Protomyces rhizobius Trail. Small root nodules on *P. annua* in Scotland.
Puccinia cognatella Bub. Yellow to dark-brown rust pustules on elongate yellow spots on *P. nemoralis* in Bohemia.
Puccinia exigua Diet. Leaf rust on *P. annua* in Brazil.
Puccinia poae-trivialis Bub. Yellow to black rust pustules on leaves of *P. trivialis* in Bohemia.
Puccinia subandina Speg. Leaf rust on *P. chorizantheae* in Argentina.
Roumegueria gangraena (Fr.) Sacc. On leaves of *P. bulbosa*, *P. nemoralis*, and *P. pratensis* in Spain, Italy, and Tripoli.
Scolecotrichum compressum Allesch. On leaves of *P. compressa* in Germany.
Septoria culmifida Lind. See *Phleum*.
Septoria macrosperma Speg. On leaves of *P. forsteri* in Chile.
Septoria nebulosa Rostr. See *Calamagrostis*.
Septoria oudemansii Sacc. On culms of *P. flexuosa* and *P. nemoralis* in Holland.
Septoria poae-annuae Bres. and var. *sepulta* Gz. Frag. On leaves and glumes of *P. annua*, *P. bulbosa*, and *P. pratensis* in Tripoli, Spain, and Germany.

POA—Continued.

- Septoria poae-trivialis** Cocc. Small subcircular yellow sunken spots on leaves and culms of *P. nemoralis* and *P. trivialis* in Russia, Italy, and Germany.
- Tilletia hordei** Schoeyen. See *Avena*.
- Uredo porophylla** Speg. Rust on leaves of *P. lanigera* in Argentina.
- Uromyces chubutensis** Speg. Leaf rust on *P. chubutensis* in Argentina.
- Uromyces poae** Rabh. Leaf rust on *P. annua*, *P. bulbosa*, *P. compressa*, *P. fertilis*, *P. nemoralis*, *P. palustris*, *P. pratensis*, *P. trivialis*, *P. violacea*, and *Agrostis alba* in Europe, Asia Minor, Nova Scotia, and north Africa. The aecial stage occurs on species of *Ranunculus*.
- Uromyces poae-alpinæ** W. Rytz. Brown to black rust pustules on leaves of *P. alpina* in Switzerland.
- Uromyces sceseli-graminis** Ed. Fisch. See *Arrhenatherum*.
- PODOCARPUS. NAGEIA** Ag. Ornamental trees grown for their evergreen foliage.
- Corynelia bispora** Fitzp. Small black stromata on lower leaf surfaces of *P. milanjiana* in central Africa.
- Corynelia brasiliensis** Fitzp. Parasitic on leaves of *Podocarpus* sp. in Brazil.
- Corynelia jamaicensis** Fitzp. Circular to elliptical black stromata on leaves of *P. purdieana* in Jamaica.
- Corynelia nipponensis** Fitzp. Black stromata on leaves of *P. macrophylla* in Japan.
- Corynelia oreophila** (Speg.) Starb. Black stromata on leaves of *P. augustifolia*, *P. sellowii*, *P. chilina*, and *P. macrostachya* in Brazil, Chile, Colombia, Bolivia, and Costa Rica.
- Corynelia portoricensis** Fitzp. Raised black stromata on leaves of *P. coriacea* in Porto Rico.
- Corynelia tropica** (Auers. and Rabh.) Starb. Irregularly elongated black stromata on leaves and stems of *P. andina*, *P. chilina*, *P. costalis*, and *P. saligna* in Chile and the Philippines.
- Corynelia uberata** Fr. Circular to irregular black stromata on leaves, stems, and fruit of *P. costalis*, *P. elongata*, *P. falcata*, *P. gracilior*, *P. latifolia*, *P. nagi*, *P. nulangianus*, and *P. thunbergii* in Japan, Philippines, New Zealand, and central Africa.
- Septoria arethusa** Penz. See *Citrus*.
- Septoria podocarpi** Niessl. Irregular ashen leaf spots with narrow dull-brown margins on *P. thunbergii* in Italy and the Union of South Africa.
- Thielaviopsis podocarpi** Petri. A doubtful species on roots of *Podocarpus* sp. in Italy.
- Tripospora tripos** (Cke.) Lind. Parasitic on leaves and green parts of stems of *P. elongata* and *P. thunbergii* in the Union of South Africa and on *P. lamberti* in Brazil.
- PODOLIPIS.** Australian herbs cultivated for their yellow, pink, or purple flower heads.
- Puccinia podolepidis** McAlp. Bright-orange and black rust sori on leaves of *P. longipedata* in Australia.
- PODOPHYLLUM. MAY APPLE.** Herbs.
- Phomopsis podophylli** Grove. On leaves of *P. peltatum* in Great Britain.
- POGOSTEMUM. PATCHOULI.** Herbs or shrubs.
- Puccinia princeps** Syd. Rust causing large tumors (1 to 3 centimeters in diameter) on branches of *Pogostemum* sp. in India.
- POINCIANA.** Royal poinciana. Flamboyant. Ornamental leguminous trees and shrubs.
- Cercospora bakeriana** Sacc. On twigs and leaves of *P. (Caesalpinia) pulcherrima* in the Philippines.
- Fomes lamaensis** Murr. See *Hevea*.
- Ravenelia humphreyana** P. Henn. Powdery brown rust pustules causing defoliation of *P. pulcherrima* in Mexico, Costa Rica, Guatemala, Cuba, Jamaica, and Porto Rico.
- Septogloeum poinciana** Syd. Indefinite yellow areas on leaves of *P. alata* in India.
- POLEMONIUM. GREEK VALERIAN. JACOB'S LADDER.** Flower-garden herbs with blue or purple flowers.
- Ascochyta polemonii** Rostr. On stems and leaves of *P. coeruleum* in Italy and Denmark.
- POLLIA.** Perennial herbs.
- Ustilago nawaschini** Radib. Smut destroying flowers of *Pollia* sp. in Java.
- POLYGALA.** Annual or perennial herbs or subshrubs.
- Aecidium palustre** Mont. Rust on leaves and stems of *Polygala* sp. in Brazil.
- Cercospora polygalae** P. Henn. On leaves of *P. paniculata* in Brazil.
- Phyllosticta chamaebuxi** A. Iesch. Ochraceous leaf spots with red margins on leaves of *P. chamaebuxus* in Germany.
- Ramularia polygala** (Schroet.) Sacc. and Syd. Light-brown spots often occupying most of leaf surface on *P. vulgaris* in Silesia.
- Septoria polygalicola** Höll. Large ochraceous leaf spots on *P. comosa* in Hungary.
- Uredo peribuyensis** Speg. Brown leaf rust on *P. americana*, *P. acicularis*, and *Polygala* sp. in Guatemala, Argentina, and Mexico.
- Uredo polygalae** Diet. Dull-yellow rust sori on yellow leaf spots on *P. japonica* and *P. ohlendoriana* in Japan.
- Uromyces polygalae** Grove. Leaf rust on *P. persicariaefolia* and *Polygala* sp. in Uganda.
- POLYGONUM. SOLOMON'S-SEAL.** Perennial herbs with simple stems and creeping rootstocks.
- Cercospora polygonati** Rostr. Subcircular pale-brown leaf spots with dark-purple margins on *P. multiflorum* in Denmark.
- Cylindrosporium komarowii** Jacz. On *P. humile* in Russia.
- Heterosporium alii** Ell. and Mart. var. **polygonati** Oud. On leaves of *P. multiflorum* in Holland.
- Mycosphaerella asteroma** (Fr.) Lind. See *Convallaria*.
- Phyllosticta polygonati** Bauml. Brown leaf spots on *P. multiflorum* and *P. officinale* in Hungary and Yugoslavia.
- Phyllosticta woronowii** Woron. On leaves of *P. officinale* in Russia.
- Rhytisma punctum** Chev. Black stromata on leaves of *P. officinale* in France.
- Sclerotinia richteriana** P. Henn. Causes a blighting of *P. multiflorum* in Germany.
- Septoria brunnea** Fr. See *Convallaria*.
- Septoria polygonati** Kab. and Bub. Subcircular to elliptical gray leaf spots with purple-brown margins on *P. multiflorum* in Bohemia.
- POLYGONUM. SMARTWEED. KNOTWEED.** Erect or twining herbs, a few species grown for ornament.
- Ascochyta polygonicola** Kab. and Bub. Circular to irregular, often confluent, brown zoned leaf spots with yellow margins on *P. lapathifolium* in Bohemia.
- Cercospora paludicola** Speg. Small angular ochraceous-brown leaf spots on *Polygonum* sp. in Argentina.
- Cercospora polygoni** P. Henn. and Shir. Circular brown leaf spots with red-brown margins on leaves of *P. multiflorum* in Japan.
- Entyloma schweinfurthii** P. Henn. Smut sori on black leaf spots on *P. monspeliense* in Egypt.
- Ovularia bistorta** (Fekl.) Sacc. Brown-gray leaf spots with purple margins on *P. bistorta* and *P. crispum* in Japan and Europe.
- Phyllosticta melanogena** Sacc. Circular to angular dark-brown leaf spots on *Polygonum* sp. in Siberia.

POLYGONUM—Continued.

Phyllosticta polygoni-aviculare Petr. Small yellowish-brown leaf spots on *P. aviculare* in Austria.

Phyllosticta polygonorum Sacc. Subcircular whitish leaf spots with reddish margins on *P. multiflorum*, *P. persicaria*, and *Pagopyrum esculentum* in Japan, Formosa, Russia, and Italy.

Phyllosticta tokutaro Speg. Small irregular or circular brown leaf spots on *P. multiflorum* in Japan.

Pseudopeziza histortae (Lib.) Fekl. On leaves of *P. bistorta* and *P. viviparum* in Greenland and Europe.

Puccinia angelicae-mamillata Kleb. See Angelica.

Puccinia astrantiae-vivipari Semad. See Astrantia.

Puccinia calumnata Syd. Leaf rust on *P. weyrichi* in Japan.

Puccinia congesta B. and Br. Leaf rust on *P. chinense* and *P. tomentosum* in China and the Philippines.

Puccinia mammillata Schroet. Leaf rust on *P. bistortum* and *P. viviparum* in Russia, Sweden, Hungary, and Germany. Reported from New Jersey.

Puccinia monticola Kom. Brown rust pustules on leaves of *P. alpinum* and *P. polymorphum* in Turkestan.

Puccinia nitida Barcl. Brown to black rust pustules on leaves of *P. amplexicaulis* and *P. chinense* in India.

Puccinia polygoni-sachalinensis Pat. and Har. Powdery brown rust pustules on leaves of *P. sachalinensis* in Japan.

Puccinia septentrionalis Juel. Leaf rust on *P. viviparum* and *Thalictrum alpinum* in Greenland, Iceland, and northern Europe. Reported from California.

Puccinia solmsii P. Henn. Leaf rust on *P. acuminatum*, *P. chinense*, *P. nipponense*, and *P. posumbu* in India, Indo-China, Japan, and Brazil.

Puccinia sommerfeltii Johans. See *Thalictrum*.

Ramularia bistortae Fekl. On leaves of *P. bistorta* in Siberia.

Septoria polygonicola (Lasch.) Sacc. Circular pale-ochraceous leaf spots on *P. convolvulum*, *P. orientale*, and *P. persicaria* in China, Russia, Yugoslavia, Austria, and Germany.

Sphaelotheca alpina Schellenb. Brown smut sori in sheaths and peduncles of *P. alpinum* in Switzerland.

Ustilago bosniaca Beck. A smut producing powdery, dark-violet spore masses in peduncles and inflorescences of *P. alpinum* in Yugoslavia.

Ustilago emodensis Berk. A smut forming clustered outgrowths up to 1 inch long on stems of *P. chinense* in Ceylon, Java, India, Australia, and the Philippines.

Ustilago koodersiana Bref. Smut in ovaries of *P. varbatum* and *Polygonum* sp. in China, Java, and the Philippines.

Ustilago marginalis (Link.) Lév. Smut sori as swellings on leaves of *P. bistorta* in central Europe.

Ustilago moelleri Bref. Smut forming swollen masses of dark-brown spores in fruit of *P. hispidum* in Brazil.

Ustilago rosulata Syd. Powdery violet smut sori in ovaries of *P. chinensis* in the Philippines.

Ustilago tuberculiformis Syd. Smut sori with powdery yellow-brown spore masses in purple-brown leaf spots on *P. chinense* and *P. runcinatum* in China and India.

POLYMNIA. Coarse viscid composites.

Phyllachora perlata Syd. Irregular black stromata on leaves of *P. glabrata* in Colombia.

Uredo banisteriicola P. Henn. Cinnamon-brown rust sori on leaves of *P. sylphoides* in Brazil.

Uromyces polymniae (P. Henn.) Diet. and Holw. Brown rust pustules on leaves of *P. glabrata*, *P. maculata*, and *P. sylphoides* in Mexico, Colombia, and Brazil.

POLYPODIUM. POLYPODY. Ferns.

Hyalopora japonica Diet. Rust on leaves of *P. senanense* in Japan.

Laestadia polypodii Sacc. and Magn. On *P. vulgare* in Italy.

Miesina dietel ana (Syd.) P. Magn. Yellow-brown rust sori on small sunken spots on leaves of *P. vulgare* in Europe.

Mycosphaerella polypodii (Rabh.) Magn. On *P. vulgare* in Denmark, Switzerland, and Germany.

Phyllosticta polypodi-australis P. Henn. Irregular pale-gray leaf spots on *P. austratum* in Kerguelen Islands.

Septoria polypodii Grove. On leaves of *P. phegopteris* in Great Britain.

Soraea maxima (B. and C.) Giesenh. Small black stromata on leaves of *P. crassifolium*, *P. phyllitidis*, *P. punctatum*, and *P. schomburgkianum* in Porto Rico, Cuba, St. Domingo, Ecuador, Venezuela, and Brazil.

POLYSTACHYA. See Orchidaceae.**POLYSTICHUM**. HOLLYFERN. Christmas fern.

Exoascus filicina (Rost.) Sacc. Small definite leaf spots on *P. spinulosum* and *Phegopteris vulgaris* in Sweden and Russia.

Magnusiella lutescens (Rost.) Sadeb. Yellow leaf spots on *P. thelypteris* in Denmark.

Metasphaeria polystichi Feltg. On petioles of *P. filix-mas* in Luxemburg.

PONGAMIA. Shrubs native of the Orient.

Cryptomyces pongamiae (B. and Br.) Sacc. Black irregular pitchlike leaf spots on *P. glabra* in India.

Fusicladium pongamiae Syd. On leaves of *P. glabra*, *P. nitida*, and *P. pinnata* in India and the Philippines.

Phyllachora pongamiae (B. and Br.) Petch. Shiny black stromata on leaves of *P. glabra* and *P. pinnata* in India, Ceylon, Java, and the Philippines.

Ravenella hobsoni Cke. Brown powdery rust pustules on leaves of *P. glabra* in India and Ceylon.

PONTEREDIA. PICKEREL WEED. Perennial aquatic herbs.

Phyllosticta pontederiae Syd. Irregular dull-brown leaf spots on *Pontederia* sp. in Brazil.

POPULUS. POPLAR. Aspen. Cottonwood. Timber, pulp, and shade trees.

Ascochyta populi Delacr. On leaves of *P. canadensis* in Italy.

Ascochyta populicola Kab. and Bub. On leaves of *P. alba* in Bohemia.

Ascochyta populina Sacc. Angular white leaf spots with dark borders on *P. nigra* in Italy.

Ascochyta tremulae Thuern. Subcircular small grayish leaf spots with dull-brown margins on *P. tremula* in Austria.

Bacillus populi Brizl. This bacterium is reported from Italy and France as the cause of a canker disease of trunks and branches of *P. alba*, *P. nigra*, and *P. tremula*.

Didymosphaeria populina Vuil. (*Fusicladium tremulae* Fr.). Tips of young shoots are killed and blackened, the seasonal recurrence of which destroys the crowns of infected trees. Blackened patches occur on the leaves, particularly at the tips and along margins. On *P. alba pyramidalis*, *P. canadensis*, *P. nigra*, *P. tremula*, and *P. tremuloides* in Russia, Sweden, France, Denmark, Austria, and Germany. Common in Canada.

POPULUS—Continued.

- Diplodia gongrogena** Tem. Causes hypertrophies of bark and wood of *P. tremula* in Germany.
- Hyalopopuli** Nypels. Said to cause a canker of *P. canadensis* in Belgium.
- Hypoderma roseum** (Pers.) Fr. On bark of *Populus* sp., *Alnus* sp., and *Juglans* sp. in Great Britain, Sweden, Italy, and Germany.
- Laestadia nesslerii** Kze. On leaves of *P. alba pyramidalis* in Germany.
- Leptosphaeria livida** Vogl. Irregular grayish-white leaf spots on *P. canadensis* in Italy.
- Leptothyrium populi** Fckl. On leaves of *P. nigra*, *P. alba pyramidalis*, and *P. tremula* in Italy, Austria, and Germany.
- Macrophoma populi-nigrae** (Allesch.) Tass. On leaves of *P. nigra* in Germany.
- Marsonia curvata** Bub. and Kab. Irregular and confluent brown leaf spots on *P. nigra* in Bohemia. Reported from Utah.
- Marsonia piriformis** (Riess.) Sacc. Dark-brown leaf spots on *P. alba* in Europe. Reported from Utah.
- Marsonia populina** Schnabl. Leaf spots on *P. nigra* in Europe.
- Melampsora alii-populina** Kleb. Yellow or golden to brown rust pustules in yellow leaf spots on *P. alba*, *P. angulata*, *P. balsamifera*, *P. canadensis*, *P. deltoides*, and *P. nigra* in Argentina and Europe. Alternate stage on *Allium ascalonicum*, *A. cepa*, *A. oleraceum*, *A. sativum*, *A. schoenoprasum*, *A. ursinum*, and *A. vinealis*.
- Melampsora ciliata** Bard. Leaf rust on *P. ciliata* in northern India.
- Melampsora laricis** R. Hartig. Yellow, then brown, rust pustules on leaves of *P. alba*, *P. balsamifera*, *P. canescens*, *P. tremula*, *Larix decidua*, *L. griffithii*, and *L. sibirica* in Europe and Japan.
- Melampsora larici-populina** Kleb. Leaf rust on *Larix decidua* and practically all species of *Populus* in Japan, Argentina, and Europe.
- Melampsora magnusiana** G. Wagn. Golden and brown rust pustules on leaves of *Chelidonium majus*, *Corydalis cava*, *C. digitata*, *C. fabacea*, *C. incisa*, *C. intermedia*, *C. laza*, *C. pallida*, *C. solida*, *Populus alba*, *P. canescens*, and *P. tremula* in Japan, China, Siberia, and Europe. Apparently not distinct from *M. tremulae* Tul., which occurs in the United States.
- Melampsora pintoerqua** Rostr. Yellow powdery aecia on *Pinus montana* and *P. silvestris*. Telia on *Populus tremuloides*, *P. tremula*, and *P. alba pyramidalis* in Russia, Denmark, Italy, and Spain.
- Melampsora pruinosa** Tranzsch. Pale-yellow to red-brown rust sori on leaves of *P. pruinosa* in Siberia.
- Melampsora pulcherrima** (Bub.) R. Maire. Leaf rust on *P. alba* in Spain.
- Melampsora rostrupii** G. Wagn. Yellow and dark-brown rust sori on leaves of *P. alba*, *P. alba pyramidalis*, *P. balsamifera*, *P. canadensis*, *P. canescens*, *P. tremuloides*, *P. nigra*, *P. tremula*, and *Mercurialis perennis* in Europe.
- Micrococcus dendroporthos** Ludw. Small gum-producing cankers on bark of young trees of *Populus* sp. in Argentina.
- Micrococcus populi** Delacr. Bacterial cankers on stems and branches of seedlings and on trunks of older trees of *P. deltoides* and *P. nigra* in France and Italy.
- Mycosphaerella macularis** (Fr.) Schroet. On leaves of *P. tremula* in Europe. Reported from Kansas.
- Myosphaerella populi** (Auersw.) Schroet. On leaves of *P. deltoides*, *P. nigra*, and *P. alba pyramidalis* in Europe.
- Napicladium asteroma** (Fckl.) Allesch. On leaves of *P. alba* in Germany.
- Phoma populi-nigrae** Allesch. On leaves of *P. nigra* in Germany.
- Phyllohendersonia alcidis** (Sacc.) Tass. On leaves of *P. nigra* in Italy.
- Phyllohendersonia foliorum** (Fckl.) Tass. On leaves of *P. nigra* in France, Italy, and Germany.
- Phyllosticta adjuncta** Bub. and Sere. On leaves of *P. euphratica* in Russia.
- Phyllosticta bacteriiformis** (Pass.) Sacc. On leaves of *P. nigra* in Italy.
- Phyllosticta cinerea** Pass. Ashen-white leaf spots on *P. alba* in Italy.
- Phyllosticta ostrospora** Sacc. See *Morus*.
- Phyllosticta populea** Sacc. Irregular white leaf spots with dull-brown margins on *P. alba*, *P. tremula*, and *Populus* sp. in Italy and China.
- Phyllosticta populina** Sacc. Angular white leaf spots with black margins on *P. nigra* and *P. alba pyramidalis* in Yugoslavia, Albania, Italy, and Denmark. Reported from Wisconsin.
- Phyllosticta populorum** Sacc. and Roum. On leaves of *P. balsamifera* and *P. canadensis* in Italy, France, Portugal, and Germany.
- Phyllosticta prominens** Oud. Brown leaf spots on *P. balsamifera* in Holland.
- Sclerotium scutellatum** A. S. See *Acer*.
- Septoria aegerina** Pass. On leaves of *P. nigra* in Italy.
- Septoria atrosanguinea** Bub. and Sere. On leaves of *P. tremula* in Russia and Poland.
- Septoria butuliformis** Bub. and Sere. On leaves of *P. euphratica* in Russia.
- Septoria candida** (Fckl.) Sacc. White leaf spots on *P. alba* in Italy, France, and Germany.
- Septoria marmorata** Kab. and Bub. Subcircular to irregular brown, then pale-yellow, leaf spots on *P. tremula* in Denmark and Italy.
- Septoria osteospora** Bri. On leaves of *P. nigra* in France.
- Septoria tremulae** Pass. Indefinite dull-brown confluent leaf spots on *P. tremula* in Italy.
- Titaesporina tremulae** (Lib.) Luyk. Anthracnose on leaves of *P. alba* and *P. tremula* in Europe. Reported in a few cases from the United States under the name *Gloeosporium tremulae* (Lib.) Pass.
- PORTULACA**. Purslane. Low, fleshy, annual or perennial herbs.
- Phyllosticta portulacae** Sacc. and Speg. Subcircular white leaf spots with raised dull-brown margins on *P. oleracea* in Italy.
- Tuberculina portulacarum** Speg. Dull-brown leaf spots on *P. plan-operculata* in Argentina.
- POSEQUERIA**. Glabrous shrubs and small trees.
- Aecidium posoquiae** Diet. Leaf rust on *P. latifolia* in Brazil.
- POTENTILLA**. CINQUEFOIL. Perennial herbs and shrubs.
- Entyloma pustulosum** Sacc. and Fautr. Leaf smut on *P. anserina* in France.
- Laestadia potentillae** Rostr. On leaves of *P. maculata* and *P. nivea* in Iceland.
- Peronospora potentillae anserinae** Güm. Downy mildew on leaves of *P. anserina* in France and Germany.
- Peronospora potentillae reptantis** Güm. Downy mildew on leaves of *P. recta*, *P. reptans*, and *P. supina* in northern Europe.
- Peronospora potentillae sterilis** Güm. As above on *P. sterilis* in Switzerland and Germany.
- Phlebospora fragariae** (Br. and Har.) Petr. See *Fragaria*.
- Phragmidium fragariastrum** (DC.) Schroet. Yellow to black rust pustules on leaves of *P. alba*, *P. alchemilloides*, *P. carnioleca*, *P. fragariastrum*, *P. hybrida*, *P. micrantha*, *P. splendens*, *P. sterilis*, *Fragaria collina*, and *F. indica* in Europe.
- Phragmidium laetanum** Barcl. Yellow rust pustules on leaves of *P. argyrophylla* in India.
- Phragmidium nepalense** Barcl. Yellow to black rust sori on leaves of *P. nepalensis* in India.

POTENTILLA—Continued.

- Phragmidium papillatum** Diet. Yellow to brown rust pustules on lower leaf surfaces of *P. striosa* in Siberia.
- Phyllosticta argentinæ** Desm. On leaves of *P. anserina*, *P. argentea*, and *Comarum palustre* in Siberia and Austria.
- Phyllosticta potentillæ** Desm. On leaves of *P. fragarioides* in Siberia.
- Phyllosticta tormentillæ** Sacc. Irregular pale ochraceous leaf spots on *P. tormentilla* in Italy.
- Physalospora potentillæ** Rostr. On stems of *P. maculata* in Greenland.
- Phyoderma vagans** Schroet. See *Ranunculus*.
- Ramularia anserina** Allesch. Subcircular, then confluent, ochraceous leaf spots on *P. anserina* in Germany.
- Ramularia martianoffiana** Thuem. Large irregular dull-violet leaf spots on *P. striosa* in Siberia.
- Septoria corcontica** Kab. and Bub. Circular to angular dull-green, then yellowish, leaf spots with brown-purple margins on *P. procumbens* and *P. tormentilla* in Bohemia.
- Septoria fragariæ** Desm. See *Fragaria*.
- Septoria tormentillæ** Rob. and Desm. Irregular, often oblong, brown leaf spots with rose-colored margins on *P. erecta*, *P. reptans*, *P. sylvestris*, and *P. tormentilla* in Russia, France, Denmark, Italy, Belgium, and Germany.
- Synchytrium globosum** Schroet. Small rough galls on leaves and stems of *P. reptans*, *Achillea millefolium*, *Cirsium oleraceum*, *Galium mollugo*, *Myosotis palustris*, *Sonchus asper*, *Veronica anagallis*, *V. beccabunga*, *V. chamaedrys*, *V. scutellata*, *Viola canina*, *V. odorata*, *V. persicifolia*, *V. silvatica*, and *V. stannina*, in northern Europe.
- Synchytrium pilificum** Thomas. Small galls on leaves and stems of *P. tormentilla* in Germany.
- Uredo nericola** Tranz. Leaf rust on *P. fragarioides* in Manchuria.
- Venturia potentillæ** (Fr.) Cke. On leaves of *P. anserina*, *P. bifurca*, *P. cinerea*, *P. reptans*, and *P. tormentilla* in Siberia and Europe.
- POTERIUM**. BURNET. Rosaceous shrubs. (See also *Sanguisorba*.)
- Ovularia bulbifera** (Fekl.) Sacc. On leaves of *Sanguisorba minor* (*P. sanguisorba*) in France.
- Phragmidium sanguisorbæ** (DC.) Schroet. Yellow to golden rust sori on leaves of *P. eriocarpum*, *P. magnolium*, *P. mauritanicum*, *P. muricatum*, *P. myriophyllum*, *P. polygamum*, *P. verrucosum*, *P. rillosum*, *Sanguisorba dictyocarpa*, *S. duria*, *S. media*, and *S. minor* in Algeria, Tunis, Asia Minor, Turkestan, and Europe.
- POURTHIAEA**. See *Photinia*.
- PRATIA**. Slender prostrate or creeping herbs, sometimes grown as ornamentals.
- Uromyces pratiae** Speg. Powdery dark-brown rust pustules on leaves and stems of *P. longiflora* and *P. repens* in South America.
- PREMNA**. Shrubs.
- Cronartium premnae** Petch. Golden-yellow to brown rust pustules on leaves of *P. cordifolia* and *P. corymbosa* in Ceylon.
- Mycosphaerella oculata** Syd. Small generally circular, leaf spots on *P. odorata* in the Philippines.
- Phyllachora premnae** Syd. Black stromata on leaves of *P. cumingiana* in the Philippines.
- Puccinia premnae** P. Henn. Powdery black rust pustules on leaves of *P. macrophylla* and *P. microphylla* in Japan.
- Uredo philippinensis** Syd. Leaf rust on *P. tomentosa* in the Philippines.
- Uredo premnae** Koord. Leaf rust on *P. cumingiana*, *P. odorata*, and *P. tomentosa* in Java and the Philippines.
- PRENANTHES**. Tall perennial herbs.
- Puccinia altaica** Syd. Black rust pustules on leaves and stems of *P. diversifolia* in central Asia.
- Puccinia prenanthis** (Pers.) Lindr. See *Lactuca*.
- Puccinia prenanthis-purpurea** (DC.) Lindr. and var. *himalensis* Barcl. Powdery yellow and brown rust pustules on leaves of *P. alba*, *P. brunoniana*, *P. purpurea*, and *P. tenuifolia* in Europe and India.
- Ramularia prenanthis** Jaap. Subcircular ochraceous or brown leaf spots, which become confluent, occupying entire leaf blades of *P. purpurea* in Austria and Switzerland.
- PRESTONIA**. Tall, climbing, pubescent or glabrous shrubs.
- Helminthosporium prestoniae** P. Henn. Circular to confluent dark-brown leaf spots on *Prestonia* sp. in Peru.
- PRIMULA**. PRIMROSE. Low herbaceous plants with attractive many-colored flowers. (Including *Auricula*.)
- Cercospora primulae** Fautr. Grayish-white leaf spots with yellow margins on *P. elatior* in France.
- Cercosporella primulae** Allesch. Subcircular ochraceous leaf spots on *P. acutis* and *P. veris* (*P. officinalis*) in Russia and Germany.
- Heterosporium auriculæ** Cke. Leaves of *P. auricula* in Great Britain are disfigured by smoky patches due to this fungus.
- Mycosphaerella primulae** (Auersw. and Heufl.) Schroet. On leaves of *P. minima* in Yugoslavia.
- Ovularia primulana** Karst. Yellowish patches on lower leaf surfaces of *P. acutis*, *P. elatior*, *P. officinalis*, *P. veris*, and *Primula* sp. in Europe.
- Peronospora oerteliana** Kuehn. Downy mildew on brown spots on lower leaf surfaces of *P. acutis*, *P. elatior*, and *P. officinalis* in Switzerland, Bohemia, Sweden, Denmark, and Germany.
- Phyllosticta primulicola** Desm. Circular white leaf spots with tawny margins on *P. acutis*, *P. elatior*, *P. officinalis*, *P. vulgaris*, and *P. veris* in Europe.
- Puccinia arctica** Lagh. Golden and yellow-brown rust sori on leaves of *P. sibirica* in Norway, Lapland, and northern Asia.
- Puccinia primulae** (DC.) Duby. Brown rust sori on leaf spots on *P. acutis*, *P. cartusoides*, *P. elatior*, *P. grandiflora*, *P. integrifolia*, *P. officinalis*, and *P. vulgaris* in Japan and Europe.
- Ramularia primulae** Thuem. Circular to angular pale-ochraceous leaf spots on *P. acutis*, *P. auricula*, *P. elatior*, *P. japonica*, *P. macrocalyx*, *P. officinalis*, *P. sinensis*, and *P. suaveolens* in Australia, Siberia, and Europe.
- Ramularia tirolensis** Maire. Subcircular, then confluent, brown leaf spots on *P. imbricata* in Tyrol.
- Septoria primulae** Buch. Circular brown leaf spots on *P. veris* and *Primula* sp. in Great Britain and Italy.
- Septoria primulicola** Rostr. Numerous ashen circular leaf spots with dull-brown margins on *P. acutis* and *P. grandiflora* in Denmark.
- Urocystis primulicola** Magn. Snout sori changing seed capsules into masses of black spores. On *P. farinosa* and *P. veris* in Great Britain, Italy, Silesia, and Germany.
- Uromyces apiosporus** Hazsl. Dark-brown rust sori on leaves of *P. minima* in Italy, Austria, Hungary, and Germany.
- Uromyces ovirensis** Jaap. Leaf rust on *P. wulfeniana* in Austria.

PRIMULA—Continued.

Uromyces primulae Fekl. Yellow and brown rust pustules on leaves of *P. auricula*, *P. balbisii*, *P. hirsuta*, *P. latifolia*, *P. minima*, *P. pedemoniana*, *P. siberica*, and *P. villosa* in Italy, Switzerland, Austria, and Germany.

Uromyces primulae-integrifoliae (DC.) Niessl. Leaf rust on *P. deorum*, *P. integrifolia*, *P. minima*, *P. muretiniana*, and *P. viscosa* in Bulgaria, Switzerland, Denmark, and Austria.

PROSOPIS. MESQUITE. Hawaiian algarroba. Leguminous trees and shrubs used for shade and forage.

Ascochyta prosopidicola Speg. Large leaf spots on *Prosopis* sp. in Argentina.

Phyllosticta prosopidicola Speg. Definite white leaf spots with purple borders on *P. alba* in Argentina.

Phyllosticta prosopidis P. Henn. Oblong white leaf spots with brownish margins on *P. ruscifolia* in Argentina.

Uncinula prosopidis Speg. Powdery mildew on leaves of *P. campestris* in Argentina.

Uredo prosopidis Jacz. Powdery brown rust pustules on leaves of *P. stephaniana* in Transcaucasia.

PROSTANTHERA. Shrubs, commonly strongly scented.

Phyllosticta prostantherae Cke. Brown leaf spots on *P. lasianthos* in Australia.

PROTEA. Tender shrubs, small trees or acaulescent perennial herbs.

Leptosphaeria protearum Syd. Subcircular, then irregular, sordid-white leaf spots on *P. madiensis* and *P. melaleuca* in Uganda and the Union of South Africa.

Oligostroma maculiformis (Wint.) Doidge. Black stromata on leaves of *P. abyssinica*, *P. nanaganii*, *P. grandiflora*, and *P. neriifolia* in the Union of South Africa.

Oligostroma proteae Syd. On *P. flanaganii* in the Union of South Africa.

Phaeosphaeria seniana Sacc. On *P. abyssinica*, *P. acutis*, and *P. melaleuca* in the Union of South Africa and Abyssinia.

Phyllachora proteae Wakef. Tar spot on leaves of *P. mellifera* in the Union of South Africa.

Pseudomonas proteamaculans Paine and Stans. This bacterium produces numerous dome-shaped blisters, 1 to 3 mm. in diameter, on upper leaf surfaces of *P. cynaroides* in Great Britain. On young leaves the areas are larger, red to reddish brown with vermilion borders and depressed.

PRUNELLA. See Brunella.

PRUNUS. PLUM. CHERRY. APRICOT. (Including *Amygdalus*, PEACH, ALMOND, and NECTARINE). Fruit and nut trees.

Ascochyta chlorospora Speg. A shot-hole disease of leaves and fruit spot of *P. armeniaca*, (*Amygdalus communis*) *P. cerasus*, *P. divaricata*, *P. domestica*, and *P. (Amygdalus) persica* in Russia and Australia.

Ascochyta crystallina McAlp. Leaf spot on *P. amygdalus* in Australia.

Ascochyta ovalispora McAlp. On leaves and twigs of *P. cerasus* and *P. domestica* in Australia.

Ascochyta pruni Kab. and Bub. Circular, often indefinite, brown, then ashen-gray, leaf spots on *P. padus* in Bohemia.

Ascospora padi Grev. Defoliates *P. cerasus* in Europe.

Auerswaidiella puccinioides (Speg.) Theiss. and Syd. Dull-black stromata on leaves of *P. argentinensis* and *P. sphaerocarpa* in Central and South America.

Bacillus spongiosus Aderh. and Ruhl. Large cankers with heavy gum exudate on trunks of *P. cerasus* in Germany. This is apparently one of the gumming diseases of *Prunus* spp., the exact causes of which have for the most part not been worked out.

Bacterium sp. (Blossom-blight.) See *Pyrus*.

Caecoma makinoui Kus. A rust attacking the buds of *P. domestica*, *P. grayana*, *P. mume*, and *P. pseudocerasus* in Japan, causing hypertrophy and distortion of the young shoots as they develop.

Camarosporium prunifolium McAlp. Elongate narrow gray leaf spots with raised margins on *P. armeniaca* in Australia.

Cercospora amygdali Ali Riza. Subcircular, often confluent, gray spots on leaves of *P. amygdalus* in Turkey.

Cercospora guliana Sacc. Subcircular yellow leaf spots on *P. amygdalus* in Malta.

Clasterosporium persicum (Sacc.) Tsuji (*Cercospora persicum* Sacc.) On leaves of *P. persica* in Japan and Italy.

Coniothecium albo-cinctum Preuss. Light gray patches on shoots of *P. domestica*, *P. persica* and *Malus* in Australia and Germany.

Coniothyrium pruni McAlp. Circular to irregular dirty-gray spots with raised dark-red margins on leaves and fruit of *P. armeniaca* and *P. domestica* in Australia.

Corticium salmonicolor B. and Br. See *Citrus*.

Cucurbitaria pruni-mahaleb. Allesch. Attacks and kills buds and small twigs, a witches'-broom effect resulting. On *P. mahaleb* in Europe. Reported from New York.

Cylindrosporium tubeufianum Allesch. Attacks green fruit of *P. avium* and *P. padus* in Germany.

Dermatella prunastri Pers. Attacks and destroys terminal twigs of *P. domestica* in Great Britain.

Diaporthe perniciosus Murchl. This fungus attacks and destroys trees of all ages, but more particularly the young ones. Slightly sunken, reddish, elongate cankers appear, increasing rapidly in size and bringing about wilting and browning of the leaves which fall prematurely. Under some circumstances there is also a rot of the fruit. The hosts are *P. armeniaca*, *P. persica*, *P. cerasus*, *P. domestica*, *Malus* (apple), and *Pyrus* (pear) in Belgium, France, and Great Britain.

Didymaria prunicola Car. Brown leaf spots on *P. domestica* in Italy, causing premature leaf fall.

Dithiorella vinosa Marchal. Forms cankers on trunks and branches, spots on leaves, and rot of fruit of *Malus*, *Prunus domestica*, *Pyrus*, and *Ribes* in Belgium.

Eoasacus andinus (Palm.) Sacc. and Trot. Indefinite reddish, often swollen, areas on leaves of *P. salicifolia* in Ecuador.

Eoasacus rostrupianus Sadeb. Forms "fruit pockets" on *P. spinosa* in Europe.

Fusarium gemmiperda Aderh. Attacks and destroys buds of *P. cerasus* in Germany.

Fusidium amygdali Duc. Small circular to angular brown leaf spots which enlarge to include the entire leaf blade, defoliation resulting. Young fruit and blossoms drop prematurely due to the weakening of the twigs. On the twigs dark-green, then brown, rough spots occur on shaded surfaces. On *P. amygdalus* in Italy and France.

Fusidium cerasi Sacc. (*Venturia cerasi* Aderh.) Olive-black minute velvet blotches on fruit of *P. cerasus* in Europe. Diseased fruits mummify. Leaves and twigs are rarely attacked. Probably not distinct from *Cladosporium carpophilum*. Reported from Ohio and New Mexico.

Fusidium pruni Duc. Gray or brown leaf spots on *P. domestica* in France.

Fusicoccum amygdali Delacr. Causes die-back of *P. amygdalus* in France.

Gloeosporium armeniacum Speg. Anthracnose on *P. armeniaca* in Argentina.

Gloeosporium cerasi Lind. Anthracnose on *P. avium* in Japan.

Glomerella mume (Hori.) Hemml. Anthracnose on *P. mume* in Japan.

Gnomonia circumscissa McAlp. Circular ochraceous to gray leaf spots on *P. amygdalus*, *P. armeniaca*, *P. cerasus*, *P. domestica*, and *P. persica* in Australia. Possibly not distinct from the following species.

PRUNUS—Continued.

- Gnomonia erythrostoma** (Pers.) Auersw. Small yellow leaf spots which become brown and much enlarged, causing defoliation in part. The fruit withers or ripens unevenly, with poor color and flavor. On *P. cerasus* in Europe and New Zealand.
- Graphium ulmi** Schwartz. See *Ulmus*.
- Haplosporella pruni** McAlp. Causes large warted swellings on branches of *P. armeniaca* and *P. domestica* in Australia.
- Helicobasidium mompa** Tan. See *Morus*.
- Helicobasidium tanakae** Miy. See *Morus*.
- Kellermannia pruni** McAlp. On leaves of *P. armeniaca* and *P. persica* in Australia.
- Laestadia circumscissa** Sacc. Subcircular to oblong red to brown leaf spots on *P. spinosa* in Italy. One of the shot-hole diseases.
- Microstoma tonellianum** Ferr. Irregular white leaf spots on *P. domestica* in Italy.
- Mycosphaerella cerasella** Aderh. (*Cercospora cerasella* Sacc.) Brown shot-hole disease on *P. arium*, *P. cerasus*, *P. iliosakura*, and *P. yamasakura* in Japan, Russia, Bulgaria, Yugoslavia, and Italy.
- Myxosporium pruni-mahaleb** Moreillon. Attacks and destroys buds and small twigs of *P. mahaleb* in Europe, witches'-brooms developing ultimately.
- Naemospora crocea** (Bon.) Sacc. Two-year-old shoots are attacked, the young leaves wilting, turning brown and dying about blossom time. The flowers also turn brown and the shoots themselves shrivel and die. On *P. persica* in Great Britain.
- Ovularia cerasi** McAlp. Scabby patches on fruit of *P. cerasus* in Australia.
- Ovularia circumscissa** Sorok. Subcircular reddish leaf spots on *Prunus* sp. in the Caucasus.
- Phoma persicae** Sacc. Grayish areas on twigs of *P. persica* in Australia, apparently causing die-back.
- Phyllosticta endersonia foliorum** (Fckl.) Tass. On leaves of *P. domestica* in Italy, France, and Germany.
- Phyllosticta albomaculans** Kab. and Bub. Irregular leaf spots, white above, yellow-brown beneath, on *P. padus* in Bohemia.
- Phyllosticta armenicula** Farneti. Irregular spots on fruit of *P. armeniaca* in Italy.
- Phyllosticta chlorospora** McAlp. Elongated irregular leaf spots dirty-white above, pale red-brown beneath, on *P. domestica* in Australia.
- Phyllosticta destruens** Desm. See *Celtis*.
- Phyllosticta macrospora** McAlp. Irregular gray leaf spots with reddish margins, the centers falling out, on *P. armeniaca*, *P. divaricata*, and *P. domestica* in Australia.
- Phyllosticta mahaleb** Thum. Circular ashen leaf spots on *P. mahaleb* in Italy and Portugal.
- Phyllosticta matthiolana** (Sacc. and Matt.) McAlp. Circular to elliptical or irregular brown leaf spots on *P. amygdalus*, *P. armeniaca*, *P. cerasus*, *P. communis*, *P. domestica*, *P. lauro cerasus*, and *P. persica* in Italy and Australia.
- Phyllosticta minutissima** Kab. and Bub. Circular to irregular, then confluent, pale ashen-brown leaf spots on *P. spinosa* in Bohemia.
- Phyllosticta padi** P. Brun. Brown or ochraceous subcircular to irregular spots on leaves of *P. padus* in Italy and France.
- Phyllosticta persicophila** Trav. and Migliardi. Marginal irregular ochraceous leaf spots with white centers on *P. persica* in Italy.
- Phyllosticta pruni-avium** Allesch. Yellow-brown leaf spots on *P. avium* in Germany.
- Phyllosticta pruni-domesticae** Vogl. Gray leaf spots, becoming white and somewhat swollen, on *P. domestica* in Italy.
- Phyllosticta serebriankowii** Bub. Circular to elliptical gray, more or less concentrically zoned, leaf spots on *P. padus* in Russia.
- Phyllostictiella pruni-spinosae** (Allesch.) Tass. Ashen leaf spots on *P. spinosa* in Italy and Germany.
- Phytophthora syringae** Kleb. See *Syringa*.
- Polystigma ochraceum** (Wahlenb.) DC. Stromata on subangular leaf spots on *P. cerasus*, *P. amygdalus*, *P. padus*, and *Malus* sp. in Japan, India, and Europe.
- Puccinia cerasi** (Bereng.) Cast. Golden-yellow to brown rust pustules on leaves of *P. avium*, *P. cerasus*, *P. domestica*, *P. persica*, and *P. spinosa* in Europe.
- Puccinia radiata** Shirai. Rust pustules on leaves of *P. pseudo-cerasus* in Japan.
- Ramularia cerasorum** Marchal. On leaves of *P. cerasus* in Belgium.
- Ramularia lata** Sacc. White leaf spots on *P. laurocerasus* in France. A doubtful species.
- Rhodoseptoria ussuriensis** Naoum. Destroys leaves and fruits of *Prunus* sp. in Russia.
- Sclerotinia kusanoi** P. Henn. Ashen-white areas on leaves of *P. pseudo-cerasus* in Japan.
- Sclerotinia laxa** Aderh. and Ruhl. Rot and mummification of fruit of *P. armeniaca* and *P. communis* in Australia, Russia, France, Hungary, and Germany.
- Septobasidium acaciae** Saw. See *Acacia*.
- Septoria amygdali** McAlp. On leaves of *P. amygdalus* and *Prunus* sp. in Australia and Russia.
- Septoria anomala** Sacc. Marginal leaf spots on *P. spinosa* in France.
- Septoria cerasi** Pass. Dark sanguineous leaf spots on *P. avium* and *P. cerasus* in France.
- Septoria disseminata** Desm. Leaf spot on *P. laurocerasus* and *P. lusitanica* in France.
- Septoria effusa** (Lib.) Desm. On leaves of *P. amygdalus* and *P. cerasus* in France, Italy, Austria, and Australia.
- Septoria lauro-cerasi** Desm. and Grog. Small brown leaf spots on *P. laurocerasus* in France.
- Septoria lauro-cerasina** Pass. On leaves of *P. laurocerasus* in Italy.
- Septoria myrobalanae** Brun. Circular to oblong brown leaf spots on *P. cerasifera* (*P. myrobalana*), in France.
- Septoria pruni-mahaleb** Therry. Small circular, then confluent, brownish-red leaf spots on *P. mahaleb* in France.
- Septoria staganosporioides** Mass. On leaves of *P. laurocerasus* in Italy.
- Sporotrichum persicae** Pöhl. White incrustations or patches on fruit of *P. persica* in Italy. The fungus in itself does not cause a rot but lowers the market value of the fruit.
- Stereum purpureum** Pers. The silver-leaf disease is considered a serious disease of *P. armeniaca*, *P. cerasus*, *P. communis*, *P. domestica*, *P. japonica*, *P. lusitanicus*, *P. persica*, *P. spinosa*, *P. serotina*, *Malus*, and *Pyrus* in Europe, New Zealand, Canada, and the Union of South Africa. It has also been reported on *Aesculus* sp., *Pernettya mucronata*, *Crataegus* spp., *Philadelphus* sp., *Ribes* sp., and *Rosa sericea*. The fungus considered as the cause of the disease is widespread throughout the United States, but apparently never occurs, or certainly only rarely, in connection with the silver-leaf type of disease, except in Oregon and Washington.
- Silver-leaf is characterized by a silverying of the foliage, which spreads gradually until the entire tree is affected. The limbs bearing these silvered leaves die back and ultimately the tree itself dies. The fungus does not attack the leaves directly, but causes a brown heart rot of roots and trunk and is supposed to secrete some deleterious substance which causes the external signs of the disease.
- Stigmata briosiana** Far. On *P. armeniaca* in Europe.

PRUNUS—Continued.

- Taphrina mexicana** Syd. Small witches'-brooms are formed on *P. microphylla* in Mexico by this fungus.
- Taphrina minor** Sadeb. Reddish spots on leaves and twigs of *P. avium*, *P. cerasus*, and *P. chamaecerasus* in New Zealand, Denmark, Holland, Russia, Great Britain, and Germany. Diseased twigs become malformed.
- Taphrina mume** Nish. On *P. armeniaca*, *P. domestica*, and *P. mume* in Japan.
- Taphrina pseudo-cerasi** Shir. Causes a witches'-broom effect on *P. subhirtella* (*P. miqueliana*) and *P. pseudo-cerasus* in Japan.
- Taphrina trunicola** S. Kus. This fungus attacks the buds of *P. incisa* and *P. pseudocerasus* in Japan. It causes hypertrophy and distortion of twigs and petioles. Leaf blades are commonly not changed, but the flowers become succulent pink masses of abnormal tissue.
- Thekopsora areolata** (Fr.) P. Magn. Dark-brown crustlike rust sori on leaves of *P. padus*, *P. serotina*, and *P. virginiana* in Japan and Europe. The aecial stage forms red-brown to dark-brown pustules on cone scales of *Picea ajanensis*, *P. excelsa*, *P. glehnii*, *P. hondoensis*, and *P. obovata*.
- Thekopsora pseudocerasi** Hiotsuka. Leaf rust on *P. pseudocerasus* in Japan.
- Uncinula prunastri** (DC.) Sacc. Powdery mildew on leaves of *P. dasycphylla*, *P. domestica*, *P. insititia*, *P. pumila*, *P. spinosa*, and *Crataegus* sp. in Europe and Transcaucasia.
- Uredo pruni-maximowiczii** P. Henn. Rust sori in young branches and petioles of *P. maximowiczii* in Japan.
- Uredo persicae** Speg. Leaf rust on *P. persica* in Brazil.
- Valsa japonica** Miy. and Henn. A wound parasite of trunks and branches of *P. cerasus*, *P. koidzumii*, *P. kurilenensis*, *P. mume*, *P. persica*, *P. sachalinensis*, *P. serrulata*, and *P. yedoensis* in Japan.
- Verticillium** sp. A wilt disease of *P. cerasus* is reported as serious in Holland, but details are lacking.
- PSEUDOTSUGA**. DOUGLAS FIR. Timber and ornamental coniferous trees.
- Lophodermium abietis** Rostr. See *Picea*.
- Myxosporium abietinum** Rostr. See *Larix*.
- Oospora abietum** Oudem. See *Abies*.
- Phomopsis pseudotsugae** Wils. This a new and apparently threatening disease of *P. taxifolia* in Great Britain and Holland. It also attacks *Larix kaempferi* (*L. leptolepis*), *L. decidua*, *Abies grandis*, *A. pectinata*, and *Tsuga heterophylla*.
- Cankers appear on the branches or stems, causing constrictions and the death of the portion of branch or stem beyond the canker. The cankers may occur along one side only, in which case death of the branch is delayed. The minute black fruiting bodies form on the cankers and enlarged gum pockets appear in the swollen portions of the limbs or trunk just above the diseased areas. Diseased trees have an unthrifty appearance, due to the yellowing of the foliage of diseased branches, and younger trees, particularly nursery stock, are killed.
- Phomopsis pithya** (Sacc.) Grove. This fungus, related to the preceding one, occurs on *Pinus*, *Pseudotsuga*, and other conifers in Europe but is generally considered as saprophytic only.
- PSIDIUM**. GUAVA. Tropical and subtropical fruit trees and shrubs.
- Atholaderma clavatisporum** Syd. On leaves of *P. guajava* in Siam.
- Catacauma goyazense** (P. Henn.) Theiss. and Syd. Tar spot on leaves of *P. guajava* in Paraguay and Brazil.
- Catacauma subcircinans** (Speg.) Theiss. and Syd. Circular black shiny stromata on leaves of *P. cattleianum* in Brazil and Argentina.
- Cercospora psidii** Rangel. On leaves of *P. araca* in Brazil.
- Linhartia hohnellii** Rehm. Small circular leaf spots on *Psidium* sp. in Brazil.
- Meliola psidii** Wint. Superficial black circular to irregular fungus patches on leaves of *P. guajava* and *P. pomiferum* in Central and South America and the West Indies.
- Phyllachora cayennensis** (DC.) Theiss. and Syd. See *Ficus*.
- Phyllachora tropicalis** Speg. Subcircular black stromata on leaves of *P. thea* in Brazil and Argentina.
- Phyllosticta psidiella** Tassi. Small irregular dirty-white leaf spots on *P. montanum* in Italy.
- Phyllosticta psidii** Tassi. Dull-brown leaf spots on *P. araca* in Italy.
- Puccinia psidii** Wint. Yellow to brown rust pustules on dull-brown or brown-purple leaf spots on *P. araca*, *P. guajava*, *P. pomiferum*, and *Eugenia* (*Caryophyllus*) *jambos* in Porto Rico, Cuba, Ecuador, and Brazil.
- PSOPHOCARPUS**. BOTOR. Ag. Tall, twining herbs with large tuberous roots.
- Woroninella psophocarpi** Racib. Forms numerous orange galls on leaves, stems, and fruit of *P. longipedunculatus*, *P. palustris*, and *P. tetragonolobus* in Java, India, the Philippines, and west tropical Africa.
- PSORALEA**. SCURF-PEA. Herbs and shrubs used somewhat in border plantings.
- Erysiphe taurica** Lévl. See *Althaea*.
- Phlytaena psoraleae** (Cast.) Karst. and Har. On stems of *Psoralea* sp. in France.
- Woroninella aequatoriensis** Syd. On leaves of *P. mutisi* in Ecuador.
- PSYCHOTRIA**. Tropical shrubs and small trees.
- Aecidium iquitosense** P. Henn. Yellow rust sori deforming leaves and stems of *Psychotria* sp. in Peru and Brazil.
- Aecidium micranthum** Syd. Leaf rust on *P. elongata* in Ceylon.
- Aecidium psychotriae** P. Henn. Leaf rust on *Psychotria* sp. in Brazil.
- Inocyclus psychotriae** Syd. On leaves of *P. luzonensis* in the Philippines.
- Phyllachora psychotriae** Rehm. Scattered black stromata on circular yellowish leaf spots on *Psychotria* sp. in Brazil.
- Phyllosticta psychotriae** P. Henn. On leaves of *Psychotria* sp. in Brazil.
- Puccinia fallaciosa** Arth. Brown leaf rust on leaves of *Palicourea crocea*, *P. riparia*, and *Psychotria patens* in Porto Rico.
- Puccinia psychotriae** P. Henn. Brown rust pustules on leaves of *Psychotria* sp. in Brazil.
- Uredo hoistii** P. Henn. Leaf rust on *Psychotria* sp. in tropical Africa.
- Uredo mkusienensis** P. Henn. Leaf rust on *Psychotria* sp. in Uganda and Tanganyika.
- Uredo psychotriae-volkensii** P. Henn. Yellow rust pustules on circular, then confluent, dull-brown leaf spots on *P. volkensii* in tropical Africa.
- Uredo psychotricola** P. Henn. Leaf rust on *Psychotria* sp. in Brazil.
- Uromyces psychotriae** P. Henn. Leaf rust on *Psychotria* sp. in Brazil.
- PTELEA**. HOP TREE. Deciduous shrubs and small trees with handsome foliage and attractive green fruit.
- Ascochyta pteleae** Kab. and Bub. Circular leather-brown, often zoned, leaf spots with narrow dark-brown margins on *P. trifoliata* in Bohemia.
- Phyllosticta pteleae** Höll. Large irregular, then confluent, brown leaf spots on *P. trifoliata* in Hungary.

PTERIS. BRAKE. A large genus of ferns, many forms cultivated.

Apelenechus olestinus Ritz. Bos. See Begonia.

Ascochyta pteridis Bres. On leaves of *Pteridium aquilinum* (*P. aquilinum*) in Germany.

Dothiella pterodophila Speg. Black stromata covering leaves of *Pteris* sp. in Brazil.

Hyalopora filicum Diet. See Asplenium.

Mycosphaerella prominula (Speg.) Lind. On leaves of *P. aquilinum* in Europe.

Phyllachora rhophographoides Wint. Doubtful species on *P. aquilinum* in Brazil.

Rhopographus malmi Starb. Linear black stromata on leaves of *Pteris* sp. in Brazil.

Ehytisma piceum Berk. Black opaque stromata on leaves of *Pteris* sp. in India.

Sestoria aquilina Pass. Brown spots on leaves of *P. aquilina* in Italy. Reported from Wisconsin.

Sestoria pteridicola Kab. and Bub. Pale-brown or dull-yellow irregular, then confluent, areas on leaves of *P. aquilina* in Italy.

Taphrina laurencia Giesenh. Causes much-branched leafy outgrowths on *P. quadriaurita* in Ceylon and India.

Taphrina rhomboidalis Syd. and Butl. Yellow to brown areas on leaves of *P. quadriaurita* in India.

Taphrina tonduziana P. Henn. Circular to angular spots on leaves of *P. aculeata* in Costa Rica.

PTEROCARPUS. Leguminous trees or woody climbers.

Aldona stella-nigra Rac. On leaves of *P. indicus* in Java and the Philippines.

Catacauma pterocarpis Syd. Small irregular black stromata on white leaf spots on *P. angolensis*, *P. indicus*, and *P. saratilis* in Java, the Philippines, and the Union of South Africa.

Helminthosporium obovatum Mass. On leaves of *P. indicus* in Malaya.

Placostroma pterocarpis (Mass.) Theiss. and Syd. Black stromata on yellow-green leaf spots on *P. indicus* in Malaya and the Philippines.

Pseudothia pterocarpis Syd. On leaves of *P. indicus* in the Philippines.

Uncinula incrassata Salm. Powdery mildew on leaves of *P. melliferus* in central Africa.

PTEROCARYA. WING NUT. Ornamental trees grown for their handsome pinnate foliage.

Phyllosticta pterocaryae Thuem. Irregular, often marginal dull-yellow to gray leaf spots on *P. caucastica* in Portugal.

PTYCHOSPELMA. See Palmae.

PUERARIA. KUDZU BEAN. Twining herbs or shrubs.

Cercospora puerariae Syd. Olive-brown leaf spots on *P. phaseoloides* in the Philippines.

Synchytrium puerariae Miy. (*Woroninella puerariae* [Henn.] Syd.). Small galls on leaves and stems of *P. hirsuta*, *P. phaseoloides*, and *P. sericea* in Japan, China, Java, India, New Guinea, and the Philippines.

PULMONARIA. LUNGWORT. Perennial herbs with creeping root-stocks, grown in gardens.

Entyloma serotinum Schroet. See Borago.

Phyllosticta pulmonariae (Fckl.) Sacc. Small, dull-brown, leaf spots on *Pulmonaria* sp. in Switzerland.

Puccinia bromina Erikss. See Bromus.

Ramularia cylindroides Sacc. Dull-yellow leaf spots on *P. mollissima*, *P. officinalis*, *P. styriaca*, *P. tuberosa* and *Verbascum* sp. in Europe.

Sestoria pulmonariae Sacc. Subcircular dark-brown leaf spots on *P. officinalis* and *Pulmonaria* sp. in Russia and Italy.

PUNICA. POMEGRANATE. Small trees with edible fruit.

Aposphaeria punicea Sacc. Found in connection with die-back of twigs of *P. granatum* in southern Europe.

Cercospora punicea P. Henn. Circular dark-fusces leaf spots on *P. granatum* in China and Japan.

Dothiorella sanninii Ciferri. Surface of fruit of *P. granatum* in Italy covered with small black patches which disfigure and lower the commercial value.

Megalonectria caespitosa Speg. Forms cankers on trunk and branches of *P. granatum* in Brazil and Argentina.

Phyllosticta granati Rangel. Circular to angular brown leaf spots on *P. granatum* in Brazil.

Phyllosticta punica Sacc. and Speg. Small subcircular white leaf spots with red margins on *P. granatum* in Italy.

PYCNOSTACHYS. Erect perennial herbs.

Uredo pycnostachydis Kalchbr. Leaf rust on *P. dawei* and *P. reticulata* in Uganda.

PYRETHRUM. See also Chrysanthemum.

Puccinia balsamitae (Strauss.) Rabh. Leaf rust on *P. (Chrysanthemum) balsamita*, *P. tanacetifolium*, and *Tanacetum* sp. in Europe.

Puccinia proximella Syd. Leaf rust on *P. millefolium* in Russia.

Sestoria pyrethri Bres. and Krieg. Irregular buff leaf spots on *P. macrophyllum* and *P. (Chrysanthemum) parthenium* in Russia and Germany.

PYROLA. Shinleaf. Low, smooth, perennial herbs with subterranean shoots.

Actinonema pyrolae Allesch. Large irregular black leaf spots on *P. secunda* in Germany.

Phyllosticta pyrolae (Ehrbg.) Allesch. Circular, often confluent, purple-brown leaf spots on *P. rotundifolia* in Germany.

Sclerotinia pyrolae Grosse. Forms black sclerotia in pods of *P. chlorantha*, *P. media*, *P. minor*, *P. rotundifolia* and *P. uniflora* in Russia.

PYRUS. PEAR. Trees and shrubs cultivated for fruit and ornament.

Ascochyta pirina Pegl. Irregular brown spots on fruit and leaves of *P. communis* in Russia, Malta, and Italy.

Bacillus sp. (Blossom Blight.) This bacterium causes a blighting of flowers and sometimes a die-back of twigs of *Pyrus*, *Malus*, and *Prunus* spp. in England. The sepals turn gray and blacken, the discoloration involves the calyx and flower stalk, and the entire blossom turns black, sometimes falling.

Bacterium nectarophilum Doidge. Causes a blackening of flowers, peduncles, and young fruit of *Pyrus* in the Union of South Africa. Distinct from fire blight.

Bacterium pyri Brzez. This bacterium is reported as the cause of a trunk and branch canker of *Pyrus* (pear) in central Europe.

Cercospora piricola K. Saw. Angular, then coalescent, brown and finally ashen leaf spots on *P. communis* and *P. sinensis* in Japan.

Cercospora porriga Speg. Leaf spot on *P. communis* in Argentina.

Colletotrichum piri Noack and f. *tirolense* Bub. Anthracnose on *Pyrus* and *Malus* in Brazil and Austria.

Coniothecium chromatoporum Oda. See Malus.

Coryneum follicolum Fckl. See Crataegus.

Diaporthe perniciosus Marchal. See Prunus.

Dothiorella vinosa Marchal. See Prunus.

Entomosporium mespill (DC.) Sacc. See Cotoneaster.

PYRUS—Continued.

- Exoascus bullatus** (B. and Br.) Fekl. Small, then confluent, reddish blisterlike swellings on leaves of *P. amygdaliformis*, *P. communis*, *P. japonica*, *P. sinensis*, *P. ussuriensis*, and *Cydonia japonica* in Australia, Chile, and Europe.
- Gymnosporangium cunninghamianum** Barcl. See *Cupressus*.
- Gymnosporangium koreanse** (P. Henn.) Jacks. See *Juniperus*.
- Gymnosporangium mespili** (DC.) Kern. See *Juniperus*.
- Gymnosporangium sabinæ** (Dicks.) Wint. See *Juniperus*.
- Gymnosporangium yamadæ** Miy. See *Juniperus*.
- Hadrotrichum piri** Montem. Small gray circular, sometimes confluent, leaf spots on *Pyrus* sp. in Italy.
- Helicobasidium mompa**. See *Morus*.
- Helicobasidium tanakæ** Miy. See *Morus*.
- Labrella picicola** Bres. and Sacc. On leaves of *P. communis* and *Malus* in Germany.
- Laestadia parmensis** Pass. On leaves of *P. communis* in Italy.
- Ochropsora sorbi** Diet. See *Sorbus*.
- Oospora picicola** Nob. Gray to white patches on leaves and branches of *Pyrus* in France.
- Phaciella discolor** (Mont. and Sacc.) A. Poteb. See *Malus*.
- Phyllosticta briardi** Sacc. See *Malus*.
- Phyllosticta mali** Prill. and Delacr. See *Malus*.
- Phyllosticta picicola** Sacc. and Speg. On leaves of *Pyrus* in Europe.
- Phyllosticta tirolensis** Bub. Small subcircular to irregular gray leaf spots on *P. communis* in Austria.
- Phytophthora syringæ** Kleb. See *Syringa*.
- Plectodiscella piri** Wor. Circular to oval grayish-white leaf spots on *P. communis* and *Malus* sp. in southern Russia.
- Ramularia magnusiana** Sacc. On leaves of *Pyrus* in Belgium.
- Septoria nigerrima** Fekl. On leaves of *P. communis* in Yugoslavia, France, and Italy. Reported from New York.
- Septoria ralfsii** B. and Br. Black irregular spots on fruit of *Pyrus* in Great Britain, with subsequent rot.
- Stagonospora mali** Delacr. On leaves of *Pyrus* and *Malus* in France.
- Sterium purpureum** Pers. See *Prunus*.
- Taphrina piri** S. Kus. Circular to irregular yellow-green leaf spots on which yellowish or white scurfy appear. These areas may spread over most of the leaf surface and cause distortion. On *P. miyabei* in Japan.
- QUAMOCLIT.** STAR GLORY. Annual or perennial twining vines.
- Phyllosticta quamoclit** Thuem. Dull-yellow to brown irregular leaf spots with gray margins on *Q. coccinea* in Portugal.
- QUERCUS.** OAK. Ornamental and timber trees.
- Cocchoarella quercicola** (P. Henn.) v. Hoeh. Shiny black circular stromata on leaves of *Q. thalassica* in Japan.
- Coccodiscus quercicola** P. Henn. Black stromata on leaves of *Q. thalassica* in Japan.
- Coccohora kusanoi** (P. Henn.) v. Hoehn. Irregular black shiny stromata on leaves of *Q. glauca* in Japan.
- Coccoidea quercicola** P. Henn. and Shir. Small black stromata on leaves of *Q. glabra* and *Q. glauca* in Japan.
- Coryneum foliolum** Fekl. See *Crataegus*.
- Coryneum umbonatum** Nees. Kills twigs and branches of *Q. castaneafolia* in France.
- Cystotheca wrightii** B. and C. Superficial brown perithecia on leaves of *Q. acuta*, *Q. glauca*, *Q. myrsinaefolia*, and *Q. vibrayana* in Japan.
- Dermatea cinnamomea** (Pers.) Rehm. Wound parasite on *Quercus* spp. in Europe.
- Diaportha taleola** (Fr.) Sacc. On branches of *Quercus* spp. in Europe.
- Exoascus krukii** Vuill. Causes a witches'-broom effect on *Q. ilex* in France and Italy.
- Gloeosporium shiratanum** Syd. Circular ochraceous to brown leaf spots with pale centers on *Q. glandulifera* in Japan.
- Gloeosporium umbrinellum** B. and Br. Small irregular brown leaf spots on *Q. robur*, *Q. sessiliflora*, and *Quercus* sp. in Russia, Switzerland, Great Britain, and Germany.
- Gnomonia quercus-iliici** Berl. On leaves of *Quercus* spp. in Italy.
- Guignardia cincta** (Desm.) Lind. Subcircular to irregular pale-rufous leaf spots with brown margins on *Q. coccifera* in France.
- Guignardia diffusa** (Crié) Sacc. and Trott. See *Castanea*.
- Guignardia punctoidea** (Cke.) Schroet. On branches of *Quercus* sp. in northern Europe.
- Hadronema orbiculare** Syd. Small circular, velvety fungus patches on leaves of *Q. glauca* and *Q. luzoniensis* in Japan and the Philippines.
- Helicobasidium mompa** Tan. See *Morus*.
- Heterosporium proteus** Starb. Brown sunken leaf spots on *Quercus* sp. in Sweden.
- Hydnum obrutans** Burt. Causes heart rot of *Quercus* sp. in Java.
- Japonia quercus** v. Hoehn. On leaves of *Q. glauca* in Japan.
- Laestadia ceris** Pass. Pale-brown leaf spots on *Q. cerris* in Italy.
- Laestadia cookeana** (Auersw.) Sacc. On leaves of *Q. ilex* and *Q. robur* in Europe.
- Laetodermia flavo-virens** Dur. and Mont. On leaves of *Quercus* spp. in Spain and Algeria.
- Macrophoma cylindrospora** (Desm.) Berl. and Vogl. See *Hedera*.
- Macrophoma fusispora** Bub. Subcircular to elongate whitish leaf spots on *Q. conferta* in Hungary.
- Macrophoma mexicana** Sacc. On leaves of *Q. glauca* in Mexico.
- Melogramma henricquetii** Br. and Cav. On *Q. suber* in southern Europe.
- Microporella quercus** v. Hoeh. On leaves of *Q. glauca* in Japan.
- Microstroma album** (Desm.) Sacc. and var. *japonicum* P. Henn. On leaves of *Q. cerris*, *Q. glauca*, *Q. grosseserrata*, and *Q. sessiliflora* in Europe, Japan, Argentina, and the Union of South Africa.
- Monochaetia pachyspora** Bub. Subcircular to irregular pale grayish-yellow leaf spots on *Q. ilex* in Austria.
- Phelodermia cuticularia** Duc. Brown leaf spots on *Q. suber* in France.
- Myceoloma cupulicola** Gz. Frag. On *Quercus* sp. in Spain.
- Phyllohendersonia foliorum** (Fekl.) Tass. On leaves of *Q. ilex* and *Quercus* sp. in Italy, France, and Germany.
- Phyllosticta associata** Bub. On leaves of *Q. robur* (*Q. pedunculata*) in Austria.
- Phyllosticta bresadoleana** Bub. and Kab. Brown or yellow-gray leaf spots with narrow brown-purple margins on *Q. pubescens* in Austria.
- Phyllosticta globulosa** Thuem. Irregular gray-white leaf spots on *Q. robur* (*Q. pedunculata*) in Austria.

QUERCUS—Continued.

- Phyllosticta hrancensis** Petr. On leaves of *Q. robur* in Bohemia.
- Phyllosticta ilicicola** Pass. Circular brown, then ashen-gray, leaf spots on *Q. ilex* in France and Italy.
- Phyllosticta ilicina** Sacc. Pale-brown leaf spots on *Q. ilex* in Italy, Portugal, and Austria.
- Phyllosticta iliciseda** Sacc. Broad, irregular leaf spots with black borders on *Q. ilex* in Italy.
- Phyllosticta macrocarpae** Monte. Brown irregular leaf spots on *Q. macrocarpa* in Italy.
- Phyllosticta phaeospora** Scalia. On leaves of *Q. ilex* in Italy.
- Phyllosticta quercicola** Oud. On leaves of *Quercus* sp. in northern Europe.
- Phyllosticta quercus-cocciferae** Bub. Circular to irregular white leaf spots on *Q. coccifera* in Yugoslavia.
- Phyllosticta roboris** Oud. Small angular, often confluent, pale-brown leaf spots on *Q. robur* in Bohemia and Holland.
- Phyllosticta shiraiana** Syd. Circular ochraceous to white leaf spots on *Q. acuta* and *Q. glauca* in Japan.
- Plowrightia noxia** (Ruhl.) Sacc. See *Castanea*.
- Rosellinia quercina** Hartig. Attacks roots of young *Quercus* seedlings in nurseries in Germany. Infected plants show an unhealthy browning of the foliage, followed by withering of young shoots and ultimate death.
- Sclerotinia candoileana** (Lév.) Fekl. Leaves of *Quercus* and *Castanea* in Great Britain, France and Germany turn brown and fall prematurely due to this fungus.
- Septoria dryophila** Sacc. White leaf spots with narrow dark margins on *Q. ilex* in Italy.
- Septoria dubia** Sacc. and Syd. Small yellowish leaf spots with red margins on *Q. pedunculata* and *Q. robur* (*Q. pubescens*) in France, Russia, Yugoslavia, and Germany.
- Septoria quercicola** Sacc. White leaf spots on *Q. humilis*, *Q. pedunculata*, and *Q. sessiliflora* in France, Italy, and Portugal. Reported from Minnesota.
- Septoria quercina** Desm. Small circular white leaf spots on *Q. robur*, *Q. sessiliflora*, and *Quercus* sp. in Japan, Russia, France, Italy, Denmark, Austria, and Germany.
- Septoria quercus** Thuem. Subcircular ochraceous leaf spots on *Q. pedunculata* and *Q. sessiliflora* in Portugal.
- Sphaerotheca kusanoi** P. Henn. and Shir. Powdery mildew on leaves of *Q. glandulifera* in Japan.
- Trabutia conzattiana** Sacc. Reddish-brown circular, sometimes confluent, leaf spots on *Quercus* sp. in Mexico.
- Typhulochaeta japonica** Ito and Hara. On leaves of *Q. glandulifera* in Japan.
- Uncinula septata** Salm. Powdery mildew on leaves of *Q. glandulifera* in Japan.
- Uredo quercus** Brond. Small circular rust pustules on lower leaf surfaces of *Quercus* spp. in Europe.
- Yoshinagaia quercus** P. Henn. Black subcarbonaceous stromata on upper leaf surfaces of *Q. glauca* in Japan.
- Yoshinagella japonica** v. Hoch. Irregular black stromata on upper leaf surfaces of *Q. glauca* in Japan.

RADICULA. HORSE-RADISH. WATER CRESS. (Including *Roripa* and *Armoracia*.)

- Ascochyta armoraciae** Fekl. Scattered pale-brown leaf spots on *Armoracia rusticana* (*R. armoracia*) in Europe.
- Ascochyta rusticana** Kab. and Bub. Ochraceous or brown leaf spots varying in size from 0.5 to 2 centimeters, on *R. armoracia* in Bohemia.
- Peronospora nasturtii aquatici** Gaum. Downy mildew on leaves of *Sisymbrium nasturtium-aquaticum* (*R. nasturtium-aquaticum*) in Germany.
- Phyllosticta armoraciae** Cke. Circular to angular dirty-white leaf spots with narrow brown margins on *R. armoracia* in Great Britain.
- Ramularia cochleariae** Cke. See *Barbarea*.
- Septoria armoraciae** Sacc. Irregular ochraceous leaf spots on *R. armoracia* in Italy.

RANDIA. Erect or climbing tropical trees or shrubs.

- Aecidium abscondens** Arth. Yellow rust pustules on leaves of *R. aculeata* in Porto Rico and Costa Rica.
- Aecidium pulverulentum** Arth. Leaf rust on *Randia* sp. in Mexico.
- Aecidium randiae** P. Henn. Yellow rust pustules on dull-brown, swollen leaf spots on *Randia* sp. in Brazil.
- Corticium salmonicolor** B. and Br. See *Citrus*.
- Endophyllum griffithiae** Rac. Yellow rust sori on circular leaf spots on *R. scandens*, *R. sinensis*, and *Peretta indica* in Java and China.
- Lembosia philippinensis** Syd. On leaves of *Randia* sp. in the Philippines.
- Pestalozzia paucisetia** Sacc. Gray leaf spots on *R. reticulata* in the Philippines.
- Taphrina randiae** Rehm. Elliptical black, swollen leaf spots on *Randia* sp. in Brazil.
- Trabutia randiae** (Rehm.) Theiss. and Syd. and var. *aculeatae* Ferd. and Winge. Black stromata on leaves of *R. aculeata* and *R. pubescens* in Bolivia and the American Virgin Islands.

RANUNCULUS. (Including *Ficaria*.) BUTTERCUP. Crowfoot. Annual and perennial herbs.

- Aphelenchus oleisitis** Ritz. Bos. See *Begonia*.
- Ascochyta carinthiaca** Jaap. On leaves of *R. thora* in Austria.
- Cercospora ranunculi** Jaap. Circular to elongate brown leaf spots on *R. muricatus* in Dalmatia.
- Cylindrosporium ficariae** Berk. On leaves of *R. ficaria* in France and Great Britain.
- Cylindrosporium ranunculi** Sacc. and *sclerati* P. Brun. On leaves of *R. acris*, *R. bulbosus*, and *R. scleratus* in France and Italy.
- Entyloma ameghinoi** Speg. Leaf smut on *R. cymbalaria* in Patagonia.
- Fabraea litigiosa** (Rob. and Desm.) Sacc. On leaves of *Ranunculus* spp. in Alaska, Europe, and north Africa.
- Fusidium eburneum** Schroet. Subcircular, often confluent, white leaf spots with yellow centers on *R. acris* and *R. repens* in Siberia and Austria.
- Heterosporium ferox** Bub. On leaves of *R. arvensis* in Bohemia.
- Leptosphaeria thorae** Jaap. On leaves of *R. thora* in Austria.
- Oularia declipensis** Sacc. On leaves of *R. acris* in Russia, Dalmatia, and Italy. Reported from New York.
- Peronospora alpicola** Gaum. Downy mildew on leaves of *R. aconitifolius*, *R. pyrenaicus*, and *R. sequieri* in France, Switzerland, Austria, and Germany.
- Peronospora ficariae** (Nees.) Tul. As above on *R. ficaria* in Europe.
- Peronospora gigantea** Gaum. Downy mildew on leaves of *R. lingua* in Denmark and Russia.
- Peronospora glacialis** (Blytt.) Gaum. Downy mildew on leaves of *R. glacialis* in Switzerland and Norway.
- Peronospora illyrica** Gaum. As above, on *R. illyricus* in Russia.
- Phyllosticta ficariae** Maire. On leaves of *R. calthaeifolius* in French north Africa.
- Phyllosticta pygmaea** Allesch. On leaves of *R. pygmaeus* in Greenland.

RANUNCULUS—Continued.

- Phyllosticta ranunculi** (Fckl.) Sacc. On leaves of *R. acris* in Germany.
- Phyllosticta ranunculeorum** Sacc. and Speg. Dull-brown leaf spots on *R. repens* in Italy.
- Physoderma vagans** Schroet. Small tubercles on leaves and stems of *R. acris*, *R. flammula*, *R. repens*, *Oenanthe phellandrium*, *Potentilla anserina*, *Selinum lineare*, *Silvaus* sp., and *Stum latifolium* in France and Germany.
- Pseudopeziza ranunculi** (Wallbr.) Fckl. Black circular fruiting bodies in dark-brown sunken leaf spots on *R. cassubicus*, *R. chaerophyllus*, *R. nemorosus*, and *R. polyanthemus* in Belgium, France, and Germany.
- Puccinia andina** Diet. and Neg. Leaf rust on *R. peduncularis* in South America.
- Puccinia hlyttiana** Lagh. Brown rust pustules on leaf blades and petioles of *R. alpestris* and *R. auricomus* in Norway and Switzerland.
- Puccinia gibberulosa** Schroet. Powdery brown rust pustules on leaf blades and petioles of *R. lancipetalus* and *Ranunculus* sp. in France and Argentina.
- Puccinia magnusiana** Koern. See Phragmites.
- Puccinia nubigena** Speg. Rust on leaf blades and petioles of *Ranunculus* sp. in Argentina.
- Puccinia ustalis** Berk. Dark-brown rust pustules on *R. hirtellus*, *R. pulchellus*, and *R. songaricus* in Mongolia, Turkestan, and India.
- Ramularia acris** Lindr. Large irregular yellow-brown to gray-brown leaf spots on *R. acris* in Finland and Sweden.
- Ramularia aequivoca** (Ces.) Sacc. On leaves of *R. auricomus*, *R. cassubicus*, *R. lanuginosus*, *R. polyanthemus*, *R. repens*, and *R. trachycarpus* in Europe. Reported from Wisconsin.
- Ramularia lapponica** Lindr. Irregular gray or dark-brown leaf spots on *R. lapponicus* in Finland.
- Ramularia repens** Oud. Elliptical to irregular leaf spots on *R. repens* in Holland.
- Ramularia sclerata** Cke. Elongate brown spots on lower leaves of *R. scleratus* in Great Britain.
- Ramularia serbica** Ranof. Irregular brown leaf spots on *R. montanus* in Yugoslavia.
- Sclerotinia ficariae** Rehm. On leaves of *R. ficaria* in Italy.
- Septoria cymbalaria** Thuem. On leaves of *R. cymbalaria* in Siberia.
- Septoria ficariae** Desm. Ashen leaf spots on *R. ficaria* and *R. vernus* in Europe.
- Septoria oreophila** Sacc. Small white leaf spots on *R. aconitifolius* in Italy.
- Septoria polaris** Karst. White leaf spots on *R. flammula* and *R. parviflorus* in Great Britain and Scandinavia.
- Septoria ranunculacearum** Lév. Brown leaf spots on *R. acris*, *R. cymbalaria*, and *R. pusillus* in France, Italy, and Siberia.
- Septoria ranunculi** West. Circular greenish-yellow leaf spots on *R. scleratus* in Belgium.
- Synchytrium andinum** Lagerh. Red-brown warty areas on leaves of *Ranunculus* sp. in Ecuador.
- Uromyces dactylidis** Oth. See Dactylis.
- Uromyces festucae** Syd. See Festuca.
- Uromyces ficariae** (Schum.) Lév. Leaf rust on *R. ficaria*, *R. grandiflora*, and *R. vernus* in Europe.
- Uromyces fischerianus** E. Mayor. Yellow-brown rust pustules on leaf blades and petioles of *R. glacialis* in Switzerland.
- Uromyces poae** Rabb. See Poa.
- Uromyces ranunculi-distichophylli** Semad. Leaf rust on *R. parnassifolius* in Switzerland.
- Uromyces ranunculi-festucae** Jaap. See Festuca.
- RAPHANUS**. RADISH. Vegetable and forage plants.
- Moniliopsis aderholdi** Ruhl. See Solanum.
- Pseudomonas destructans** Potter. See Brassica.
- Puccinia isiacae** (Thuem.) Wint. See Arundo.
- RAUWOLFIA**. Ornamental trees or shrubs.
- Aecidium rauwolfiae** F. Henn. Leaf rust on *Rauwolfia* sp. in the Congo.
- RAVENALA**. TRAVELER'S-TREE. Large *Musa*-like plants.
- Catacauma ravenala** (Pat. and Har.) Theiss. and Syd. Black stromata on leaves of *R. madagascariensis* in Madagascar.
- Zukalia stuhlmanniana** P. Henn. See Palmae.
- RENANTHERA**. Tall climbing epiphytic orchids. See Orchidaceae.
- RENEALMIA**. Perennial aromatic herbs.
- Catacauma renealmiae** (Rehm.) Theiss. and Syd. Circular black stromata on brown leaf spots on *Renealmia* sp. in Brazil.
- Endothella renealmiae** (Rehm.) Theiss. and Syd. Circular to irregular black stromata on leaves of *Renealmia* sp. in Brazil.
- Septoria renealmiae** Tassi. Large brown leaf spots on *R. cinnamomea* in Italy.
- RESEDA**. MIGNONETTE. Garden annuals.
- Peronospora crispula** Fckl. Downy mildew on leaves of *R. lutea* and *R. luteola* in France, Denmark, Switzerland, and Germany.
- Phyllosticta resedae** Petch. On leaves of *R. odorata* in Ceylon.
- RETINOSPORA**. See Chamaecyparis.
- RHAMNUS**. BUCKTHORN. Woody plants grown for their foliage and some species for their attractive fruit and medicinal bark.
- Asechyta frangulina** Kab. and Bub. On leaves of *R. cathartica* and *R. frangula* in Russia and Bohemia.
- Endophyllum elegans** (Diet.) Pole-Evans. Leaf rust on *R. prinoides* in the Union of South Africa.
- Mycosphaerella vogelii** Syd. On leaves of *R. cathartica* in Germany.
- Phyllosticta advena** Sacc. Oval to indefinite ashen-white leaf spots with dull-brown margins on *R. (Sageretia) corymbosa* in Italy.
- Phyllosticta alaterni** Pass. Circular grayish-white leaf spots with dark-brown margins on *R. alaternus* and *R. frangula* in France and Italy.
- Phyllosticta cathartici** Sacc. Subcircular ochraceous leaf spots with dull-red margins on *R. cathartica* in Russia, Italy, Denmark, and Austria.
- Phyllosticta desmazieri** Tass. On leaves of *R. cathartica* in France.
- Phyllosticta jahnlana** Petr. and Sacc. Subelliptical to irregular black leaf spots on *R. cathartica* in Bohemia.
- Phyllosticta osteospora** Sacc. See Morus.
- Phyllosticta rhamnicola** Desm. Circular ashen leaf spots on *R. alpina* in France and Belgium.
- Phyllostictella rhamnigena** (Sacc.) Tass. Subcircular dull-white leaf spots with dull-brown margins on *R. alaternus* and *R. cathartica* in Portugal, Italy, and France.
- Puccinia himalensis** (Barcl.) Diet. Brown to black rust pustules on leaves and twigs of *R. dahurica* in India. Aerial stage on leaf blades and sheaths of *Brachypodium silvaticum*.

RHAMNUS—Continued.

Puccinia schweinfurthii (P. Henn.) P. Magn. Brown crustlike rust pustules deforming the leaves of *R. prinoides* and *R. stado* in Abyssinia and east Africa.

Ramularia alaterni Thuem. Large gray spots on leaves of *R. alaternus* in Spain and France.

Rostrupia schweinfurthii P. Henn. On leaves of *Rhamnus* sp. in Abyssinia.

Septoria alaterni Pass. and var. *hispanica* Gz. Frag. Gray leaf spots on *R. alaternus* in Spain and Italy.

Septoria cathartica Pass. Irregular gray leaf spots on *R. cathartica* and *R. saxatilis* in Denmark and Italy.

Septoria frangulae Guep. Small dark-rufous leaf spots on *R. frangula* in France, Bohemia, Silesia, Poland, and Germany.

RHAPHIDOPHYLLUM. See *Palmae*.**RHAPONTICUM**. See, also, *Centaurea*.

Puccinia rhapsontici Syd. Powdery dark-brown rust pustules on leaves of *R. acaule* and *R. pusillum* in north Africa.

RHEUM. *RHUBARB*. Herbs, grown for their foliage and edible leaf stocks.

Bacterium rhapsonticum Millard. This bacterium is reported as the cause of crown rot of *R. rhapsonticum* in Great Britain. A soft brown rot attacks the crowns and sheathing leaves are discolored and swollen at the base where joined to the crowns. The leaves turn dark-brown and in the absence of terminal buds only spindling shoots are produced from lateral buds.

Cercospora rhei Grog. On leaves of *R. officinale* in France.

Peronospora jaipiana Magn. Downy mildew forming large irregular gray patches on leaves of *R. officinale*, *R. palmatum*, and *R. undulatum* in Switzerland and Germany.

Ramularia rhei Allesch. Circular, then confluent, red-brown leaf spots on *R. undulatum* and *Rheum* (cult.) in Russia, Denmark, and Germany.

Uredo rhei-undulati Diet. Brown leaf rust on *R. undulatum* in Japan.

RHIPHALIS. Epiphytic succulent plants.

Phaeospora caticola Stevens. On *R. cassytha* in Porto Rico.

RHODODENDRON. (Including *Azalea*.) Woody plants cultivated for their flowers and foliage.

Ascochyta rhododendri Lind. Angular brown leaf spots on *Rhododendron* (cult.) in Denmark.

Cercospora handelii Bub. Irregular, then confluent, ochraceous spots destroying leaves of *R. ponticum* in Asia Minor.

Chrysomyxa dietelii Syd. Yellow rust pustules on leaves of *R. arboreum* in India.

Chrysomyxa expansa Diet. Yellow rust pustules on red-brown leaf spots on *R. metternichii* in Japan.

Chrysomyxa himalaensis Barcl. Rust on leaf blades and petioles of *R. arboreum*, *R. campanulatum*, and *R. hodgsonum* in India.

Chrysomyxa rhododendri (DC.) De By. Yellow to red-brown rust pustules on *R. brachycarpum*, *R. dauricum*, *R. ferrugineum*, *R. hirsutum*, *R. intermedium*, *R. kotschyum*, *R. myrsifolium*, and *R. suave* in Japan, Siberia, and Europe. The aecal stage occurs on *Picea abies*, *P. obovata*, and *P. pungens*.

Exobasidium butleri Syd. Flethy growths on lower leaf surfaces of *R. arboreum* in India.

Exobasidium hemisphaericum Shir. Flethy pyriform to globose galls on lower leaf surfaces of *R. metternichii* and *R. pentamerum* in Japan.

Exobasidium japonicum Shir. Flethy, smooth, globose, greenish, then reddish-white, galls on leaves of *R. indicum* in Japan.

Exobasidium vexans Mass. See *Thea*.

Exobasidium yoshinagai P. Henn. Circular to effuse dull-brown leaf spots on *R. tosaense* and *Azalea* sp. in Japan.

Exobasidium zeylanicum Petch. Large spherical white or greenish-white, then powdery white galls on lower leaf surfaces of *R. arboreum* in Ceylon.

Gloeosporium rhododendri Br. and Cav. Large irregular blotches on leaves of *R. ponticum* in Italy and Australia.

Gloeosporium succineum Sacc. Anthracnose on leaves of *R. chrysanthemum* in Siberia.

Leptosphaeria rhododendri P. Henn. Large brown, then gray, leaf spots on *R. falconeri* in Germany.

Melasmia rhododendri P. Henn. and Shir. On leaves of *R. indicum*, *R. kaempferi*, and *R. tschonoskii* in Japan.

Mycosphaerella occulta Bub. Circular gray leaf spots on *R. ponticum* in Bohemia.

Naemosphaera japonica Sacc. and Syd. On branches of *R. indicum* in Japan.

Phacidium falconeri P. Henn. Gray leaf spots with broad red-brown margins on *R. falconeri* in Germany.

Phyllohendersonia bicolor (Pat.) Tass. On *Rhododendron* sp. in China.

Phyllohendersonia rhododendri (Thuem.) Tass. On leaves of *R. hirsutum* in Italy.

Phyllosticta berolinensis P. Henn. Dull-brown to gray leaf spots on *R. falconeri* in Germany.

Phyllosticta cunninghami Allesch. On leaves of *R. cunninghami* in Germany.

Phyllosticta falconeri P. Henn. Dull-brown to gray circular leaf spots with dark-red margins on *R. falconeri* in Germany.

Phyllosticta rhododendri Brun. Large oblong to irregular ashen-gray leaf spots with narrow brown margins on *Rhododendron* sp. in France.

Phyllosticta rhododendri-flavi Bub. and Kab. Large circular to irregular red-brown, then dark-brown, leaf spots on *R. flavum* in Bohemia.

Phyllosticta rhodorae (Cke.) Tass. On leaves of *Rhododendron* sp. in England.

Phyllosticta saccardoi Thuem. Brown leaf spots on *R. ponticum* in Russia, France, and Portugal. Reported from New York on *Rhododendron* sp.

Physalospora alpina Speg. On leaves of *R. ferrugineum* in Italy.

Physalospora rhododendri Naum. Die-back of twigs and browning of leaves of *Rhododendron* sp. in Germany.

Placosphaeria rhododendri P. Henn. Circular black stromata on leaves of *Rhododendron* sp. in Japan.

Puccinia rhododendri Fekl. Leaf rust on *R. ferrugineum* in Austria.

Rhytisma rhododendri Fr. Black irregular areas on leaves of *Rhododendron* sp. in Siberia.

Sclerotinia rhododendri Fisch. Forming sclerotia in fruit of *R. ferrugineum* and *R. hirsutum* in Europe.

Septoria azaleae Vogl. Die-back of twigs and browning of leaves of *Azalea indica* in Italy and Germany.

Septoria azaleae-indicae Maubl. Circular white leaf spots with black margins on *A. indica* and *Azalea* sp. in Brazil and Germany.

RHODOMYRTUS. *ROSE MYRTLE*. Trees or shrubs native to southern Asia.

Helminthosporium rhodomirti Syd. On leaves of *R. tomentosa* in China.

RHODOTYPOS. *JET BEAD*. Shrubs grown chiefly for their large white flowers and bright-green foliage

Septoria rhodotypi Höll. Irregular rufous leaf spots on *R. kerrioides* in Hungary.

RHUS. SUMAC. Ornamental woody plants, some species of economic importance.

Aecidium foetidum Diet. Leaf rust on *Rhus* sp. in Japan.

Aecidium inecassatum Syd. Leaf rust on *R. sylvestris* in Japan.

Bagnisiopsis rhoina Syd. and Hara. Dull-black circular to elliptical stromata on leaves of *R. sylvestris* in Japan.

Colletotrichum rhoinum F. Tassi. Anthracnose attacking young shoots and leaf blades and petioles of *R. javanica* (*R. semi-alata*) and *R. verniciflua* in Japan. The lesions are long, elliptical, black, and sunken.

Exobasidium hesperidium Maire. Leaf spots, brown-orange on the lower surfaces, yellow-green, then dark-brown, on the upper, depressed and causing rolling of the leaves of *R. oxyacantha* in Algeria.

Helicobasidium mompa Tan. See Morus.

Phylachora marginalis Pat. Irregular black stromata on lower leaf surfaces of *Rhus* sp. in Ecuador.

Phylosticta glabra Brun. Brown angular spots on leaves of *R. glabra* in France.

Phylosticta rhois West. Angular to irregular pale brown leaf spots on *R. coriaria* and *R. cotinus* in Denmark, France, and Belgium.

Pileolaria dieteliana Syd. Leaf rust on *R. hypoleuca* in China.

Septonema henningsii Bresad. On leaves of *R. abyssinica* in Abyssinia.

Septoria acuriana P. Henn. Yellow leaf spots on *R. retinorrhæa* in Abyssinia.

Septoria cotini C. Mass. Irregular, then confluent, reddish leaf spots on *R. cotinus* in Italy.

Septoria rhois Lévl. Dull-brown leaf spots on *R. hirta* (*R. typhina*) in Russia.

Uncinula verniceifera P. Henn. Powdery mildew on leaves of *R. javanica* (*R. semi-alata*) and *R. verniciflua* in Japan.

Uredo rhoina Syd. Yellow-brown leaf rust on *Rhus* sp. in central Africa.

Uromyces barbeyanus P. Henn. Dark-brown rust pustules on leaves of *R. falcata* in Abyssinia.

Uromyces klugkistanus Diet. Leaf rust on *R. javanica* and *R. succedanea* in Japan.

Uromyces propinquus Syd. Powdery brown leaf rust on *R. mollis* in Mexico.

Uromyces shiraianus Diet. and Syd. Powdery brown rust pustules on leaves of *R. silvatica*, *R. succedanea*, *R. sylvestris*, and *R. trichocarpa* in Japan.

RHYNCHOSIA. **DOLICHOLUS** Ag. ROSARYBEAN. Twining, prostrate, rarely erect leguminous herbs and shrubs.

Aecidium erosematis P. Henn. Leaf rust on *Rhynchosia* and *Eriosema* sp. in Brazil.

Uredo pamparum Speg. Powdery dark-ochraceous rust pustules on leaves of *Rhynchosia* sp. in Argentina.

Woroninella dolichi (Cke.) Syd. See Dolichos.

RIBES. CURRANT. (Including *Grossularia*, GOOSEBERRY.) Shrubs cultivated for fruit and ornament. **Alternaria ribis** Bub. and Ranj. Effuse gray leaf spots with brown margins on *R. rubrum* in Yugoslavia.

Aphelenchus ribes (Taylor.) Goodey. This nematode attacks the buds of *R. nigrum* in Great Britain. The buds are destroyed and adjacent unfolding leaves may also be attacked.

Ascochyta monachorum Bub. On *Grossularia* in Bulgaria.

Ascochyta ribesia Sacc. and Fautr. Numerous circular to angular white leaf spots with dull-purple margins on *R. nigrum* and *Grossularia* sp. in Great Britain, France, and Austria.

Ascochyta ribis Massal. On *R. aureum* and *R. rubrum* in Italy and Russia.

Bacterium sp. A blossom-blight of *Grossularia* sp. due to a bacterium is reported from Great Britain.

Cercospora magellanica Speg. Small reddish leaf spots on *R. magellanicum* in Chile.

Colletotrichum grossulariae Jacz. Brown spots on fruit of *Grossularia* sp. in Russia and Italy.

Coniothyrium vagabundum Sacc. Said to cause premature fall of leaves of *Grossularia* sp. in Europe.

Cronartium ribicola Fisch. See Pinus.

Cytosporina ribis Magn. Causes a dying back and stem cankering of *R. grossularia*, *R. nigrum*, and *R. rubrum* in England and Holland.

Dothiorella vinosa Marchal. See Prunus.

Dothidella ribesia (Pers.) Theiss. and Syd. On twigs of *R. nigrum*, *R. rubrum*, *R. vulgare*, and *Ribes* sp. in Great Britain, middle Europe, and Alaska.

Helicobasidium tanakae Miy. See Morus.

Marsonia grossulariae Oud. Black spots on branches of *Grossularia* sp. in Europe.

Melampsora ribesii-purpureae Kleb. See Salix.

Melampsora ribesii-viminalis Kleb. See Salix.

Phylosticta pallidocarpa Bub. and Sereb. On leaves of *R. aureum* in Russia.

Phylosticta ribesicida Speg. Definite circular white leaf spots on *R. rubrum* in Argentina.

Phylosticta ribicola Sacc. Large yellowish leaf spots on *R. alpinum*, *R. aureum*, *R. nigrum*, *R. rubrum* and *Grossularia* sp. in Europe.

Phylosticta ribiseda Bub. and Kab. Circular, zoned, red-brown, to gray leaf spots on *R. rubrum* in Bohemia.

Phylosticta ribis-rubri Vogl. Circular, then confluent, chestnut-brown leaf spots on *R. rubrum* in Italy.

Puccinia depressa Diet. and Neg. Brown rust pustules on leaf blades and petioles of *R. glandulosum* in Chile.

Puccinia jaffueliana Speg. Leaf rust on *R. punctatum* in Chile.

Puccinia ribesii-caricis Kleb. Brown leaf rust on *R. aureum* in Europe.

Puccinia ribis-japonici P. Henn. Brown rust pustules on leaves of *R. japonicum* in Japan.

Septoria grossularicola C. Mass. Irregular brown leaf spots on *Grossularia* sp. in Italy.

Stereum purpureum Pers. See Prunus.

Venturia grossulariae (Awd. and Fleisch.) Sacc. On leaves of *Grossularia* sp. in Germany.

RICINUS. CASTOR BEAN. Tall annuals, or small trees in the Tropics.

Colletotrichum ricini Bub. and Frag. Anthracnose on leaves of *R. communis* in Spain.

Macrosporium cavaræ Parisi. Circular dark-yellow to brown spots on leaves and cotyledons preventing development of plants of *R. communis*, *R. gibsoni*, and *R. viridis* in Italy.

Melampsorella ricini (Biv.-Bern.) Det. Powdery orange-yellow rust pustules on small circular yellow leaf spots on *R. communis* and *R. spectabilis* in India, Ceylon, Madeira, north, south, and east Africa, and southern Europe.

Phylosticta ricini Rostr. Subcircular brown leaf spots with purple borders on *R. communis* in Denmark.

Physalospora propinqua Sacc. On *R. communis* in the Philippines.

Phytophthora melongenæ K. Saw. See Solanum.

RICINUS—Continued.

Phytophthora parasitica Dastur. Circular dull-green infection areas over both surfaces of cotyledons, destroying them and spreading down the stems causing damping off. Irregular spots occur on older leaves. On *R. communis*, *Clarkia* sp., *Pagopyrum* sp., *Gilia* sp., *Lycopersicon esculentum*, *Salpiglossis* sp., *Sesamum indicum*, *Solanum melongena*, *S. tuberosum*, and *Vinca rosea* in India.

A leaf-stalk rot of *Cocos nucifera* is attributed to this fungus in Jamaica. Dark-brown, somewhat sunken spots, one to several inches in diameter, occur on the leaf stalks causing a yellowing and wilting of the leaves, progressing successively until all are attacked and drooping. An internal brown rot of the leaves follows the spotting. The bud and central tissues as a rule are not affected. The larger nuts drop as a result of loss of support from the leaves.

This species is now considered identical with *P. terrestris* Sherb. which occurs in the southern United States and the West Indies as the cause of a foot rot or crown rot of *Citrus*, *Lycopersicum*, and other economic plants.

Sclerotinia ricini Godf. This serious blight of *R. communis*, characterized by a rot of the various parts of the plant, was introduced into the United States on seeds during the late war. It also occurs in Ceylon and India.

RIVINA. ROUGE PLANT. Erect tropical herbs.

Endophyllum rivinae (B. and C.) Arth. Yellow to brown rust pustules on branches, inflorescences and leaf blades and petioles of *R. humilis* and *Trichostigma octandrum* in Cuba and Argentina.

Phyllosticta rivinae Speg. Circular grayish-white leaf spots on *R. humilis* in Argentina.

Puccinia rivinae (B. and C.) Speg. Brown rust pustules on stems and leaf blades and petioles of *R. humilis* and *Trichostigma octandrum* in Porto Rico, Cuba, St. Thomas, and Argentina.

Septoria rivinae Pat. Brown, then white, irregular leaf spots on *R. humilis* and *Trichostigma octandrum* in Guadeloupe.

ROBINIA. LOCUST. Timber and ornamental trees cultivated for flowers and graceful foliage.

Ascochyta robiniae Sacc. and Speg. Grayish-white leaf spots on *R. pseudoacacia* in Italy.

Mycosphaera robiniae Siem. On leaves of *R. pseudoacacia* in Russia.

Oidium orbiculare Nannizzi. Powdery mildew on leaves of *R. hispida* in Italy.

Phleospora robiniae (Lib.) v. Hoeh. (*Septoria robiniae* Desm.) Leaf spot on *R. pseudoacacia* in Europe.

Phyllosticta advena Pass. Oval to vague ashen-white leaf spots on *R. corymbosa* and *R. pseudoacacia* in France, Italy, and Bohemia.

Phyllosticta robiniae Sacc. Irregular rufous leaf spots on *R. pseudoacacia* in Italy.

Phytophthora fagi R. Hart. See *Fagus*.

Septoria curvata (Rbh. and Braun.) Sacc. Circular to oblong brown leaf spots on *R. pseudoacacia* in Russia, Italy, and Germany. Reported from Ohio.

ROLLINIA. Tropical American fruit trees and shrubs.

Ovulariopsis monospora (Pass.) Sacc. and D. Sacc. Ashen leaf spots on *R. hispida* in Italy.

ROSE. **Uredo cherimoliae** Lagerh. See *Annona*.**ROMULEA**. Crocuslike bulbs.

Phyllosticta romuleae Gz. Frag. Leaf spots on *R. bifrons* in Spain.

RONDELETIA. Evergreen tropical trees and shrubs.

Phyllosticta rondeletiae Fl. Red, then ashen-white, leaf spots on *R. speciosa* in Italy.

ROSA. **Uredo rondeletiae** Arth. and Holw. Leaf rust on *R. cordata* in Guatemala.**RORIPA**. See *Radicula*.**ROSA**. ROSE. Ornamental shrubs.

Aecidium rosae-abysynicae P. Henn. Yellow rust pustules on leaves of *R. abyssinica* in Abyssinia.

Apiospora rhodophila Sacc. On branches and spines of *R. alpina* in Italy.

Apiospora rosae Oud. On branches of *R. canina* in Holland.

Ascochyta rosicola Sacc. This fungus produces more or less circular brown spots with distinct red margins on the upper surfaces of leaves of *R. muscosa* in Italy.

Botryosphaeria dothidea (Fr.) Ces. and de Not. (*B. diploia* Moug.) This disease, sometimes known as brier scab, has been epidemic at times in England on certain varieties of roses. Large, slightly raised black scabs are formed on the stems, which crack more or less concentrically. The hosts are *R. arvensis*, *R. canina*, and *R. centifolia* in Great Britain, continental Europe, and South America.

Catacauma dothidea (Moug.) v. Hoeh. Black stromata on twigs of *Rosa* sp. in Albania and Yugoslavia.

Cercospora hypophylla Cav. Causes a leaf spot of *R. gallica* in Italy.

Cercospora rosae (Fekl.) v. Hoehn. Reported as causing a leaf spot of *R. agrestis*, *R. alpina*, *R. arvensis*, *R. indica*, and *R. pendulina* in Madeira, Italy, Switzerland, and Austria.

Cercospora rosae-alpinae C. Mass. Causes a spotting of the under sides of leaves of *R. alpina* in Italy.

Coniothyrium wernsdorffiae G. Koch. This species, which is said by several writers to differ from *C. fuckelii*, causes a serious canker disease of canes of *Rosa* spp. in Denmark, Austria, and Germany. Diseased areas are more or less oval to circular, frequently girdling the stem, and brown in color with golden-brown margins.

Cryptosporium minimum Lan. Forms yellow-gray to brown spots, 1 to 2 centimeters in diameter on stems of *R. multiflora* in Germany.

Cryptostictis caudata (Preuss.) Sacc. Said to form cankers on branches of *Rosa* spp. in Europe.

Cryptostictis cynosbati (Fekl.) Sacc. Found on fruit of *R. pimpinellifolia* in Germany.

Diaporthe incarcerationa (B. and Br.) Nke. (*Phoma incarcerationa* [Nke.] Sacc.) Forms dead areas on twigs of *R. banksiae*, *R. canina* and *R. centifolia* in Europe.

Dicoccum rosae Bon. Leaf spot of *Rosa* spp. in Europe.

Didymella seipicolaeformis (DeNot.) Sacc. Minute fruiting bodies on dead areas on branches of *R. alpina* in Italy.

Diplodia rosarum Fr. (*D. centrophila* Pass.) Attacks the branches and spines of *R. banksiae*, *R. canina*, and *R. centifolia* in France and Austria.

Exosporium rosae Fekl. Forms spots on the under sides of living leaves of *Rosa* sp. in Switzerland and Germany.

Guignardia rosae (Auersw.) Petr. (*Laestadia rosae* Auersw.) Brown spots on the upper surfaces of leaves of *R. canina*, *R. centifolia*, and *R. rubiginosa* in Germany.

Gymnoconia rosae (Barcl.) Liro. (*Puccinia rosae* Barcl.) Powdery red-brown to dark-brown rust pustules on both sides of the leaves and on young twigs. Since the mycelium is perennial in the twigs, inspection of dormant plants would fail to reveal the presence of the disease. The hosts are *R. acicularis*, *R. cinnamomea*, *R. macrophylla*, *R. pimpinellifolia*, *R. platyacantha*, and *R. rugosa* in northern Europe, Turkey, Kamechatka, Caucasus, India, and Japan.

Hendersonia canina Brun. (*H. lichenicola* [Cda.] Fr.) On branches of *R. canina* in France.

Hendersonia rosicola Petch. On twigs of *Rosa* sp. in Ceylon.

ROSA—Continued.

- Hendersonia vulgaris** Desm. var. *rosae* Vesteg. Produces irregular spots on leaves of *R. alpina* and *R. pimpinellifolia* in Sweden.
- Kuehneola japonica** Diet. (*Phragmidium japonicum* Diet.) Brown rust pustules on both leaf surfaces of *R. lucia*, *R. multiflora*, and *R. wichuriana* in Japan.
- Kuehneola rosae** Sawada. A disease characterized by reddish-orange rust pustules on both leaf surfaces and on stipules, young twigs, and fruit of *R. indica* var. *formosana* in Formosa.
- Leptosphaeria coniothyrium** (Fckl.) Sacc. On *Rosa* sp. in Tasmania.
- Metasphaeria canina** Pass. (*M. constricta* Bres.) Attacks branches and spines of *R. canina* in Italy.
- Metasphaeria persistens** (B. and Br.) Sacc. On stems of *Rosa* sp. in England.
- Mycosporium rosae** Fckl. Produces blackened areas on branches of *R. canina* in Austria and Germany.
- Phoma aculeorum** Sacc. Reported on twigs of *R. canina* in Denmark, France, Italy, Switzerland, Austria, and Germany.
- Phoma canina** P. Brun. (*P. sepincola* [Kichx.] Sacc.) On *R. canina* in France, Switzerland, Austria, and Germany.
- Phoma rubiginosa** Brun. On fruit of *R. canina* and *R. rubiginosa* in Denmark and France.
- Phomopsis incarceration** v. Hoeh. (*Phoma incarceration* Sacc.) On branches of *R. canina* and *R. banksiae* in Great Britain. See also *Diaporthe incarceration*.
- Phomopsis rosae** Grove. (*Phoma rosae* Schulz. and Sacc.) On *R. canina* in Great Britain, Albania and Yugoslavia.
- Phragmidium butleri** Syd. Yellow to deep-brown or black rust pustules on the lower leaf surfaces of *R. macrophylla* in the Himalaya Mountains in India.
- Phragmidium devastatrix** Sor. Brown to black powdery rust pustules on both leaf surfaces and on petioles and young twigs of *R. lutea* in central Asia.
- Phragmidium egenulum** Syd. and Butl. Yellow or dark brown rust pustules on the leaves of *R. webbiana* in Kashmir (India.)
- Phragmidium fusiforme** Schroet. (*P. rosae alpinae* Wint.) Powdery rust pustules on the under sides of leaves and on petioles and fruits. *P. rosae-acicularis* on *R. acicularis* in Japan is said to be identical. The hosts are *R. acicularis*, *R. alpina*, *R. canina*, *R. nipponensis*, *R. pseudulina*, and *R. sinatica* in Europe.
- Phragmidium rosae-lacerantis** Diet. Yellow rust pustules changing to brown on the lower sides of leaves of *R. lacerans* in Persia.
- Phragmidium rosae-moschatae** Diet. Yellow to brown rust pustules on the under surfaces of leaves of *R. moschata* and *R. webbiana* in India.
- Phragmidium rosae-multiflorae** Diet. Small yellow swellings occur on the petioles and larger leaf veins with yellow rust pustules on the lower leaf surfaces of *R. laevigata* and *R. multiflora* in Japan.
- Phragmidium rosae-rugosae** Kasai. This rust forms gall-like pustules on petioles, fruit and twigs and often deforms the leaf blades of *R. rugosa* in Japan. The pustules form on the lower leaf surfaces and are yellow at first, becoming brown to black.
- Phragmidium rosae-sempervirentis** Maire. Orange-yellow rust pustules on the leaves and black powdery spore masses in cankered areas on the stems of *R. sempervirens* in Algeria.
- Phragmidium rufanjense** Bub. and Ranaj. Bright-yellow to deep brown rust pustules on the lower leaf surfaces of *R. spreta* in Albania and Yugoslavia.
- Phragmidium tuberculatum** J. Mueller. Brown to black rust pustules on *R. arvensis*, *R. canina*, *R. cinnamomea*, *R. inodora*, *R. lucida*, *R. micrantha*, *R. mollissima*, *R. rubiginosa*, *R. rubrifolia*, *R. scandens*, *R. sempervirens*, *R. sepium*, *R. spinosissima*, *R. tomentosa*, and *R. trachyphylla* in Europe, Siberia, and north Africa.
- Phragmidium yezoense** Kasai. Brown to black rust pustules on the lower surfaces of leaves and on petioles of *R. rugosa* in Japan.
- Phyllosticta argillaceae** Bres. Causes a leaf spot of cultivated roses in Europe.
- Phyllosticta rosarium** Pass. A dark-red to black leaf spot of *R. canina* and other cultivated roses in Russia, Italy, Switzerland, Austria, New South Wales, and Germany.
- Physoctenaria rosicola** (Fckl.) Sacc. On branches of *R. canina* in Germany.
- Protoventuria rosae** (DeNot.) Berl. On branches and twigs of *R. alpina* in Italy.
- Ramularia banksiana** (Pass.) Sacc. On leaves of *R. banksiae* in Europe.
- Septoria rosae** Desm. This widespread European disease, known as leaf scorch, is characterized by pale-brown or dirty-white spots, irregular in shape, bounded by a broad purple margin. The centers of the diseased areas tend to fall out. The disease is especially serious on nursery stock, causing practical defoliation, and when this occurs over several seasons, the cumulative effect is serious. The hosts are *R. canina*, *R. pomifera*, *R. pumila*, *R. sancta*, *R. scandens*, and other species of cultivated roses in Europe, New Zealand, Abyssinia, and Porto Rico.
- Septoria rosae-arvensis** Sacc. Causes a leaf spot of *R. arvensis* and *R. sempervirens* in Spain and other European countries. Probably synonymous with the preceding.
- Septoria rosarium** West. Produces numerous white rounded spots with broad, purple borders on the upper surfaces of leaves of *R. cinnamomea*, *R. collina*, and *R. corifolia*. Reported from Russia, Belgium, Italy, Great Britain, India, and Brazil. Probably not specifically distinct from *S. rosae* Desm.
- Stereum purpureum** Pers. See *Prunus*.
- Silbum** sp. This fungus produces "small purplish spots on the rose stems which gradually increase in size, sometimes even girdling the branch. The bark becomes shrivelled and sunken, and in the later stages is covered with the bright red stalks of the fungus surmounted by red globules." Cultivated roses in Trinidad are subject to this disease.
- Ucinula simulans** Salm. A powdery mildew on the leaves and young twigs of *R. multiflora* in Japan.
- ROSMARINUS.** ROSEMARY. Hardy evergreen shrubs with aromatic leaves.
- Ascochyta rosmarinii** Tassi. Circular to angular white leaf spots with red-brown borders on *R. officinalis* in Italy.
- ROTTBOELLIA.** Annual or perennial robust tropical grasses.
- Cintractia densa** McAlp. Smut sori in compact, dark-brown masses along rachides, destroying the florets of *R. compressa* in Australia.
- Meliola parenchymatica** Gaill. Black, superficial fungus patches on leaves of *R. exaltata* in the Philippines.
- Phyllachora rottioidiae** Syd. and Butl. Dull-black stromata on leaves of *R. exaltata* in India and the Philippines.
- Puccinia cacao** McAlp. Chocolate-brown to dark-brown rust pustules on leaves of *R. compressa* and *R. japonica* in India, Japan, and Australia.

ROTTBOELLIA—Continued.

- Puccinia microspora** Diet. Brown rust pustules on purple to brown leaf spots on *R. compressa* in Japan.
- Puccinia rottboelliae** Syd. Linear powdery black rust sori on leaves of *R. arundinacea* in Abyssinia and the Congo.
- Schizothyrum congoensis** Beeli. On *Rottboellia* sp. in the Congo.
- Uromyces rottboelliae** Arth. Leaf rust on *R. compressa* and *R. speciosa* in India and Japan.
- Ustilago flagellata** Syd. and Butl. Elongate powdery dark-brown smut sori in spikes and rachides of *R. exaltata* and *Rottboellia* sp. in the Philippines and the Congo.
- Ustilago rottboelliae** Syd. and Butl. Black powdery smut sori, deforming and destroying rachides and flowers of *R. compressa* in China, Japan, and India.

ROUPALA. Tropical American trees.

- Catacauma rhopalinum** (Mont.) Theiss. and Syd. Black stromata on leaves of *Roupala* sp. in Cayenne, Brazil and Peru.
- Melasma roupalae** Allesch. On leaves of *Roupala* sp. in Brazil.

ROYENA. Evergreen trees and shrubs.

- Aecidium royenae** C. and M. Leaf rust on *R. pallens* in the Union of South Africa.

ROYSTONEA. See *Palmae*.**RUBIA**. Hardy herbs.

- Aecidium archaevaletae** Speg. Yellow stem rust on *Rubia* sp. in Uruguay.
- Aecidium rubiae** Diet. Leaf rust on *R. cordifolia* in Japan.
- Ascochyta rubiae** Rabh. Gray circular leaf spots with purple-brown areoles on *R. peregrina* in Yugoslavia.
- Mycosphaerella peregrina** (Cke.) Lind. On stems and leaves of *R. peregrina* in England.
- Peronospora rubiae** Gäum. Downy mildew on leaves of *R. tinctorum* in Holland, France and Dalmatia.
- Pseudopeziza autumnalis** (Fr.) Karst. See *Galium*.
- Pseudopeziza komarovii** Jacz. Brown fruiting disks on yellow-brown sunken leaf spots on *R. cordifolia* in Manchuria.
- Puccinia algerica** Pat. Brown to black rust pustules on leaves of *R. laevis* in Algeria.
- Puccinia collettiana** Barcl. Leaf rust on *R. cordifolia* in India.
- Puccinia dimorpha** Syd. Chestnut-brown rust pustules on leaves and stems of *R. petiolaris* in the Union of South Africa.
- Puccinia rubicola** Syd. Brown rust pustules on leaves of *R. discolor* in Abyssinia.
- Puccinia rubivora** P. Magn. Leaf rust on *R. fruticosus* in the Canary Islands.
- Thecopora rubiae** Kom. Yellow and dark-brown rust pustules on leaves of *R. cordifolia* in Japan, China, Manchuria, Ceylon, Siberia, and Russia.

RUBUS. BLACKBERRY. RASPBERRY. DEWBERRY. Herbs and shrubs cultivated for their fruit.

- Ascochyta fenilleaubsiana** Sacc. and Roum. White leaf spots with narrow dark borders on *Rubus* sp. in France.
- Ascochyta rubi** Sacc. Small subcircular deep red-brown leaf spots on *R. caesius*, *R. fruticosus* and *R. saxatilis* in Siberia, Russia, Italy, and Germany.
- Cercospora garbiniana** C. Mass. Grayish-white leaf spots on *R. fruticosus* in Italy.
- Cercospora rubicola** Thuem. Subcircular dull yellow-brown leaf spots with purplish margins on *R. fruticosus* in Russia and Portugal.
- Cladosporium** sp. Said to cause a stunting of plants of *Rubus* sp. (raspberry) in Switzerland, due to cankers formed on the stems. Fruit fails to form and the leaves are discolored.
- Coccomyces rubi** (Fr.) Karst. On leaves of *R. caesius*, *R. fruticosus*, and *R. idaeus* in Europe.
- Coleroa chaetomium** (Kze.) Rabh. On leaves of *Rubus* spp. in Europe.
- Coniothyrium tumaeafaciens** Gueiss. Causes galls as large as walnuts on canes of *R. fruticosus* in Great Britain.
- Coryneum foliocolum** Fckl. See *Crataegus*.
- Didymella applanata** (Niessl.) Sacc. Irregular gray cankered areas on canes of *Rubus* spp. (raspberry and loganberry), causing wilting as well as witches'-brooms in Great Britain and continental Europe.
- Gerwasia rubi** Rac. Golden yellow rust pustules on leaves of *Rubus* sp. in Java.
- Gnomoniella cercosporae** Pass. On leaves of *R. glandulosus* in Italy.
- Hamaspora acutissima** Syd. Leaf rust on *R. moluccanus*, *R. nantoensis*, *R. rolfeii*, and *Rubus* sp. in Java, Formosa, Australia, and the Philippines.
- Hamaspora engleriana** (Diet.) Syd. Yellow rust pustules on leaves of *R. volkensianus* in East Africa.
- Hamaspora gedecana** Rac. Yellow leaf rust on *R. alpestris* in Java.
- Hamaspora longissima** (Theum.) Koern. Yellow leaf rust on *R. moluccanus*, *R. pinnatus* and *R. rigidus* in India, Australia, Uganda, and the Union of South Africa.
- Hapalosphæria deformans** Syd. Attacks flowers of *R. dumetorum* in Germany.
- Helminthosporium cymbispermum** Pat. On leaves of *Rubus* sp. in Ecuador.
- Hendersonia rubi** West. White patches on canes, causing sterility and finally wilting of *R. fruticosus*, *R. idaeus*, and *R. vitifolius* in Great Britain, France, Italy, and Belgium.
- Hyalothales dimerosperma** Speg. On leaves of *R. urticaefolius* in Brazil.
- Hypomyces rubi** (Osterw.) Wr. Attacks roots of *R. idaeus* in Switzerland.
- Kuehneola andicola** Diet. Leaf rust on *R. geoides* in Chile.
- Kuehneola uleana** Syd. Rust on leaves of *Rubus* sp. in Brazil.
- Mycosphaerella chamacemori** (Karst.) Lind. On leaves of *R. chamacemorus* in Finland.
- Mycosphaerella minoensis** Syd. On leaves of *Rubus* sp. in Japan.
- Mycosphaerella winteri** (Pass.) Sacc. On leaves of *R. corylifolius* in Italy.
- Ovularia rubi** Bub. Circular to irregular, often confluent, brown leaf spots on *Rubus* sp. in Hungary.
- Phragmidium arcticum** Lagh. Yellow and black rust pustules on leaves of *R. arcticus* in Sweden and Finland.
- Phragmidium assamense** Syd. Leaf rust on *R. lasiocarpus* in India.
- Phragmidium barclayi** Diet. Yellow and brown rust pustules on red-brown leaf spots on *R. koehleri* and *R. lasiocarpus* in India.
- Phragmidium barnardi** Plowr. and Wint. Rust on leaf blades and petioles and on peduncles of *R. parvifolius* in Australia.
- Phragmidium burmanicum** Syd. Leaf rust on *R. lasiocarpus* in Burma.
- Phragmidium griseum** Diet. Yellow and black rust sori on irregular leaf spots on *R. incisus* in Japan.
- Phragmidium heterosporium** Diet. Leaf rust on *R. trifidus* in Japan.
- Phragmidium incompletum** Barcl. Yellow leaf rust on *R. paniculatus* in India.
- Phragmidium nambrianum** Diet. Powdery black rust pustules on sunken violet-brown leaf spots on *R. kinashii* and *R. occidentalis* in Japan.

RUBUS—Continued.

- Phragmidium oecoloculare** Barcl. Black leaf rust on *R. rosaeifolius* in Japan.
- Phragmidium orientale** Syd. Golden rust pustules on leaves of *R. ellipticus* and *R. moluccanus* in India and Ceylon.
- Phragmidium pauciloculare** Syd. Powdery rust pustules on leaf blades and petioles and on peduncles of *R. parvifolius*, *R. phoenicolasius*, and *R. triphyllus* in Japan.
- Phragmidium quinqueloculare** Barcl. Rust on leaves and stems of *R. biflorus* in India.
- Phragmidium rubi** (Pers.) Wint. Yellow and powdery black rust pustules on lower leaf surfaces of *Rubus* spp. (nearly 40 species are recorded as hosts) in Japan and Europe.
- Phragmidium rubi-fraxinifolii** Syd. Leaf rust on *R. fraxinifolius* and *R. taiwanianus* in Formosa.
- Phragmidium rubi-idaei** (DC.) Karst. Yellow to black rust pustules on leaves of *R. idaeus* in Japan, Alaska, Siberia, and Europe. Probably occurs in the United States.
- Phragmidium rubi-japonici** Kasal. Black rust pustules on leaves of *R. japonicus* in Japan.
- Phragmidium rubi-saxatilis** Liro. Yellow and powdery black rust pustules on leaves of *R. saxatilis* in northern Europe.
- Phragmidium rubi-sieboldii** Kaw. (*Hamasporea rubi-sieboldii* [Kaw.] Diet.) Leaf rust on *R. sieboldii* in Japan.
- Phragmidium rubi-thunbergii** Kus. Leaf rust on *R. idaeus* and *R. thunbergii* in Japan.
- Phragmidium violaceum** (Schultz.) Wint. Golden-yellow and black rust pustules on leaves of *Rubus* spp. (60 species are listed as hosts) in China, Indo-china, Egypt, and Europe.
- Phragmidium yoshinagai** Diet. Leaf rust on *R. crataegifolius*, *R. morifolius*, and *R. sorbifolius* in Japan.
- Phragmidium zeylanicum** Petch. Leaf rust on *R. lasiocarpus* in Ceylon.
- Phyllosticta argillacea** Bres. On leaves of *R. idaeus* in Bohemia and Silesia.
- Phyllosticta cryptocarpa** Kab. and Bub. Brown, often confluent, leaf spots with purple margins on *R. glandulosus* in Bohemia.
- Phyllosticta fusco-zonata** Thuem. Subcircular dull-brown leaf spots on *R. idaeus* in Algeria, Russia, and Italy.
- Phyllosticta pallor** (Berk.) Oud. On *Rubus* (raspberry) in Great Britain and Holland.
- Phyllosticta rubi** P. Henn. Circular leaf spots with red-brown margins on *Rubus* sp. in Brazil.
- Phyllosticta rubicola** Rabenh. Subcircular small white leaf spots with blood-red margins on *R. caesius* and *R. idaeus* in Russia, Dalmatia, Italy, and Germany.
- Phyllosticta rubi-odorati** Bub. and Kab. On leaves of *R. odoratus* in Bohemia.
- Rhabdospora ramealis** (Desm. and Rob.) Sacc. and var. **macrospora** App. and Laub. Causes small purple-red areas on stems of *Rubus* spp. in Europe.
- Sclerotinia rubi** Carm. White, then brown, sclerotia on leaves of *R. fruticosus* in Great Britain.
- Septoria campoi** Speg. On leaves of *R. sanctus* in Chile.
- Uredo chinensis** Diet. Leaf rust on *R. buergeri* and *R. reflexus* in Japan.
- Uredo imperialis** Speg. Golden-yellow leaf rust on *R. imperialis* in Argentina.
- Uredo sinensis** Diet. Brown rust pustules on leaves of *R. reflexus* in China.
- Uromyces arthuri** Syd. Leaf rust on *R. guyanensis* and *R. schiedeanus* in Guatemala.
- Uromyces cundimarcensis** Mayor. Leaf rust on *R. peruvianus* in Colombia.
- Uromyces lagerheimii** P. Magn. Yellow leaf rust on *R. glaucus* and *Rubus* sp. in Ecuador and Colombia.
- Uromyces loesenerianus** (P. Henn.) Syd. Yellow leaf and stem rust on *R. bogotensis*, *R. brazilensis*, *R. trichomallus*, and *R. urticaefolius* in Central and South America.
- Uromyces pittierianus** P. Henn. Yellow leaf rust on brown sunken spots on *R. adenotrichos* and *Rubus* sp. in Costa Rica.
- Uromyces quitensis** Lagh. Leaf rust on *Rubus* sp. in Ecuador.
- Uromyces rubi** Diet. and Holw. Rust on sunken leaf spots on *R. adenotrichos*, *R. glaucus*, *R. laxus*, *R. poliphyllus*, *R. pringlei*, and *R. trichomallus* in Mexico, Guatemala, and Costa Rica.
- Uromyces rubi-urticifolii** Mayor. Rust on leaves of *R. urticaefolius* in Colombia.
- Uromyces variabilis** Mayor. Leaf rust on *Rubus* sp. in Colombia.
- Venturia kunzei** Sacc. On leaves of *R. caesius*, *R. idaeus*, and *R. pedatus* in Europe and Alaska.
- RUDGEA**. Shrubs or small trees.
- Phyllachora rudgeae** Syd. Black stromata on yellow-brown leaf spots on *R. coriacea* in Brazil.
- RUELLIA**. Acanthaceae herbs or shrubs.
- Accidium tuberosa** P. Henn. Leaf rust on *R. tuberosa* in Peru.
- Puccinia mussonii** McAlp. Yellow and dark-brown rust sori on leaves of *R. australe* in Australia.
- Puccinia paranahybae** P. Henn. Dark-brown leaf rust on *R. longifolia* in Brazil and Argentina.
- Puccinia ruelliae-bourgaei** Diet. and Holw. Powdery yellow and brown rust pustules on leaves of *R. bourgaei* in Mexico.
- Uromyces ruelliae** Holw. Brown to black rust pustules on leaves of *Ruellia* sp. in Mexico.
- RUMEX**. DOCK. SORREL. Perennial herbs, mostly weeds.
- Ascochyta rumicis** Bub. and Malk. On *Rumex* sp. in Bulgaria.
- Ascochyta vicinia** Sacc. On stems of *R. acetosella* and *Salvia officinalis* in Spain, France, and Italy.
- Cylindrosporium pulchrum** Speg. On leaves of *R. obtusifolius* and *R. pulcher* in Argentina.
- Depazea acetosa** Op. On leaves of *R. acetosa* in Siberia, Austria, and Germany.
- Mycosphaerella insulana** Bub. and Syd. On leaves of *Rumex* sp. in Germany.
- Mycosphaerella rumicis** Desm. Circular brown leaf spots on *R. conglomeratus*, *R. crispus*, *R. hydro-lapathum*, and *R. nemolapathum* in China and Europe.
- Ovularia canaegracola** P. Henn. Pale-yellow or brown leaf spots with white centers on *R. hymenosepalus* in Germany.
- Ovularia rubella** (Bon.) Sacc. On leaves of *R. aquaticus* in Germany.
- Ovularia rumicis** Eliasson. Circular to elliptical ochraceous leaf spots on *R. crispus* in Sweden.
- Pellicularia chilensis** Speg. Leaf blight of *R. crispus* in Chile.
- Peronospora rumicis** Cda. Downy mildew on leaves, stems, and panicles of *R. acetosa*, *R. acetosella*, *R. arifolius*, *R. curviculatus*, *R. crispus*, *R. lunaria*, *R. scutatus*, *R. thyrsoiflorus*, and *R. vesicarius* in India, Tunis, and Europe.
- Phyllosticta acetosae** Sacc. Subcircular brown leaf spots on *R. acetosa* in Italy and Russia.
- Phyllosticta straminea** Bres. On leaves of *R. acetosa* in Germany.
- Physoderma acetosellae** Rostr. Spore masses in deformed ovaries of *R. acetosella* in Denmark.
- Puccinia bififormis** Lagh. Dark-brown rust pustules on leaves and stems of *R. bucephalophorus* in Portugal.
- Puccinia dissiliens** Cke. Brown leaf rust on *Rumex* sp. in northern India.
- Puccinia ludwigii** Tepper. Rust on leaves of *R. brownii* in Australia.
- Puccinia nepalensis** Barcl. and Diet. Brown rust pustules on leaves of *R. nepalensis* in India.
- Puccinia pachyphloea** Syd. Black rust pustules on leaf blades and petioles and stems of *R. tuberosus* in Kurdistan.
- Puccinia rumesicicola** Gz. Frag. Leaf rust on *R. papillaris* in Spain.

RUMEX—Continued.

- Puccinia rumicis-scutati** (DC.) Wint. Brown leaf and stem rust on *R. scutatus* in Europe and possibly Australia.
- Puccinia trillii** Plowr. See Phragmites.
- Ramularia rumicis** K. and Cke. Large dull-brown leaf spots on *R. obtusifolius* in the Union of South Africa.
- Ramularia rumicis-scutati** Allesch. Circular leaf spots with white centers and broad ochraceous margins on *R. scutatus* in Spain, Switzerland, and Germany.
- Schroeteria alpinus** (Schroet.) P. Magn. Brown rust pustules on leaves of *R. alpinus* in central Europe.
- Septoria acetosae** Oud. Brown leaf spots with purple borders on *R. acetosa* and *Rumex* sp. in Denmark and Holland.
- Septoria glacialis** Ferr. Ochraceous circular zoned leaf spots on *R. scutatus* in Italy.
- Stigmataea rumicis** (Desm.) Schroet. On leaves of *R. alpinus* in Switzerland.
- Uredo purpurascens** Diet. Leaf rust on *R. romassa* in Chile.
- Uromyces acetosae** Schroet. Yellow and brown rust pustules on leaf blades and petioles of *R. acetosa*, *R. acetosella*, and *R. arifolius* in Europe.
- Uromyces argaeus** Maire. Powdery brown rust pustules on leaves of *R. tuberosus* in Asia Minor.
- Uromyces borealis** R. Leaf rust on *R. arifolius* in Finland.
- Uromyces crassipes** Diet. and Neg. Brown leaf rust on *R. cuneifolius* in South America.
- Uromyces tingitanus** P. Henn. Leaf rust on *R. tingitanus* in Algeria and Tunis.
- Urophlyctis rubsaameni** Magn. Tuberous excrescences on roots of *R. scutatus* in Switzerland and Germany.
- Ustilago domestica** Bref. Dull-violet masses of smut spores in stems, leaves, and inflorescences of *R. domesticus* in Norway.
- Ustilago kühneana** (Wolf) F. v. W. Smut sori in leaves and panicles of *R. acetosa*, *R. acetosella*, and *R. tuberosus* in north Africa and Europe.
- Ustilago lagerheimii** Bref. Powdery dark-violet masses of smut spores in stems and petioles of *Rumex* sp. in Ecuador.
- Ustilago warmingii** Rostr. Leaf smut on *R. crispus* and *R. domesticus* in Finland.
- Venturia rumicis** (Desm.) Wint. On leaves of *R. alpinus*, *R. crispus*, *R. nemorosus*, and *R. obtusifolius* in French North Africa, Denmark, and Bulgaria.

RUSCUS. BUTCHER'S-BROOM. Erect shrubs.

- Colletotrichum erumpens** Sacc. On *R. aculeatus* in France.
- Leptosphaeria rusci** (Walbr.) Sacc. On stems of *R. aculeatus* and *R. hypoglossum* in Switzerland, Spain and Italy.
- Phyllosticta danoë** Pass. Large irregular gray-white spots with brown margins on pseudo leaves of *R. racemosus* in France.
- Phyllosticta ruscicola** Dur. and Mont. Subcircular white leaf spots with brown margins on *R. aculeatus*, *R. androgyneus*, *R. hypophyllum*, and *R. hypoglossum* in Spain, Portugal, Italy, and France.
- Septoria hypoglossi** Massal. Leaf spots on *R. hypoglossum* in Italy.
- Urocystis jaaplana** Sacc. Smut on *R. aculeatus* in France, Spain, and Italy.

RUSSELIA. CORAL BLOW. Shrubs.

- Phyllosticta armitageana** Sacc. On leaves of *R. juncea* in Malta.

RTTA. RUE. Perennial glaucular herbs.

- Aecidium rutae** Har. Yellow rust on leaf blades and petioles of *R. chalepensis* in France and Sardinia.
- Oulariopsis haplophylla** (P. Magn.) Trav. On leaves of *R. graveolens* in Italy.

SABAL. See Palmetto.**SACCHARUM**. Sugar cane. Tall-growing, perennial, sugar-yielding grasses.

- Allantopora radiculicola** Wakk. On roots of *S. officinarum* in Java and Hawaii.
- Bacillus flavidus** Fawc. Reported as one of the causes of top rot or "polvillo" of sugar cane (*S. officinarum*) in Argentina.
- Bacillus sacchari** Speg. This bacterium is considered by Spegazzini as the cause of the "polvillo" disease of sugar cane (*S. officinarum*) in Argentina. Fawcett, however, in the most recent work on the disease, assigns *B. flavidus* and two other unnamed species as the causes. The disease is typical of diseases discussed under the head of top rot.
- Bacterium** sp. Causes a leaf-scald disease on *S. officinarum* in Australia and probably New Guinea.
- Bacterium vasculare** Cobb. The gumming disease due to this bacterium is a serious sugar-cane (*S. officinarum*) trouble in New South Wales, Queensland, Mauritius, New Guinea, Borneo, Fiji, Reunion, Java and possibly in Brazil. What is apparently the same disease has appeared in Porto Rico within the last few years. Externally diseased canes are stunted, the leaves show red longitudinal stripes, and there may be top rotting. On cutting across infected canes the bundles show discolored and a yellow gumlike substance oozes out. Many of the bundles are stained red, the discoloration extending out into the leaves. The gumming disease of Java is probably not due to *B. vasculare*, since the typical gummy exudate from the cut end of infected canes is entirely absent. Etiological studies have not yet been carried out.
- Bakerophoma sacchari** Died. Small brown leaf spots with red margins on leaves and leaf sheaths of *S. officinarum* in the Philippines.
- Bunchy top**. See Musa.
- Cephalosporium sacchari** Butl. This fungus causes a wilt of sugar cane in India, the Philippines, British West Indies, and the Union of South Africa. Affected stools become stunted, the leaves drying up and stems becoming light and hollow. Diseased canes internally are diffuse purple or dirty-red in color. These areas are distinct from the red-bordered white patches characteristic of red rot (*Colletotrichum falcatum*).
- Cercospora acerosum** Dick. and Hein. Black spots on leaves of *S. officinarum* in Java and the Philippines.
- Cercospora kopkei** Krueg. Dull-yellow, often confluent, leaf spots becoming red at the centers on *S. edule*, *S. glongong*, *S. officinarum*, and *S. spontaneum* in Java, Japan, Reunion, Formosa, Queensland, the Philippines, Argentina, and Cuba.
- Cercospora longipes** Butl. Small oval red leaf spots on *S. officinarum* in Porto Rico and India. The spots increase in size, becoming brown with yellow areolae and finally straw-colored, with oval, deep brown rings surrounding.
- Cintractia pulverulenta** Cke. and Mass. Smut in inflorescences of *S. arundinaceum* in India.
- Cytospora sacchari** Butl. The leaf sheaths take on a dark dull-red color and are bound firmly together by the mycelium of the fungus. The leaf blades turn brown and hang parallel to the stalk. Young stalks are killed outright and on older ones which have formed cane, brown lesions appear on the internodes, triangular at first when originating at the nodes, irregular if starting from wounds or cracks in the internodes. The fungus fruits on the outer leaf sheaths, the black, elongated necks of the pycnidia being rough to the touch. The disease occurs on *S. officinarum* in Porto Rico, Barbados, Formosa, India, and Brazil.

SACCHARUM—Continued.

Dinemasporium sacchari P. Henn. Circular to oblong brown leaf spots with red-brown margins on *S. officinarum* in Peru.

Diplodia cacaoicola P. Henn. Common as a wound parasite of cuttings and on dead canes and occasionally causing damage to susceptible varieties of *S. officinarum*. Affected canes wither and die, and are internally reddened, turning brown on exposure to air. See also *Theobroma*.

Eriosphaeria sacchari (v. Breda) Went. Dark-red circular leaf spots up to 1 centimeter in diameter on *S. officinarum* in Java and Trinidad.

Eurychora sacchari Avena. Stromata on leaves of *S. officinarum* in Brazil.

Fiji disease. Linear galls are produced on the leaves and leaf sheaths from 2 millimeters to 6 centimeters long, light-green in color at first and finally brown, bursting open, exposing a brownish mass. Diseased leaves are darker green than normal, as well as smaller and more slender. Infected plants, due to the shortening of the internodes are small, stunted, and have a bunched growth of leaves at the top. The roots are small, bunched, and slightly rotted. New shoots arising from the bases of heavily infected plants have a yellow streak running down the unfolded leaves which become distorted and wrinkled. A brown rot sometimes occurs in advanced cases. The cause of the disease is uncertain, but has recently been attributed to a protozoan (*Phytamoeba sacchari* McWhorter). The disease occurs in Australia, New Guinea, Fiji, and the Philippines on *S. officinarum*.

Gnomonia liiau Lyon. Liiau disease of sugar cane (*S. officinarum*) is characterized by a binding of the leaf sheaths firmly together by the mycelium of the causative fungus. The leaf blades die, turn brown, and hang down. The leaf sheaths are pinkish-brown in color and the stalks, if any cane is produced at all, are deep bluish-gray. The disease commonly attacks young shoots and prevents their further development. The imperfect or Melanconium stage of the fungus is produced internally in leaf sheaths and stalks, but the perithecia occur on the outer leaf sheaths, the beaks projecting. The disease is prevalent in Hawaii and is known also from limited areas in Louisiana.

Helminthosporium sacchari Butler. The eye-spot disease occurs as linear brown leaf spots with definite margins on *S. officinarum* in Porto Rico, Jamaica, Cuba, Barbados, St. Domingo, Hawaii, the Philippines, Java, Reunion, India, and the Union of South Africa. On some varieties the spots are oval, small, zoned, and yellow-brown in color. Susceptible varieties are checked in their growth and often stunted. The disease is known from Florida and Georgia.

Hendersonia sacchari Butl. This fungus causes withering of *S. officinarum* in India and Ceylon. The leaves die from the center outward and downward. Individual leaves die back from the tip and along the margins. Internally the diseased canes are dry and pithy, with a cavity in each internode. The roots are rotted.

Leptosphaeria sacchari v. B. H. Numerous oval dull-gray spots with definite red or red-brown margins occur on the leaves of *S. officinarum*. There may be an indistinct yellow halo surrounding each spot. In many varieties of cane the fungus causes premature withering and death of the leaf blades. The fungus is known from practically all the sugar-cane growing countries of the world. It is reported from Alabama and Florida.

Leptosphaeria saccharicola P. Henn. Pale-brown leaf spots with darker margins on *S. officinarum* in Brazil. Probably not distinct from the preceding species.

Leptosphaeria spegazzinii Sacc. and Syd. and var. *minor* Speg. On leaf blades and sheaths of *S. officinarum* in Argentina.

Leptosphaeria tucumanensis Speg. On leaves of *S. officinarum* in Argentina.

Linospora sacchari Avena. On leaves of *S. officinarum* in Brazil.

Lophodermium sacchari Lyon. Black linear fruiting bodies on midribs and sheaths of "sereh" cane in Hawaii, and "may be mildly parasitic."

Marasmius sacchari Wakker. Root disease is one of the most serious, if not the most serious, disease of sugar cane (*S. officinarum*) and occurs in every sugar-cane growing country. There is a very extensive literature on the subject, the disease being attributed in large part to the above fungus. Most recent studies indicate, however, that other fungi, such as *Rhizotonia* spp., *Pythium*, etc., are the initial agents and *Marasmius*, *Odontia*, and other hymenomyces secondary forms, only mildly parasitic at best. *M. sacchari* is commonly found in connection with root disease, the mycelium binding the lower leaf sheaths together by typical mycelial fans. The small white sporophores are produced in abundance in and around diseased stools during wet weather. Other species occur less commonly in similar manner.

Melanconium sacchari Mass. The rind disease of sugar cane (*S. officinarum*) occurs in practically every cane-growing country. The fungus causes a souring of the juice and a soft white rot of the tissues, affected stalks withering and drying up. Fruiting pustules form in great numbers on rotted stalks and at the base of leaf sheaths as small black areas from which the spore mass exudes, often as long, threadlike processes. The fungus is for the most part a wound parasite and seldom serious, except on overmature cane or cane first weakened by other agencies. Occurs as a saprophyte in Louisiana and Florida.

Meliola arundinis Pat. Superficial black fungus patches on leaves of *S. officinarum* in China and the Philippines.

Meliola sacchari Syd. Similar to the preceding species, in the Philippines.

Mosaic (*Motting*). The mosaic or yellow-stripe disease of sugar cane (*S. officinarum*) occurs in Porto Rico, Cuba, Jamaica, Guadeloupe, Barbados, Trinidad, Santo Domingo, Argentina, Egypt, Hawaii, the Philippines, Java, China, and Formosa. The disease has been found also on other grasses, including *Chaetochloa magna*, *Brachiaria platyphylla*, *Holcus sorghum*, *Paspalum boscinum*, *Pennisetum glaucum*, *Syntherisma sanguinale*, and *Zea mays*. It has been especially serious on maize (*Zea*). By inoculation other grasses including *Miscanthus sinensis*, *Saccharum narenga*, *Chaetochloa lutescens*, *C. magna*, and *Echinochloa crusgalli* have been proved susceptible. This disease was introduced into Louisiana some years ago and has been disseminated from there into Florida, Georgia, Alabama, and Mississippi, and has already caused considerable losses.

The disease is characterized by a motting of the leaves, varying in shading and patterns, but always very characteristic. In early infections there are no other outstanding symptoms. As the disease progresses in susceptible varieties, and generally after one or two seasons, cankers appear on the stalks, at first as discolored or water-soaked patches or streaks on the internodes, later sunken and brown to grayish white in color. On very susceptible varieties the cankers are very numerous and cause the canes to dry out. In such cane there is also a marked shortening of the joints and premature root development beneath the leaf sheaths. Resistant varieties may show no more than the leaf symptoms over long periods, with relatively slight reductions in yield, while on the other hand fields of susceptible types may be rendered worthless within two seasons.

Mycosphaerella sacchari Speg. On leaves of *S. officinarum* in Cuba, Porto Rico, and Argentina.

Mycosphaerella striatiformans Cobb. This fungus is said to attack the leaf tissue between the veins, causing a characteristic yellow striping. In the final stage the leaves break up into as many strings as there are stripes, the resulting shreds hanging down, while the mid-rib remains erect. The disease occurs in Hawaii on *S. officinarum*.

SACCHARUM—Continued.

Odontia saccharicola Burt. The granular leaf-sheath fungus binds together the lower leaf sheaths and develops fruiting areas as uniform thin white patches with a granular, somewhat powdery surface encircling the stalk from the ground level to a height of about a foot. The stellate-crystal fungus (*Himantia stellifera* Johnston), which often occurs in connection with root disease, is undoubtedly the vegetative stage of this fungus. The exact relation of this fungus to root disease is entirely problematical and its status is approximately the same as *Marasmius*. *O. saccharicola* is reported from Porto Rico and Santo Domingo, although the stellate crystal form is known from Cuba, Barbados, St. Croix, British Guiana, Trinidad, Hawaii, Zululand, and the Union of South Africa on *S. officinarum*.

Pestalozzia fuscescens Sor. var. *sacchari* Wakk. Irregular gray areas with brown margins on leaves of *S. officinarum* in Java and the Philippines.

Phoma heterospora Speg. Pale-brown indefinite spots on leaves of *S. officinarum* in Argentina.

Phyllachora sacchari P. Henn. Linear dull-black stromata on leaves of *S. officinarum* and *S. spontaneum* in Java, India, and the Philippines.

Phyllachora sacchari-aegyptiaci Br. and Cav. Oblong to irregular, black stromata on leaves of *S. aegyptiacum* in Italy.

Phyllachora sacchari-spontanei Syd. Black stromata on yellowish to red-violet leaf spots on *S. spontaneum* in India and the Philippines.

Phyllosticta hawaiiensis Caum. On leaves of *S. officinarum* in Hawaii.

Phyllosticta sacchari Speg. Linear gray-white leaf spots with purple margins on *S. officinarum* in Porto Rico and Argentina.

Phyllosticta saccharicola P. Henn. Oblong gray-white leaf spots with dull-brown margins on *S. officinarum* in the Congo.

Plasmodiophora vascularium Matz. This organism is said to inhabit the vascular areas of cane stalks (*S. officinarum*) causing a stunting and top rot in Porto Rico and Barbados.

Polvillo (*Gangrena humida*). See *Bacillus sacchari*.

Rhizoctonia ferruginea Matz. Isolated from diseased roots of *S. officinarum* in Porto Rico.

Rhizoctonia pallida Matz. Found on roots of *S. officinarum* in Porto Rico, as well as on *Capsicum* sp. and *Zea mays*.

Sclerospora sp. Oogonial stages of *Sclerospora* occur in the Philippines on *S. officinarum*, *S. spontaneum*, and *Miscanthus japonicus*, the relationships of which to the conidial forms on *Zea* (q. v.) and other hosts are uncertain.

Sclerospora philippinensis Weston. See *Zea*.

Sclerospora sacchari Miy. This downy mildew of sugar cane (*S. officinarum*) is very serious in Formosa, Australia, and Fiji and has recently been found in the Philippines. Yellow stripes parallel to the veins appear on the leaves, increasing in number until the greater part of the leaf becomes pale and is marked by long, reddish-brown spots, on the under surfaces of which fluffy fungus outgrowths occur. Diseased leaves die and dry up, often becoming much torn or shredded before dying. Leaf sheaths are also affected. With some varieties the joints are abnormally long and greater in number than in healthy canes. If the cane is attacked while young, shoots arise from buds along the stalk, giving a broomlike appearance. It has been possible to infect *Euchlaena* and *Zea mays*.

Sclerospora spontanea Weston. See *Zea*.

Sclerotium sp. Yellowish to white irregular patches, broader than long, and with narrow purple or red-brown surrounding zones on leaves of *S. officinarum*. These patches are separated by narrow green or brown gaps, the result being a variegated appearance which is very striking. Irregular white to brown sclerotia form on the diseased areas. The disease is prevalent and serious in Java, India, and the Philippines.

Sclerotium griseum Stevenson. Rounded gray sclerotia on leaf sheaths of *S. officinarum* in Porto Rico. This form has recently been connected with root disease in common with one or more species of *Rhizoctonia*.

Sereh. Sereh has been a serious disease of sugar cane (*S. officinarum*) in Java, although it is now fairly well controlled by the use of resistant varieties and by growing the seed cane in high altitudes. The disease or one very similar is known from Malacca, Borneo, Siam, Hawaii, Malaya, and the Philippines. The cause remains unknown, although the disease is probably to be classed with the infectious mosaics.

Diseased stools fail to produce upright stems and the shortening of the internodes close to the ground gives the stunted plants a fan-shaped appearance. The leaves are smaller than normal and yellow-striped, dying in severe cases. The roots are weakly developed and soon die. The vascular bundles are reddened and show some gumming. The disease is readily carried in seed pieces without its presence being in evidence.

Sphaeronema adiosum Butl. A weak parasite of cane (*S. officinarum*) cuttings in India and Java. Diseased cuttings are dull-red within, finally drying up, with a black fungus growth on the exterior.

Sphaerulina sacchari P. Henn. Circular to oblong, then confluent, pale brown leaf spots with red-brown margins on *S. officinarum* in Peru.

Stigmata sacchari Speg. On leaves of *S. officinarum* in Argentina.

Thielaviopsis paradoxa (DeSeyn.) v. Hoeb. The pineapple or black rot disease of sugar cane is a disease of cuttings only, acting as a wound parasite when the plantings are made in wet soils or under other unfavorable conditions. Diseased cuttings fail to germinate or young shoots die back, the interior of the cutting turning black and giving off an odor of rotting pineapple. The fungus also causes the common soft rot or shipping rot of pineapples (*Ananas*) which has caused heavy losses to pineapple growers in the West Indies. Circular to irregular brown sunken leaf spots also occur on the pineapple plant due to this fungus. The stem-bleeding disease of coconut (*Cocos nucifera*) and betel palm (*Areca catechu*) is likewise attributed to it. In this instance it acts as a wound parasite, discoloring the tissues as it advances and causing a gummy exudate. The fungus is known to occur in the West Indies from Cuba to Trinidad, British Guiana, Hawaii, the Philippines, Mauritius, Ceylon, Java, Malaya, India, and Gold Coast. It has been isolated from coconut and sugar cane (*S. officinarum*) from Florida and probably occurs on other hosts in the southern United States.

Top rot. Top rot, characterized by the death of the terminal bud and inclosing leaves, is a common disease of sugar-cane (*S. officinarum*) in all cane-growing countries. A number of fungi and probably bacteria are undoubtedly involved, some at least of which do not occur in the United States. The "polvilla" of Argentina and dry top rot (*Plasmodiophora*) of Porto Rico and Cuba are two diseases of this class.

Tylenchus similis Cobb. See *Musa*.

Uromyces kühni Krueg. Elongated brown rust pustules on long orange spots on both leaf surfaces of *S. arundinaceum*, *S. fuscum*, *S. officinarum*, and *S. spontaneum* in Java, Japan, India, Burma, Ceylon, the Philippines, and Australia.

SACCHARUM—Continued.

- Ustilago noidea sacchari-narengae** K. Saw. Top-shaped dark-olive fruiting bodies in ovaries of *S. narenga* in Japan, the protecting membranes of which rupture at maturity, exposing the dark-olive spore masses.
- Ustilago sacchari** Rabh. From the growing point there is produced a long, whiplike, black shoot without leaves, often several feet in length and much curved or twisted. In the earlier stages these smutted shoots are protected by silvery-white membranes, which rupture, exposing dense black spore masses. Secondary lateral shoots may also be infected. On *S. cylindricum*, *S. officinarum*, *S. spontaneum*, and *Erianthus asper* in British Guiana, Trinidad, Mauritius, Reunion, Union of South Africa, the Philippines, Japan, Formosa, China, India, Java, Queensland, and Italy.
- Ustilago sacchari-ciliaris** Bref. Powdery black masses of smut spores in ovaries of *S. ciliare* in India.
- SAGERETIA**. Armed or unarmed shrubs.
- Aecidium sageretiae** P. Henn. Leaf rust on *Sageretia* sp. in China.
- SAGINA**. PEARLWORT. Annual or perennial herbs.
- Septoria nivalis** Rostr. On peduncles of *S. nivalis* in Greenland.
- Septoria velenovskyi** Bub. Small circular yellow leaf spots on *S. tenuifolia* in Yugoslavia.
- SAGITTARIA**. ARROWHEAD. Hardy perennial aquatic herbs.
- Cercospora macrospora** Bres. Subcircular dull-brown leaf spots on *S. sagittifolia* in Germany.
- Doassansia horiana** P. Henn. Black smut sori deforming leaf blades and petioles of *S. sagittifolia* in Japan.
- Doassansia limosellae** (Kze.) Schroet. Smut sori in circular brown leaf spots on *S. sagittifolia* and *Limosella aquatica* in Europe.
- Mycosphaella sagittariae** Tass. On leaf blades, petioles and sheaths and on stems of *S. montenidensis* in Italy.
- Phyllosticta sagittifoliae** Brun. Subcircular to oblong ochraceous leaf spots with dark brown margins on *S. sagittifolia* in Siberia and France.
- Ramularia sagittariae** Bres. Small dull-brown leaf spots on *S. sagittifolia* in Germany.
- SALIX**. WILLOW. Dioecious trees and shrubs.
- Ascochyta martianoffiana** Thuem. Subcircular ochraceous to gray leaf spots on *S. pyrolaeifolia* in Siberia.
- Ascochyta salicicola** Pass. Angular reddish to dull brown leaf spots on *S. alba* in France.
- Ascochyta translucens** Kab. and Bub. Circular to angular, then confluent, broad gray leaf spots on *S. caprea* in Bohemia.
- Ascochyta vitellinae** Pass. On leaves of *A. vitellina* in Europe.
- Bacillus hari Hori**, and Miy. Bacteriosis of *Salix* sp. in Japan.
- Bacterium salicis** Day. This bacterium is reported as the cause of the "watermark" disease of *S. alba* and *S. caerulea* in Great Britain. Leaves of infected trees wither and die, turning brown and remaining attached. New shoots and finally entire branches are infected, the disease ultimately reaching the trunk and killing the tree after one or more years. Infected trees develop adventitious shoots and a gummy exudate from dying branches. There is an internal gray staining of the wood.
- Botryosphaeria gregaria** Sacc. Causes the bark to turn brown, crack and die on twigs and branches of *Salix* sp. (basket willow) in Ireland.
- Dinemasprium purpurascens** Rich. On branches of *Salix* sp. in France.
- Diplotinea salicina** Cke, and Mass. Causes a die-back of *Salix* spp. in Great Britain.
- Discula carbonacea** (Fr.) B. and Br. Small black cankers causing die-back of *Salix* Sp. in Holland and Sweden.
- Entyloma salicis** Karst. Leaf smut on *S. caprea* in Finland.
- Fusiculadium salsiperdum** (All. and Tub.) Tub. This disease attacks *S. alba*, *S. aurita*, *S. caprea*, *S. cinerea*, *S. cuspidata*, *S. fragilis*, *S. mollissima*, *S. nigricans*, and *S. pentandra* in Scotland, Russia, Denmark, Holland, and Germany. *Venturia chlorospora* (Ces.) Karst. is considered the perfect stage. Dark olive, velvety patches occur on the leaves and spread down along the petioles to the twigs which as a result die back. Long black patches, sharply demarked from the healthy yellow bark, form on the branches.
- Gloeosporium deformans** (Schroet.) J. Lind. On catkins of *S. caprea* and *S. caprea-cinerea* in Sweden, Finland, Denmark, Silesia, and France.
- Graphium ulmi** Schwarz. See Ulmus.
- Haplotheclium amenti** (Rostr.) Theiss. and Syd. On catkins of *S. reticulata* in Norway.
- Helicobasidium mompa** Tan. See Morus.
- Helicobasidium tanakae** Miy. See Morus.
- Hendersonia foliorum** Fekl. Parasitic on leaves of *S. caprea* in Europe.
- Marsonia didyma** (Fekl.) Sacc. Small dull-brown leaf spots on *S. daphnoides*, *S. fragilis*, and *S. amygdalina* (*S. triandra*) in Russia, Poland, France, and Germany.
- Marsonia kriegeria** (Bres.) P. Magn. Circular black leaf spots on *S. amygdalina* and *S. triandra* in Germany.
- Marsonia obscura** Rom. Dull-red leaf spots on *Salix* sp. in Scandinavia.
- Marsonia pyrenaica** Gz. Frag. On leaves of *S. caprea* in Spain.
- Marsonia salicicola** (Bres.) Magn. Indefinite rufous leaf spots on *S. alba* and *S. caprea* in Bohemia, Denmark, and Germany.
- Marsonia salicis** Trail. On leaves of *Salix* sp. in Norway.
- Marsonia santonensis** (Pass.) Bub. On leaves of *S. pentandra* in Austria.
- Melampsora abietis-capraearum** Tub. Yellow and brown rust pustules on needles of *Abies alba* and *A. nordmanniana* and on the leaves of *S. caprea* in Denmark, France, Spain, and Germany.
- Melampsora allii-fragilis** Kleb. Yellow leaf rust on *Allium ascalonicum*, *A. cepa*, *A. oleraceum*, *A. porrum*, *A. sativum*, *A. schoenoprasum*, *A. ursinum*, *A. vineale*, *Salix fragilis*, *S. pentandra*, *S. triandra*, and *S. viminalis* in Europe.
- Melampsora amygdalinae** Kleb. Leaf rust on *S. amygdalina*, *S. hippophaefolia*, and *S. pentandra* in Europe.
- Melampsora coleosporioides** Diet. Yellow to brown rust pustules on leaves of *S. babylonica* and *S. glandulosa* in Japan.
- Melampsora epiphylla** Diet. Yellow to dark-brown rust pustules on leaves of *S. shikokiana* in Japan.
- Melampsora euonymi-capraearum** Kleb. Yellow to brown rust pustules on sunken, yellow circular to irregular leaf spots on *S. aurita*, *S. caprea*, *S. cinerea*, *S. incana*, *S. viminalis*, *Euonymus europaeus*, *E. latifolia*, and *E. verrucosa* in Europe and Japan.
- Melampsora galanthi-fragilis** Kleb. See Galanthus.
- Melampsora humboldtiana** Speg. Leaf rust on *S. humboldtiana* in Uruguay and Argentina.
- Melampsora humilis** Diet. Yellow to brown rust pustules on leaves of *S. multinervis* in Japan.
- Melampsora lapponum** Lindf. Leaf rust on *S. lapponum* and *Viola epipsila* in Sweden.

SALIX—Continued.

- Melampsora larici-caprearum** Kleb. Yellow leaf rust on *S. aurita*, *S. caprea*, *S. lapponum*, *S. pentandra*, *S. phylicifolia*, *S. pirolaeifolia*, *S. repens*, *S. sieboldiana*, *S. smithiana*, *Larix decidua*, *L. occidentalis*, and *L. sibirica* in Europe and Japan.
- Melampsora larici-epitea** Kleb. Leaf rust on *Salix* spp. (33 species listed) and *Larix decidua* in Siberia, India, Ceylon, and Europe.
- Melampsora larici-pentandrae** Kleb. Yellow to brown leaf rust on *S. amygdalina*, *S. fragilis*, *S. humboldtiana*, *S. jaspidea*, *S. pentandra*, *S. purpurea*, *Larix decidua*, and *L. sibirica* in Uruguay and Europe.
- Melampsora larici-urbaniana** Mat. Leaf rust on *S. urbaniana* and *Larix* sp. in Japan.
- Melampsora microscora** Diet. Yellow to brown rust pustules on leaves of *S. nipponica* in Japan.
- Melampsora repentis** Plowr. See *Orchidaceae*.
- Melampsora reticulatae** Blytt. Leaf rust on *S. reticulata* and *Saxifraga aizoides* in Sweden, Norway, Switzerland, Austria, and Hungary.
- Melampsora ribesii-purpureae** Kleb. Rust on leaves of *Salix* spp. and *Ribes* spp. in Europe, Japan, and Siberia. Reported from Utah and Colorado on *Ribes*.
- Melampsora ribesii-viminalis** Kleb. Golden-yellow to brown rust pustules on leaves of *S. viminalis*, *purpurea*, *Ribes alpinum*, *R. aureum*, *R. nigrum*, *R. rubrum*, *R. sanguineum*, and *Grossularia* in Europe.
- Melampsora salicis-albae** Kleb. Yellow and brown rust pustules on leaves of *Allium cepa*, *A. porrum*, *A. schoenoprasum*, *A. ursinum*, *A. vinealis*, and on leaves and twigs of *Salix alba* in India and Europe.
- Melampsora salicina** Lév. Black stromata on leaves of *S. purpurea* and *S. viminalis* in Italy.
- Mycosphaerella genutifusa** Auersw. On leaves of *S. alba* in Germany.
- Mycosphaerella salicicola** (Fr.) Fekl. On leaves of *S. caprea*, *S. nigricans*, and *S. triandra* in France, Italy, and Germany.
- Phoma intricans** Schwarz. Causes a die-back of twigs, with resulting death of leaves of *S. alba* var. *vitellina-pendula* in Holland.
- Phyllo Hendersonia foliorum** (Fekl.) Tass. On *S. caprea* in France, Italy, and Germany.
- Phyllosticta iserana** Kab. and Bub. Circular to angular brown-gray leaf spots on *S. fragilis* in Bohemia.
- Phyllosticta salicicola** Thuem. Subcircular gray-white leaf spots on *S. alba* and *S. triandra* in Denmark, Russia, Poland, and France.
- Phyllosticta translucens** Bub. and Kab. Angular gray-white leaf spots with yellow-brown margins on *S. nigricans* in Austria.
- Physalospora miyabeana** Fuku. Elliptical whitish-gray stem lesions, 5 to 30 millimeters long surrounded by black borders, on *S. purpurea* in Japan. These lesions may unite and girdle infected branches. Circular to irregular, dark-brown, often concentrically zoned areas occur on the leaves.
- Podosphaera schlechtendalii** Lév. Powdery mildew on leaves of *S. alba* and *S. viminalis* in France and Russia.
- Ramulaspera salicina** (Vestergr.) Lind. Violet leaf spots on *S. cinerea* and *S. hastata* in Sweden, Finland, and Austria.
- Septobasidium acaciae** Saw. See *Acacia*.
- Septoria blennorioides** (Karst.) Berl. and Vogl. On leaves of *S. pendula* in Finland.
- Septoria capreae** West. Angular gray-white leaf spots with rufous borders on *S. caprea*, *S. cinerea*, and *S. atrocinerea* in Belgium and Portugal.
- Septoria jeniiseica** Thuem. Circular, then confluent, dull-brown to gray leaf spots on *Salix* sp. in Siberia.
- Septoria salicicola** (Fr.) Sacc. Circular brown leaf spots on *S. caprea*, *S. cinerea*, *S. grandifolia*, *S. pyrolaeifolia*, and *S. viminalis* in Russia, Siberia, Denmark, Sweden, France, Italy, and Austria.
- SALOA**. See *Blumenbachia*.
- SALPICHRUA**. COCKSEGGS. Herbs and shrubs, sometimes under cultivation.
- Puccinia pampeana** Speg. Rust on leaves and stems of *S. rhomboidea* in Argentina, Brazil, and Uruguay.
- Septoria bonariensis** Speg. Circular to irregular gray-white leaf spots on *S. rhomboidea* in Argentina.
- SALPIGLOSSIS**. Flower-garden annuals.
- Phytophthora parasitica** Dast. See *Ricinus*.
- SALVIA**. SAGE. Herbs and subshrubs, with some economic and ornamental species.
- Aphelenchus oleisistis** Ritz. Bos. See *Begonia*.
- Ascochyta vicina** Sacc. See *Rumex*.
- Cercospora salviae** Pat. Brown leaf spots on *Salvia* sp. in Ecuador.
- Coleosporium salviae** Diet. Orange rust pustules on leaves of *S. bipinnata* and *S. japonica* in Japan.
- Puccinia badia** Holw. Leaf rust on *S. albicans* and *S. chrysanth* in Mexico.
- Puccinia bithynica** P. Magn. Leaf rust on *S. bithynica*, *S. grandiflora*, and *S. pomifera* in Russia and Crete.
- Puccinia conspersa** Diet. Brown rust pustules on small yellow sunken leaf spots on *Salvia* sp. in Brazil.
- Puccinia delavayana** Pat. and Har. Black rust sori on leaves of *Salvia* sp. in China.
- Puccinia delicatula** (Arth.) Sacc. and Trott. Leaf rust on *S. cinnabarina*, *S. elegans*, *S. holwayi*, and *S. pulchella* in Mexico and Guatemala.
- Puccinia diutinum** Mains. and Holw. Brown rust pustules on leaves of *S. chrysanth*, *S. pittieri*, and *S. scordoniaefolia* in Mexico and Costa Rica.
- Puccinia filioli** Mains. and Holw. Leaf rust on *S. involucreta* and *S. pulchella* in Guatemala.
- Puccinia gentilis** Arth. Cinnamon-brown to dark-brown rust pustules on leaves of *S. alamosana* in Mexico.
- Puccinia gilliesii** Speg. Brown rust pustules on leaf blades and petioles of *S. gilliesii* in Argentina.
- Puccinia griseola** Lagh. Leaf rust on *Salvia* sp. in Ecuador.
- Puccinia impedita** Mains. and Holw. Leaf rust on *S. hyptoides*, *S. occidentalis*, and *S. tiliaefolia* in Guatemala, Costa Rica, and Trinidad.
- Puccinia infrequens** Holw. Brown rust pustules on leaves of *S. cinnabarina* in Mexico and Guatemala.
- Puccinia mitrata** Syd. Dark-brown powdery rust pustules on leaves of *S. fluriatilis*, *S. mexicana*, *S. polystachya*, *S. purpurea*, *S. sessiliflora*, and *S. tiliaefolia* in Mexico and Guatemala.
- Puccinia nigrescens** Kirchn. Brown to black rust pustules on elongated sunken spots on leaves and petioles of *S. judaica*, *S. verticillata*, and *S. virgata* in Europe and Palestine.
- Puccinia nipponica** Diet. Leaf rust on *S. nipponica* in Japan.
- Puccinia nevadensis** Syd. Leaf rust on *S. lavandulaefolia* in Spain.
- Puccinia nivea** Holw. Brown rust pustules on leaves of *S. purpurea* in Mexico.
- Puccinia paramensis** Mayor. Leaf rust on *S. cernua* in Colombia.

SALVIA—Continued.

- Puccinia prospera** Arth. Brown rust pustules on leaves of *S. microphylla* in Mexico.
Puccinia roestliiformis Lagh. Yellow and brown rust pustules on leaves of *Salvia* sp. in Ecuador.
Puccinia salviae Ung. Brown leaf rust on *S. glutinosa* and *S. lavandulifolia* in China and Europe.
Puccinia soledadensis Mayor. Leaf rust on *S. pauciserrata* in Colombia.
Septoria salviae Pass. Dull-brown leaf spots on *S. hymninum*, *S. pratensis*, *S. sylvestris*, and *S. sclarea* in Russia, Italy, Dalmatia, and France.
Ustilago betonica Beck. Anther smut of *Betonica* spp. and *Salvia* sp. in Europe.
- SAMBUCUS**. ELDER. Shrubs grown for their foliage, flowers, and black and red berries.
Ascochyta fernandi Bub and Malk. Circular to irregular gray-brown leaf spots on *S. ebulus* in Bulgaria.
Ascochyta sambucella Pass. On branches of *S. ebulus*, *S. nigra*, and *S. racemosa* in Germany and Italy.
Ascochyta sambuci (Sacc.) Tass. On leaves of *S. nigra* in Italy and France.
Cercospora ticinensis Cav. Gray-brown leaf spots on *S. nigra* in Italy.
Exosporium depazeoides Desm. On leaves of *S. nigra* in France.
Gloeosporium ebuli Allesch. Subcircular to irregular, then confluent, brown leaf spots on *S. ebulus* in Germany.
Helicobasidium mompa Tan. See Morus.
Marsonia sambuci E. Rostr. Large brown leaf spots on *S. nigra* in Europe.
Mycosphaerella ebulina Petr. On leaves of *S. ebulus* in Bohemia and Austria.
Phyllosticta ebuli Fekl. On leaves of *S. ebulus* and *S. racemosa* in France, Italy, Hungary, and Germany.
Phyllosticta sambucina Allesch. On leaves of *S. nigra* in Switzerland.
Septoria ebuli Desm. and Rob. Small subcircular pale-rufous to purple, often confluent, leaf spots on *S. ebulus* and *S. nigra* in Europe.
- SANCHEZIA**. Strong, erect herbs or half shrubby plants grown for their flowers and foliage.
Pyrenochaete vexans Syd. Large confluent brown leaf spots on *S. nobilis* in Brazil.
- SANGUISORBA**. See also Poterium. BURNET. Perennial herbs grown in hardy borders and as salad plants.
Marsonia sennensis Gz. Frag. On leaves of *S. officinalis* in Spain.
Ovularia bulbifera (Fekl.) Sacc. Ochraceous leaf spots with dark-purple margins on *S. minor* in Alaska, Switzerland, and Germany.
Peronospora sanguisorbae Güm. Downy mildew on leaves of *S. officinalis* in central Europe.
Phragmidium carbonarium (Schlecht.) Wint. Golden and black rust pustules on leaves of *S. canadensis*, *S. carnea*, *S. officinalis*, and *S. tenuifolia* in Japan, Siberia, and Europe.
Phragmidium minor (Arth.) Syd. Dark-brown rust pustules on leaf blades and petioles of *S. latifolia* in Alaska.
Phragmidium sanguisorbae (DC.) Schroet. See Poterium.
- SANSEVIERIA**. BOWSTRING HEMP. Herbaceous perennials cultivated as ornamentals and sometimes as fiber plants.
Chaetophoma sansevieriae Tass. On leaves of *S. zeylanica* in Italy.
Leptosphaeria baldraffiana Bacc. Small brown spots on leaves of *S. ehrenbergia* in Abyssinia.
Phyllachora pappiana Bacc. Small subcircular black stromata on leaves of *S. ehrenbergia* in Abyssinia.
- SEPTOGLOUM concentricum** Syd. On leaves of *S. guineensis* in Abyssinia and central Africa.
- SANTALUM**. SANDALWOOD. Evergreen trees with aromatic properties.
Spik disease. This disease, characterized by die-back, unthriftiness, and final death of infected trees, is a serious disease of sandalwood (*S. album*) in India. The cause is unknown, but the disease is apparently one of the infectious mosaics.
- SANTOLINA**. LAVENDER COTTON. Shrubs and herbs, sometimes cultivated.
Puccinia chamaecyparissi Trott. Rust on leaves and stems of *S. chamaecyparissus* in Italy.
- SAPINDUS**. SOAPBERY. Trees and shrubs cultivated for ornament and economic value.
Mycosphaerella reyesii Syd. On leaves of *S. saponaria* in the Philippines.
Phyllosticta patouillardii Sacc. and D. Sacc. Circular to angular, bright-brown leaf spots with dull-brown margins on *Sapindus* sp. in Algeria.
Phyllosticta raimundi Sacc. Leaf spot on *Sapindus* sp. in the Philippines.
Phyllosticta sapindi P. Henn. Circular pale-brown leaf spots on *S. saponaria* in Brazil.
Uredo cristata Speg. Leaf rust on *S. bifoliatus* and *Sapindus* sp. in Ceylon and Argentina.
- SAPIUM**. CHINESE TALLOW TREE. Tropical trees and shrubs cultivated for their economic products.
Cercospora micromera Syd. On leaves of *S. sebiferum* in China.
Cercospora sapicola Speg. Angular brown leaf spots on *S. aucuparium* and *S. mannianum* in Argentina and Uganda.
Exosporium lateritium Syd. On leaves of *S. abyssinicum* in Tanganyika.
Helmintosporium sapii Miy. On leaves of *S. sebiferum* in China.
Uromyces cisnerosianus Speg. Dark-brown to black rust pustules on leaves of *S. aucuparium*, *S. biglandulosum*, and *S. salicifolium* in Argentina and Paraguay.
Uromyces globosus Diet. and Holw. Powdery black rust pustules on leaves of *S. biglandulosum* in Mexico.
Uromyces vestitus Diet. Brown rust pustules on leaves and stems of *Sapium* sp. in Brazil.
- SAPONARIA**. SOAPWORT. Hardy, often coarse, annual or perennial, erect or decumbent herbs, sometimes cultivated.
Ascochyta dianthi Berk. See Dianthus.
Ascochyta saponariae Fekl. Small dull-brown leaf spots on *S. officinalis* in Germany.
Cylindrosporium saponariae Roum. Large circular, often confluent, gray leaf spots on *S. officinalis* in France.
Phyllosticta saponariae (Fekl.) Sacc. On leaves of *S. officinalis* in Belgium, Italy, and Germany.
Septoria saponariae (DC.) Sav. and Becc. Pale-brown circular to irregular leaf spots on *S. officinalis* and *Silene latifolia* in Bulgaria, Yugoslavia, Great Britain, Denmark, and Belgium.
Uromyces gypsophila Cke. See Gypsophila.
- SARCANTHUS**. See Orchidaceae.
- SASA**. See Bambuseae.
- SATYRIUM**. See Orchidaceae.
- SAURURUS**. LIZARD'S-TAIL. Perennial marsh herb.
Uromyces saururi P. Henn. Leaf rust of *S. loureiri* in Japan.
- SAUSSUREA**. Annual, biennial, or perennial herbs, sometimes planted in gardens.
Aecidium saussureae-affinis Diet. Leaf rust on *S. affinis* in Japan.
Bremia saussureae Saw. Downy mildew on pale-yellow polygonal leaf spots on *S. carthamoides* in Formosa.

SAUSSUREA—Continued.

- Coleosporium saussureae** Thuern. Golden-yellow rust pustules on leaves of *S. grandifolia*, *S. japonica*, *S. latifolia*, *S. mazimowiczii*, *S. triangulata* and *S. ussuriensis* in Siberia, Japan, and China.
- Erysiphe taurica** Lévl. See *Althaea*.
- Phyllosticta saussureae** Thuern. On leaves of *S. salicifolia* in Siberia.
- Puccinia rupestris** Juel. Rust on leaves of *S. alpina* and *Carex rupestris* in Sweden and Norway.
- Puccinia saussurea** Thuern. Powdery dark-brown rust pustules on leaves of *S. glomerata* in Siberia and India.
- Puccinia saussureae-alpinae** Lindr. Brown rust sori on leaves of *S. alpina* in Finland.
- Puccinia vaginatae** Juel. Brown to black rust pustules on leaves of *S. alpina*, *Carex panicea*, and *C. vaginata* in Sweden.
- Puccinia vomica** Thuern. Leaf rust on *S. tanakae* and *Saussurea* sp. in Siberia.
- Ramularia saussureae** Thuern. Subcircular gray leaf spots on *S. glomerata* in Siberia.
- Septoria saussureae** Thuern. Circular gray leaf spots with dull-brown margins on *S. glomerata* in Siberia and China.
- Thecaphora trailii** Cke. Powdery purple-brown smut sori in flowers of *S. alpina* and *Carduus heterophyllus* in Scotland.
- Uromyces saussureae** Karst. Leaf rust on *S. circa*, *S. loureiri*, *S. nipponica*, and *Saussurea* sp. in Japan and Siberia.
- SAXIFRAGA**. SAXIFRAGE. Perennial herbs cultivated in gardens and rockeries.
- Caecoma cernuae** Th. Lind. Yellow leaf rust on *S. cernua* in Lapland.
- Cylindrosporium microspermum** (Speg.) Sacc. Indefinite circular leaf spots on *S. rotundifolia* in Italy.
- Diptochorella stromatica** (Rehm.) Theiss. and Syd. On leaves of *Saxifraga* sp. in Austria.
- Entyloma schinzianum** (P. Magn.) Bub. Leaf smut on *S. heuffeliana* and *S. rotundifolia* in Asia Minor, Switzerland, Yugoslavia, Austria, and Hungary.
- Leptosphaeria monotis** Rehm. On stems of *Saxifraga* sp. in Austria.
- Melampsora hirculi** Lindr. Rust on stems and leaves of *S. hirculus* in Russia, Finland, and Switzerland.
- Melampsora reticulata** Blytt. See *Salix*.
- Melampsora vernalis** Niessl. Golden, then dark-brown, rust pustules on leaves of *S. granulata* in Europe.
- Peronospora minima** G. W. Wils. Downy mildew on leaves of *S. cernua* in Norway.
- Peronospora saxifragae** Bub. Downy mildew on leaves of *S. granulata* in Bohemia and Germany.
- Phyllosticta groenlandica** Allesch. On leaves of *S. aizoides*, *S. oppositifolia*, and *S. stellaris* in Greenland.
- Phyllosticta saxifragae** P. Brun. Circular gray-white leaf spots with brown margins on *S. geum* in France.
- Phyllosticta saxifragicola** P. Brun. Circular leaf spots with brown margins on *S. cordifolia* in France.
- Puccinia fischeri** Crut. and Mayor. Leaf rust on *S. biflora* in Switzerland.
- Puccinia laurentiana** Trel. Leaf rust on *S. neglecta stolonifera* in Alaska.
- Puccinia huteri** Syd. Powdery brown leaf rust on *S. mutata* in Austria.
- Puccinia juliana** Diet. Dark-brown rust pustules on leaves of *S. aizoides* and *S. oppositifolia* in Norway and Austria.
- Puccinia pazschkei** Diet. Powdery dull-brown rust sori on leaves of *S. aizoon*, *S. elatior*, *S. longifolia*, and *S. mutata* in Switzerland and Austria.
- Puccinia saxifragae** Schlecht. Powdery brown leaf rust on *Saxifraga* spp. in Japan and Europe. Also reported from Oregon.
- Puccinia saxifragae-ciliatae** Barcl. Brown rust pustules on circular brown sunken leaf spots on *S. ciliata* and *S. lingulata* in India.
- Puccinia saxifragae-micranthae** Barcl. Brown rust sori on leaves of *S. micrantha* in India.
- Puccinia saxifragae-tricuspidatae** P. Henn. Yellow-brown leaf rust on *S. tricuspidata* in Greenland.
- Ramularia saxifragae** Syd. Large brown leaf spots on *S. granulata* in Europe.
- Ramularia sennensis** G. Frag. On leaves of *S. rotundifolia* in Spain.
- Septoria saxifragae** Pass. and forma *arctica* Allesch. On leaves of *S. aizoides*, *S. hirculus*, and *S. rotundifolia* in Greenland and Italy.
- Synchytrium rubrocinctum** P. Magn. Small galls on leaves of *S. granulata* in Germany.
- Synchytrium groenlandicum** Allesch. On leaves of *S. cernua* in Greenland.
- SCABIOSA**. Annual or perennial herbs, cultivated in gardens for their showy flowers. Sometimes called mourning bride.
- Aphelenchus oleisistus** Ritz. Bos. See *Begonia*.
- Cercospora scabiosicola** Rangel. On leaves of *S. atropurpurea* in Brazil.
- Peronospora knautiae** Fekl. Downy mildew on leaves of *S. columbaria*, *Knautia arvensis*, and *K. silvatica* in Switzerland, France, Scandinavia, Austria, and Germany.
- Peronospora violaceae** Berk. See *Dipsacus*.
- Ramularia bosniaca** Bub. Small circular to oblong, often confluent, dark-violet, then gray-white, leaf spots on *S. columbaria* in Yugoslavia.
- Ramularia knautiae** (Mas.) Bub. On leaves of *S. columbaria* and *S. lucida* in Dalmatia and Austria.
- Ramularia scabiosa** Lind. (*R. scabiosae* Jaap.) Subcircular leaf spots on *S. columbaria* and *S. lucida* in Denmark and Switzerland. Jaap has also described what is apparently the same species under the same name.
- Septoria scabiosicola** Desm. Circular purple-brown leaf spots on *S. arvensis*, *S. atropurpurea*, *S. balansae*, *S. columbaria*, *S. kladnikiana*, *S. ochroleuca*, and *S. succisa* in Europe.
- Stigmatea scabiosae** Lebed. On leaves of *S. ucranica* in Russia.
- Ustilago intermedia** Schroet. A smut destroying flowers of *S. arvensis* and *S. columbaria* in Denmark, Belgium, Great Britain, and Germany.
- Ustilago scabiosa** (Sow.) Wint. A smut forming powdery spore masses in anthers of *S. columbaria* and *Knautia arvensis* in Europe.
- SCAEVOLA**. Herbs or shrubs, sometimes cultivated.
- Phyllosticta scaevolae** E. and E. Circular dull-brown leaf spots on *S. chamissoniana* in Hawaii.
- Ramularia scaevolae** Rac. Circular pale-brown leaf spots on *S. koenigii* in Java.
- Uromyces puccinioides** Berk. Rust on leaves and flower stalks of *Scaevola* sp. and *Selliera radicans* in Queensland.
- SCANDIX**. Umbelliferous annual herbs cultivated for their flowers and foliage.
- Uredo petroselinii** DC. Leaf rust on *S. pecten-venensis* in France and Great Britain.
- SCHINUS**. PEPPER TREE. Dioecious, resinous, ornamental trees.
- Cercospora schinii** Syd. On leaves of *S. dependens* in Argentina.
- Meliola ludibunda** Speg. Black fungus patches on leaves of *S. dependens* in Brazil.
- Phyllosticta schinii** Thuern. Large irregular ochraceous to dull-brown leaf spots on *S. latifolius* and *S. molle* in Italy and Portugal.
- Septoria schinii** Tassi. Ashen leaf spots with black borders on *S. latifolius* in Italy.

- SCHIZANTHUS.** BUTTERFLY FLOWER. Erect, half hardy, annual herbs grown for their showy flowers.
- Cronartium flaccidum** (Alb. and Schw.) Wint. See Paonia.
- SCHIZOCODON.** Fringe bell. Glabrous perennial herbs.
- Puccinia schizocodonis** Pat. Brown to black rust pustules on leaves of *S. soldanelloides* in Japan.
- SCHIZOLOBIUM.** Tall leguminous trees.
- Phylachora schizolobii** P. Henn. Black stromata on leaves of *S. parahybium* (*S. excelsum*) in Brazil.
- SCHIZONOTUS.** See Sorbaria.
- SCHIZOPHRAGMA.** Ornamental vines grown for their handsome green foliage and showy clusters of white flowers.
- Mycosphaella exigua** Syd. On leaves of *S. hydrangeoides* in Japan.
- SCHIZOSTACHYUM.** See Bambuseae.
- SCILLA.** SQUILL. Perennial, bulbous plants with blue, rose, or white flowers. (Including *Endymion*)
- Aecidium doidgei** Syd. Leaf rust on *S. saturata* in the Union of South Africa.
- Aecidium scillae** Fekl. Yellow rust pustules on leaves of *S. bifolia* in England, France, and Switzerland. Said to be distinct from *Uromyces scillarum*, a common rust on Scillas.
- Aecidium scellinum** Dur. and Mont. This rust, reported on leaves of *S. autumnalis*, is probably only one stage of one of the *Scilla* rusts recorded elsewhere.
- Physoderma debeauxii** (Bub.) Syd. Ellipsoidal spots on the leaves of *S. maritima* (*Urginea scilla*) in Italy. The leaves are injured and bulb development is prevented.
- Puccinia ficulhoana** Lagh. Elongated black rust pustules on both leaf surfaces of *S. campanulata* (*Endymion campanulatum*) in Portugal.
- Puccinia lilacearum** Duby. See Ornithogalum.
- Puccinia rossiana** (Sacc.) Lagh. Rufous-brown rust pustules on both leaf surfaces of *S. bifolia* and *S. cernua* in Switzerland, Italy, Russia, and Hungary.
- Sclerotinia tuliparum** (Wakk.) Rehm. See Tulipa.
- Septoria scillae** West. See Muscari.
- Urocystis colchici** (Schlecht.) Rab. See Colchicum.
- Uromyces algeriensis** Syd. Circular to irregular cinnamon-colored rust pustules on both leaf surfaces of *S. obtusifolia* and other species of *Scilla* in Tunis and Algeria.
- Uromyces circinalis** Kalchbr. and Cke. This rust produces broad spots from 2 to 8 centimeters in diameter in which powdery, cinnamon-colored pustules form. On both leaf surfaces of *S. prasina* in the Union of South Africa.
- Uromyces scillarum** (Grev.) Wint. See Hyacinthus.
- Ustilago vaillantii** Tul. See Hyacinthus.
- SCOLYMUS.** Spanish oyster plant. Fleshy, edible roots.
- Phyllosticta scolyi** Pat. Angular ashen leaf spots on *S. hispanicus* in Tunis.
- Puccinia scolyi** Syd. Powdery brown rust sori on leaves of *S. grandiflorus* and *S. hispanicus* in north Africa, Syria, Spain, and Sicily.
- Ramularia cynaræ** Sacc. Gray spots with brown margins on leaves of *S. hispanicus* in Chile, France, and French North Africa.
- Ustilago scolyi** Roum. Powdery black masses of smut spores in flowers and receptacles of *S. grandiflorus* and *S. hispanicus* in Algeria and Tunis.
- SCOPOLIA.** SCOPOLINA. Ag. Erect, hardy, perennial herbs cultivated for their lurid-purple flowers.
- Aecidium pascheri** Bub. Rust on large round gray leaf spots on *S. japonica* in Japan.
- Ramularia scopalia** Vos. Irregular brown spots on leaves of *S. atropoides* in Austria.
- SCOPOLINA.** See Scopolia.
- SCORPIURUS.** Hardy stemless or decumbent herbs.
- Cercospora scorpiuri** Thum. Subcircular reddish-gray leaf spots with dull-brown margins on *S. muricata* in Portugal.
- SCORZNERA.** SERPENT ROOT. BLACK SALSIFY. Herbs cultivated for the fleshy, edible roots.
- Ascochyta scorzonerae** Rostr. Irregular brown-leaf spots on *S. hispanica* in Denmark.
- Cercospora scorzonerae** v. Hoeh. Oblong pale-brown leaf spots with broad purple margins on *S. humilis* in Austria.
- Phyllosticta scorzonerae** Pass. Small brown leaf spots on *S. humilis* in France and Bohemia.
- Puccinia angustifolia** McAlp. Yellow and dark-brown rust pustules on leaves of *S. angustifolia* in Australia.
- Puccinia scorzonerae** (Schum.) Jacky. Golden to brown rust pustules on leaves of *S. austriaca*, *S. hispanica*, *S. humilis*, *S. purpurea*, and *S. rosea* in China and Europe.
- Ramularia scorzonerae** Jaap. Circular to elongate dark-brown leaf spots on *S. aristata* and *S. humilis* in Austria and Spain.
- Sporidesmium scorzonerae** Aderh. Attacks stems and leaves of *Scorzonera* sp. in Germany.
- Ustilago receptaculorum** Tul. A smut destroying the flowers of *Scorzonera* sp. in France.
- Ustilago scorzonerae** (Alb. and Schw.) Schrot. Dark-violet smut spores in flowers of *S. aristata*, *S. hispanica*, *S. humilis*, and *S. purpurea* in Europe.
- SCROPHULARIA.** FIGWORT. Herbs, occasionally cultivated.
- Ascochyta scrophulariae** Kab. and Bub. Round to irregular zoned brown leaf spots on *S. nodosa* in Bulgaria and Bohemia.
- Phyllosticta ehrhartii** Sacc. Gray-white sinuous leaf spots on *S. aquatica* and *S. ehrhartii* in Portugal and Italy.
- Phyllosticta scrophulariae** Sacc. Ochraceous leaf spots on *S. nodosa* in Denmark, Russia, and Italy.
- Phyllosticta scrophulariae-bosniacae** Bub. Subcircular to ellipsoid yellow-brown leaf spots on *S. bosniaca* in Yugoslavia.
- Phyllosticta scrophularinea** Sacc. Small angular white leaf spots on *S. kakudensis* and *S. nodosa* in Japan and Italy.
- Ramularia nicolai** Bub. (*Ovularia carneola* Sacc.). Angular to irregular leaf spots on *S. bosniaca*, *S. ehrhartii*, *S. nodosa*, and *S. vernalis* in Yugoslavia, Switzerland, Spain, Denmark, and Austria.
- Ramularia scrophulariae** Fautr. and Roum. Purple leaf spots on *S. aquatica* and *S. nodosa* in France and Bulgaria.
- Septoria rohlenae** Bub. Angular, often confluent, gray leaf spots with brown margins on *S. scopoli* in Yugoslavia and Albania.
- Uromyces scrophulariae** (DC.) Fekl. Yellow and brown rust pustules on leaf blades, petioles, and stems of *S. aquatica*, *S. bosniaca*, *S. ebullifolia*, *S. ehrhartii*, *S. nodosa*, and *S. scorodonia* in Europe.
- SCUTELLARIA.** SKULLCAP. Annual and perennial herbs.
- Aecidium scutellariae** Syd. Leaf rust on *S. angulosa* and *S. repens* in India.
- Aecidium scutellariae-indicae** Diet. Leaf rust on *S. indica* in Japan.
- Erysiphe taurica** Lévy. See Althaea.

SEBASTIANA. Tropical shrubs.

Exoascus sebastianae Sadeb. Fruiting on lower surfaces and often deforming leaves of *S. brasiliensis* in Brazil.

Phyllachora subtropica Speg. Black stromata on leaves of *Sebastiania* sp. in Brazil and Paraguay.

Puccinia sebastianae Syd. Brown rust pustules on leaves of *S. klotzschiana* in Brazil.

Uredo sebastianae Wint. Leaf rust on *S. microstachys* in Brazil.

SECALE. Rye.

Acremoniella occulta Cav. Brownish-yellow flakes on culms of *Secale* and *Triticum* in Great Britain and Italy.

Bacillus cerealinus Gntner. See *Hordeum*.

Dilophia graminis (Fekl.) Sacc. See *Triticum*.

Fusarium nivale (Fr.) Sor. (*Calometria graminicola* B. and B.) Attacks the culms of *S. cereale*, *Hordeum*, *Lolium*, and *Triticum* in Europe, causing the so-called "snow disease."

Gloeosporium secalis Rostr. Gray-white leaf spots on *S. cereale* in Denmark.

Leptosphaeria eustoma (Fr.) Sacc. On culms and peduncles of *S. cereale*, *Holcus sorghum*, and *Phragmites* sp. in Europe.

Leptosphaeria herpotrichoides DeN. See *Triticum*.

Mycosphaerella basicola Frank. See *Hordeum*.

Ophiobolus cariceti (B. and Br.) Sacc. See *Triticum*.

Ophiobolus herpotrichoides (Fr.) Sacc. See *Triticum*.

Sclerotinia temulenta Prill and Delacr. (*Hymenoscypha temulenta* P. and D.) Forms sclerotia in place of grain in *S. cereale* in France. Flour made from grain contaminated with these sclerotia is poisonous.

Septocylindrium secales Oud. Pale-brown leaf spots on *S. cereale* in Holland.

Sphaeroderma damnosum Sacc. See *Triticum*.

Tilletia secalis (Cda.) Kueh. Stinking smut of rye (*S. cereale*) in Europe. The smut sori take the place of the grain and are composed of a dense black mass of spores. Reported from Washington and New York.

Tylenchus dipsaci Kuehn. See *Narcissus*.

Tylenchus tritici Bast. See *Triticum*.

SECHUM. CHAYOTA Ag. CHAYOTE. Cucurbitaceous tropical vines grown for their edible single-seeded fruit.

Cercospora sechiae Stevenson. Yellow subcircular, then irregular and indefinite, areas on leaves of *S. edule* (*C. edulis*) in Porto Rico and Cuba.

Helminthosporium sechicolum Stevenson. Small angular brown spots on leaves of *S. edule* (*C. edulis*) in Porto Rico.

Phyllosticta sechii Young. Irregular, then confluent, yellowish-white to gray-white leaf spots on *S. edule* (*C. edulis*) in Porto Rico.

SECURINEGA. Shrubs with bright-green foliage.

Notharavenella japonica Diet. Brown rust pustules on leaves of *S. flueggeoides* and *Flueggea japonica* in Japan and China.

SEDUM. STONECROP. Hardy, perennial, succulent herbs grown in rockeries.

Accidium koreaense P. Henn. Leaf rust on *S. maximowiczii* in Chosen.

Accidium sedi-alzooitis Tranzsch. Leaf rust on *S. aizoon* in Russia.

Ascochyta sedi Allesch. On stems of *S. affine* and *S. maximum* in Austria and Germany.

Ascochyta telephii Vesterg. Circular to irregular brown leaf spots with dark-purple margins on *S. telephium* in Sweden.

Endophyllum sedi (DC.) Lév. Brown leaf rust on *S. acre*, *S. nicaense* (*S. altissimum*), *S. boloniense*, *S. maximum*, *S. pruinatum*, *S. purpureum*, *S. reflexum*, and *S. sesangulare* in Europe.

Euryachora thoracella (C. Rusfr.) Schroet. Shiny black stromata, circular on the leaves and elongate on the stems of *S. fabaria*, *S. maximum*, *S. purpureum*, and *S. telephium* in Italy, Switzerland, Austria, Hungary, and Germany.

Helminthosporium rhopaloides Fres. and f. *sedi* P. Brun. On stems of *S. fabaria* in France.

Mycosphaerella sedi Jacz. On leaves of *S. telephium* in Russia.

Phyllosticta aizoi Cke. Small subcircular brown leaf spots on *S. aizoon* in Great Britain.

Puccinia australis Koern. Leaf rust on *S. boloniense*, *S. cepaea*, *S. mite*, *S. reflexum*, and *Molinia serotina* in Dalmatia, Italy, and Austria.

Puccinia benkel Kusano. Leaf rust on *S. alboroseum* and *S. telephium* in Japan.

Puccinia longissima Schroet. Leaf rust on *S. acre*, *S. boloniense*, *S. maximum*, *S. purpurascens*, *S. reflexum*, and *Koeleria* spp. in Sweden, Austria, Hungary, and Germany.

Puccinia sedi Koern. Dark-brown rust pustules on leaves and stems of *S. elegans* and *S. glanduliferum* in Italy, France, Switzerland, Belgium, and Germany.

SELAGINELLA. CLUB MOSS.

Helminthosporium filicicola P. Henn. See *Lygodium*.

Phyllosticta helvetica Sacc. On leaves of *S. helvetica* in Dalmatia.

Phyllosticta selaginellae Sacc. Indefinite pale-brown leaf spots on *S. helvetica* in Russia.

Sclerotium selaginellae Peyl. On leaves and stems of *Selaginella* sp. in Bohemia.

Synchytrium selaginellae Sorok. Doubtful species on *Selaginella* sp. in Russia.

SELINUM. Umbelliferous branched perennial herbs.

Accidium selini Lindr. Yellow leaf rust on *S. lineare* in Finland.

Physoderma vagans Schroet. See *Ranunculus*.

Puccinia apii Desm. See *Apium*.

SEMELE. CLIMBING BUTCHER'S-BROOM. Ornamental shrubs.

Ascochyta semeles Sacc. Gray-white oblong leaf spots with rufous margins on *S. androgyna* in Italy.

Phyllosticta semeles Ohl. Numerous white leaf spots with red borders on *S. androgyna* in Russia. Possibly only a stage of the preceding species.

SEMPERVIVUM. HOUSELEEK. Perennial stemless herbs or subshrubs with thick fleshy leaves.

Ascochyta sempervivi Fautr. On leaves and stems of *S. tectorum* in Denmark and France.

Endophyllum sempervivi (A. and S.) De By. Rust on leaves of *Sempervivum* spp. (about 20 species so far reported) in Europe. Introduced into New York City, where it has done considerable damage in public plantings.

SENECIO. GROUNDSEL. Composites with yellow flower heads.

Accidium dubiosum Syd. Leaf rust on *S. mikanioides* in the Union of South Africa.

Accidium halophilum Speg. Leaf rust on *S. alcornis* in Chile.

Accidium huaitinum Speg. Yellow leaf rust on *S. gilliesii*, *S. glabrum*, *S. halorragis*, *S. hieracioides*, *S. huaitata*, and *S. thermarum* in Chile and Argentina.

Accidium killmadscharicum P. Henn. Yellow leaf rust on *S. multicymbosus* in central Africa.

Accidium margeryanum Maire. Rust on leaves of *S. aquaticus* in France.

Accidium permulatum Syd. Leaf rust on *S. inornatus* in the Union of South Africa.

Accidium senecionis-stenocephali Diet. Leaf rust on *S. stenocephalus* in Japan.

SENECIO—Continued.

- Aecidium serrae* Syd. Leaf rust on *S. serra* in the Union of South Africa.
- Ascochyta senecionis* Fekl. On leaves of *S. sarracenicus* in Germany.
- Baedromus holwayi* Arth. Leaf rust on *S. cinerarioides* in Mexico.
- Baedromus senecionis* Syd. Leaf rust on *Senecio* sp. in Ecuador.
- Bremia tulasnei* (Hoffm.) Syd. Downy mildew on leaves of *S. vulgaris* in Germany.
- Cercospora jacquiniana* Thum. Irregular ochraceous leaf spots on *S. jacquinianus* in Germany.
- Colosporium senecionis* Fr. Yellow rust pustules on leaves of *Senecio* spp. and *Cineraria* spp. in Europe, Brazil, Uruguay, Chile, Siberia, Manchuria, Japan, and the Canary Islands. Collected once in Rhode Island. The aecial state occurs on needles of *Pinus austriaca*, *P. insignis*, *P. echinata*, *P. halepensis*, *P. mughus*, *P. montana*, *P. pinaster*, and *P. sylvestris*.
- Cylindrosporium eleonorae-reginae* Bub. and Malk. On leaves of *S. transilvanicus* in Bulgaria.
- Peronospora senecionis* Fekl. Downy mildew on leaves of *S. cordatus* in Germany.
- Phyllachora nidulans* Pat. Dull-black stromata on leaves of *Senecio* sp. in Ecuador.
- Phyllosticta albo-brunnea* Bub. and Wrobl. On leaves of *S. umbrosus* in Russia and Poland.
- Phyllosticta hualtatae* Speg. Subcircular gray-white leaf spots on *S. hualtata* in Argentina.
- Phyllosticta jacobaeae* Sacc. Indefinite ochraceous leaf spots on *S. jacobaea* in Italy.
- Phyllosticta senecionis-cordati* Allesch. Dull-yellow leaf spots on *S. cordatus* and *S. fuchsii* in Bohemia, Silesia, and Germany.
- Puccinia glomerata* Grev. Brown rust pustules on brown leaf spots on *S. jacobaea* in Great Britain and Belgium.
- Puccinia ligericae* Syd. Leaf rust on *S. viscosus* and *Carex ligerica* in Germany.
- Puccinia oedipus* Cke. Yellow-brown to black rust pustules on leaves of *S. pandurifolius* in the Union of South Africa.
- Puccinia schoeleriana* Plowr. and Magn. Leaf rust on *S. jacobaea* and *Carex* spp. in Europe.
- Puccinia senecionicola* Arth. Brown rust sori on leaves of *Cacalia amplifolia*, *C. anapullacea*, *C. calotricha*, *C. pringlei*, *C. obtusiloba*, *C. sinuata*, *Senecio angulifolius*, *S. petastioides*, *S. sinuatus*, and *S. warszewiczii* in Mexico and Guatemala.
- Puccinia senecionis* Lib. Brown to black rust pustules on leaves of *Senecio* spp. in central Europe and Australia.
- Puccinia senecionis-acutiformis* Hasl. and Mayor. Leaf rust on *Senecio* spp. and *Carex* spp. in Switzerland.
- Puccinia senecionis-ochrocarpi* Bacc. Leaf rust on *S. ochrocarpus* in Abyssinia.
- Puccinia silvatica* Schroet. See Taraxacum.
- Puccinia tasmanica* Diet. Black rust pustules on leaves and stems of *S. vulgaris* in Tasmania.
- Puccinia uralensis* Tranzs. Leaf rust on *S. fuchsii*, *S. nemorensis* and *S. scandens* in Ceylon, Yugoslavia, Russia, and Hungary.
- Ramularia chlorina* Bres. On leaves of *S. fuchsii* in Germany.
- Ramularia coeosporii* Sacc. See Campanula.
- Ramularia pruinosa* Speg. Small circular ochraceous leaf spots which finally involve the entire leaf area of *S. jacobaea* in Great Britain and Italy.
- Ramularia senecionis* (B. and Br.) Sacc. On leaves of *S. carniolicus*, *S. jacobaea*, *S. nebrodensis*, *S. (Cineraria) palustris*, and *S. subalpinus* in Great Britain, the Balkans, and Austria. Reported from Colorado.
- Septoria anaxaea* Sacc. Ochraceous leaf spots on *S. praealtus* in Italy.
- Septoria martiniae* Cke. Gray confluent leaf spots on *S. bedfordii* in Victoria.
- Septoria selioi* Speg. Small circular gray-white leaf spots on *S. selioi* in Argentina.
- Septoria senecionis* West. Circular to irregular gray-black leaf spots with white centers on *Senecio* spp. in Siberia and Europe. Reported from California.
- Uromyces ameghinoi* Speg. Leaf rust on *S. laevicaulis* in Patagonia.
- Uromyces araucanus* Diet. and Neg. Brown leaf rust on circular spots on *S. otites* in Chile.
- Uromyces kurtzii* P. Henn. Rust deforming leaves and stems of *S. salsus* in Argentina.
- Uromyces psamathophilus* Speg. A rust deforming leaves, branches, and inflorescences of *Senecio* sp. in Argentina.
- Uromyces senecionicola* Arth. Brown leaf rust on *S. roldana* in Mexico and Guatemala.
- SERJANIA.** Climbing or twining tropical shrubs.
- Aecidium serjaniae* P. Henn. Ochraceous rust pustules on yellow leaf spots on *S. fulla* and *Serjania* sp. in Brazil and Argentina.
- Dothidella serjaniae* P. Henn. Brown stromata on stems of *Serjania* sp. in Brazil.
- Phyllachora duplex* Rehm. Shiny black stromata on brown spots on leaf blades and petioles of *S. caracasana* in Brazil and Argentina.
- Phyllachora intermedia* Speg. Small brown galls on leaves and stems of *S. glabrata* in Brazil. The fungus is said to be more properly a *Valsaria*.
- Phyllachora subrepens* Speg. Black stromata on leaves of *Serjania* sp. in Paraguay and Brazil.
- Physalospora serjaniae* Rehm. Circular red to dull-brown leaf spots on *Serjania* sp. in Brazil.
- SEBRATULA.** Thistlelike perennial herbs with purple or violet flower heads.
- Ovularia serratulae* Sacc. Leaf spot on *S. tinctoria* in Spain.
- Puccinia heterophyllae* Cke. Powdery brown rust pustules on leaves and stem of *S. cerinthefolia* and *S. heterophylla* in Kurdistan and Asia Minor.
- Puccinia miurae* Syd. Leaf rust on *S. atriplicifolia* in Japan.
- Puccinia schirajewskii* Tranzsch. Leaf rust on *S. pallida* in India.
- Puccinia schroeteriana* Kleb. Leaf rust on *S. radiata*, *S. tinctoria*, and *Carex* spp. in Russia and central Europe.
- Puccinia serratulae* Thum. Brown leaf rust on *S. tinctoria* and *Serratula* sp. in Siberia and Russia.
- Puccinia serratulae-oligocephalae* Syd. Powdery brown to black rust pustules on leaves of *S. oligocephala* in Syria.
- Puccinia serratulae-pinnatifidae* Gz. Frag. Leaf rust on *S. pinnatifida* in Spain.
- Puccinia tinctoricola* P. Magn. Leaf rust on *S. centauroides*, *S. coronata*, *S. heterophylla*, *S. sloanei*, and *S. tinctoria* in Siberia and Europe.
- Septoria cirsi* Niessl. Subcircular brown leaf spots on *S. glauca* and *Cirsium arvense* in Italy.
- Septoria tinctoriae* Brun. On leaves of *S. tinctoria* in Germany.
- SESAMUM.** SESAME. Tropical herbs producing oil-bearing seeds.
- Ascochyta* sp. On *S. orientale* in south Russia.
- Bacillus sesami* Malkoff. Attacks *S. orientale* (*S. indicum*) in connection with *Pseudomonas sesami* in Bulgaria.
- Cercospora sesami* A. Zimm. Numerous small light-brown spots with dark-brown margins on leaf blades, petioles, stems, and capsules of *S. orientale* (*S. indicum*) in Porto Rico, Cuba, Brazil, Tanganyika, the Philippines, Ceylon, China, and Japan. Reported from Florida.
- Fusarium* sp. Causes a serious wilt of *S. orientale* in India. The disease is seed-borne.

SESAMUM—Continued.

Helminthosporium sesami Miy. On leaves of *S. orientale* in China.

Hypochnus cucumeris Frank. See Cucumis.

Phytophthora parasitica Dast. See Ricinus.

Pseudomonas sesami Malkoff. Produces brown to black spots on leaves and stems of *S. orientale* in Bulgaria. Diseased plants turn black and rot.

Rhizoctonia sp. See Vigna.

SESBANIA. Leguminous herbs or shrubs.

Cercospora agatidis Foex. On leaves of *S. grandiflora* in Indo-China.

Cercospora sesbaniae P. Henn. Effuse yellowish leaf spots on *S. grandiflora* in India and the Congo.

Uredo fimbriata Speg. Brown leaf rust on *Sesbania* sp. in Brazil.

Uredo sesbaniae P. Henn. Rust pustules on yellow leaf spots on *S. aegyptiaca* and *Sesbania* sp. in India and the Congo.

SESEL. Perennial and biennial herbs.

Puccinia phymatospora Lindr. Brown to black rust pustules on leaves of *S. coloratum* in Italy, France, Belgium, and Switzerland.

Septoria desciscens Sacc. On leaves of *S. strictum* in Siberia.

Septoria seseli Höll. On leaves of *S. glaucum* in Hungary.

Uromyces graminis (Niessl.) Diet. Powdery brown to black rust pustules on leaves of *S. glaucum*, *Laserpitium siler*, and *Melica ciliata* in Europe.

Uromyces seseli-graminis Ed. Fisch. See Arrhenatherum.

SETARIA. **CHAETOCHELOA** Ag. MILLET. Annual and perennial grasses, mostly weeds. See, also *Chaetochloa*.

Acrothecium lunatum Wak. See Panicum.

Balanisia claviceps Speg. See Pennisetum.

Phyllachora evansii Syd. Dull-black stromata on yellowish leaf spots on *S. aurea*, *S. nigrirostris*, and *S. sulcata* in the Union of South Africa.

Phyllachora setariaeacola Speg. Black stromata on leaves of *Setaria* sp. in Brazil and Ecuador.

Phyllosticta crastophila Sacc. See Phalaris.

Phyllosticta glumarum-setariae P. Henn. Pale-brown spots on glumes of *S. aurea* in the Congo.

Phyllosticta setariae Ferr. Irregular gray-white leaf spots with purple margins on *S. glauca* in Italy.

Puccinia atra Diet. and Holw. Black rust sori on leaves of *S. grisebachii* in Mexico.

Puccinia cameliae (Mayor) Arth. Leaf rust on *S. scandens* and *Chaetochloa setosa* in Porto Rico and Colombia.

Puccinia setariae Diet. and Holw. Brown rust pustules on leaves of *S. caudata* and *S. imberbis* in Chile and Mexico.

Scirrhia setariae (Sacc.) Theiss. and Syd. On *S. glauca* in Italy.

Sorisorium setariae McAlp. Ovaries converted into black powdery masses of smut spores. On *S. glauca* in Queensland.

Sphacelia grisea Speg. On culms of *Setaria* sp. in Argentina.

Sphaelotheca pamparum (Speg.) Clint. Smut sori in the ovaries as oblong bodies protected by false membranes which break irregularly, disclosing dusty olive-black spore masses. On *S. geniculata* and *S. imberbis* in Mexico, Cuba, Bahamas, and Chile.

Sporotrichum peribebuense Speg. On leaves of *Setaria* sp. in Brazil.

Tolyposporium pampeanum Speg. Dark powdery smut sori in spikelets of *Setaria* sp. in Argentina.

Tolyposporium setariicolum Syd. Smut on *S. aurea* in Central Africa.

Uromyces setariae-italicae (Diet.) Yosh. Cinnamon-brown to black rust pustules on leaves of *S. glauca*, *S. intermedia*, *S. italica*, *S. verticillata*, and *S. viridis* in India, Japan, Ceylon, China, Manchuria, the Philippines, and north Africa.

Ustilago inoidea setariae Bref. Powdery smut sori in ovaries of *S. crus-ardea* in Brazil.

Ustilago evansii P. Henn. Smut on *S. aurea* in Africa.

Ustilago heterospora P. Henn. See Panicum.

Ustilago setaria Rbh. A smut destroying the panicles of *S. glauca*, *S. viridis*, and *Syntherisma sanguinale* in Argentina and Germany.

Ustilago setariae-aureae P. Henn. Black powdery smut sori destroying the ovaries of *S. aurea* in the Congo and the Union of South Africa.

Ustilago verruculosa Wakef. Smut on *S. aurea* in tropical Africa.

SHAWIA. See Olearia.**SHOREA**. Tropical trees.

Corticium salmonicolor B. and Br. See Citrus.

Fomes lignosus Klotzsch. See Hevea.

Polyporus shoreae Wakef. Rot of trunk and branches of *S. robusta* in India.

SIDA. Malvaceous herbs or shrubs, mostly weeds, one species cultivated for fiber.

Cercospora densissima Speg. Large indefinite pale-brown leaf spots on *S. rhombifolia* in Porto Rico and Argentina.

Cercospora sidae P. Henn. Pale-brown leaf spots on *S. cordifolia* in the Congo.

Dietelia verruciformis P. Henn. Yellow-brown rust pustules on leaves of *S. macrodon* in Argentina.

Phyllosticta sidaeacola Cke. Irregular, then confluent, brown leaf spots on *S. napaea* in Great Britain.

Septoria heterochroa Desm. See Malva.

SIDEROXYLON. Tropical trees and shrubs, some of value for timber.

Phyllachora baumii P. Henn. Black stromata on gray to brown leaf spots on *Sideroxylon* sp. in the Union of South Africa.

Puccinia johnstonii Arth. Brown leaf rust on *Dipholtis salicifolia* and *S. foetidissimum* in Cuba.

SILENE. CAMPION. CATCHFLY. Erect annual, biennial, or perennial herbs.

Marsonia delastrei (Delacr.) Sacc. See Lychnis.

Myosphaerella silenii v. Hoeh. Numerous small ochraceous spots on leaves and calyces of *S. latifolia* in Austria.

Peronospora vexans Gäum. Downy mildew on leaves of *Silene* spp. in Europe.

Phacidium pumilum Desm. On stems and leaves of *S. conica* and *S. rubella* in France and Tunis.

Phyllosticta nebulosa Sacc. Pale-brown leaf spots on *S. armeria*, *S. pendula*, and *S. splendens* in north Africa, Denmark, and Italy.

Phyllosticta otites Brun. Olivaceous leaf spots on *S. otites* in France.

Phyllosticta zahlbruckneri Baenil. Subcircular gray-white leaf spots on *S. latifolia* and *S. nutans* in Russia, Hungary, and Austria.

Puccinia silenens Schroet. Brown rust pustules on leaves of *Silene* spp. and *Melandryum* sp. in Siberia and Europe.

Ramularia didymarioides Br. and Sacc. Indefinite ochraceous leaf spots on *S. latifolia* in Spain and France.

SILENE—Continued.

- Ramularia lychnicola** Cke. See *Lychnis*.
Ramularia silenes Karst. Ochraceous leaf spots on *Silene* sp. in Finland.
Septoria apetalae P. Magn. On leaves of *S. apetalae* in Palestine.
Septoria dimera Sacc. Indefinite ochraceous leaf spots on *S. dichotoma*, *S. latifolia*, *S. livida*, and *S. nutans* in Europe.
Septoria doehlii Syd. Circular gray-white leaf spots on *S. nutans* in Germany.
Septoria dominii Bub. and f. *calycicola* Gz. Frag. Small circular dull-yellow leaf spots on *S. latifolia* in Yugoslavia and Spain.
Septoria silenes-nutantis C. Mass. Subcircular yellow-white leaf spots on *S. nutans* in Italy.
Septoria saponariae (DC.) Sav. and Becc. See *Saponaria*.
Sorosporium saponariae Rud. See *Cerastium*.
Uromyces behensis (DC.) Ung. Dark-brown to black rust pustules on leaves of *Silene* spp. in Europe.
Uromyces inaequalis Lasch. Yellow to dark-brown rust pustules on leaves of *Silene* spp. in Europe, Siberia, Asia Minor, and Japan.
Ustilago major Schroet. Dark-violet smut sori in anthers of *S. otites* in Europe.
- SILYBUM. MARIANA** Ag. Erect thistlelike plants.
Aecidium marianum Gz. Frag. Leaf rust on *S. marianum* in Spain.
Hypochnus fuciformis (Berk.) McAlp. See *Lolium*.
Ramularia cynarae Sacc. See *Cynara*.
Septoria silybi Pass. Subcircular dull-brown to gray leaf spots on *S. marianum* in Italy.
Ustilago cardui F. v. Wald. Powdery brown-violet masses of smut spores replacing ovaries of *S. marianum* and *Carduus acanthoides* in Germany.
- SIPHOCAMPYLUS.** Herbs or shrubs, sometimes climbing.
Uredo siphocampyli P. Henn. Powdery brown rust pustules on leaves of *Siphocampylus* sp. in Argentina.
Uromyces cystopiformis Lagerh. Leaf rust on *Siphocampylus* sp. in Ecuador.
- SIRMUELLERA.** See *Banksia*.
- SISYRINCHIUM. BLUE-EYED GRASS.** Iridaceous fibrous-rooted perennials.
Aecidium bunsteri Neg. Rust on leaves of *S. andinum* in Chile.
Phyllosticta chilensis Allesch. Leaf spots on *S. pedunculatum* in Chile.
Puccinia angulata Diet. & Neg. Yellow rust pustules, becoming dark-brown to black, on both leaf surfaces and on scapes of *S. pedunculatum* in Chile.
Puccinia sisyrinchii Mont. (*Uromyces sisyrinchii* Mont.) Produces brown to black rust pustules on both leaf surfaces and on scapes of *S. graminifolium* and *S. palmifolium* in Chile and Argentina. Reported from Washington.
Puccinia straminea Diet. Brown to dark-brown powdery rust pustules on both surfaces of leaves of *Sisyrinchium* sp. in Brazil.
Septoria sisyrinchii Speg. Leaf spot on *S. bonariense* in Argentina.
Uredo nominata Arth. Leaf rust on *S. bermudianum* in Bermuda.
- SMILACINA. VAGNERA** Ag. FALSE SOLOMON'S SEAL. Perennial herbs.
Protomyces purpureo-tingens Mass. Elongated or broadly effused red or purple patches on cotyledons and young leaves of *Smilacina* sp. in Great Britain.
- SMILAX. GREENBRIER. CATBRIER.** Herbaceous or shrubby climbers.
Blastospora smilacis Diet. Yellow rust pustules on small sunken leaf spots on *S. herbacea* and *S. sieboldii* in Japan.
Cercospora miyakei P. Henn. Circular to angular, then confluent, leaf spots on *S. herbacea* in Japan.
Cercospora smilacina Speg. Small brown leaf spots on *Smilax* sp. in Brazil. Saccardo has given the same name to a species on leaves of *S. aspera* and *S. mauritanica* in Italy, Spain, Portugal, and the United States.
Dothidella smilacicola Rehm. Black stromata on leaves of *Smilax* sp. in Brazil.
Gloeosporium thuementi Sacc. See *Anthurium*.
Marsonia smilacina Thuem. Irregular ochraceous leaf spots on *S. mauritanica* in Spain and Portugal.
Phaeochorella clypeata (Wint.) Theiss. and Syd. Shiny black stromata on leaves of *Smilax* sp. in Brazil.
Phyllosticta smilacina Speg. Rufous leaf spots on *S. campestris* in Argentina.
Puccinia citrina Syd. Leaf rust on *S. gaudichaudiana*, *S. stenopetala* and *Smilax* sp. in China, Japan, and the Philippines.
Puccinia ferruginea Lév. Dull brown rust pustules on leaves of *S. aspera* and *S. zeylanica* in Macao and Ceylon.
Puccinia henryana Syd. Leaf rust on *S. china*, *S. herbacea*, and *S. menispermoides* in Japan and China.
Puccinia kraussiana Cke. Brown rust pustules on leaves of *S. kraussiana* in central and south Africa.
Puccinia merrillii P. Henn. Leaf rust on *S. bracteata*, *S. latifolia*, and *S. vicaria* in the Philippines.
Puccinia prainiana Barcel. Brown rust pustules on leaves of *S. aspera*, *S. elegans*, and *S. maculata* in India.
Puccinia smilacis-chinae P. Henn. Leaf rust on *S. chinense* and *S. herbacea* in Japan and the Philippines.
Sphenospora pallida (Wint.) Diet. Waxy rust pustules on leaves of *Smilax* sp. in Guatemala, Costa Rica, and South America.
Uredo yurimaguasensis P. Henn. Leaf rust on *Smilax* sp. in Peru.
- SOBRALIA.** See *Orchidaceae*.
- SOJA. SOYBEAN.** Herbaceous leguminous plants, with a wide range of economic uses. (Including *Glycine*.)
Aecidium glycines P. Henn. Leaf rust on *S. max* in Tanganyika and Uganda.
Asochyta sp. On *S. max* in Japan.
Bacterium sp. A leaf-spotting disease of *S. max*, said to be due to a bacterium distinct from other species on this host, is reported from Japan.
Cercospora daizii Miura. Leaf spot on *S. max* in Manchuria.
Cercospora glycines Cke. Definite dark-brown leaf spots on *Glycine clandestina* in Australia.
Colletotrichum glycines Hori. Anthracnose on stems and pods of *S. max* in Japan and Chosen.
Fusarium sp. Causes ring spot disease of *S. max* in Manchuria.
Hypochnus centrifugus Tul. Causes cankers on stems, blighting infected plants of *S. max* in Manchuria.
Hypochnus cucumeris Frank. See *Cucumis*.
Mosaic. Mosaic disease of *S. max* reported from Japan is probably the same as the mosaic and crinkling reported from the United States.

SOJA—Continued.

Mycosphaerella phaseolarum Siem. See *Phaseolus*.

Peronospora manshurica (Naoum.) Syd. (*P. trifoliorum* De B. var. *manshurica* Naoum.) Downy mildew on leaves of *S. maz* in Siberia, India, Formosa, and Manchuria, causing premature leaf fall.

Phyllosticta sojaecola Massal. (*Pharosphaerulina sojaecola* Miura.) Subcircular, dull-brown leaf spots on *S. maz* in Japan, Russia, Manchuria, and Italy.

Pseudomonas glycines Nak. Circular yellow leaf spots, becoming brown to dark-brown with yellow margins, on *S. maz* in Japan. The spots may be as numerous as 70 to 80 per square centimeter, causing death of infected leaves.

Septoria glycines T. Hem. Brown leaf spots on *S. maz* in Japan and Manchuria.

Septoria sojae v. Thum. Irregular yellowish leaf spots on *S. maz* in Italy, Japan, and Austria.

Troteria venturioides Sacc. Black mildew on leaves of *S. maz* in the Philippines.

Uredo vignae Bres. See *Vigna*.

Uromyces sojae (P. Henn.) Syd. Brown rust pustules on leaves of *S. maz* in Japan, Java, China, India, Formosa, Manchuria, and the Philippines.

SOLANUM. NIGHTSHADE. POTATO. EGGPLANT. Temperate and tropical herbs and shrubs, many species of economic, ornamental, or medicinal value.

Actinomyces spp. A number of species of *Actinomyces* have been described from Germany, causing various types of scab on potato (*S. tuberosum*) tubers. Whether these forms differ from *A. scabies* the common American species, is problematical as yet. These forms include *A. albus* (K. D.) Gasp. (girdle scab); *A. incanescens* Wr. (deep scab); *A. intermedius* (Krug.) Wr. (shallow scab); *A. nigricans* (Krug.) Wr. (girdle scab); *A. aeruginus* Wr. (knobby scab), and *A. tricolor* Wr. (shallow scab).

Aecidium habuense P. Henn. Leaf rust on *S. melongena* in India.

Aecidium solani Beeli. Leaf rust on *Solanum* sp. in Belgian Congo.

Aecidium solani-unguiculati P. Henn. Leaf rust on *S. unguiculatum* in Abyssinia.

Ascochyta dulcamara Bub. On leaves of *S. dulcamara* in Bohemia and Germany.

Ascochyta solani-nigri Died. Circular to ovate white leaf spots with dark margins on *S. nigrum* in Germany.

Bacillus spp. Bacteria, the occurrence of which is unknown in the United States, cause the rotting of potato (*S. tuberosum*) tubers in India and Malaya.

Bacillus caulivorus Prill, and Delacr. Said to cause cankers on the stems of *S. tuberosum*, *Lycopersicon esculentum*, and *Nicotiana tabacum* in Italy and France.

Bacillus tubifex Dale. Irregular brown spots, generally along veins or margins of leaves of *S. tuberosum* in Great Britain.

Bacterium sp. An undescribed species of *Bacterium* is reported from Morocco as causing a serious potato (*S. tuberosum*) disease. The disease begins at the apex of the plant as brown patches on the leaflets, which rapidly turn black and wilt. The disease works down the petioles into the stems and finally to the tubers. The tops of the plants are covered with elongate black patches and finally die. Small violet-black spots appear on the tubers, enlarging until the entire tuber surface is involved, a soft rot ensuing.

Cercospora aratali Speg. Circular ashen-brown leaf spots on *S. glaucum* in Argentina.

Cercospora dulcamaricola Höll. Circular brown leaf spots on *S. dulcamara* in Hungary.

Cercospora heterosperma Bres. On leaves of *S. tuberosum* in Poland.

Cercospora incarnata P. Henn. On leaves of *Solanum* sp. in Brazil.

Cercospora melongena Welles. Irregular, then confluent, zoned, chlorotic areas on leaves of *S. melongena* in the Philippines.

Cercospora solanacea Sacc. and Berl. Subcircular brown leaf spots on *S. verbascifolium* in Australia.

Cercospora tosenensis P. Henn. Circular, then confluent, dull-brown spots on leaves of *S. biflorum*, *S. nigrum*, and *Solanum* sp. in the Philippines and Japan.

Chrysophlyctis endobiotica Schilb. (*Synchytrium endobioticum* [Schilb.] Perc.) The potato (*S. tuberosum*) wart disease or potato canker is one of the most serious diseases of this important crop. It occurs in Great Britain, Holland, Norway, Poland, Luxemburg, the Union of South Africa, and Germany. In the United States the disease also occurs in limited areas in Maryland, West Virginia, and Pennsylvania. The disease is characterized by wartlike outgrowths on the tubers, which vary in size and distribution. They commonly develop from the eyes of tubers, more rarely on stolons or base of stems. The warts are at first whitish, turning black with age. Exposed warts take on a green color. In size they vary from small nodules to masses larger than the tubers on which they are borne. The organism causing the disease can persist in the soil for many years and is readily carried from place to place in this medium. The tomato (*Lycopersicon esculentum*) has been found capable of harboring the disease. By inoculation *S. dulcamara* and *S. nigrum* have been found susceptible.

Didymopora solani Diet. Leaf rust on *Solanum* sp. in Brazil.

Didymopora solani-argentei (P. Henn.) Diet. Rust pustules on circular dull-brown leaf spots on *S. argenteum* in Brazil.

Guignardia nervisequa (Rehm.) Sacc. and Syd. On leaves of *S. auriculatum* in Brazil.

Helminthosporium solani McAlp. On leaf blades and petioles and stems of *S. viride* in New South Wales.

Helminthosporium solanum Sacc. and Syd. Leaf spots on *S. argenteum* in Brazil.

Leptosphaeria circinans (Fckl.) Sacc. See *Asparagus*.

Monilopsis aderholdi Ruhl. The "vermerbrungs Pilze" attacks the tubers of *S. tuberosum* and also causes a damping-off of seedlings of many species of cultivated plants, including *Beta vulgaris*, *Brassica* spp., *Cinchona* sp., and *Raphanus* in Europe and Java. The fungus forms black sclerotia similar to those of *Rhizoctonia solani*, with which it is sometimes considered identical.

Oospora pustulans Owen and Wakef. The skin spot disease of potatoes (*S. tuberosum*) occurs in Great Britain and in limited areas in Washington and British Columbia. Numerous small circular, slightly raised areas occur on the tubers. The possible connection of this disease with immature stages of powdery scab (*Spongospora*) has not yet been definitely settled.

Phellomyces sclerotiphorus Frank. Reported as the cause of a scab and dry rot of potato (*S. tuberosum*) tubers in Russia and Great Britain.

Phoma solanicola Prill, and Delacr. See *Nicotiana*.

Phylachora fluminensis Theiss. Shiny black stromata on brown leaf spots on *Solanum* sp. in Brazil.

Phyllohendersonia dulcamaræ (Sacc.) Tass. On leaves of *S. dulcamara* in Italy.

Phyllosticta concentrica Th. On leaves of *Solanum* sp. in Brazil.

Phyllosticta dulcamaræ Sacc. Brown, often marginal, leaf spots on *S. dulcamara* in France, Italy, and Denmark. Reported from Kansas.

Phytophthora arecae (Colem.) Pethybr. See *Palmae*.

SOLANUM—Continued.

Phytophthora erythroseptica Pethybr. This downy mildew fungus causes a wilting of potato (*Solanum tuberosum*) plants and a moist rot of the tubers in Ireland, Switzerland, Holland, and Dutch East Indies. Cut surfaces of infected tubers turn pink rapidly upon exposure to air, becoming purplish-brown to black after several hours. Diseased tubers remain firm, but of rubbery consistency, and if pressed exude a quantity of juice, becoming finally completely rotted. *Atropa belladonna* is reported as an additional host in Poland.

Phytophthora melongenae K. Saw. Downy mildew attacking primarily fruit of *S. melongena* in Japan but also capable of infecting *S. tuberosum*, *Areca catechu*, *Hibiscus esculentus*, *Lycopersicon esculentum*, *Nicotiana* sp., and *Ricinus* sp.

Phytophthora parasitica Dastur. See *Ricinus*.

Pseudomonas solaniolens Paine. This bacterium has been described as the cause of internal brown spot of potato (*S. tuberosum*) tubers in Europe.

Puccinia araucana Diet. and Neg. Yellow and brown rust pustules on deformed leaves and branches of *S. cryptopodium* and *S. valdivianum* in Chile.

Puccinia hieronymi P. Henn. Leaf rust on *S. incisum* and *S. nodiflorum* in Argentina.

Puccinia huallagensis P. Henn. Cinnamon-brown rust sori on dark-brown leaf spots on *S. torrum* in Peru and Colombia.

Puccinia imitans Syd. Leaf rust on *Solanum* sp. in Ecuador.

Puccinia nigeriana Diet. Dark-brown rust pustules on leaf blades and petioles of *S. furcatum* in Chile.

Puccinia pitteriana P. Henn. Numerous brown rust pustules on lower leaf surfaces of *S. tuberosum* and *Lycopersicon esculentum* in Costa Rica and Colombia. This rust appears to be capable of causing heavy damage to the host plants through drying and premature fall of leaves.

Puccinia solanacearum Sacc. and Syd. Black rust pustules on leaves of *Solanum* sp. in India.

Puccinia solanita (Schw.) Arth. Brown leaf rust on *Solanum* sp. and *S. donnellsmithii* in Trinidad, Panama, and Surinam.

Puccinia solani-tristis P. Henn. Powdery black rust pustules on leaves of *S. triste* in Brazil.

Puccinia tijibodensis Gáum. Brown leaf rust on *S. biflorum* in Java.

Puccinia tubulosum (Pat. and Gaill.) Arth. See *Paspalum*.

Puccinia weberbaueri P. Henn. Dark-brown rust pustules on yellow-brown leaf spots on *Solanum* sp. in Peru.

Pucciniosira solani Lagerh. Yellow leaf rust on *Solanum* sp. in Ecuador.

Rhizoctonia destruens Tassi. This fungus attacks its hosts at the ground level, setting up a rot which destroys the plants, the numerous brown sclerotia appearing on the rotted areas. Potato tubers are soft rotted, thick white strands of mycelium growing over the surface. The fungus is reported from Italy, India, and Java, on *S. melongena*, *S. tuberosum*, *Amorphophallus campanulatus*, *Arachis hypogaea*, *Delphinium decorum*, *D. grandiflorum*, *D. magnificum*, *D. machayanum*, *D. montanum*, *Dianthus* sp., *Hibiscus rosa-sinensis*, *Lobelia laxiflora*, *Medicago sativa*, *Oryza sativa*, *Piper betel*, and other hosts. The fungus is not distinct from *Sclerotium rolfsii*, but may represent a strain not present in the United States.

Sarcinella raimundoi Sacc. On leaves of *S. melongena* in the Philippines.

Septoria dulcamaræ Desm. Small subcircular brown, then gray, leaf spots on *S. dulcamara* in Europe.

Septoria pseudo-quinae Pat. Dull-brown to gray leaf spots with black borders on *S. pseudo-quina* in Ecuador.

Septoria solanina Speg. Small circular dull-brown leaf spots on *S. gracile* in Argentina.

Septoria solani-nigri Scalia. Circular, then confluent, dull-brown leaf spots on *S. nigrum* in Italy.

Septoria solanophila Speg. Circular gray-white leaf spots with purple margins on *S. verbascifolium* in Argentina.

Spongospora subterranea (Walls.) Lang. Powdery scab is a disease introduced within recent years into the United States from Europe, and now established in the northern potato (*S. tuberosum*) growing regions. It does not seem to be able to persist south of the New England and Lake States. It occurs in northern Europe, Algeria, northern South America, and New Zealand. Other hosts are *S. citiatum*, *S. commersoni*, *S. haematocladum*, *S. mammosum*, *S. marginatum*, *S. warszewiczii*, and *Lycopersicon esculentum*.

The disease closely resembles common scab, but is distinguished by the smaller, more definite lesions, which on maturity break open, revealing brown, powdery masses, surrounded by the torn skin of the tubers.

Sporidesmium exitiosum Kuehn. var. *solani* Fekl. On stems of *S. tuberosum* in Russia. See also *Brassica*.

Sporidesmium melongenae Thuem. On leaves of *S. melongena* in Portugal and Brazil.

Tylenchus dipsaci Kuehn. A biological strain of this nematode attacks potato (*S. tuberosum*) tubers in Great Britain. Brown patches appear on the skin followed by internal rot. See also *Narcissus*.

Uredo minitans Speg. Brown rust pustules on circular leaf spots on *S. commersoni* in Argentina.

Urocystis hieronymi Schroet. Ochraceous smut sori on branches and peduncles of *Solanum* sp. in Bolivia.

Uromyces solani Diet. and Holw. Yellow-brown to dark-brown rust pustules on leaves of *S. appendiculatum* and *S. nudum* in Mexico.

Vermicularia capsici Syd. See *Capsicum*.

Vermicularia varians Duc. The black dot disease or "dartrose" is considered a serious disease of *S. tuberosum* in France and some other European countries. It is also reported from the Union of South Africa, Australia, Brazil, and Canada. The fungus undoubtedly occurs in the United States, but has not been associated with a definite disease of the potato. In Europe the disease attacks the stems, roots, and tubers. The fungus also occurs on *Lycopersicon esculentum* and *Physalis peruviana*. It is more properly referable to *Colletotrichum*.

SOLARIA. Bulbous plants.

Uromyces solariae Diet. Chestnut colored powdery rust pustules on both leaf surfaces of *S. miersioides* in Chile.

SOLDANELLA. Small, glabrous, perennial herbs with short rhizomes.

Puccinia soldanelleae (DC.) Fekl. Yellow and dark-brown rust pustules on leaves of *S. alpina*, *S. minima*, *S. montana*, and *S. pusilla* in Europe.

Septoria soldanelleae Speg. and var. *pyrolaeifoliae* Vogl. Brown spots, often occupying the entire leaf area, on *S. alpina*, *S. montana*, and *S. pyrolaeifolia* in Italy.

Septoria versicolor Bub. Circular to irregular yellow-white leaf spots on *S. montana* in Bohemia.

Thecaphora hyalina Fingerh. See *Convolvulus*.

SOLIDAGO. GOLDENROD. Perennial herbs.

Cercospora fulvescens Sacc. Small yellow-brown leaf spots on *S. virgaurea* in Denmark, Italy, and Japan. Reported from Colorado and Kansas.

Mycosphaerella virgaureae Krieg. On leaves of *S. virgaurea* in Germany.

Ovularia virgaureae (Thum.) Sacc. On leaves of *S. virgaurea* in Denmark, Switzerland, and Germany.

Phyllosticta solidaginis Bres. On leaves of *S. virgaurea* in Germany.

Puccinia solidaginicola Diet. Leaf rust on *Solidago* sp. in Chile.

Puccinia solidaginis-microglossae Diet. Chestnut-brown rust pustules on yellow or brown leaf spots on *S. microglossa* in Argentina.

Puccinia virgaureae (DC.) Lib. Black rust pustules on leaves of *S. microglossa* and *S. virgaurea* in Japan and Europe.

Septoria virgaureae Desm. On leaves of *S. virgaurea* in Yugoslavia, Italy, and Switzerland.

Uromyces komarovii Bub. Brown rust pustules on leaves of *S. virgaurea* in Japan and Manchuria.

SONCHUS. Sow THISTLE. Composites, mostly weeds, some species used for foliage effects.

Bremia sonchi K. Saw. Downy mildew causing small polygonal spots on stems and leaves of *S. oleraceus* in Formosa.

Coleosporium sonchi Lévl. Leaf rust on *S. asper*, *S. oleraceus*, *S. palustris*, and *S. uliginosus* in Europe. The aecial stage occurs on *Pinus sylvestris*. Introduced into Wisconsin and found locally on *S. asper* and *Pinus sylvestris*.

Phyllosticta sonchi Sacc. Gray-white leaf spots on *S. oleraceus* in Russia and Italy.

Puccinia sonchi Rob. Yellow to black rust pustules on leaves of *Sonchus* spp. in Ceylon, Japan, Algeria, Canary Islands, and Europe.

Puccinia sonchina Syd. Brown rust pustules on leaves of *S. oleraceus* in Japan and Portugal.

Ramularia sonchi-arvensis Fautr. On leaves of *S. oleraceus* in Spain and France.

Septoria modonia Sacc. Small subcircular spots on leaves and stems of *S. arvensis* in France.

Septoria sonchi Sacc. Oblong greenish-gray leaf spots on *S. oleraceus* in Italy.

Septoria sonchicola Höll. Circular ochraceous leaf spots with black borders on *S. uliginosus* in Hungary.

Septoria sonchina Thum. Leaf spots on *S. oleraceus* in Siberia.

Synchytrium globosum Schroet. See *Potentilla*.

SOPHRONITIS. See *Orchidaceae*.**SOPHORA.** Shrubs or herbs grown for their attractive flowers and handsome foliage.

Aecidium sophorae Kus. Leaf rust on *S. platycarpa* in Japan.

Ascochyta sophorae Allesch. On branches of *S. japonica* in Germany.

Macrosporium sophorae Turconi. Circular brown leaf spots on *S. japonica* in Italy.

Phyllosticta sophoricola Höll. Irregular ashen leaf spots with dull-brown borders on *S. japonica* in Hungary.

Uromyces shikokianus Kus. Leaf rust on *S. shikokiana* in Japan.

Uromyces sophorae-flavescentis Kus. Brown leaf rust on *S. flavescentis* in Japan.

Uromyces sophorae-japonicae Diet. Brown rust pustules on leaf blades and petioles of *S. japonica* in Japan.

Uromyces trunicola P. Henn. and Shir. Dark-brown rust pustules on branches and trunks of *S. japonica* in Japan.

SORBARIA. SCHIZONOTUS. Ag. FALSE SPIREA. Ornamental woody plants related to *Spiraea*.

Leptosphaeria spiraeae Karst. On branches of *S. sorbifolia* in France.

Phyllosticta spiraeina Brun. See *Spiraea*.

SORBUS. MOUNTAIN ASH. Trees grown for their handsome foliage, white flowers and ornamental red fruit.

Cercospora ariae Fekl. On leaves of *S. aria* in Germany.

Cercospora kriegeriae Bres. Rufous-brown leaf spots on *S. aucuparia* in Germany.

Fusiculadum orbiculatum (Desm.) Thum. Brown scabby areas on leaves of *S. aria*, *S. aucuparia*, *S. domestica*, and *S. torminalis* in Europe.

Gloeosporium aucupariae P. Henn. Dark-brown anthracnose spots on fruit of *S. aucuparia* in Russia.

Gloeosporium sorbi Rostr. Numerous subcircular ashen leaf spots on *S. fennica* in Finland.

Gymnosporangium miyabei Yam. and Miy. See *Chamaecyparis*.

Gymnosporangium solenoides (Diet.) Kern. See *Chamaecyparis*.

Gymnosporangium terminali-juniperinum Ed. Fisch. See *Juniperus*.

Laestadia radiata (Walbr.) Sacc. On leaves of *S. torminalis* in Germany.

Leptosphaeria sorbi Jacc. Subcircular gray leaf spots on *S. aucuparia* in Siberia.

Marsonia sorbi P. Magn. Indefinite, yellow-brown leaf spots on *S. aria* in Europe.

Mycosphaerella cinerascens (Fekl.) Vgr. On leaves of *S. aria* and *S. scandica* in Denmark and Germany.

Mycosphaerella topographica (Sacc.) Vgr. On leaves of *S. aucuparia* and *S. torminalis* in Denmark and Italy.

Ochropsora ariae (Fekl.) Syd. Yellow leaf rust on *S. americana*, *S. amicana*, *S. aria*, *S. aucuparia*, *S. fennica*, *S. intermedia*, *S. latifolia*, *S. scandica*, *S. torminalis*, *Amelanchier canadensis*, *Aruncus sylvestris*, *Pyrus communis*, and *Malus* in Europe and Formosa. The aecial stage occurs on *Anemone nemorosa*.

Phyllobendersonia terminalis-ariae (Br. and Har.) Tass. On *S. aria* and *S. torminalis* in France and Italy.

Phyllosticta aucupariae Thum. Subcircular ashen leaf spots with dull-brown margins on *S. aucuparia* in Siberia, Russia, Italy, and Hungary.

Phyllosticta sorbi West. Subcircular, then confluent, ashen leaf spots with dark-purple margins on *S. aucuparia*, *S. domestica*, *S. lanuginosa*, and *S. scandica* in Belgium and Portugal. Reported from Missouri.

Sclerotinia aucupariae Woron. Brown rot of fruit, which finally are mummified. On *S. aucuparia* in Finland.

Septoria aucupariae Bres. Dull-yellow leaf spots on *S. aucuparia* in Denmark and Germany.

Septoria aucuparicola Oud. Circular to irregular, then confluent, dull-brown to ashen leaf spots on *S. aucuparia* in Holland.

Septoria hyalospora (Mont. and Ces.) Sacc. On leaves of *S. aria*, *S. domestica*, and *S. torminalis* in Italy and Germany.

Septoria inaequalis Sacc. and Roum. On *S. aucuparia* in France.

Septoria sorbi Lasch. Small circular brown spots on leaves of *S. aucuparia* in Austria.

Septoria sorbi-hybridae Pass. On leaves of *S. hybrida* in Italy.

SORGHUM. See *Helcus*.

- SPARAXIS.** WAND FLOWER. Iridaceous, bulbous or cormous plants.
Uromyces sparaxidis Syd. Yellow to dark-brown rust pustules on both leaf surfaces of *S. lineata* in Natal.
- SPARGANIUM.** BUR REED. Marsh or aquatic plants.
Ascochyta quadriguttulata Kab. and Bub. Irregular linear, then confluent, yellow-brown leaf spots on *S. ramosum* in Bohemia.
Cylindrosporium aquaticum (Fautr. and Roum.) Sacc. On leaves of *Sparganium* sp. in France.
Physoderma sparganii-ramosi (Buesg.) Schroet. On *S. ramosum* in Germany.
Ramularia frutescens Kab. and Bub. On leaves of *S. ramosum* in Bohemia.
Ramularia sparganii Lindr. Circular brown leaf spots on *S. glomeratum* and *S. simplex* in Sweden, Finland, Silesia, and Austria.
Septoria flexuosa Oud. On leaves of *S. ramosum* in Holland.
Septoria sparganii Pass. On leaves of *S. erectum* and *S. ramosum* in Italy.
- SPARTINA.** Perennial marsh grasses.
Phyllosticta spartinae Brun. Subcircular small white spots with brown margins on leaf sheaths of *S. stricta* in France.
Uredo spartinae-strictae Pat. and Har. Linear brown rust pustules on leaves of *S. stricta* in France.
- SPARTIUM.** WEAVERS'-BROOM. Ornamental hardy shrubs grown for their bright-yellow flowers.
Phylospora euganea Sacc. (*P. albanica* Petr.) On stems of *S. junceum* in Albania and Yugoslavia.
Septoria spartii Rob. and Desm. Circular olivaceous leaf spots on *S. junceum* in France.
Uromyces spartii-juncei Syd. Powdery brown rust sori on leaves of *S. junceum* in Dalmatia, France, and Portugal.
- SPATHOGLOTTIS.** See Orchidaceae.
- SPECULARIA.** VENUS'S LOOKING-GLASS. Erect or decumbent annual herbs.
Ramularia prismatocarpi Oud. Forms grayish areas on lower leaf surfaces of *S. speculum* in Dalmatia and Holland.
Septoria prismatocarpi Desm. On leaves of *S. hybrida* and *S. speculum* in Italy and France.
- SPERGULA.** SPURRY. Annual herbs.
Mycosphaerella alsines Pass. On stems and leaves of *S. campestris* in Denmark.
Peronospora lepigoni Fekl. Downy mildew on leaves of *S. campestris* and *S. marina* in north Europe.
Peronospora vernalis Gäum. Downy mildew on leaves of *S. pentandra* and *S. vernalis* in Germany.
Puccinia spergulae DC. Brown to dark-brown rust pustules on leaf blades and petioles and stems of *S. arvensis*, *S. pentandra*, *S. vernalis*, and *Spergularia rubra* in Europe. Reported from New York.
Septoria spergulae West. On leaves and stems of *S. arvensis* in Sweden and Belgium.
Uromyces sparsus (Kze. and Schm.) Lévl. Leaf and stem rust on *S. marginata*, *S. media*, *S. (Spergularia) rubra*, and *S. salina* in French North Africa and Europe.
- SPHAERALCEA.** PHYMOSIA Ag. GLOBE MALLOW. Shrubs and herbs.
Aecidium sphaeralcearum (Speg.) Sacc. and Troit. Leaf rust on *S. tonariensis* and *S. cisplatina* in Argentina.
Uromyces costesianus Speg. Leaf rust on *S. velutina* in Chile.
Uromyces platysporus Speg. Dark-brown leaf rust on *S. miniata* and *S. rhombifolia* in Argentina.
- SPIGELIA.** PINKROOT. Annual and perennial herbs.
Coleosporium spigeliae Arth. Yellow powdery rust pustules on leaves of *S. humboldtiana* in Salvador and Guatemala.
Puccinia spigeliae Syd. Leaf rust on *Spigelia* sp. in Brazil.
Septoria spigeliae P. Henn. Subcircular dull-brown leaf spots with dark-brown borders on *S. anthelmintica* in Jamaica and Brazil.
- SPILANTHES.** Herbs, mostly weeds, a few species cultivated for their yellow flower heads.
Entyloma spilanthi Speg. Smut sori on leaves of *S. leptophylla* in Argentina.
Puccinia africana Cke. Dark-brown rust sori on brown sunken spots on *S. acmella*, *S. africana*, and *S. oleracea* in Abyssinia, Madagascar, and tropical and south Africa.
Puccinia barranquillae Mayor. Leaf rust on *S. urens* in Colombia.
Puccinia melampodii Diet. and Holw. See Zinnia.
Puccinia spilanthicola Mayor. Leaf rust on *S. americana* and *S. ciliata* in Colombia.
Puccinia spilanthi P. Henn. Dark-brown rust sori on leaves of *S. ciliata*, *S. oleracea*, *S. salzmanni*, and *S. uliginosa* in Martinique, Brazil, Argentina, and Peru.
- SPINACIA.** SPINACH. Annual herbs, cultivated for "greens."
Aecidium spinaciae Rostr. Leaf rust on *S. tetrandra* in Russia.
Alternaria spinaciae Allesch. and Noack. On living leaves of *S. oleracea* in Brazil.
Ascochyta chenopodii Rostr. See Atriplex.
Cercospora spinaciae Oud. Pale greenish or yellowish, often confluent, leaf spots on *S. oleracea* in Holland.
Gloeosporium spinaciae Ell. and Fautr. Subcircular, then confluent, brown leaf spots on *S. oleracea* in France.
Phyllosticta spinaciae A. Zimm. Large irregular leaf spots on *S. oleracea* in Belgium and Austria.
Ramularia spinaciae Nypels. Brown leaf spots on *S. oleracea* in Belgium.
Septoria spinaciae West. Circular leaf spots on *S. oleracea* in France, Russia, and Belgium.
- SPIRAEA.** SPIREA. Bridal wreath. Shrubs grown for their profuse, handsome white or pink flowers.
Aphelenchus olesistus Ritz. Bos. See Begonia.
Cercospora spiraeae Thuem. Irregular, often confluent, dull-brown leaf spots on *S. chamaedryfolia* and *Opulaster opulifolius* (*S. opulifolia*) in Siberia.
Cercospora callosa Allesch. Subcircular brown, then ashen-white, leaf spots with brown margins on *S. japonica* (*S. callosa*) in Germany.
Cylindrosporium filipendulae Thuem. See Filipendula.
Excipula spiraeae Thuem. On leaves of *S. thalictroides* in Siberia.
Gloeosporium spiraeae Bres. Indefinite dull-brown leaf spots on *Opulaster opulifolius* (*S. opulifolia*) in Germany.
Heterosporium spiraeae Syd. Dull-brown irregular, then confluent, spots on leaves of *Spiraea* sp. in Austria.
Ochropsora ariae (Fekl.) Syd. See Sorbus.
Phyllosticta ariaefolia Allesch. Subcircular to irregular dull-brown leaf spots on *S. (Holodiscus) ariaefolia* in Germany.
Phyllosticta crenatae Brun. Oblong to angular leaf spots, black above, grayish-brown beneath, on *S. crenata* in France.
Phyllosticta spiraeae-salicifoliae Kab. and Bub. Circular to oblong dull-yellow to dark-brown leaf spots on *S. salicifolia* in Bohemia.

SPIRAEA—Continued.

Phyllosticta spiracina Brun. Subcircular brown leaf spots with red-brown margins on *S. bumalda* and *Sorbaria sorbifolia* in France and Germany.

Septoria ascochytoidea Sacc. Subcircular to oblong ochraceous to dull-brown leaf spots on *S. chamaedryfolia* and *S. decumbens* in Siberia.

Triphragmium filipendulae (Lasch.) Pass. See *Filipendula*.

SPIRANTHES. LADY'S-TRESSES. See *Orchidaceae*.**SPONDIAS**. YELLOW MOMBIN. JOBO. SPANISH PLUM. Tropical fruit trees.

Ascochyta spondiacearum A. L. Sm. White leaf spots with dull-brown margins on *S. mombin* in Angola.

Endothia havanensis Bruner. See *Eucalyptus*.

Fomes lamaensis Murr. See *Hevea*.

Kuehneola aliena Syd. and Butl. Leaf rust on *S. mangifera* in India and Ceylon.

Uredo spondiadis Petch. Leaf rust on *S. mangifera* in Ceylon.

SPOROBOLUS. DROPSEED. Grasses, a few species cultivated for forage.

Entyloma majus Har. and Pat. Smut sori on small gray leaf spots on *S. spicatus* in the Congo.

Phylachora sporoboli Pat. Elongate black stromata on *S. pungens* in Algeria and the Philippines.

Puccinia spegazziniella Sacc. and Trav. Dark-brown rust pustules on leaves of *S. asperifolius* in Argentina.

Tilletia zonata Bref. Black powdery smut masses in flowers of *S. ligularis* in Ecuador.

Uredo egenula Arth. Brown powdery leaf rust on *S. argutus* in Jamaica.

Uromyces major Arth. Brown leaf rust on *S. diander* and *S. indicus*, in India, Ceylon, and Trinidad.

Uromyces tenuiculis McAlp. Yellow to brown rust pustules on leaves and stems of *S. indicus* in Australia and Japan.

Ustilaginoides ochracea P. Henn. In spikes of *S. elongatum* in North Borneo.

Ustilago schlechteri P. Henn. Black smut sori distorting the panicles and splitting the stems of *Sporobolus* sp. in the Union of South Africa.

STACHYS. BETONY. Tall perennial herbs or diffuse winter annuals.

Ovularia betonicae Mass. On leaves of *S. alopecuroides* in Austria.

Ovularia stachydis Bres. On leaves of *S. palustris* in Germany.

Peronospora stachydis Syd. Downy mildew on leaves of *S. arvensis* in Europe.

Phyllosticta stachydis Brun. and var. **arvensis** Allesch. Angular brown leaf spots on *S. arvensis* and *S. sylvatica* in France and Germany.

Phyllosticta varicolor Bub. Irregular brown, then yellow-white, leaf spots on *S. germanica* in Hungary.

Puccinia aethiopica Kalchbr. and Cke. Brown leaf rust on *S. aethiopica* and *S. grandifolia* in the Union of South Africa.

Puccinia albida Diet. and Neg. Leaf rust on *S. grandidentata* and *S. mayorti* in Chile.

Puccinia harti Lagh. Yellow and dark-brown rust pustules on leaves of *S. setifera* and *S. spectabilis* in Persia and Anatolia.

Puccinia pallidissima Speg. Brown leaf rust on *S. arvensis* and *S. lindeni* in Argentina, Brazil, Ecuador, and Guatemala.

Puccinia stachydis DC. Brown rust pustules on leaves, stems, and calyxes of *S. recta* in Russia, France, Italy, Austria, and Germany.

Puccinia vossii Koern. Powdery brown rust pustules on leaves of *S. recta* and *S. setifera* in Europe.

Ramularia cardiaca Sacc. and Penz. Oblong to irregular spots on leaves and bracts of *S. annua* and *S. recta* in Italy and Switzerland.

Ramularia stachydis (Pass.) Mass. On leaves of *S. annua* and *S. recta* in Yugoslavia.

Ramularia stachydis-alpinae Allesch. Irregular ochraceous-brown leaf spots on *S. alpina* in Germany.

Septoria stachydis Rob. and Desm. Irregular olivaceous, then pale-brown, leaf spots on *S. annua*, *S. palustris*, and *S. sylvatica* in Europe.

Ustilago betonicae Berk. Smut in anthers of *A. alopecuroides* in Austria.

STACHYTARPHETA. **VALERIANOIDES** Ag. Tropical and subtropical herbs and shrubs.

Cercospora stachytarphetae E. and E. Small circular white leaf spots on *S. jamaicensis* in the Bahamas and Porto Rico.

Endophyllum stachytarphetae (P. Henn.) W. and O. Small yellow rust pustules on leaves of *S. cayennensis* and *S. dichotoma* in Porto Rico, Brazil, and Colombia.

Puccinia urbaniana P. Henn. Dark-brown rust pustules on small circular brown spots on *S. cayennensis* and *S. jamaicensis* in Porto Rico, Jamaica, and Trinidad.

STACHYRUS. Ornamental shrubs grown for their early flowers and handsome foliage.

Uredo stachyuri Diet. Brown leaf rust on *S. praecox* in Japan.

STANGERIA. A subtropical cycad.

Placosphaeria stangeriae (Zimm.) Bub. Large irregular brown leaf spots on *S. paradoxa* in Bohemia.

STANHOPEA. See *Orchidaceae*.**STAPHYLEA**. BLADDER NUT. Shrubs cultivated for their white flowers, handsome foliage, and inflated podlike fruit.

Ascochyta staphyleae Syd. Circular brown leaf spots with purple margins on *S. trifolia* in Germany.

Laestadia staphyleae Hazsl. Attacks young branches of *S. pinnata* in Hungary.

Phyllosticta osteospora Sacc. var. **staphyleae** C. Mass. Irregular brown leaf spots on *S. pinnata* in Italy and Austria.

Phyllosticta staphyleicola Oud. On leaves of *S. pinnata* in Holland.

Septoria staphyleae Pass. Rufous leaf spots on *S. pinnata* in Italy.

STATICE. THRIFT. Sometimes called "sea pink." Annuals, biennials, and perennials cultivated for their flowers and foliage. See also *Armeria*.

Ascochyta statidis P. Nag. On leaves of *Statice* sp. in Russia.

Phyllosticta statidis Petr. Large brown leaf spots, lighter at the center, and with red-brown margins on *Statice* sp. in Bohemia.

Ramularia statidis E. Rostr. Large circular to oblong ochraceous leaf spots with dull-brown margins on *S. (Limonium) behusensis* and *S. gmelini* (*Limonium gmelini*) in Russia and Norway.

Uromyces guayaurei Speg. Leaf rust on *S. brasiliensis* (*Limonium brasiliense*) in Argentina.

STELLARIA. **ALSINE** Ag. Annual or perennial herbs, mostly weeds.

Aecidium stellariae Kirch. Leaf rust on *S. graminea* in Bohemia.

Ascochyta stellariae Fautr. On leaves of *S. graminea* in France.

Fabraea cerastiorum (Fr.) Rehm. See *Cerastium*.

Isariopsis alborosella (Desm.) Sacc. On leaves of *S. aquatica*, *S. graminea*, *S. media*, *S. nemorum*, *Cerastium pumilum*, and *C. vulgatum* in Japan and Europe.

STELLARIA—Continued.

- Isariopsis stellariae** Trail. Indefinite brown spots on *S. graminea* in Scotland.
- Mycosphaerella isariphora** (Desm.) Johan. On leaves of *Stellaria* spp. in Siberia and Europe.
- Omphalospora stellariae** (Lib.) Theiss. and Syd. Shiny black stromata on leaves and stems of *S. holostea*, *S. moehringia*, and *S. nemorum* in Siberia and Europe.
- Ovularia stellariae** (Rabh.) Sacc. On leaves of *S. nemorum* in Russia, France, Switzerland, Yugoslavia, and Germany.
- Peronospora media** Gäum. Downy mildew on leaves of *S. media* and *S. nemorum* in Europe and Japan.
- Peronospora parva** Gäum. Downy mildew on leaves of *S. graminea*, *S. holostea*, and *S. uliginosa* in Europe.
- Phyllosticta holostea** Allesch. On leaves of *S. holostea* and *S. uliginosa* in Great Britain and Germany.
- Phyllosticta holosteicola** Oud. Gray leaf spots on *S. holostea* in Holland.
- Placosphaeria stellariae** (Lib.) Sacc. On leaves and stems of *S. holostea* and *Dianthus monspessulanus* in Denmark, France, Italy, and Germany.
- Puccinia detonsa** Arth. and Holw. Leaf rust on *S. ovata* in Costa Rica and Guatemala.
- Ramularia stellariae** Rab. On leaves of *S. nemorum* in France.
- Septoria henslowiana** Sacc. Indefinite pale-ochraceous leaf spots on *S. media* in Malta.
- Septoria paraphysoides** Speg. On leaves and stems of *S. debilis* in Chile.
- Synechitrium stellariae** Fckl. Galls on leaves and stems of *S. media* and *S. nemorum* in Brazil, Argentina, Dalmatia, Denmark, and Germany. Reported from Mississippi.
- Uromyces stellariae** Syd. Dark-brown leaf rust on *S. kotschyana* in Persia.
- Ustilago duriaana** Tul. See *Cerastium*.
- STENANDRUM. GERARDIA** Ag. Stemless or short-stemmed acanthaceous herbs.
- Aecidium cystopoides** Speg. Leaf rust on *S. dulce* in Argentina.
- Puccinia stenandrii** Diet. and Neg. Golden and brown rust pustules on leaves of *S. dulce* in Chile.
- Uredo stenandrii** Speg. Leaf rust on *S. trinerve* in Argentina.
- STENOCARPUS. CYBELE** Ag. Tropical trees.
- Phyllosticta stenocarpi** Tassi. Irregular brown leaf spots on *S. sinuatus* in Italy.
- STENORRHYNCHUS.** See *Orchidaceae*.
- STEPHANIA.** Climbing shrubs.
- Trabutia stephaniae** Rac. Circular shiny black stromata on leaves of *S. capitata* in Java.
- STEPHANOTIS.** Twinning, glabrous tropical shrubs.
- Phyllosticta stephanotidis** Grove. Circular to irregular gray-white leaf spots with dark-brown margins on *S. floribunda* in Great Britain.
- STERCULIA.** BOTTLE TREE. Ornamental and economic trees.
- Ascochyta sterculiae** Tassi. Angular to irregular gray-white to dull-brown leaf spots on *S. diversifolia* (*Brachychiton diversifolium*) in Italy.
- Dothidasteroma maculosum** (B. and Br.) v. Hoeh. Circular brown stromata on leaves of *S. alata* (*Pterygata alata*) in Ceylon.
- Gloeosporium rhodospermum** Delac. Brown leaf spots on *S. acuminata* in central Africa.
- Guignardia sterculiae** Rehm. On *S. foetida* in the Philippines.
- Phyllosticta sterculiae** Wint. Angular to irregular ochraceous leaf spots on *S. acrifolia* (*Brachychiton acrifolium*) and *S. heterophylla* in Queensland and Portugal.
- Phyllosticta sterculicola** Trav. Large gray-white leaf spots on *S. frondosa* in Italy.
- Rosellinia bunodes** B. and Br. See *Citrus*.
- STEREOSPERMUM.** Tropical evergreen trees with large begonialike flowers.
- Phakopsora erythraea** Sacc. Black leaf rust on *S. dentatum* in Abyssinia.
- Uredo stereospermi** Syd. Leaf rust on *S. chelonioides* in Ceylon.
- STEVIA.** Tropical shrubs.
- Aecidium steviae** P. Henn. Leaf rust on *S. urticifolia* in Brazil.
- Coelosporium reichel** Diet. Leaf rust on *Stevia* sp. in Mexico.
- Coelosporium steviae** Arth. Golden rust pustules on leaves of *S. lucida*, *S. monardaefolia*, *S. reglensis*, *S. rhombifolia*, *S. salicifolia*, *S. subpubescens*, *S. trachelioides*, and *S. viscida* in Mexico and Guatemala.
- Puccinia rosea** Roth. See *Ageratum*.
- STIGMAPHYLLON.** AMAZON VINE. Tropical woody vines.
- Mycosphaerella stigmaphylli** Rangel. On leaves of *S. ciliatum* in Brazil.
- Phyllosticta stigmaphylli** Speg. Circular brown leaf spots on *S. littorale* in Argentina.
- Puccinia inflata** Arth. Brown leaf rust on *S. lingulatum*, *S. periplocifolium*, and *S. sagraeanum* in Porto Rico and Cuba.
- Puccinia insueta** Wint. Brown to black rust pustules on leaves of *S. jatrophaeifolium* and *S. littorale* in Brazil, Uruguay, and Ecuador.
- STILLINGIA.** CHINESE TALLOW TREE. Tropical American shrubs and herbs.
- Cercospora sebiferae** Pat. On leaves of *S. sebifera* (*Sapium sebiferum*) in Indo-China.
- STIPA.** FEATHER GRASS. Perennial grasses.
- Puccinia flavescens** McAlp. Brown to black rust pustules on leaves of *S. flavescens*, *S. pubescens*, and *S. scabra* in Australia.
- Puccinia hierochloae** S. Ito. See *Hierochloa*.
- Puccinia oligocarpa** Syd. and Butl. Leaf rust on *Stipa* sp. in India.
- Puccinia stipae-sibiricae** S. Ito. Leaf rust on *S. sibirica* in Japan.
- Puccinia volgensis** Nawasch. Powdery brown rust pustules on leaves of *S. barbata* and *S. pennata* in Russia.
- Septoria stipae** Died. On leaves of *S. capillata* in Germany.
- Sorosporium tumefaciens** McAlp. Olivaceous to dark-brown smut sori involving entire panicles of *S. pubescens* and *Stipa* sp. in Australia.
- Sphaelotheca macrochloae** (Pat.) Maire. Smut on *S. tenacissima* in Tripoli.
- Tilletia hypsophila** Speg. Powdery brown masses of smut spores in ovaries of *S. caespitosa* and *S. tenuissima* in Argentina.
- Uredo pencana** Diet. and Neg. Brown leaf rust on *S. manicata* in Chile.
- Urocystis stipae** McAlp. Smut sori as elongated dark-colored streaks on leaves, sheaths, and stems of *S. luehmanni* in Australia.
- Uromyces argentinus** Speg. Yellow to cinnamon-brown rust pustules on leaves of *S. neesiana* in Argentina.
- Uromyces mussooriensis** Syd. Brown leaf rust on *S. sibirica* in India.
- Ustilago comburens** Ludw. Powdery black smut sori in culms and inflorescences of *Stipa* sp. in Australia.
- Ustilago macrochloae** Pat. Inflorescences of *S. tenacissima* in Algeria converted into masses of smut spores.
- Ustilago nummularia** Speg. Black smut sori in inflorescences of *Stipa* sp. in Argentina.

STIPA—Continued.

Ustilago stipae-barbatae Maire. Smut on *S. barbata* and *S. gigantea* in French North Africa.

Ustilago stiparum Speg. Oligaceous masses of smut spores in panicles of *Stipa* sp. in Chile.

Ustilago stipicola Speg. Powdery black smut sori in panicles of *S. filiculmis* and *S. setigera* in Argentina and Uruguay.

STIZOLEBIUM. See *Mucuna*.**STOBÆA**. Shrubs or herbs.

Puccinia stobæae McO. and var. *woodii* Syd. Yellow and dark-brown rust pustules on leaves of *S. membranifolia* and *S. speciosa* in the Union of South Africa.

STRANVAESIA. Shrubs grown for their handsome foliage and attractive fruit and flowers.

Acidium stranyaesiae Syd. Leaf rust on *S. glaucescens* in India.

STRATIOTES. WATER SOLDIER. Perennial aquatic herbs.

Leptosphaeria stratiotis Oud. On leaves of *S. aloides* in Holland.

Phyllosticta aloidis Oud. On leaves of *S. aloides* in Holland.

Phyllosticta stratiotis Oud. Dull-brown leaf spots on *S. aloides* in Holland.

Septoria stratiotis Oud. On leaves of *S. aloides* in Holland.

STRELITZIA. BIRD-OF-PARADISE FLOWER. Perennial herbs.

Endodothella strelitziae (Cke.) Theiss. and Syd. Circular dull-black stromata on leaf spots which are wine-red beneath, on *S. regina* in the Union of South Africa.

STROBILANTHES. CONEHEAD. Acanthaceous ornamental herbs and shrubs.

Acidium strobilanthæ Barch. Leaf rust on *Strobilanthes* sp. in the Philippines and India.

Corticium salmonicolor B. and Br. See Citrus.

Puccinia aggregata Syd. Brown rust pustules on sunken brown leaf spots on *S. barbatus* in India.

Puccinia polliniae Barch. Yellow and brown to black rust pustules on leaf blades and petioles and stems of *S. cuspidatus*, *S. dalhousianus*, and *Pollinia nuda* in India and Indo-China.

STROPHANTHUS. Tropical shrubs.

Hemileia strophanthi Rac. Yellow rust pustules on *S. dichotomus* and *S. hispidus* in Gold Coast Colony.

STRYCHNOS. Shrubs and trees of economic importance, cultivated for their fruit and medicinal properties.

Meliola steniospora Hint. Subcircular black superficial fungus patches on leaves of *S. nux-vomica* in India.

Phyllosticta strychni Allesch. Subcircular ochraceous leaf spots with dark-brown margins on *S. stuhlmanni* in tropical Africa.

Puccinia lindaviana P. Henn. Dark-brown rust pustules on leaves of *S. henningsii* in tropical Africa and the Union of South Africa.

Uredo strychni P. Henn. Yellow leaf rust on *Strychnos* sp. in the Congo.

STYLIIDUM. CANDOLLEA Ag. Herbs or subshrubs.

Puccinia styliidi McAlp. Orange-yellow and dark-brown to black rust pustules on leaves of *S. graminifolium* in Tasmania.

STYRAX. SNOWBELL. Ornamental woody plants grown for their flowers.

Atichia millardetii Racib. See Cinnamomum.

Monopus pulverulentus (B. and C.) Theiss. and Syd. Gray-black stromata on leaves of *Styrax* sp. in Cuba.

Pucciniastrum stryacinum Hirats. Leaf rust on *S. japonica* and *S. obassia* in Japan.

Schneepia guaranitica Speg. On *S. leprosum* and *S. parvifolia* in Brazil.

SUTHERLANDIA. Tropical shrubs grown for their flowers.

Stigmatula sutherlandiae (K. and Cke.) Syd. On leaves of *Sutherlandia* sp. in the Union of South Africa.

SWERTIA. Erect perennial and annual herbs grown in borders and rock gardens.

Albugo swertiae Berl. and Kom. Small white powdery pustules on leaves of *S. connata* in Manchuria and Russia.

Septoria swertiae Pat. Circular rufous leaf spots on *Swertia* sp. in China.

Uredo ophellae Syd. Brown rust on leaves and stems of *S. angustifolia* in India.

SWIETENIA. MAHOGANY. Tropical timber trees.

Tylenchus mahogani Cobb. This nematode is found in the bark tissues of *S. mahogani* in Barbados, probably parasitic and capable of causing injury.

SYMPHORICARPOS. SNOWBERRY. Ornamental shrubs.

Ascochyta grandispora Kab. and Bub. Subcircular ochraceous to brown leaf spots on *S. orbiculatus* in Bohemia.

Ascochyta symphoricae Kab. and Bub. Circular to irregular spots on leaves of *Symphoricarpos* sp. in Bohemia and France. *A. symphoricae* Br. and Har. has priority.

Ascochyta symphoricarpi Pass. On branches of *S. albus* (*S. racemosus*) in Italy and France.

Heterosporium symphoricarpi Ranoj. On branches of *S. albus* (*S. racemosus*) in Yugoslavia.

Phyllosticta symphoricae Sacc. and March. Leaf spots on *S. cinerascens* in Belgium.

SYMPHYSTEMUM. Bulbous plants.

Uromyces symphyostemi Speg. Brown linear rust pustules on leaves of *S. narcissoides* in Patagonia.

SYMPHYTUM. COMFREY. Erect, often hispid, herbs.

Acidium asperifolii Pers. Leaf rust on *S. tuberosum* in Austria.

Acidium symphyti Thuem. Leaf rust on *S. tuberosum* in Spain and Dalmatia.

Cercospora agnostoica Speg. Subcircular pale-brown leaf spots on *S. asperum* in Brazil.

Cylindrosporium myosotidis Sacc. See Myosotis.

Eutyloma serotinum Schroet. See Borago.

Melampsorella symphyti Bub. Powdery brown rust pustules on leaves of *S. bulbosum*, *S. cordatum*, *S. ibericum*, *S. officinale*, *S. tauricum*, and *S. tuberosum* in Europe. The aecial stage occurs on the needles of *Abies pectinata*.

Ovularia asperifolii Sacc. and var. *symphyti-tuberosi* Allesch. On leaves of *S. officinale* and *S. tuberosum* in Germany.

Ovularia farinosa (Bon.) Sacc. On leaves of *S. officinale* in Germany and Russia.

Peronospora symphyti Gäum. Downy mildew on leaves of *S. bulbosum*, *S. cordatum*, and *S. tuberosum* in southern Europe.

Phyoderma speciosum Schroet. Brown areas on leaf blades and petioles and stems of *S. officinale* in Silesia and Germany.

Puccinia bromina Eriks. See Bromus.

Ramularia symphyti-tuberosi (Allesch.) Jaap. On leaves of *S. tuberosum* in Dalmatia. Possibly the same as *Ovularia farinosa*.

- SYMPLOCOS.** SWEETLEAF. Shrubs grown chiefly for their attractive flowers and fruits.
- Coccenia placenta** (B. and Br.) Sacc. Dull-black stromata on brown leaf spots on *S. spicata* and *Panella indica* in Ceylon.
- Exobasidium indicum** Syd. and Butl. Sordid white galls on leaves of *S. theaeifolia* and *Symplocos* sp. in India and Ceylon.
- Exobasidium symptloci-fasciculatae** Rac. Circular to ovate spots on *S. fasciculata* in Java.
- Exobasidium symptloci-japonicae** Kus. Distorts young buds of *S. japonica*, *S. lucida*, and *Maba buxifolia* in Japan.
- Fomes pseudoferreus** Wakef. See Hevea.
- Mycosphaerella bhouria** (Cke.) Lind. On leaves of *S. spicata* in India and Malaya.
- Phacidium symptlocinum** Syd. On leaves of *Symplocos* sp. in India.
- Phyllachora symptloci** Pat. Black stromata on leaves of *S. japonica*, *S. lucida*, *S. neritifolia*, and *S. thwaitesii* in Indo-China, Japan, and Australia.
- Physalospora symptloci** Rac. Circular to irregular yellow leaf spots on *S. fasciculata* in Java.
- Rhytisma austro-caledonicum** Crie. Black irregular stromata on leaves of *S. arborea*, *S. caerulea*, *S. gracilis*, *S. lenormandiana*, *S. mandiana*, and *S. vielillardii* in New Caledonia.
- Rosellinia arcuata** Petch. See Thea.
- Septoria sydowii** Henn. and Sacc. Small subcircular to irregular gray-white leaf spots on *S. crataegoides*, *S. myrtacea*, and *S. prunifolia* in Japan and Brazil.
- Septoria symptloci** Allesch. and P. Henn. On *Symplocos* sp. in Brazil.
- Trochila symptloci** P. Henn. On leaves of *S. japonica* in Japan.
- Uredo inouyei** P. Henn. and Shir. A rust deforming leaves and branches of *Symplocos* sp. in Japan.
- SYRINGA.** LILAC. Shrubs cultivated for their flowers.
- Ascochyta orientalis** Bond. Confluent gray leaf spots with dark-brown margins on *S. vulgaris* in Russia.
- Ascochyta syringae** Bres. On leaves of *S. chinensis* and *S. vulgaris* in Russia, Denmark, Italy, and Germany.
- Ascochyta syringicola** Bub. and Kab. Circular to angular dull-yellow leaf spots on *S. vulgaris* in Bohemia.
- Ascochyta syringae** Jaap. On fruit capsules of *S. vulgaris* in Germany.
- Gloeosporium syringae** Allesch. Irregular ochraceous to brown leaf spots, often marginal, or involving entire leaf blades on *S. alba* and *S. vulgaris* in Russia and Germany.
- Helminthosporium syringae** Kleb. Irregular brown areas on leaves, gradually involving entire area of the blades on *S. vulgaris* in Great Britain and Germany.
- Heterosporium syringae** Oud. Large irregular gray-brown blotches on leaves of *S. vulgaris* in Spain, Italy, France, Holland, and Germany.
- Phyllosticta superflua** Oud. Circular, then confluent, pale ashen-brown leaf spots on *S. vulgaris* in Holland.
- Phyllosticta syringophila** Oud. On leaves of *S. vulgaris* in Holland.
- Phytophthora syringae** Kleb. This downy mildew causes a leaf-blight and die-back of *S. persica*, *S. vulgaris*, and other woody plants in Europe, including *Acer* sp., *Aesculus* sp., *Alnus* sp., *Corylus* sp., *Crataegus oxyacantha*, *Forsythia viridissima*, *Jasminum nudiflorum*, *Ligustrum vulgare*, *Malus*, *Prunus cerasus*, *P. domestica*, *Pyrus communis*, and *Tilia* sp.
- Pseudomonas syringae** v. Hall. Large black blotches on young leaves and twigs of *S. vulgaris* in Holland, Great Britain, and Germany. The organism is said to be able to attack also *Malus*, *Populus*, *Prunus*, and *Pyrus*.
- Septoria syringae** Sacc. and Speg. Dull-yellow leaf spots on *S. chinensis* and *S. vulgaris* in Italy and France.
- SYZGIUM.** Tropical trees and shrubs.
- Atichia millardetii** Racib. See Cinnamomum.
- Lizoniella syzgi** (Rac.) Sacc. and Trott. Subcircular black leaf spots on *Syzgium* sp. in Java.
- TABEBUIA.** Ornamental evergreen trees grown for their showy flowers.
- Mycosphaerella tabebuiae** Miles. Amphigenous small white transparent leaf spots with brown-purple margins on *T. haemantha* in Porto Rico.
- Phyllachora sordida** Speg. Black subcircular stromata on pale, indefinite leaf spots on *Tabebuia* sp. in Brazil.
- Phyllachora tabebuiae** (Rehm.) Theiss and Syd. Shiny black stromata on small dark-brown leaf spots on *T. leucoxydon* in Brazil.
- Physalospora tabebuiae** Rehm. On leaves of *T. leucoxydon* in Brazil.
- Prospodium bahamense** Arth. Leaf rust on *T. bahamensis* and *T. leucoxydon* in the Bahamas.
- Prospodium suppressum** Arth. Leaf rust on *Tabebuia* sp. in Trinidad.
- TABERNAEMONTANA.** CRAPE JASMINE. Evergreen trees and shrubs.
- Aecidium ceraceum** B. and Br. Leaf rust on *T. dichotoma* in Ceylon, Brazil, and Argentina.
- Aecidium ochraceum** Speg. Indefinite pale-ochraceous leaf spots on *T. australis* in Paraguay.
- Caecoma apocyni** McAlp. Brownish-yellow rust pustules on leaves of *T. orientalis* in Australia.
- Cercospora tabernaemontanae** Syd. On leaves of *T. pandocahui* and *Tabernaemontana* sp. in Malaya and the Philippines.
- Fomes pectinatus** (Kl.) Cke. Attacks *T. sphaerocarpa* in Java.
- Gloeosporium holstii** P. Henn. Anthracnose on leaves of *T. dichotoma* in Ceylon.
- Hemileia jurensis** Syd. Pale-yellow leaf rust on *Tabernaemontana* sp. in Brazil.
- Puccinia cookei** DeT. Yellow to brown rust pustules on leaves of *T. ventricosa* in India and the Union of South Africa.
- Puccinia engleriana** P. Henn. Leaf rust on *T. heyneana* and *T. pandocahui* in India and the Philippines.
- Puccinia tabernaemontanae** B. and Br. Leaf rust on *T. dichotoma* in Ceylon.
- Seynesia fusco-paraphysata** P. Henn. On leaves of *Tabernaemontana* sp. in Tanganyika.
- Uredo manilensis** Syd. Brown leaf rust on *T. coronaria* and *T. polygama* in the Philippines.
- TACCA.** Perennial herbs with tuberous or creeping rhizomes.
- Cercospora** sp. On leaves of *T. pinnatifida* in Fiji.
- Cercosporina taccae** Syd. Circular brown leaf spots on *T. palmata* in the Philippines.
- TACSONIA.** Tendril-climbing vines resembling *Passiflora*.
- Seynesia lagerheimii** Rehm. On leaves of *Tacsonia* sp. in Ecuador.
- TAGETES.** MARIGOLD. Garden annuals.
- Puccinia tagetica** Diet. and Holw. Powdery brown to black rust pustules on leaves of *T. erecta*, *T. filifolia*, *T. lucida*, *T. micrantha*, *T. patula*, *T. microglossa*, and *T. tenuifolia* in Porto Rico, Mexico, Costa Rica, and northern South America.
- TAINIA.** See *Orchidaceae*.

- TAMARINDUS.** TAMARIND. Tropical leguminous tree. Cultivated for ornament and fruit.
Exosporium tamarindi Syd. On leaves of *T. indica* in India.
Gloeosporium tamarindi P. Henn. Anthracnose on leaves of *T. indica* in Tanganyika.
Mycosphaerella tamarindi P. Henn. Dull-brown leaf spots on *T. indica* in Tanganyika.
- FAMARIX.** Ornamental shrubs or trees, cultivated for their graceful foliage and showy clusters of white or pink flowers.
Pyrenopeziza tamaricis (Roum.) Sacc. See Myricaria.
- TAMONEA.** Tropical shrubs.
Gloeosporium cyanophylli P. Henn. Dull-brown leaf spots on *T. (Cyanophyllum) magnifica* in Germany.
- TAMUS.** Black bryony. Herbaceous perennials with tuberous roots related to Dioscorea.
Cercospora scandens Sacc. and Wint. and var. *longissima* Gz. Frag. Small subcircular dull-brown leaf spots with yellow margins on *T. communis* in Italy, Spain, and Russia.
Cercospora tamicola Lamb. and Fautr. Irregular gray leaf spots on *T. communis* in Italy and France.
Phyllosticta tami Sacc. Subcircular to oblong brown leaf spots with red margins on *T. communis* in Italy.
Septoria sarmeticia Sacc. On stems of *T. communis* in France.
Septoria tami West. Subcircular to angular brown leaf spots with purple-brown margins on *T. communis* in Belgium.
- TANACETUM.** TANSY. Annual or perennial herbs with small yellow flower heads.
Peronospora tanacetii Güm. Downy mildew on leaves of *T. vulgare* in Europe.
Puccinia balsamitae (Strauss.) Rabh. See Pyrethrum.
Puccinia seriata Syd. Black rust sori on leaves and stems of *Tanacetum* sp. in Kurdistan.
Puccinia tanacetii DC. Dark-brown to black rust pustules on leaves of *T. corymbosum* and *T. vulgare* in Europe.
Ramularia tanacetii Lind. Brown spots, occupying most of leaf area of *T. vulgare* in Denmark and Spain.
Septoria tanacetii Niessl. Indefinite irregular often confluent, dull-brown leaf spots on *T. vulgare* in Bulgaria, Bohemia, and France.
Septoria tanacetii-macrophylli Bub. Elongate or irregular, often confluent, dull-yellow to brown leaf spots on *T. macrophyllum* in Central Europe.
- TARAXACUM.** LEONTODON. Ag. DANDELION. Weedy perennials, sometimes cultivated as a vegetable crop.
Cercospora augustana Ferr. Irregular ochraceous leaf spots on *T. officinale* in Italy.
Erysiphe taurica Lévy. See Althaea.
Phyllosticta taraxaci Höll. Irregular ashen black-bordered leaf spots on *T. officinale* in Hungary.
Protomyces pachydermus Thuem. On peduncles and leaves of *T. officinale* in Japan, Scandinavia, Italy, Switzerland, Austria, and Germany.
Puccinia decipiens Mass. Powdery black rust pustules on leaves of *T. canescens* and *T. montanum* in Persia and Kurdistan.
Puccinia kurdistanii Cke. Brown leaf rust on *T. glaucum* in Kurdistan.
Puccinia silvatica Schroet. Leaf rust on *T. dens-leonis*, *T. officinale*, *Crepis biennis*, *Lappa officinalis*, *Senecio juchsi*, *S. nemorensis*, *S. sarraenicum*, and *Carex* spp. in Siberia and Russia.
Puccinia taraxaci-bithynici R. Maire. Brown leaf rust on *T. bithynicum* in Asia Minor.
Puccinia variabilis Grev. Yellow to dark-brown rust pustules on leaves of *T. collinum*, *T. officinale* and *T. palustre* in Siberia and Europe.
Ramularia lampsana (Desm.) Sacc. See Cichorium.
Septoria taraxaci Höll. Circular ashen, black-bordered leaf spots on *T. officinale* in Hungary.
 Sydow has given the same name to a species on *T. officinale* in Japan.
- TAXUS.** YEW. Ornamental coniferous trees.
Gloeosporium eavarae (Mont.) Sacc. and Syd. On needles of *T. baccata* in Italy.
Gloeosporium taxicolum Allesch. On needles of *T. baccata* in Denmark, France, and Germany.
Lophodermium abietis Rostr. See Picea.
Macrophoma taxi Berl. and Vogl. Causes leaves of *T. baccata* in Europe to turn brown.
Phyllosticta taxi Höll. Ochraceous spots on needles of *T. baccata* in Hungary.
- TECOMA.** TRUMPET BUSH. Ornamental tropical shrubs and small trees grown for their showy flowers.
 See also Tabebuia.
Aecidium simplicius Arth. and Johnst. Leaf rust on *T. pentaphylla* in Cuba.
Cercospora leprosa Speg. On leaves of *T. arborea* in Brazil.
Cercospora stenolobicola Speg. Subcircular definite dull-rufous leaf spots on *T. (Stenolobium) stans* in Argentina.
Cylindrosporium aureum Speg. Indefinite dull-brown leaf spots on *Tecoma* sp. in Brazil.
Munkia guaranitica Speg. On leaves of *Tecoma* sp. in Brazil and Paraguay.
Ovalariopsis obelavata Wakef. On leaves of *T. leucozyllon* in Barbados.
Phyllosticta erysipoides Sacc. On leaves of *T. (Bignonia) radicans* in France.
Phyllosticta henriquesii Thuem. On leaves of *T. (Bignonia) radicans* in Italy.
Phyllosticta tecomae Sacc. Dull-brown to ochraceous leaf spots on *T. (Bignonia) radicans* in Italy and Portugal.
Prospodium appendiculatum (Wint.) Arth. Leaf rust on *T. mollis* and *T. (Stenolobium) stans* in Porto Rico, Cuba, Costa Rica, Mexico, and Trinidad.
Prospodium bahamensis Arth. See Tabebuia.
Prospodium plagiopus (Mont.) Arth. Brown rust pustules on leaves of *T. lepidota* in Cuba.
Puccinia extiosa Syd. and Holw. Brown leaf rust on *T. mollis* in Mexico.
Puccinia maligna Diet. Leaf rust on *T. stans* in Mexico.
Uredo longicauleata P. Henn. Powdery dull-brown to ochraceous rust pustules on leaves of *Tecoma* sp. in Brazil.
- TECOPHILAEAE.** CHILIAN CROCUS. Bulbous plants.
Puccinia roseanae Arth. Leaf rust on *T. roseana* in Peru.
- TECTARIA.** Ferns.
Uredo gymnogrammes P. Henn. See Dryopteris.
- TECTONA.** TEAK. Valuable timber trees.
Aecidium effusum Niessl. Leaf rust on *T. grandis* in India.
 Heart-rot. A heart-rot of *T. grandis* due to an undetermined fungus is reported from Java. The heartwood is stained dark brown and finally rots. The tops of diseased trees die.
Uncinula tectonae Salm. Powdery mildew on upper leaf surfaces of *T. grandis* in India.
Uredo tectonae Raeb. Brown leaf rust on *T. grandis* in Java, Indo-China, Ceylon, and India.
- TELLIMA.** FALSE ALUMROOT. Erect, hardy, herbaceous perennials.
Phyllosticta tellimae Tassi. Subcircular, brown leaf spots on *T. grandiflora* in Italy.

- TEPHROSIA. CRACCA** Ag. HOARY PEA. Perennial herbs or shrubs, some species cultivated for forage.
- Corticium salmonicolor** B. and Br. See Citrus.
- Fomes pseudoferreus** Wakef. See Hevea.
- Ophiodothella edax** (Br. and B.) v. Hoeh. Shiny black stromata on leaves of *T. suberosa* in Ceylon and the Union of South Africa.
- Poria hypobrunnea** Petch. See Hevea.
- Ravenelia caulicola** Arth. Brown to black powdery rust pustules on stems of *T. cinerea* in the Bahamas.
- Ravenelia irregularis** Arth. Brown rust pustules on leaves of *T. macrantha* in Mexico.
- Ravenelia stictica** B. and Br. Yellow to brown rust pustules on leaves of *T. suberosa* in India, Ceylon, and the Union of South Africa.
- Ravenelia talpa** (Long) Arth. Brown leaf rust on *T. talpa* in Mexico.
- Ravenelia tephrosiae** Kalchbr. Yellow-brown to dark-brown rust pustules on leaves of *T. macropoda* in the Union of South Africa.
- Rosellinia arcuata** Petch. See Thea.
- Uredo tephrosiae** Rabb. Leaf rust on *T. purpurea* in Ceylon.
- Uredo tephrosicola** P. Henn. Brown leaf rust on *T. toxicaria* and *Tephrosia* sp. in Peru and Brazil.
- TERAMNUS.** Tropical trees.
- Cercospora maricaoensis** Young. Diffuse, brown leaf spots on *T. uncinatus* in Porto Rico.
- Physopella concors** Arth. See Phaseolus.
- Uredo teramni** Mayor. Leaf rust on *T. uncinatus* in Colombia.
- Uromyces cologaniae** Arth. Brown leaf rust on *T. uncinatus* in Porto Rico.
- Uromyces vanderystii** P. Henn. Yellow-brown leaf rust on *T. labialis* in Congo.
- TERMINALIA.** INDIA ALMOND. ALMENDRO. SPANISH ALMOND. Tropical trees with many economic uses.
- Cercospora catappae** P. Henn. Circular dark-brown leaf spots on *T. catappa* in Zanzibar and the Philippines.
- Fomes lamaoensis** Murr. See Hevea.
- Gnomoniella catappae** Koord. Large indefinite angular dull-brown leaf spots on *T. catappa* in Java.
- Phyllosticta terminaliae** P. Henn. Circular brown to olivaceous leaf spots with red margins on *T. baumii* and *T. catappa* in Brazil and the Union of South Africa.
- Ramularia catappae** Rac. Circular ashen leaf spots with dull-red margins on *T. catappa* in the Philippines.
- Uredo terminaliae** P. Henn. Ochraceous rust pustules on leaves of *T. argentea* in Brazil.
- TETRAGONIA.** NEW ZEALAND SPINACH. Herbs cultivated as vegetables.
- Cercosporina tetragoniae** Speg. Subcircular leaf spots on *T. expansa* in Argentina and Russia.
- Puccinia tetragoniae** McAlp. Yellow to brown rust pustules on leaves of *T. implexicoma* in Australia.
- Tubercularia caca** McAlp. On leaf blades and petioles of *T. implexicoma* in Australia.
- TETRAGONOLOBUS.** Leguminous shrubs related to Lotus.
- Ovularia deusta** (Fockl.) Sacc. See Lathyrus.
- Peronospora tetragonolobi** Gäum. Downy mildew on leaves of *T. siliquosus* in central Europe.
- Uromyces genistae-tinctoriae** Pers. See Caragana.
- TEUFLANA.** See Fatsia.
- TEUCRIUM.** GERMANDER. Herbs and shrubs.
- Ascochyta teucrii** Lasch. On leaves of *T. scordium* in Germany.
- Erysiphe taurica** Lévy. See Althaea.
- Ovulariopsis teucrii** Jaap. Diffuse yellow areas on leaves of *T. chamaedrys* in Dalmatia.
- Peronospora teucrii** Gäum. Downy mildew on leaves of *T. botrys* in Bohemia.
- Phyllosticta infusca** Wint. Subcircular to irregular dull-brown leaf spots on *T. scordium* in Portugal.
- Phyllosticta obliqua** Tassi. Dull-brown zoned leaf spots on *T. flavum* and *T. pulverulentum* in Italy.
- Phyllosticta teucrii** Sacc. and Speg. Grayish-white leaf spots on *T. chamaedrys* in Italy.
- Puccinia annularis** (Strauss.) Schlecht. Brown rust pustules on leaves of *Teucrium* spp. in Europe.
- Puccinia constricta** (Lagh.) Bub. Brown leaf rust on *T. montanum* in Russia and Germany.
- Puccinia istriaca** Syd. Powdery dark-brown rust pustules on leaves and twigs of *T. aureum* and *T. polium* in Spain and the Balkans.
- Puccinia majoricensis** R. Maire. Leaf rust on *T. capitatum* in the Balearic Islands.
- Puccinia teucrii** Biv. Bernh. Dark-brown rust pustules on leaves of *T. fruticans* and *T. polium* in Morocco, Algeria, Syria, Spain, and Italy.
- Ramularia microspora** Thuem. Brown spots on leaves of *T. chamaedrys* in Austria and Italy.
- Septoria scorodoniae** Pass. Small subcircular reddish to ochraceous leaf spots on *T. scorodonia* in Belgium, France, and Italy.
- Septoria teucrii** Sacc. Circular brown to gray leaf spots with dark-purple margins on *T. chamaedrys* in France.
- Taphrina candicans** Sacc. On leaves of *T. chamaedrys* in Europe.
- THALIA.** Perennial marsh or aquatic herbs.
- Puccinia cannae** (Wint.) P. Henn. See Canna.
- Puccinia thaliae** Diet. Powdery brown to black rust pustules on leaves of *T. dealbata* in Brazil.
- THALICTRUM.** MEADOW RUE. Herbs, sometimes cultivated.
- Cercospora thalictri** Thuem. Irregular, often confluent, violet-purple leaf spots on *T. flavum*, *T. glaucum*, *T. jacquinianum*, and *T. speciosum* in Portugal.
- Haplobasidium thalictri** Eriks. Dark-brown leaf spots on *T. flavum* in Sweden.
- Marsonia clematidis** All. See Clematis.
- Oedemium thalictri** Jaap. Gray spots on leaves of *T. minus* in Germany.
- Phyllosticta leucosticta** C. Mass. Small angular spots on leaves of *T. aquilegifolium* in Italy.
- Phyllosticta thalictri** West. Angular to irregular leaf spots on *T. flavum* and *T. minus* in Belgium and Germany.
- Plasmopara alpina** (Johans.) Blytt. Downy mildew on leaves of *T. alpinum* in Scandinavia.
- Pseudopeziza repanda** Sow. On *T. minus* in Japan.
- Puccinia borealis** Juel. See Agrostis.
- Puccinia castagnei** Schroet. Leaf rust on *T. angustifolium* in France, Belgium and Switzerland.
- Puccinia rhytismoides** Johans. Black rust pustules on leaves and stems of *T. alpinum* in Sweden and Norway.
- Puccinia septentrionalis** Juel. See Polygonum.
- Puccinia sommerfeldii** Johans. Dark-brown, powdery rust pustules on leaves of *T. alpinum* and *Polygonum viviparum* in Alaska, Newfoundland, and Europe.

THEA. TEA. Broad-leaved evergreen shrubs cultivated for their leaves and flowers.

Aglaospora aculeata Petch. Causes blight of stems of *Thea* (cult.) in Ceylon.

Ascochyta theae Hara. On stems of *Thea* (cult.) in Japan and Ceylon.

Bacillus theae Hori and Bokura. The "red scaled" disease of *Thea* (cult.) is known only from Japan. On fully developed leaves small pale-brown circular spots occur, which rapidly increase in size to form large confluent, irregular, reddish-brown to black patches. Infected leaves wither and fall. The disease spreads from the leaves to the buds and twigs. The buds turn red-brown and then black. Green twigs turn black, wither, and die back for a length of 4 to 5 inches.

Cercospora theae v. B. de H. Circular purplish-red leaf spots with indefinite yellow-green borders on *Thea* (cult.) in Java, India, and Ceylon. The spots finally become white with narrow purple-red borders.

Cercospora theae Petch. Numerous, small, dark-brown to black, subcircular then irregular spots on young tea leaves. The spots coalesce, destroying the leaves. On older leaves large diffuse chocolate-brown, finally gray, patches occur with narrow purplish-black margins. Sunken purple patches appear on green twigs. Additional hosts are *Acacia decurrens*, *Eucalyptus dealbata*, *E. diversicolor*, *E. melanoxylon*, *E. paniculata*, and *E. robusta* in Ceylon.

Chaetophoma penzigi Sacc. On leaves of *Thea* (cult.) in Italy and the Caucasus.

Colletotrichum camelliae Mass. Brown blight or anthracnose on leaves and twigs of *Thea* (cult.) in Japan, India, Ceylon, Malaya, Java, Formosa, Caucasia, and Uganda. The leaf spots are at first yellowish-green, becoming red-brown or dark brown with first yellow-green margins, and finally gray. Young leaves are blackened and sometimes soft rotted. Twigs of unthrifty plants are at times attacked, becoming hard, gray, and stunted.

Corticium sp. Black rot of tea (*Thea*) in Ceylon is caused by a *Corticium* distinct from *C. theae* of Java. The Ceylon fungus or similar fungi also attack *Calophyllum burmanii*, *Hemidesmus indicus*, *Ozyanthus tubiflorus*, *Pithecolobium dulce*, and *Zinnia* sp.

Corticium salmonicolor B. and Br. See *Citrus*.

Corticium theae Bern. Black rot of tea (*Thea*) in Java, India, and Sumatra is characterized by the drying out and blackening of the leaves and die-back of the twigs due to the fungus, which forms brown fungus strands over infected organs. The disease is one of the thread-blight diseases.

Didymostilbe coffeae Zimm. Causes a die-back of plucked *Thea* branches in Ceylon.

Diplodia cacaocicola P. Henn. See *Theobroma*.

Dioscisia theae Cav. Circular definite, often confluent, black leaf spots on *Thea* (cult.) in Italy and the Caucasus.

Exobasidium camelliae Shir. and var. *gracilis* Shir. Leaves and leaf-shoots are attacked, becoming thick and fleshy or gall-like. Inflorescences are also deformed and reduced to irregular spherical masses. On *T. (Camellia) sasanqua* and *T. japonica* in Japan.

Exobasidium reticulatum Ito and Saw. Attacks *Thea* (cult.) in Japan and Formosa in a similar manner to the following species, except that the leaves are not blistered.

Exobasidium vexans Mass. The blister blight is a serious disease of cultivated *Thea* in India and probably in Formosa. It has also been found on *Rhododendron* spp. Numerous translucent spots appear on the leaves from one-fifth to 1 inch in diameter, red below, pale-green, yellowish, or pink above. The upper surfaces gradually become depressed into shallow cavities. The upper concave surfaces are smooth, shining, and pale-green, the lower surfaces dull-gray to white and powdery. Petioles and stems may also be attacked, the latter dying back. As many as 20 blisters may occur on a single leaf, remaining separate or at times coalescing.

Fomes lamaoensis Murr. See *Hevea*.

Fomes lignosus Klotzsch. See *Hevea*.

Fomes pseudoferreus Wakef. See *Hevea*.

Fusicladium theae K. Hara. Velvety-black fungus patches on leaves of *T. sinensis* in Japan.

Gloeosporium theae Zimm. Large red-brown, then ashen, leaf spots on *Thea* (cult.) in Tanganyika.

Gloeosporium theae-sinensis Miy. Large reddish-brown, then gray, leaf spots on *Thea* (cult.) and *T. sasanqua* in Japan. Probably identical with *Colletotrichum camelliae*.

Guignardia camelliae (Cke.) Butler. (*Laestadia theae* Rac.) This fungus causes the so-called "copper blight" of tea (*Thea*) in India, Ceylon, Java, and Malaya. In the early stages of the disease a peculiar coppery-colored sheen forms on the lower surfaces of the leaves, leaves so affected bending over, with the undersides outermost. Yellow-brown patches then appear, becoming finally gray and very brittle. Possibly the perfect stage of *Colletotrichum camelliae* Mass.

Helicobasidium tanakae Miy. See *Morus*.

Helminthosporium gigasporum B. and Br. On *Thea* (cult.) in Ceylon.

Hendersonia theae K. Hara. Dark-brown, then gray, leaf spots on *Thea* (cult.) in Japan.

Hendersonia theicola Cke. On leaves of *Thea* in Ceylon and the Caucasus.

Hypodermopsis theae K. Hara. Light reddish-brown circular to irregular, often confluent, patches on trunk and branches of *Thea* (cult.) in Japan.

Leptosphaeria hottai K. Hari. Circular to irregular large brown spots on trunks of *T. sinensis* in Japan.

Macrophoma theae Speeschn. Subcircular gray-brown leaf spots on *Thea* (cult.) in the Caucasus.

Macrophoma theicola Petch. This fungus is reported as the cause of a canker of *Thea* (cult.) in Ceylon. Small, slightly sunken, dark patches first appear on the branches and develop into irregular sunken cankers.

Marasmius spp. Horsehair blight. See *Theobroma*.

Massaria theicola Petch. This fungus attacks the branches of the tea (*Thea*) plant in Ceylon, killing them gradually one by one. Individual branches may die slowly, the leaves withering and falling, or because of girdling a branch may be killed suddenly.

Mycosphaerella theae K. Hara. Large irregular dark-brown, then ashen, leaf spots on *Thea* (cult.) in Japan.

Nectria caneri Rutg. A wound parasite causing callused cankers on twigs, branches, and trunks of *Thea* sp. in India and Java.

Neottispora theae Saw. Irregular brown to ashen leaf spots with raised purplish-black margins on *Thea* (cult.) in Formosa.

Pestalozzia theae Saw. The gray blight of tea (*Thea sinensis*) occurs in India, Ceylon, Java, Formosa, and the Caucasus. The fungus produces subcircular to oval leaf spots which vary in size from one centimeter to almost the entire area of the leaf. The spots are dark-brown and concentrically zoned above, brighter beneath, with narrow greenish-yellow surrounding zones. Old spots become gray. Very young leaves when attacked are blackened and distorted. Twigs are also attacked at times, a die-back resulting.

Phaeosphaerella theae Petch. Small yellow leaf spots, circular, then angular, on *Thea* (cult.) in Ceylon. The centers of the spots turn gray or gray-brown and fall out, leaving irregular holes bordered by narrow brown zones.

Phoma theicola Petch. Subcircular to oval or angular bright red-brown leaf spots on *Thea* (cult.) in Ceylon.

THEA—Continued.

Phyllohendersonia theicola (Oke.) Tass. On leaves of *Thea* (cult.) in Caucasasia.

Phyllosticta camelliae West. See *Camellia*.

Phyllosticta theae Speshchn. Subcircular gray-white leaf spots on *T. assamica* and *Thea* (cult.) in Caucasasia.

Physalospora neglecta Petch. Causes cankers on branches of *Thea* (cult.) in Ceylon.

Poria hypobrunnea Petch. See *Hevea*.

Protomyces theae Zimm. On roots of *Thea* (cult.) in Java.

Pseudocommis theae Speshchn. Indefinite, often confluent, gray-brown leaf spots on *T. viridis* (*Thea sinensis*?) in Russia.

Rhizoctonia lamellifera Small. See *Grevillea*.

Rosellinia arcuata Petch. This fungus causes a rotting of the bark around the collar and on the larger roots, forming a white mycelial layer between the bark and wood. Above ground the fungus forms a layer of purple-gray, then black, mycelium around the stems up to a height of about 6 inches. Some plants, although girdled, continue growth, resulting in peculiar enlargements of stems down to the point of girdling. Diseased plants often wilt and die. The hosts in addition to tea (*Thea*) are *Capsicum annuum*, *Cinnamomum camphora*, *Erythrina* sp., *Grevillea robusta*, *Manihot glaziovii*, *Panax fruticosum*, *Strobilanthes* sp., *Symplocos obtusa*, and *Tephrosia candida* in Ceylon and India.

Rosellinia bunodes B. and Br. See *Citrus*.

Sclerotium zeylanicum (B. and Br.) Petch. Attacks seedlings of *Thea* (cult.), *Caladium* spp., and other economic plants in Ceylon.

Septobasidium acaciae Saw. See *Acacia*.

Septoria theae Cav. On leaves of *T. viridis* and *Thea* (cult.) in Italy and Russia.

Sillia theae K. Hara. Small dark-pink or gray spots appear on trunks and branches of *T. sinensis* in Japan. These spots increase until the infected trunk or branch is girdled. The disease is known as the "shark-skin" disease because of the shape of the stromata scattered over the cankered area.

Sphaerostilbe repens B. and Br. See *Hevea*.

Stilbella theae Bern. (*Stilbum nanum* Mass.) Causes a die-back of branches of *Thea* (cult.) in Java.

Valsa theae K. Hara. Acts as a wound parasite of *Thea* (cult.) in Japan.

Venturia speshnewii Sacc. and D. Sacc. Indistinct brown leaf spots on *Thea* (cult.) in Caucasasia.

White-stem blight. See *Hevea*.

Zukalia nantoensis K. Saw. This species of sooty mold and others, including *Z. theae* Saw., *Scorias capitata* Saw., and *Capnodium footii* Berk. and Desm., occur on tea (*Thea*) leaves, injuring the commercial value of the leaves.

THELYMITRA. See *Orchidaceae*.**THEOBROMA.** CACAO OR CHOCOLATE TREE. Tropical trees.

Aspergillus delacroixii Sacc. and Syd. Hastens the hardening and rot of pods of *T. cacao* in Colombia and the Philippines.

Ceratocarpia theobromae von Faber. Thick black fungus crust on leaves of *T. cacao* in Cameroon.

Colletotrichum brachytrichum Delacr. Broad gray-white leaf spots with brown margins on *T. cacao* in Trinidad and Ceylon.

Colletotrichum cradwickii Bancr. Anthracnose of pods of *T. cacao* in Jamaica and Porto Rico. Probably not distinct from *C. gloeosporioides*.

Colletotrichum incarnatum Zimm. Anthracnose on pods of *T. cacao* in Ceylon, Java, Belgian Congo, and Uganda, following *Phytophthora*.

Colletotrichum luxifum v. Hall. and Drost. This fungus was at one time assigned as the cause of the Surinam witches'-broom disease. See *Marasmius perniciosus*.

Colletotrichum theobromae Appel. and Str. Small black sunken areas on pods of *T. cacao* in central Africa.

Colletotrichum theobromicolum Delacr. Anthracnose on pods of *T. cacao* in the West Indies and Uganda. Most, if not all, of the various species of *Colletotrichum* reported as attacking leaves and pods of *T. cacao* can be referred to *C. gloeosporioides*. They are all more or less secondary in their parasitism, attacking trees weakened by adverse cultural conditions or following primary parasites such as *Phytophthora*.

Corticium salmonicolor B. and Br. See *Citrus*.

Diplodia cacaoicola P. Henn. This fungus is found on a wide range of tropical and subtropical plants and has been described under many names, among which are *D. rapax* Mass., *Botryodiplodia theobromae* Pat., *B. elastica* Petch., *Lasiodiplodia theobromae* (Pat.) Griff. and Maubl., *L. nigra* App. and Laub., and *Chaetodiplodia vanillae* Maubl. *Thyridaria tarda* Bancr. has been reported as the perfect stage. Among the many hosts are *T. cacao*, *Hevea brasiliensis*, *Vanilla planifolia*, *Ananas sativa*, *Albizia* sp., *Carica papaya*, *Cajanus indicus*, *Citrus* spp., *Castilleja elastica*, *Cocos nucifera*, *Erythrina* sp., *Cinnamomum camphora*, *Coffea arabica*, *Ficus elastica*, *Grevillea robusta*, *Indigofera* spp., *Mangifera indica*, *Manihot utilissima*, *Nicotiana tabacum*, *Saccharum officinarum*, and *Thea sinensis*. Countries reporting the fungus are Porto Rico, Cuba, French and British West Indies, Trinidad, Brazil, northern South America, Java, India, Ceylon, Malaya, the Philippines, Fiji, and central Africa.

Circular brown spots appear on cacao pods, most frequently at the basal end, and, spreading rapidly, soon bring about complete rot of the pods. The fungus acts as a wound parasite for the most part. Twigs and branches are also attacked, causing a typical die-back and finally stag-headed condition of infected trees. In this phase of the disease, also, the fungus is not virulently parasitic, attacking through wounds or infecting trees weakened by overshadowing or other unfavorable cultural conditions. Where the progress of the fungus is slow, cankers of various sizes form. The action of the fungus on other hosts is the same as on *Theobroma*, a brown rot or soft rot (stem-end rot of *Citrus* fruits) of the fruit and a die-back or cankering of twigs and branches. *D. natalensis* Evans, which causes a typical stem-end rot and die-back of *Citrus* spp. in the Union of South Africa, Porto Rico, Cuba, and southern Florida, is probably not distinct.

Epicoccium theobromae Petch. On leaves of *T. cacao* in Ceylon.

Eutypa erumpens Mass. Causes a rot of bark and wood of branches and trunks of *T. cacao*. *Ficus* spp., *Mangifera indica*, and *Myristica* sp. in Trinidad and the British West Indies. The fungus forms black shining carbonaceous raised stromatic patches on the rotted areas. Probably a wound parasite only.

Exoascus bussei v. Fab. This fungus causes a witches'-broom disease of *T. cacao* in Cameroon (Africa). The brooms arise from an infected bud, which forms an abnormal fleshy branch with shortened internodes. From other buds at the base of this branch other abnormal branches arise, forming the broom. Such leaves as form are small and soon wither and die, remaining attached to the twigs. The brooms may persist for years, never flowering.

Exoascus theobromae Ritz. Eos. Assigned at one time as the cause of the Surinam witches'-broom disease. See *Marasmius perniciosus*.

Fomes lamaensis Murr. See *Hevea*.

Fomes lignosus Klotzsch. See *Hevea*.

Fomes pseudo-ferreus Wakef. See *Hevea*.

Gloeosporium theobromicolum Vinc. Anthracnose on leaves and pods of *T. cacao* in Brazil¹¹

THEOBROMA—Continued.

- Helminthosporium theobromae** Turc. On leaves of *T. cacao* in Italy.
- Macrophoma vestita** Prill. and Delacr. Said to attack the roots of *T. cacao* in tropical America.
- Marasmius** spp. The so-called "horse-hair" blights and thread blights occur on *T. cacao*, *Thea* (cult), *Hevea brasiliensis*, *Myristica fragrans*, and many other tropical shrubs and trees. This type of disease has been reported from Ceylon, west Africa, the West Indies, Trinidad, British Guiana, and Brazil, and doubtless occurs in all tropical regions. Among the species of fungi involved are *M. equicrinis* Mull., *M. pulcher* (B. and Br.) Fetch, and *M. sarmentosus* Berk. Most of these fungi form thin (hairlike) polished black mycelial cords which grow over branches and leaves, adhering at certain points. They are only weakly parasitic, if at all, in contrast to fungi such as *Corticium koleroga* (see *Coffea*), with which class of parasites they are often confused in the literature.
- Marasmius pernicius** Stahel. The "krulloten" or Surinam witches'-broom disease which has made the growing of *T. cacao* commercially almost impossible in Surinam is now attributed by Stahel to the above fungus, *Colletotrichum* and *Exoascus* having been previously assigned as causes by other workers. The disease has also been found in British Guiana. The disease is characterized by somewhat curved broomlike growths brought about by an excessive development of lateral shoots, together with a shortening of the internodes of infected twigs. Diseased shoots are thicker than normal and the surfaces ridged. The leaves are always small and soft and pliant, while the petioles are always much enlarged. The witches'-brooms die in from two to three weeks and fall from the trees. Very few pods are formed and these remain small and misshapen or become very hard and black. Continued attacks of the disease not only prevent production of healthy pods but eventually destroy the trees.
- Melanomma henriquesianum** Bres. and Roum. On bark of *T. cacao* in San Thomé and Kamerun.
- Meliola guianensis** Stev. and Dowell. Black fungus patches on brown leaf spots, 3 to 10 millimeters in diameter on *T. cacao* in British Guiana.
- Meliola theobromae** von Faber. Superficial black fruiting patches on leaves of *T. cacao* in Cameroon.
- Monilia** sp. The watery or *Monilia* pod disease of *T. cacao* occurs in Ecuador, having been found there also on the wild species, *T. bicolor* and *T. balaensis*. The disease is first noted on young pods as small protuberances or discolored areas varying from yellow to brown or even bluish. A white powder consisting of conidia of the fungus, often appears on the diseased areas. The interior of diseased pods is dark-brown, dark-brown or black strands and spots also occurring throughout the husk itself. A watery fluid forms in the interior in considerable abundance. Seeds in diseased pods are rendered worthless.
- Nectria** spp. A number of species of *Nectria*, including *N. bainii* Mass., *N. camerunensis* App. and Str., *N. jungeri* P. Henn., *N. theobromae* Mass., *N. ditissima* Tul., and *N. striatopora* Zimm. have been reported from the various cacao-growing countries as attacking branches, trunks and fruit of *T. cacao* as well as other tropical economic woody plants. There is a voluminous literature describing these fungi and the symptoms of the diseases resulting from their activities, but recent studies indicate that the Nectrias are for the most part saprophytic or at best weak parasites following other fungi, notably *Phytophthora faberi* Maubl. Some of the species may act as wound parasites.
- Phyllosticta theobromae** d'Alm. and S. da Cam. Large irregular leaf spots on *T. cacao* in Portuguese St. Thomas, Java, and Ceylon.
- Phyllosticta theobromicola** Vinc. Irregular brown areas on leaves of *T. cacao* in Brazil.
- Phyalospora affinis** Sacc. On branches of *T. cacao* in the Philippines.
- Phyalospora theobromae** Turc. On leaves of *T. cacao* in Italy.
- Phytophthora faberi** Maubl. This downy mildew fungus causes the most serious known disease of *T. cacao*. Other tropical economic plants, including *Artocarpus integrifolia*, *Carica papaya*, *Cocos nucifera*, and *Hevea brasiliensis* are also attacked. The fungus is reported from Cuba, Jamaica, Dominican Republic, Guadeloupe, Lesser Antilles, Trinidad, Surinam, Ceylon, Java, India, the Philippines, Malaya, Fiji, Siam, Samoa, New Guinea, Gold Coast, Uganda, Cameroon, and Belgian Congo.
- The pod-rot phase of the disease begins as brown spots at the end of the pods, which rapidly spread, soon involving all of the rind and seeds in a brown rot. Diseased pods may be covered with a white mycelium and may either fall to the ground or harden and hang on the tree for a long time. The disease is more serious in moist, shady situations. Seedlings are blighted and destroyed by the fungus. Cankers very similar to those described for *Hevea* occur on the trunks and branches and are often extensive enough to kill the trees. See also *Hevea*.
- Phytophthora meadii** McRae. See *Hevea*.
- Rosellinia bunodes** B. and Br. See *Citrus*.
- Rosellinia paraguayensis** Starb. Found in connection with root disease of *T. cacao* in the British West Indies.
- Rosellinia pepo** Pat. See *Citrus*.
- Ramularia necator** Mass. White mold appears on the cotyledons of seedlings of *T. cacao* in the British West Indies and west Africa. A damping-off results.
- Sphaeronema** sp. This fungus is reported as the cause of a black spot and bark rot disease of *T. cacao* in Ecuador. In the first phase of the disease small dark spots appear on nearly mature pods and increase in size up to 4 centimeters. Diseased pods are invaded by a dry rot and are lighter in weight than normal ones. The fungus gains entrance through wounds and causes large cankers, from which a dark wine-colored liquid exudes in damp weather. The diseased bark tissues become yellow or reddish, a dark line separating healthy tissue from diseased.
- Stachyliidium theobromae** Turc. On leaves of *T. cacao* in Italy.
- Trachysphaera fructigena** Tabor and Bunting. See *Coffea*.
- THERMOPSIS.** Hardy perennial herbs.
- Phyllosticta thermopsisidis** Thuem. Large irregular dull-brown leaf spots on *T. lanceolata* in Siberia.
- Uredo thermopsisidis** Thuem. Leaf rust on *T. lanceolata* in Siberia.
- THESPESIA.** See *Hibiscus*.
- THEVETIA.** Small, glabrous trees and shrubs.
- Aecidium thevetiae** Sacc. Leaf rust on *T. cuneifolia* in Mexico.
- THLASPI.** PENNYCRESS. FRENCH WEED. Annual or perennial herbs.
- Aecidium thlaspiarum** Syd. Leaf rust on *T. ceratocarpon* in Austria.
- Peronospora thlaspeos alpestris** Gäum. Downy mildew on leaves of *T. alpestre* in central Europe.
- Peronospora thlaspeos arvensis** Gäum. As above on *T. arvense* in Europe.
- Peronospora thlaspeos perfoliati** Gäum. As above on *T. perfoliatum* in Yugoslavia, Austria, and Germany.
- Puccinia thlaspeos-glaucophylli** P. Henn. Brown rust pustules on leaves of *T. glaucophyllum* in Chile.
- Ustilago thlaspeos** (Beck.) Lagh. See *Arabis*.
- THRINAX.** THRATCH PALM. See *Palmae*.
- THUJA.** ARBORVITAE. Timber trees, also grown as ornamentals.
- Caecoma deformans** (B. and Br.) Tub. A rust causing witches'-brooms, 4 to 6 inches in diameter. On *T. japonica* and *Thuopsis dolabrata* in Japan.
- Helicobasidium mompa** Tan. See *Morus*.

- THUJOPSIS.** FALSE ARBORVITAE. Ornamental coniferous trees and shrubs grown for their handsome foliage and formal habit.
- Caecoma deformans** (B. and Br.) Tub. See Thuja.
- THUNBERGIA.** CLOCK VINE. Acanthaceous, tall, perennial climbers, with a profusion of variously colored flowers.
- Puccinia thunbergiae** Cke. Brown leaf rust on *T. natalensis* in Japan.
- Puccinia thunbergiae-alatae** P. Henn. Powdery brown rust pustules on leaves of *T. alata* in tropical Africa.
- THYMUS.** THYME. Small shrubs and subshrubs, grown in borders and rockeries.
- Puccinia caulincola** Schneid. Dark-brown rust pustules on leaf blades, petioles, and stems of *T. angustifolium*, *T. chamaedrys*, *T. pannonicum*, *T. serpyllum*, and *Origanum vulgare* in Europe.
- THYRSACANTHUS.** Acanthaceous erect herbs.
- Uromyces hartioides** Lagerh. Leaf rust on *T. strictus* in Costa Rica and Ecuador.
- TIARELLA.** FOAMFLOWER. Slender, hardy, perennial herbs.
- Puccinia asiatica** (Kom.) Syd. See Mitella.
- TIBOUCHINA.** GLOBY BUSH. Melastomaceous shrubs, some species climbers.
- Dothidina peribueyensis** (Speg.) Chardon. See Miconia.
- Physalospora tibouchinae** P. Henn. On *Tibouchina* sp. in Brazil.
- TILIA.** LINDEN. Sometimes called basswood and lime. Timber and ornamental trees.
- Actiononema tillae** Allesch. Causes premature leaf fall and swelling of buds in autumn of *Tilia* sp. in Belgium, Austria, and Germany.
- Ascochyta tillae** Kab. and Bub. Subcircular to irregular, often confluent, yellow-gray, then ashen, leaf spots with purple-brown margins on *T. dasystyla* in Bohemia.
- Ectostroma tillae** Fr. On leaves of *Tilia* sp. in Sweden.
- Gloeosporium tillae** Oud. and var. **maculicolum** Allesch. On leaves of *T. cordata*, *T. platyphyllos*, and *T. cordata* (*T. ulmifolia*) in Europe.
- Gloeosporium tilaeolum** Allesch. Irregular yellow to yellow-brown patches on leaf blades, petioles, and young twigs of *T. platyphyllos*, *T. parvifolia*, and *T. cordata* (*T. ulmifolia*) in Great Britain and Germany.
- Gloeosporium vogelii** Syd. On leaves of *T. cordata* (*T. ulmifolia*) in Germany.
- Helminthosporium tillae** Fr. On leaves of *T. platyphyllos* and *T. cordata* in France.
- Leptosphaeria vagabunda** Sacc. On branches of *Tilia* sp. in Europe.
- Phyllosticta bacterioides** Vuill. Dark-brown leaf spots on *T. cordata* in France.
- Phyllosticta bractephila** Ferr. Small ashen-white leaf spots on *T. platyphyllos* in Italy.
- Phyllosticta praetervisa** Bub. Numerous small angular leaf spots on *T. cordata* (*T. parvifolia*) in Bohemia.
- Phyllosticta tillicola** Oud. Small angular brown leaf spots on *T. cordata* (*T. ulmifolia*) in Holland.
- Phyllosticta vogelii** (Syd.) Died. On leaves of *T. platyphyllos* in Poland.
- Phytophthora syringae** Kleb. See Syringa.
- Placopsphaeria tillae** Bub. Circular to elliptical brown, then yellowish, leaf spots on *T. cordata* in Hungary.
- Pucciniastrum tillae** Hirats. Yellow to brown rust pustules on leaves of *T. cordata*, *T. manshurica*, and *T. miqueliana* in Japan, China, Manchuria, and possibly Russia.
- Pyrenochaeta filarskyi** Bub. Circular to irregular yellow leaf spots on *T. cordata* in Hungary.
- Pyrenochaete pubescens** E. Rostr. Circular to oblong purplish, then ashen, areas on branches of *Tilia* sp. in Denmark.
- Septoria tillae** West. On leaves of *T. europaea* and *T. cordata* (*T. ulmifolia*) in Europe.
- Uncinula miyabei** Salm. Powdery mildew on leaves of *T. miqueliana* in Japan.
- TILLANDSIA.** SPANISH MOSS. Epiphytic bromeliads.
- Phyllosticta tillandsiae** Speg. On leaves of *Tillandsia* sp. in Argentina.
- Ustilago tillandsiae** Patters. Powdery black smut sori destroying the inflorescences of *T. leiboldiana* and *Tillandsia* sp. in Mexico and Costa Rica.
- TINNEA.** Tall perennial herbs and subshrubs.
- Aecidium tinneae** P. Henn. Yellow leaf rust on *T. eriocalyx* in the Union of South Africa.
- TIPUANA.** TIPU TREE. Ornamental leguminous trees.
- Phyllosticta tipuanae** Tassi. On leaves of *T. tipu* (*T. speciosa*) in Argentina.
- TITHONIA.** Tropical composite herbs.
- Puccinia lithoniae** Diet. and Holw. Brown to black rust pustules on leaves of *T. rotundifolia*, *T. scaberrima*, and *T. tubaeformis* in Mexico, Costa Rica, and Guatemala.
- TODDALIA.** Ornamental subtropical shrubs.
- Aecidium toddaliae** Petch. Yellow leaf rust on *T. aculeata* in Ceylon.
- Bacterium citri** Hasse. See Citrus.
- Puccinia kentaniensis** Pole-Evans. Rust on *T. lanceolata* in the Union of South Africa.
- Puccinia tectae** Pass. Black leaf rust on *T. nobilis* in Abyssinia.
- Puccinia toddaleae** Racib. Leaf rust on *T. aculeata* in Java.
- TOFIELDIA.** Perennial rhizomatous herbs.
- Phyllosticta tofieldiae** Rostk. Leaf spot on *T. borealis* in Europe.
- TOLPIS.** Composite herbs with showy yellow flower heads.
- Puccinia heribaudiana** Har. Leaf rust on *T. barbata* in France.
- TORENIA.** Annual or perennial herbs, sometimes grown in gardens.
- Cercospora toreniae** P. Henn. On leaves of *Torenia* sp. in Brazil.
- Puccinia toreniae** Rac. Brown leaf rust on *T. asiatica* in Java.
- TORRESIA.** See Hierochloa.
- TOXYLON.** See MacLura.
- TRADESCANTIA.** SPIDERWORT. Perennial hardy herbs.
- Kordyana tradescantiae** (Pat.) Rac. On leaves of *T. virginiana* and *Tradescantia* sp. in Java and Ecuador.
- Uromyces commelinae** (Speg.) Cke. See Commelina.
- TRAGOPOGON.** VEGETABLE OYSTER. Erect biennial or perennial herbs with narrow grasslike leaves and heads of yellow or purple flowers.
- Ascochyta tragopogonis** Boud. On leaves and stems of *T. major* in Russia.
- Entyloma tragopogi** Lagerh. Smut sori in elongate yellow, then brown, leaf spots on *Tragopogon* sp. in France.
- Puccinia tragopogi** (Pers.) Cda. Yellow and powdery dark-brown rust pustules on leaves and stems of *T. coloratus*, *T. dubius*, *T. floccosus*, *T. major*, *T. orientalis*, *T. palastinus*, *T. porrifolius*, and *T. pratensis* in Asia Minor, Algeria, and Europe. Reported from Idaho.
- Ustilago tragopogonis-pratensis** (Pers.) Wint. Smut sori distorting flower heads of *T. major*, *T. orientalis*, and *T. parvifolius* in Argentina and Europe.

TRAPA. WATER CHESTNUT. Floating aquarium plants.

Septoria trapae-nantatis Wisn. Subcircular leaf spots with dark-purple margins on *Trapa* sp. in Galicia.

TREMANDRA. Australian shrubs.

Puccinia tremandrae B. and Br. Brown leaf rust on *T. oppositifolia* and *T. stelligera* in Australia.

TRICALYSIA. Erect or climbing shrubs.

Hemileia evansii Syd. Leaf rust on *Tricalysia* sp. in the Union of South Africa.

Phyllosticta tricalysiae A. L. Sm. Large gray leaf spots with red-brown margins on *T. griseiflora* in Angola.

TRICHILIA. Ornamental trees and shrubs.

Campostriehum cladosporioides Sacc. Large subcircular leaf spots with dark-olive margins on *T. emetica* in tropical Africa.

Coccinia concentrica Syd. Circular crustaceous black stromata on leaves of *T. emetica* in Portuguese East Africa.

Mycosphaerella asunciensis Starb. On leaves of *Trichilia* sp. in Paraguay.

Phyllachora explanata (Lév.) Sacc. Circular black stromata on leaves of *T. havanensis* in Cuba.

Phyllosticta pertundens Sacc. Subcircular dull-yellow leaf spots on *T. emetica* in Africa.

Uredo trichilliae Arth. Leaf rust on *T. pallida* and *T. trinitensis* in Porto Rico and Trinidad.

TRICHOLOBI. Perennial grasses.

Puccinia trichloridis Speg. Linear black rust sori on leaves of *T. mendocina* in Argentina.

TRICHOSANTHES. SNAKE GOURD. Annual or perennial climbing herbs.

Irene confragosa Syd. Superficial black patches on leaves of *T. quinqueangulata* in the Philippines.

Macrophoma trichosanthis Syd. Causes a fruit rot of *T. anguina* in the Philippines.

Uredo trichosanthes Petch. Leaf rust on *T. palmata* in Ceylon.

TRICHOSTEMA. BLUECURLS. Herbs, sometimes cultivated in gardens or rockeries.

Cercospora trichostemmatis P. Henn. Pale-brown irregular leaf spots on *T. volkensii* in Tanganyika.

TRICHOSTIGMA. Ornamental climbing shrubs.

Cercospora trichostigmae Stevens. Leaf spot on *T. octandra* in Porto Rico.

Endophyllum rivinae (B. and C.) Arth. See Rivina.

Puccinia rivinae (B. and C.) Speg. See Rivina.

Septoria rivinae Pat. See Rivina.

TRIDAX. Hardy perennial composite herbs.

Puccinia melampodii Diet. and Holw. See Zinnia.

TRIENTALIS. STAR FLOWER. Small, glabrous perennials.

Aecidium trientalis Tranzsch. Leaf rust on *T. europaea* in Russia.

Puccinia karelica Tranzsch. Leaf rust on *T. europaea* and *Carex limosa* in Russia. Reported from New York.

Ramularia magnusiana (Sacc.) Lind. On leaves of *T. europaea* in Denmark, Austria, and Germany.

Septoria trientalis (Lasch.) Sacc. Small circular gray-white leaf spots on *T. europaea* and *Trientalis* sp. in Russia, Sweden, Denmark, and Germany.

Sorosporium trientalis Woron. Smut on *T. europaea* in Finland.

Tubercinia trientalis (B. and Br.) Wor. Thickened leaf spots and crustlike growths on stems of *T. europaea* in Japan and northern Europe.

TRIFOLIUM. CLOVER. Herbaceous perennials grown for lawns and as forage crops.

Ascochyta trifolii A. Bond and Trus. On *T. pratense* in Russia.

Ascochyta violartii Bub. On *T. repens* in Bulgaria.

Bacillus trifolii Vogl. Attacks stems, leaves, and peduncles of *T. repens* in Italy.

Bacterium sp. A leafcurl and wilt of stems of *T. pratense* in Italy is thought to be due to a *Bacterium*.

Botrytis anophila A. Bond. Gray mold destroying flowers and preventing seed production of *T. pratense* in Russia.

Cercospora strobiliana Magn. Yellow swollen leaf spots on *T. repens* in Bohemia and Austria.

Hypochnus cucumeris Fr. See Cucumis.

Leptosphaeria circinans (Fckl.) Sacc. See Asparagus.

Mycosphaerella carinthiaca Jaap. (*Ramularia trifolii* Jaap). Leaf spots, brown above, gray-green below, on *T. medium* in Great Britain, Switzerland, Austria, and Germany.

Ovularia exigua (W. Sm.) Sacc. On leaves of *Trifolium* sp. in Great Britain.

Peronospora pratensis Syd. Downy mildew on leaves of *T. incarnatum*, *T. medium*, and *T. pratense* in Europe.

Peronospora trifolii alpestris Gäum. Downy mildew on leaves of *T. alpestre* in Switzerland, Denmark, and Germany.

Peronospora trifolii arvensis Syd. As above, on *T. arvense* in Switzerland, Russia, Denmark, and Germany.

Peronospora trifolii hybridi Gäum. Downy mildew on leaves of *T. hybridum*, *T. spadiceum*, and *T. striatum* in Europe.

Peronospora trifolii minoris Gäum. As above on *T. agrarium*, *T. badium*, *T. minus*, *T. patens* and *T. procumbens* in Europe.

Peronospora trifolii repentis Syd. Downy mildew on leaves of *T. monatum*, *T. repens*, and *T. rubens* in central and northern Europe.

Phyllosticta trifolii Rich. On leaves of *T. pratense* and *T. repens* in Sweden and Russia.

Pseudovularia trifolii Speg. Indefinite subcircular brown spots with subshen centers on *T. platensis* in Argentina.

Rhabdospora alexandrina Chrest and Maire. Irregular black spots on stems and circular brown leaf spots on *T. alexandrinum* in Egypt.

Septoria trifolii Cav. On leaves of *T. repens* in Italy.

Sphaerulina trifolii E. Rostr. Circular brown leaf spots with pale-purple surrounding zones on *T. pratense* and *T. repens* in Denmark and Germany.

Tylenchus dipsaci Kuehn. See Narcissus.

Typhula trifolii Rostr. Fom small black sclerotia in rotted areas in stems and leaves of *T. pratense*, *T. repens*, *Anthyllis vulneraria*, and *Medicago lupulina* in northern Europe. The parasitism of this fungus is uncertain.

Uromyces flectens Lagh. Brown powdery rust pustules on leaf blades and petioles of *T. repens* in Japan and Europe. Also reported from Wisconsin.

Urophlyctis trifolii (Pass.) Magn. Pustules on leaf blades and petioles of *T. montanum*, *T. pratense*, and *T. repens* in Great Britain, Denmark, Italy, and Germany.

TRIGONELLA. FENUGREEK. Annual or perennial herbs.

Cercospora traversiana Sacc. Subcircular brown leaf spots on *T. foenum-graecum* in Italy.

Cercospora trigonellae A. Maubl. On leaves of *T. foenum-graecum* in Brazil.

Peronospora trigonellae Gäm. Downy mildew on leaves of *T. foenum-graecum* and *T. polycerata* in Algeria, Russia, and India.

Uromyces anthyllidis (Grev.) Schroet. See Anthyllis.

Uromyces trigonellae Pass. Leaf rust on *T. foenum-graecum* and *T. occulta* in Egypt, India, Italy and France. Said to be synonymous with the above species.

TRIPETALEIA. Shrubs cultivated for their flowers.

Uredo tripetaleiae Diet. Leaf rust on *T. bracteata* in Japan.

TRIPSACUM. GAMA GRASS. Perennial grasses with stout culms, used as fodder plants.

Puccinia pallescens Arth. Brown powdery rust pustules on leaves of *T. dactyloides*, *T. latifolium*, *T. lanceolatum*, and *Zea mays* in Mexico, Porto Rico, Guatemala, Nicaragua, Salvador, and Trinidad.

Ustilago dieteliana P. Henn. Dark, olivaceous smut sori in panicles of *T. dactyloides* in Mexico.

TRISTETUM. Hardy perennial grasses.

Puccinia triseti Erikss. Yellow-brown to black rust pustules on leaves of *T. barcinonensis*, *T. deyeuzioides*, *T. flavescescens*, *T. neglectum*, and *T. ovatum* in Guatemala and Europe. Also reported from Utah.

Septoria caballeri Gz. Frag. On *T. ovatum* and *T. panicum* in Spain.

Septoria nebulosum Rostr. See Calamagrostis.

Septoria triseti Speg. See Agrostis.

TRISTANIA. Evergreen trees or shrubs.

Ascochyta brunnea Cke. and Mass. Ochraceous to dull-brown leaf spots on *T. conferta* in Australia.

Helminthosporium puccinioides Sacc. and Berl. On leaves of *T. laurina* in Australia.

Mycosphaerella tristaniae Wakef. On leaves of *T. griffithii* in the Federated Malay States.

Phyllosticta tristaniae D. Sacc. Subcircular gray-white leaf spots with red margins on *T. conferta* in Italy.

Septoria banburyana Tassi. On leaves of *T. nereifolia* in Italy.

Septoria tristaniae P. Henn. Brown leaf spots on *T. laurina* in Germany.

TRITELEIA. Bulbous plants.

Uromyces reichii Diet. Brown rust pustules on leaves of *T. gaudichaudiana* in Chile.

Uromyces triteleiae Diet. and Neg. Rust on leaves and scapes of *T. porrifolia* in Chile.

TRITHINAX. South America fan palms. See Palmae.**TRITICUM.** WHEAT.

Acremoniella occulta Cav. See Secale.

Acremoniella verrucosa Togn. See Avena.

Baeillius cerealinum Gentner. See Hordeum.

Dilophospora graminis Desm. (*Mastigosprium album* Reiss.) (*Dilophia graminis* Fekl.) Small light-colored circular to oblong spots appear on both leaf surfaces, becoming larger and dark in color to form blotches and stripes. Infected leaves soon assume a light-yellow color and die. Leaf sheaths are also attacked, particularly the upper ones, and generally before heading out, so that the growing point is firmly inclosed by the young leaves bound together by a mass of mycelium. The continued growth of the tip causes characteristic distortion or buckling of the stems. When infection occurs after heading out deformation of the spikes may be partial or complete, affected areas covered at first with a dirty-white fungus growth which later becomes black and gives a charred appearance. This disease occurs in connection with the nematode *Tylenchus tritici*. Many grasses are attacked, including *Triticum* spp., *Agrostis alba*, *A. stolonifera*, *A. vulgaris*, *Alopecurus agrestis*, *A. pratensis*, *Arrhenatherum elatius*, *Anthraxanthum odoratum*, *Calamagrostis arundinacea*, *C. canadensis*, *C. lanceolata*, *C. epigeios*, *Dactylis glomerata*, *Festuca ovina*, *F. pratensis*, *F. rubra*, *Holcus lanatus*, *H. mollis*, *Phleum pratense*, *Poa trivialis*, and *Secale cereale*. In England, Holland, France, Belgium, Switzerland, Norway, Denmark, and Germany. Reported on *Calamagrostis canadensis* from Wisconsin.

Fusarium nivale (Fr.) Sor. See Secale.

Gibellina cerealis Pass. Round or elongate black patches on leaf sheaths of *Triticum* in Italy and Hungary. Infected plants turn yellow, wilt, and finally die.

Helminthosporium tritici P. Henn. Brown areas on leaves, culms, and spikes of *Triticum* in Tanganyika.

Leptosphaeria culmicola (Fr.) Karst. See Bromus.

Leptosphaeria herpotrichoides De N. Brown spots appear on the leaf sheaths at the base of infected plants, enlarging so as to encircle the culms and penetrating to the stem. Mycelial mats form between and bind the leaf sheaths together. Diseased plants fail to produce grain and infected stems lodge. The hosts are *Triticum*, *Andropogon* sp., and *Koeleria cristata* in Argentina, France, Italy, Finland, Sweden, Denmark, and Germany. One of the foot-rot or take-all group of diseases. See Ophiobolus.

Macrophoma hennebergii (Kuehn) Berl. and Vogl. Brown spots on glumes and leaves of *Triticum* in Italy, Russia, Sweden, and Germany. The disease causes the grain to shrivel up.

Micrococcus tritici Prill. Grains develop poorly, shrivel, and turn red. On *Triticum* (cult.) and *Hordeum vulgare* in Brazil, Argentina, Italy, France, and Denmark.

Mycosphaerella exitialis Mor. See Hordeum.

Mycosphaerella hordicola Hara. See Hordeum.

Myrosporum abrodens Neum. Dark patches on the lower nodes and leaves of *Triticum* in France. Diseased plants fail to form grain.

Nigrospora panici Zimm. Black fungus patches on stems and leaf sheaths of *Triticum*, *Oryza sativa*, and *Zea mays* in Java.

Ophiobolus cariceti (B. and Br.) Sacc. (*O. graminis* Sacc.) The take-all, pietin, or foot-rot disease of cereals has been the cause of heavy losses for many years. A number of fungi have been found associated with the disease, but *O. cariceti* is apparently the most common and most virulent species present. The hosts are *Triticum*, *Secale cereale*, *Agropyron* sp., *Avena sativa*, *Agrostis palustris*, *Hordeum sativum*, *H. murinum*, *Oryza sativa*, *Bromus mollis*, *B. sterilis*, *Elymus*, *Festuca*, *Lolium*, *Hystriz*, and *Phalaris* in Australia, Japan, Argentina(?), Great Britain, Sweden, Holland, France, Belgium, Italy, and Germany. The disease has also been found recently in a number of the States.

The symptoms of the disease are very characteristic. There is a browning or blackening of the base of the stems with the development of an enveloping brown mycelial mat in which the perithecia develop. The roots turn black and rot off. Diseased plants turn yellow and finally die, always failing to set grain. The disease kills out irregular patches in the field which may become very extensive.

Ophiobolus herpotrichus (Fr.) Sacc. This species attacks the stems and leaves at or near the base of the plant, causing a disease similar to the take-all described above. It attacks *Triticum*, *Hordeum* sp., *Cynodon* sp., and *Agrostis* sp. in Italy, France, Denmark, Belgium, Holland, Hungary, and Germany.

TRITICUM—Continued.

- Phoma hennebergii** Kuehn. Large irregular gray-brown spots on glumes of *Triticum* in Sweden.
- Phyllachora tritici-gracilis** (Cast.) Sacc. Black stromata on leaves of *T. gracile* in France.
- Pseudomonas tritici** Hutch. The inflorescences and stems are covered with a bright primrose-yellow slime which forms sticky layers between the glumes and between the stems and sheaths. Such heads as form are distorted. The disease is readily carried on seed grain. On *Triticum* in India.
- Puccinia brachypus** Speg. See Bromus.
- Puccinia megalopotamica** Speg. Powdery brown rust pustules on leaves and sheaths of *Triticum* in Argentina.
- Pyroctonum sphaericum** Prunet. Attacks all parts of *Triticum* plants, causing a yellowing and shriveling of the leaves, abortion of the grain, and dwarfing of the plants in France.
- Rhizoctonia napi** West. See Brassica.
- Sclerospora macrospora** Sacc. This downy mildew gives infected plants a scalded appearance, the leaves dry up, and the heads abort or are deformed. A white mold growth appears on lower leaf surfaces. In addition to *Triticum*, *Alopecurus* sp., *Avena sativa*, *Festuca elatior*, *Lolium temulentum*, *Oryza sativa*, *Phragmites communis*, *Phalaris canariensis*, and *Zea mays* are attacked in Australia, Japan, Abyssinia, Russia, France, Spain, and Italy. The disease is present in the United States in a few localities.
- Septoria agrestis** Sacc. On leaves of *Triticum* in Russia and France.
- Septoria cristati** Höll. Leaf spots on *T. (Agropyron) cristatum* in Hungary.
- Septoria neglecta** Sacc. On leaves of *Triticum* in France.
- Septoria nymniana** Sacc. Elongate ochraceous areas on *Triticum* in Malta.
- Sphaeroderma damnosum** Sacc. The fungus develops a white mycelium beneath the sheaths and stems at the base of infected plants, causing a yellowing and dwarfing of the plants and abortion of the spikes. On *Triticum* and *Secale* in Italy. Considered by some workers as identical with *Fusarium culmorum*.
- Tilletia controversa** Kuehn. Black powdery, fetid masses of smut spores in ovaries of *T. vulgare*, *T. (Agropyron) repens*, *T. glaucum*, and *Elymus orolensis* in Turkestan, Russia, Denmark, Bohemia, Italy, Albania, Austria, and Germany.
- Tylenchus tritici** Bast. This nematode attacks young wheat (*Triticum*) plants, causing a wrinkling and distortion of the upper leaves. Such plants may die or remain alive and produce dwarfed, diseased heads. In place of normal grain dark hard galls form, which are shorter and thicker than wheat grains and filled with the parasites. The nematode affects *Triticum* primarily, but by inoculation has been found capable of attacking spelt, emmer, oats, and rye. The disease is known from Brazil, India, China, Turkestan, Australia, and Europe. In the United States it is reported from Virginia, Maryland, West Virginia, Georgia, and California.
- Typhula graminum** Karst. See Lolium.
- Urocystis tritici** Koern. The flag smut of wheat (*Triticum*) is characterized by long gray sori on the leaves and sometimes on the sheaths and stems, which finally rupture, exposing black powdery spore masses. Diseased leaves and stems become twisted, assume a drooping habit, and finally wither. If grain is formed at all, it shrivels up and is valueless. Diseased plants are much stunted. The disease occurs in India, Japan, China, Australia, Spain, Italy, and the Union of South Africa. Within recent years it has been found in Illinois, Missouri, and Kansas in limited areas.
- Ustilago trebouxii** Syd. See Melica.
- TRITONIA.** Cormous or bulbous plants.
- Septoria tritoniana** Pat. Elongate ashen-colored spots on stems of *Tritonia* sp. in Ecuador.
- Uromyces bonae-palis** Bub. Yellow to chestnut-brown rust pustules on both leaf surfaces of *T. scillarum* and *Acidanthera pallida* in the Union of South Africa.
- Uromyces transversalis** Thuem. See Gladiolus.
- TROCHODENDRON.** Evergreen trees with aromatic bark and foliage.
- Phyllosticta trochodendri** Alm. and Cam. Leaf spot of *T. aralioides* in Portugal.
- TBOLLIIUS.** GLOBEFLOWER. Hardy herbaceous perennials.
- Aecidium trollii** Blytt. Leaf rust on *T. europaeus* in Russia.
- Ascochyta trollii** Thuem. Irregular gray-white leaf spots with dark-brown margins on *T. asiaticus* in Siberia.
- Didymaria trollii** Jacz. On leaves of *T. europaeus* in Russia.
- Metasphaeria trollii** Karst. On stems of *T. europaeus* in Finland.
- Phyllosticta trollii** Trail. Irregular brown leaf spots on *T. europaeus* in Great Britain, Italy, and Spain.
- Puccinia dietrichiana** Tranz. Leaf rust on *T. europaeus* in Russia.
- Puccinia trollii** Karst. Brown rust pustules in irregular sunken spots on leaf blades and petioles of *T. europaeus* in Italy, Switzerland, France, Belgium, Norway, Lapland, and Germany.
- Ramularia trollii** (Jacz.) Lind. On leaves of *T. europaeus* in Spain, Switzerland, Russia, and Finland.
- Septoria trollii** Sacc. and Wint. Ochraceous leaf spots on *T. europaeus* in Yugoslavia, Switzerland, Spain, and the Union of South Africa.
- TROPAEOLUM.** NASTURTIUM. Climbing or erect herbs, cultivated in gardens.
- Colosporium tropaceoli** Palm. Rust on leaves of *T. peregrinum* in Germany.
- Cronartium flaccidum** (Alb. and Schw.) Wint. See Paeonia.
- Phyllosticta tropaceoli** Sacc. and Speg. On leaves of *T. major* in Italy, France, Portugal, and Austria.
- Uredo tropaceolum** Desm. Bright-orange rust pustules on lower leaf surfaces of *Tropaceolum* sp. in Europe.
- Uromyces tropaceoli** Ranoj. Leaf rust on *T. major* in Yugoslavia.
- TSUGA.** HEMLOCK. Coniferous timber and ornamental trees.
- Ascochyta piniperda** Lindau. See Picea.
- Phomopsis pseudotsugae** Wils. See Pseudotsuga.
- TULIPA.** TULIP. Many-colored spring-flowering hardy bulbs.
- Aecidium tulipae** Komarov. Yellow rust pustules on the leaves of *T. turkestanica* in central Asia. Possibly one stage of *Uromyces tulipae* Diet.
- Helmintosporium tenuissimum** Nees. Dead spots on scapes of *T. heteropetala* in Siberia.
- Puccinia prostrii** Moug. This rust produces brown oblong convex rust pustules on both surfaces of the leaves of *T. celsiana* and *T. silvestris*, including many cultivated varieties in Italy, Great Britain, and France. The disease checks flowering or may even prevent it entirely.
- Puccinia tulipae** Schroet. Brown leaf rust on *T. gesneriana* and *T. suaveolens* in Austria and Germany.
- Rhizoctonia tuliparum** (Kleb.) Whet. and Arth. (*Sclerotium tuliparum* Kleb.) This disease has been very destructive at times to tulip plantings in Holland and has also been reported from England, Switzerland, and Germany. There are several reports of occurrence in America. Infected plants rot at the ground level, wither, and die. Diseased bulbs show a dry grayish or reddish-gray rot internally, with abundant grayish-white mycelium externally and between the diseased scales. Numerous more or less globose, dark-brown to black sclerotia occur on the rotted scales and neck of the bulb and occasionally between the scales.

TULIPA—Continued.

Sclerotinia tuliparum (Wakk.) Rehm. This fungus causes the so-called black rot of hyacinth bulbs in Italy, Great Britain, Holland, France, and Germany. Bulbs of *Crocus*, *Scilla* spp., and *Tulipa* spp. are also attacked.

Sclerotium spp. (*Botrytis* spp.) Several *Sclerotium* and *Botrytis* diseases on tulips and other bulbs have been listed and studied by various workers. It is far from clear how many distinct diseases are involved or which ones have reached the United States. *Botrytis tulipae* (Lib.) Hopk., the so-called *Botrytis* blight of tulips, is widespread in this country, having been introduced times without number on Dutch bulbs. See *Rhizoctonia*, *Sclerotinia*.

Synchytrium laetum Schroet. See Gagea.

Urocystis colchici (Schlecht.) Rab. See Colchicum.

Uromyces tulipae Diet. A rust known only from Japan causing yellow to brown rust pustules on both surfaces of leaves of *T. edulis*.

Ustilago tulipae Wint. Elliptical convex pustules appear on the leaves and split longitudinally exposing the black dusty spore masses. *Tulipa* spp., including *T. sylvestris* in France, Austria, and Germany, are the hosts. *Ustilago heufferi* Fekl., which is considered synonymous by some workers with the above, occurs in America on *Erythronium americanum*.

TUMBOA. See Welwitschia.**TUNICA**. TUNIC FLOWER. Annual or perennial hardy herbs.

Septoria dianthicola Sacc. See Dianthus.

Urocystis purpurea Hazl. See Dianthus.

TURRAEA. Ornamental trees and shrubs.

Aecidium ugandense Syd. Rust pustules in circular to irregular yellow to yellow-brown leaf spots on *Turraea* sp. in British tropical Africa.

Pseudographis volkensii P. Henn. On branches of *T. volkensii* in tropical Africa.

TUSSILAGO. COLTSFOOT. Acaulescent perennial herbs with solitary yellow flower heads.

Ascochyta tussilaginis Oud. On leaves of *T. farfara* in Holland.

Coleosporium tussilaginis (Pers.) Lévl. See Pinus.

Phyllosticta caballeri Gz. Frag. On leaves of *T. farfara* in Spain.

Phyllosticta farfarae Sacc. and var. **major** P. Brun. Irregular gray-white leaf spots on *T. farfara* in Italy, France, and Denmark.

Ramularia purpurascens Wint. Grayish-white leaf spots with indefinite purplish surrounding zones on *T. fragrans* and *Nardosmia fragrans* in Portugal.

Septoria farfarae Pass. Gray-brown leaf spots on *T. farfara* in Italy.

Septoria tussilaginis West. Irregular brown to olivaceous, finally gray-white, leaf spots on *T. farfara* and *T. fragrans* in Italy, Belgium, and Germany.

Stagonospora tussilaginis (Fekl.) Died. On leaves of *T. farfara* in Italy, Denmark, Austria, and Germany.

TYPHA. CAT-TAIL. Hardy perennial marsh herbs.

Halmesia corallina Sacc. and Fautr. On leaves of *T. latifolia* in France.

Heterosporium typharum Cke. and Mass. On leaves of *T. angustifolia* in Great Britain.

Ophiobolus typhae Feltg. On leaves of *T. latifolia* in Luxemburg.

Phyllosticta corallibola Bub. and Kab. Irregular elongate ochraceous to brown leaf spots on *T. angustifolia* in Bohemia.

Phyllosticta typha (Pass.) Oud. On leaves of *T. angustifolia* in Holland.

Phyllosticta typhina Sacc. and Malbr. Oblong reddish leaf spots on *T. japonica* and *T. latifolia* in Japan and Europe.

Septoria filispora (Cke.) Sacc. On leaves of *T. latifolia* in France.

Septoria menispora B. and Br. On leaves of *T. latifolia* in Belgium and Great Britain.

Ustilago grandis Fr. See Phragmites.

ULMARIA. See Spiraea.**ULMUS**. ELM. Timber and ornamental trees.

Acremonia pallida Cke. and Mass. On *Ulmus* sp. in Europe.

Actinonema ulmi Allesch. Irregular yellow, then brown, leaf spots on *U. campestris* in Austria.

Ascochyta ulmella Sacc. Pale-brown leaf spots on *U. campestris* in Italy.

Asteroma fuckelii Sacc. On leaves of *Ulmus* sp. in Europe.

Asteroma ulmi Klotzsch. On leaves of *U. campestris*, *U. parvifolia*, and *U. glabra*, (*U. scabra*) in Japan, Great Britain, France and, Portugal.

Cercospora ulmicola v. Hoeh. Subcircular ashen-brown leaf spots with dark margins on *Ulmus* sp. in Austria.

Cucurbitaria naucosa Fr. Causes swellings on the branches of *U. glabra* in Switzerland, followed by withering of the leaves and death of infected branches.

Exoascus campester Sacc. Ochraceous leaf spots on *U. campestris* and *U. glabra* (*U. montana*) in France.

Exosporium ulmi Eriks. Found in connection with die-back of twigs of *U. campestris*, *U. pedunculata* (*U. effusa*), and *U. glabra* in Sweden.

Laetstadia comedens (Pass.) Sacc. Subcircular to irregular leaf spots on *U. campestris* in Italy.

Leptosphaeria ulmicola Mass. Irregular brown leaf spots on *U. campestris* in Italy.

Mycosphaera insularis Walbr. On leaves of *Ulmus* sp. in Germany.

Mycosphaerella oedema (Fr.) Fekl. On leaves of *U. campestris* and *U. effusa* in central Europe.

Phyllosticta bellunensis Mart. Large irregular brown leaf spots on *U. campestris* (*U. elliptica*) and *U. glabra* in Russia, Poland, and Italy.

Phyllosticta lacerans Pass. Gray confluent leaf spots, the centers of which tear and drop out, on *U. campestris* in Russia and Italy.

Phyllosticta ulmi West. Subcircular, then confluent, ashen leaf spots on *U. americana*, *U. montana* and *U. fulva* in Italy, Bulgaria, and Belgium.

Placosphaeria ulmi P. Henn. Circular to angular yellow leaf spots on *U. parvifolia* in Japan.

Septoria ulmicolum (Biv. Bern.) Elenk. and Ohl. On *Ulmus* sp. in Russia.

Sphaeropsis nervisequa Lang. Causes a die-back of twigs of *U. glabra* in Germany. The leaves are first attacked, the fungus working down through the petioles.

Sporidesmium ulmi Fekl. On leaves of *U. pedunculata* in Germany.

Stagonospora ulmifolia (Pass.) Sacc. Small leaf spots on *U. campestris* in Italy.

Systremma ulmi (Schleich.) Theiss. and Syd. (*Phyllachora ulmi* [Duv.] Fekl.) Subcircular scattered, raised black stromata on leaves of *U. campestris*, *U. glabra*, *U. pedunculata*, and *U. suberosa* in Algeria and Europe. Often confused with a similar-appearing American elm-leaf disease caused by *Gnomonia ulmea* (Schw.) Thum.

Taphrina ulmi (Fekl.) Johans. Pellucid, blisterlike spots on leaves of *U. alata*, *U. campestris*, and *U. montana* in Europe.

Teratosperma singulare Syd. Black fruiting pustules on leaves of *U. parvifolia* in Japan.

ULMUS—Continued.

Uncinula candelina (Biv. Bern.) Schroet. Powdery mildew on leaves of *U. campestris*, and *U. montana* in Japan, Algeria and Europe.

A serious disease of elms (*U. americana*, *U. campestris*, *U. pedunculata*, *U. monumentalis* and *U. rupestris*), characterized by a dying back of the twigs and branches, occurs in Holland and adjacent countries. Young trees frequently die while older trees are permanently affected, the leaves remaining small and falling prematurely. A number of fungi, including *Graphium ulmi* Schwartz have been found in connection with the disease, but their parasitic relationships have not yet been worked out. Brusseff attributes the disease to bacteria and considers the fungi secondary.

UMBELLIFERAE. Herbs related to *Cotyledon*.

Aecidium umbilici Trott. Leaf rust on *U. erectus* and *U. pendulinus* in Portugal and Spain.

Phyllosticta umbilici Brun. Indefinite brown leaf spots on *U. erectus* and *U. pendulinus* in Spain, Portugal, and France.

Puccinia umbilici Guep. Powdery red-brown rust pustules on leaf blades and petioles of *U. erectus* and *U. pendulinus* in Great Britain, Belgium, France, Spain, and Portugal.

Septoria umbilici Maire. On leaves of *U. horizontalis* in French North Africa.

UNIFOLIUM. See *Maianthemum*.**URCEALES**. Tropical and subtropical twining vines.

Phyllosticta glaucispora Delacr. On leaves of *U. lutea* in Porto Rico.

URGINEAE. SEA ONION. Bulbous herbs.

Aecidium urginiae P. Henn. and Evans. Yellow rust pustules on leaves of *Urginea* sp. in the Union of South Africa.

Cladochytrium urginiae Pat. and Trab. Oval or subcircular spots, 1 to 2 centimeters in diameter on leaves of *U. maritima* in Algeria.

Septoria urginiae Pass. and Beltr. A leaf spot of *U. scilla* in Sicily.

Uromyces scillarum (Grev.) Wint. See *Hyacinthus*.

Ustilago vaillantii Tul. See *Hyacinthus*.

UROSTIGMA. See *Ficus*.**UTRICULARIA**. BLADDERWORT. Aquatic or terrestrial plants.

Doassansia utriculariae P. Henn. Smut sori in subcircular, then confluent, black leaf spots on *U. reniformis* in Brazil.

UVARIA. Tropical climbers.

Actinonema uvariae (Lév.) Sacc. On leaves of *U. parviflora* in central Africa.

Aecidium uvariae-rufae P. Henn. Circular red-brown leaf spots on *U. rufa* in the Philippines.

Phyllachora incarcerata (Berk.) Sacc. Shiny black stromata on leaves of *Uvaria* sp. in Ceylon.

Phyllosticta uvariae Berk. Subcircular to irregular brown leaf spots on *U. triloba* in Great Britain.

VACCINIUM. BLUEBERRY. CRANBERRY. Shrubs cultivated for their fruit.

Gloeosporium myrtilli Allesch. Anthracnose on *V. myrtilus* in Europe.

Mycosphaerella stemmatea (Fr.) Rom. On leaves of *V. vitis-idaea* in Sweden, Denmark, Esthonia, and Austria.

Phacidium vaccinii Fr. On leaves of *V. vitis-idaea* in Europe.

Phyllachora japonica Cke. and Mass. Doubtful species on *V. japonicum* in China.

Phyllosticta lepidica (Fr.) Allesch. On leaves of *V. vitis-idaea* in France.

Phyllosticta vaccinii-hirti P. Henn. Effuse red-brown leaf spots, ashen-brown beneath, on *V. hirtum* in Japan.

Physalospora leptidea (Hanzl.) Sacc. and Trott. On leaves of *V. vitis-idaea* in Hungary.

Physalospora vitis-idaea Rehm. On leaves of *V. vitis-idaea* in Germany.

Ramularia tumescens (Fckl.) Sacc. On branches of *V. uliginosum* in Belgium and Germany.

Sclerotinia baccharum (Schroet.) Rehm. Fruit rot of *V. myrtilus* in Russia, Switzerland, Austria, and Germany.

Sclerotinia heteroica Wor. See *Ledum*.

Sclerotinia urnula Rehm. White mildew on stems, leaves, and fruit of *V. vitis-idaea* in Finland, Switzerland, and Germany. The fruits become hard and dry.

Septoria lagerheimi Pat. On leaves of *Vaccinium* sp. in Ecuador.

Septoria oleae-vaccinii P. Henn. Circular gray-white leaf spots with dark-brown margins on *V. bracteatum* in Japan.

Septoria stemmatea (Fr.) Berk. Subcircular dull-brown leaf spots on *V. uliginosum* in Siberia, Sweden, Great Britain, Russia, and Italy.

Septoria vaccinii P. Henn. On leaves of *V. bracteatum* and *V. hirtum* in Japan.

Sphaerodothis circumscripta (Berk.) Theiss. and Syd. Irregular black shiny stromata on leaves of *Vaccinium* spp. in Colombia and Peru.

VAGNERA. See *Smilacina*.**VALERIANA**. VALERIAN. Hardy herbs or shrubs, grown in gardens.

Aecidium valdivianum Sacc. and Syd. Rust on leaf blades and petioles of *V. valdiviana* in Chile.

Cylindrosporium valerianae Speg. On leaves of *V. heterophylla* in Siberia.

Endophyllum valerianae-tuberosae R. Maire. Yellow rust sori on leaves of *V. tuberosa* in France and Macedonia.

Peronospora valerianae Trail. Indefinite violet-gray downy-mildew patches on leaves of young plants of *V. morisonii*, *V. officinalis*, and *V. olitoria* in Scotland, France, Esthonia, and Switzerland.

Puccinia commutata Syd. Golden and brown rust pustules on leaf blades, petioles, and stems of *V. officinalis*, *V. sambucifolia*, and *V. tripteris* in Europe. Reported from New York and Oregon.

Puccinia valerianae Carst. Dark-brown rust pustules on leaves of *V. capitata*, *V. celtica*, *V. officinalis*, *V. sambucifolia*, *V. saxatilis*, *V. tripteris*, and *Centranthus calcitrapa* in Alaska, Italy, Switzerland, and Austria.

Septoria valerianae Sacc. and Fautr. Ovate to angular dull-rufous leaf spots on *V. dioica*, *V. major*, *V. montana*, *V. officinalis*, and *V. tripteris* in France, Switzerland, Denmark, and Austria.

Uredo valerianae-wallichii Diet. Brown leaf rust on *V. leschenaultii* and *V. wallichii* in India.

Uromyces valerianae (Schum.) Fckl. Brown leaf rust on *Valeriana* spp. in Europe, Siberia, and the Union of South Africa.

VALERIANELLA. Herbs grown in rock gardens, one species a salad plant.

Aecidium velenovskii Bub. Leaf rust on *V. membranacea* in Bulgaria.

Aecidium valerianellae Biv. Leaf rust on *V. campanulata*, *V. discoidea*, *V. eriocarpa*, *V. microcarpa*, *V. olitoria*, *V. pumila*, and *V. truncata* in Europe and north Africa.

Peronospora valerianellae Fckl. Downy mildew on leaves of *V. carinata*, *V. dentata*, *V. morisonii*, *V. olitoria*, and *V. ramosa* in Europe.

VALERIANOIDES. See *Stachytarpheta*.**VANDA**. See *Orchidaceae*.**VANDOPSIS**. See *Orchidaceae*.

VANGUERIA. Tropical shrubs or trees.

Aecidium vangeriae Cke. and var. **abyssinica** P. Henn. Leaf rust on *V. edulis*, *V. infausta*, and *V. latifolia* in Abyssinia, Uganda, and the Union of South Africa.

Hemileia woodii Kalchbr. and Cke. Orange-yellow rust pustules on leaves of *V. edulis*, *V. euonymoides*, *V. infausta*, *V. latifolia*, *V. madagascarensis*, and *V. pubescens* in Abyssinia, central, and south Africa.

VANILLA. Climbing orchids. Cultivated for their aromatic pods. See Orchidaceae.**VERATRUM.** FALSE HELLEBORE. Hardy perennial herbs grown in gardens.

Cylindrosporium veratrinum Sacc. and Wint. On leaves of *V. album* in Switzerland. Reported from Utah and New York.

Gloeosporium veratrinum Allesch. Anthracnose on leaves of *V. album* (*V. lobelianum*) in Germany.

Mycosphaerella veratri v. Hoeh. On *V. nigrum* in Europe.

Phyllachora melanoplaca Sacc. See *Convallaria*.

Phyllosticta albina Bub. and Kab. Oblong dark-brown leaf spots on *V. album* and *V. lobelianum* in Hungary and Austria.

Phyllosticta macrotheca (Thuem.) Gz. Frag. On leaves of *V. album* in Spain.

Phyllosticta melanoplaca Thuem. Linear black leaf spots on *V. album* and *V. nigrum* in Siberia, Russia, Switzerland, Italy, and Bulgaria. Reported from Utah.

Phyllosticta veratrina Sacc. and Paol. On leaves and stems of *V. album* in Siberia.

Puccinia pachycephala Diet. Brown to black rust pustules on brown or yellow leaf spots on *V. maximowiczii* and *V. nigrum* in Japan and Russia.

Septoria sublineolata Thuem. Linear gray-white leaf spots with dull-brown margins on *V. album* and *V. nigrum* in Siberia.

Uromyces veratri (DC.) Schroet. Yellow and powdery chestnut-brown rust pustules on leaves of *V. album*, *V. nigrum*, *V. stamineum*, *Adenostyles albisfrons*, *A. alpina*, *Cacalia delphinifolia*, *C. krameri* and *Homogyne alpina* in Japan, Siberia, and Europe.

VERBASCUM. MULLEIN. Hardy biennial herbs, mostly weeds.

Asochyta verbasci Sacc. and Speg. Ochraceous leaf spots on *V. phlomoides* and *V. sinuatum* in Italy.

Erysiphe taurica Lévl. See *Althaea*.

Peronospora verbasci Gäum. Downy mildew on leaves of *V. lychnites*, *V. montanum*, *V. nigrum*, *V. phlomoides*, *V. thapsiforme*, and *V. thapsus* in central and northern Europe.

Phyllosticta banatica Bub. Large irregular leaf spots on *V. banaticum* in Hungary.

Phyllosticta verbasci Sacc. Small, pale-brown leaf spots on *Verbascum* sp. in France. Reported from New York and New Jersey.

Ramularia cylindroides Sacc. See *Pulmonaria*.

Septocylindrium heliocense C. Mass. and Sacc. On leaves of *V. nigrum* in Italy.

Uromyces thapsi (Opiz.) Bub. Yellow and black rust pustules on leaves of *V. lychnites*, *V. neilreichii*, *V. nigrum*, *V. orientale*, *V. phlomoides*, *V. schraderei*, *V. thapsus*, and *V. thapsiformis* in Europe.

VERBENA. Annual or perennial herbs or subshrubs, cultivated in gardens.

Aecidium spegazzinianum Sacc. and Trott. Leaf rust on *V. tenera* in Argentina.

Cronartium flaccidum (Alb. and Schw.) Wint. See *Paeonia*.

Microsphaera ferruginea Erikss. Powdery mildew on leaves of *V. hybrida* in Sweden.

Phyllosticta verbenae Sacc. Ashen leaf spots on *V. hastata* and *V. officinalis* in France and Denmark.

Puccinia elongata Speg. Dark-brown rust pustules on stems of *V. litoralis* in Argentina.

VERBESINA. CROWNBEARD. Annual or perennial herbs.

Cercospora tubercularioides Speg. On leaves of *Verbesina* sp. in Argentina.

Coleosporium anceps Diet. and Holw. Golden rust pustules on leaves of *V. sphaerocephala* in Mexico.

Coleosporium verbesinae Diet. and Holw. Golden rust pustules on leaves of *Verbesina* spp. in Mexico, Costa Rica, Guatemala, and Jamaica.

Puccinia affinis Syd. Powdery black rust pustules on leaves of *V. trilobata* and *V. perymentoides* in Mexico and Guatemala.

Puccinia cundimarcensis Mayor. Leaf rust on *V. verbascifolia* in Colombia.

Puccinia diaziana Arth. Chestnut-brown rust pustules on leaves of *V. encelioides* in Mexico.

Puccinia ferox Diet. and Holw. Powdery brown rust pustules on *V. diversifolia* and *V. myriocephala* in Mexico and Costa Rica.

Puccinia involuta H. S. Jack. Leaf rust on *V. montanoifolia* in Mexico.

Puccinia irregularis Diet. Brown leaf rust on *V. pallens* and *V. subcordata* in Nicaragua and Brazil.

Puccinia spegazziniana Det. Leaf rust on *V. montevidensis* in Argentina.

Puccinia verbesinicola Diet. Leaf rust on *Verbesina* sp. in Ecuador.

Septoria balansae Speg. Indefinite leaf spots on *Verbesina* sp. and *Bidens helianthoides* in Brazil.

Sphaerodothis pirifera (Speg.) Theiss. and Syd. Small, black stromata on leaves of *Verbesina* sp. in Brazil and Venezuela.

Uredo verbesinae-dentatae Syd. Brown rust sori on circular leaf spots on *V. dentata* in Ecuador.

VERNONIA. IRONWEED. Perennial herbs, shrubs, and trees; some ornamentals.

Aecidium banosense Syd. Leaf rust on *Vernonia* sp. in the Philippines.

Aecidium tarapotense P. Henn. Dull-brown rust sori on circular leaf spots on *Vernonia* sp. in Peru.

Aecidium vanderystianum P. Henn. Leaf rust on *Vernonia* sp. in Congo.

Aecidium vernoniae P. Henn. Yellow leaf rust on *Vernonia* sp. in Brazil.

Aecidium vernoniae-cinerea Patch. Rust on leaves of *V. cinerea* in Ceylon.

Aecidium vernoniae-hookerianae Patch. Yellow leaf rust on *V. hookeriana* in Ceylon.

Cercospora sublaterita P. Henn. Leaf spot on *Vernonia* sp. in Congo.

Endophyllum vernoniae Arth. Powdery ochraceous rust pustules on leaves of *Vernonia* sp. in Mexico.

Melanospora ruspollana P. Henn. Yellow-brown rust sori on leaves of *Vernonia* sp. in central Africa.

Phyllachora vernonicola P. Henn. and var. **microspora** Theiss. and Syd. Small, scattered shiny black stromata on brown leaf spots on *V. tweediana* and *Vernonia* sp. in Brazil.

Puccinia arthuriana Jacks. Leaf rust on *V. albicaulis*, *V. arbuscula*, *V. bahamensis*, *V. boricuensis*, *V. canescens*, *V. phillostachys*, and *V. sericea* in Porto Rico, the Bahamas, and Costa Rica.

Puccinia beeki Mayor. Leaf rust on *V. cotoneaster* and *V. divaricata* in Colombia and Jamaica.

Puccinia discreta Jacks. and Holw. Brown leaf rust on *V. deppeana* and *V. stellaris* in Guatemala and Costa Rica.

Puccinia egregia Arth. Leaf rust on *V. uniflora* in Mexico.

Puccinia erratica Jacks. and Holw. Leaf rust on *V. schiedeana* in Guatemala.

VERNONIA—Continued.

- Puccinia fraterna* Jacks. Rust on leaves of *V. plurialis* in Jamaica.
- Puccinia fuscella* Arth. and Johnst. Brown leaf rust on *V. menthaefolia* in Cuba.
- Puccinia hyalina* Diet. Leaf rust on *V. scariosa* in Ceylon.
- Puccinia idonea* Jacks. and Holw. Brown leaf rust on *V. triflosculosa* in Guatemala and Costa Rica.
- Puccinia inaequata* Jacks. and Holw. Rust on leaves of *V. patens* in Guatemala.
- Puccinia insulana* (Arth.) Jacks. Brown rust pustules on leaves of *V. albicaulis*, *V. arborescens*, *V. divaricata*, and *V. longifolia* in Porto Rico, St. Croix, Antigua, Guatemala, and Jamaica.
- Puccinia kuntzii* Jacks. Leaf rust on *V. kuntzei* in Bolivia.
- Puccinia le-testii* Maubl. Leaf rust on *Vernonia* sp. in central Africa.
- Puccinia lorentzii* P. Henn. Yellow-brown to dark-brown rust pustules on leaves of *V. cinerea*, *V. lorentzii*, *V. mollissima*, and *V. scorpioides* in Brazil, Argentina, and Ceylon.
- Puccinia membranacea* Diet. Yellow and cinnamon-brown rust pustules on leaves of *V. cauloni* in Brazil.
- Puccinia notha* Jacks. and Holw. Leaf rust on *V. leiocarpa* and *V. shannoni* in Guatemala.
- Puccinia pinguis* Diet. Powdery chestnut-brown rust pustules on leaves of *V. platens* in Brazil.
- Puccinia rota* Jacks. and Holw. Brown leaf rust on *V. leiocarpa* in Guatemala.
- Puccinia rotundata* Diet. Cinnamon-colored rust pustules deforming leaf blades, petioles, and stems of *V. patens*, *V. scabra*, and *V. tuesidiana* in Costa Rica, Panama, Colombia, and Brazil.
- Puccinia semisculpta* Arth. Powdery cinnamon brown to dark-brown rust sori on leaves of *V. alamanii*, *V. dictyophlebia*, *V. karvinskiana*, *V. serratuloides*, and *V. umbellifera* in Mexico.
- Puccinia tonduziana* Speg. Leaf rust on *V. triflosculosa* in Costa Rica and Guatemala.
- Puccinia vernoniae-mollis* Mayor. Leaf rust on *V. mollis* in Colombia.
- Puccinia vernoniae-scariosae* Petch. Brown leaf rust on *V. scariosa* in Ceylon.
- Puccinia veroniciola* P. Henn. Ochraceous rust pustules on leaves of *Vernonia* sp. in tropical Africa.
- Puccinia veroniphila* Speg. Leaf rust on *V. flexuosa* in Argentina.
- Uredo pachystegia* Diet. Brown leaf rust on *Vernonia* sp. in Brazil.
- Uredo vernoniae* P. Henn. Dull-brown rust pustules on circular leaf spots on *Vernonia* sp. in Congo.
- Uredo vernoniae-hookerianae* Petch. Leaf rust on *V. hookeriana* in Ceylon.
- Uredo veroniciola* Petch. Brown leaf rust on *V. cinerea*, *V. setigera*, and *V. wightiana* in Ceylon.
- Uromyces pressus* Arth. and Holw. Leaf rust on *V. deppiana* in Guatemala and Costa Rica.
- VERONICA.** SPEEDWELL. Annual or biennial herbs or shrubs.
- Aecidium disciforme* McAlp. Rust distorting and thickening leaves of *V. calycina* and *V. gracilis* in Australia and Tasmania.
- Aecidium veronicae* Berk. Yellow rust pustules on lower leaf surfaces of *V. sibirica* and *Veronica* sp. in Japan and Australia.
- Ascochyta veronica* Rostr. On leaves of *V. saxatilis* in Denmark.
- Clypeostroma hemisphaericum* (Berk.) Theiss. and Syd. Small black stromata on leaves of *V. elliptica* and *V. odora* in New Zealand.
- Gloeosporium pruinum* Baenl. and var. *tirolense* Kab. and Bub. Small subcircular, then confluent, brown spots on leaves, stems and capsules of *Veronica* spp. in Spain, Bulgaria, Denmark, and Austria.
- Laestadia veronicae* Rostr. On leaves of *V. alpina* in Iceland.
- Olpidiaster radices* (Willd) Pascher. See Linum.
- Ovularia chamaedrys* Lindr. Subcircular gray-brown, then black, leaf spots on *V. chamaedrys* in Finland.
- Ovularia veronicae* (Fekl.) Sacc. and var. *microsticta* Sacc. On leaves of *Veronica* spp. in Ceylon, Sweden, Denmark, Italy, Esthonia, and Germany.
- Peronospora agrestis* Gäum. Downy mildew on leaves of *V. agrestis* and *V. polita* in Switzerland, Galicia, and Russia.
- Peronospora arvensis* Gäum. Downy mildew on leaves of *V. hederifolia* and *V. triphyllus* in Europe.
- Peronospora grisea* Unger. Downy mildew on leaves of *V. beccabunga* in Europe.
- Peronospora palustris* Gäum. Downy mildew on leaves of *V. scutellata* in Europe and Switzerland.
- Peronospora saxatilis* Gäum. Downy mildew on leaves of *V. fruticans* in Europe.
- Peronospora silvestris* Gäum. Downy mildew on leaves of *V. officinalis* and *V. urticifolia* in Europe.
- Phyllosticta prostrata* Brun. Gray-white leaf spots with dull-brown margins on *V. prostrata* in France.
- Phyllosticta veronicae* Thuem. On leaves of *V. longifolia* in Siberia.
- Puccinia rhaetica* Ed. Fisch. Powdery brown rust pustules on leaves of *V. bellidioides* in Switzerland. Reported from Washington.
- Puccinia veronicae* Schroet. Yellow-brown to brown rust pustules on leaves of *V. lutea* and *V. montana* in Europe.
- Ramularia anagallidis* Lindr. Circular to oblong pale-brown leaf spots on *V. anagallis* and *V. moretti* in Italy, Spain, Denmark, and Finland.
- Ramularia beccabungae* Fautr. Circular leaf spots on *V. beccabunga* in Denmark, France, and Austria.
- Ramularia caruianiana* Sacc. Subcircular, often concentrically zoned, leaf spots on *V. anagallis* in Malta.
- Ramularia coccinea* (Fekl.) Vester. (*R. pygmaea* Lindr.) Rufous-brown leaf spots on *V. chamaedrys*, *V. pygmaea*, *V. officinalis*, and *V. serpyllifolia* in Finland, Denmark, Sweden, Esthonia, Austria, and Germany.
- Ramularia nivea* Kab. and Bub. Subcircular, often confluent, dull yellow to brown leaf spots on *V. anagallis* in Bohemia.
- Ramularia pseudococcinea* Liro. Rufous to brown leaf spots with dark margins on *V. chamaedrys* in northern Europe and Spain. Said to be the same as *R. coccinea*.
- Septoria exotica* Speg. and var. *andersoni* Tassi. Confluent brown leaf spots on *V. andersoni*, *V. elliptica*, *V. salicifolia*, and *V. speciosa* in New Zealand, Argentina, Ecuador, Italy, and France.
- Septoria triphylli* Höll. On leaves of *V. triphyllus* in Hungary.
- Septoria veroniciola* Karst. On leaves of *V. formosa* in Italy.
- Sorosphaera veronicae* Schroet. Small irregular or cylindrical smut sori as outgrowths from leaves of *V. chamaedrys*, *V. hederifolia*, and *V. triphyllus* in Sweden, Denmark, and France.
- Synchytrium globosum* Schroet. See Potentilla.
- VESICARIA.** Annual or perennial herbs cultivated in gardens.
- Septoria ceresperma* Rostr. On leaves of *V. arctica* in Greenland.
- VIBURNUM.** Ornamental shrubs grown for their attractive flowers, fruit, and foliage.
- Aecidium viburni* Henn. and Shir. Leaf rust on *V. burceaticum*, *V. dilatatum*, *V. formosanum* and *V. sargentii* in Russia and Japan.
- Aecidium viburnophilum* Syd. Leaf rust on *V. opulus* in Japan.
- Ascochyta tini* Sacc. Ochraceous leaf spots on *V. tinus* in Italy.

VIBURNUM—Continued.

- Ascochyta viburni* (Roum.) Sacc. Brown spots with brown-purple margins on branches of *V. lantana* and *V. opulus* in France, Denmark, Austria, and Germany.
- Cercospora penicillata* Fekl. On leaves of *V. opulus* in Siberia, Italy, Portugal, and Germany.
- Cercospora tineae* Sacc. On leaves of *V. tinus* in Portugal.
- Laetadiala tuscula* Pass. Subcircular to irregular gray-white leaf spots on *V. tinus* in Italy.
- Meliola viburni* Syd. Superficial black fungal patches on leaves of *V. odoratissimum* in China and the Philippines.
- Mycosphaerella crepidophora* (Mont.) Rehm. Circular dull-brown leaf spots on *V. tinus* in Corsica, Dalmatia, Portugal, and Tunis.
- Mycosphaerella viburni* (Nitsch) Fekl. On leaves of *V. lantana* and *V. opulus* in Italy and Germany.
- Phyllosticta lantanæ* Pass. See *Lantana*.
- Phyllosticta lantanicola* Sacc. Irregular ochraceous, then gray-white, leaf spots on *V. lantana* in Bohemia.
- Phyllosticta opuli* Sacc. On leaves of *V. opulus* in Europe.
- Phyllosticta roumeguerii* Sacc. Indefinite gray leaf spots on *V. opulus* and *V. tinus* in Europe.
- Phyllosticta tineae* Sacc. Subcircular to irregular gray-white leaf spots on *V. tinus* in Italy.
- Phyllosticta viburni* Pass. Large irregular gray or rose-gray leaf spots on *V. tinus* in France and Portugal.
- Piasosphaeria viburni* P. Henn. Black subcarbonaceous stromata on leaves of *V. dilatatum* in Japan.
- Pucciniastrum miyabeanum* Hirats. Yellow rust pustules on leaves of *V. furcatum* in Japan.
- Rhytisma viburni* P. Henn. Black stromata on leaves of *V. dilatatum* and *V. luzonicum* in the Philippines.
- Septoria tui* (Arc.) Sacc. Irregular ashen-brown leaf spots with purple margins on *V. tinus* in Italy.
- Septoria viburni* West. On leaves of *V. lantana* and *V. opulus* in Great Britain, Denmark, Belgium, and Italy.

VICIA. VETCH. Annuals cultivated for forage and flowers.

- Ascochyta bolthauseni* Sacc. See *Phaseolus*.
- Ascochyta fabae* Speg. Circular to irregular, then confluent, dull-tan leaf spots on *V. faba* in Argentina.
- Ascochyta phaseolorum* Sacc. See *Phaseolus*.
- Ascochyta viciae-lathyroides* Syd. Large gray-white areas on leaves of *V. lathyroides* in Germany.
- Ascochyta viciae-pisiformis* Bub. Subcircular to irregular, then confluent, dull-yellow leaf spots with rufous margins on *V. pisiformis* in Bohemia.
- Cephalotheca francisci* Sacc. On stems of *V. faba* in Italy.
- Cercospora fabae* Fautr. Dark-purple, zoned, then confluent, leaf spots with gray centers on *V. faba*, *V. gigantea*, and *V. narbonensis* in China, Russia, Italy, Bohemia, and France.
- Cercospora zonata* Wint. On leaves of *V. faba* and *Phaseolus vulgaris* in Brazil, Portugal, Russia, Italy, and Germany.
- Cercospora viciae* Siem. Leaf spot on *V. aurantia* in Poland.
- Didymaria lindaviana* Jaap. Large pale-brown leaf spots on *V. cracca* in Germany.
- Erysiphe taurica* Lévy. See *Althaea*.
- Gloeosporium tricolor* Lind. Circular dark-brown leaf spots with paler margins on *V. cracca* in Sweden and Denmark.
- Hypochnus cucumeris* Frank. See *Cucumis*.
- Microsphaera bäumieri* P. Magn. Powdery mildew on leaves of *V. cassubica* and *V. sylvatica* in Great Britain, Russia, Italy, Austria, Hungary, and Germany.
- Ovularia fallax* (Bon.) Sacc. On leaves of *V. cracca*, *V. cassubica*, *V. pyrenaica*, and *V. villosa* in Spain and Germany.
- Ovularia schwarziana* Magn. Brown leaf spots on *V. villosa* in Denmark and Germany.
- Ovularia sphaeroidea* Sacc. See *Lotus*.
- Ovularia viciae* (Frank.) Sacc. Brown leaf spots on *V. cassubica* and *V. tenuifolia* in Denmark and Germany.
- Ovularia villiana* P. Magn. On leaves of *V. cassubica* in Germany.
- Peronospora sepium* Gäum. Downy mildew on leaves of *V. sepium* in Scotland, central and northern Europe.
- Peronospora viciae* (Berk.) Gäum. Downy mildew on leaves of *V. grandiflora*, *V. hirsuta*, *V. lathyroides*, *V. pisiformis*, *V. tenuifolia*, and *V. tetrasperma* in Europe.
- Phyllachora lathyri* (Lév.) Theiss. and Syd. See *Lathyrus*.
- Phyllosticta fabae* West. Dull-brown leaf spots on *V. faba*, *V. sativa*, and *V. villosa* in Russia, Sweden and Belgium.
- Phyllosticta viciae* (Lib.) Cke. Subcircular to oblong ochraceous to brown leaf spots with dark margins on *V. faba* and *V. sepium* in Russia, France, and Belgium.
- Rhizoctonia napi* West. See *Brassica*.
- Rhytisma onobrychidis* DC. See *Onobrychis*.
- Septoria silvestris* Pass. See *Lathyrus*.
- Septoria viciae* West. Dull-brown leaf spots on *V. angustifolia*, *V. pyrenaica*, and *V. sativa* in Denmark, Russia, Spain, and Italy.
- Stagonospora carpathica* Bauml. Small circular to subcircular red-brown spots appear on the leaves, petioles, stems, and pods, changing to dark-brown with red-brown margins, and finally to gray-brown. Infected plants often die. The hosts are *Vicia faba* in Japan and *Melilotus alba* and *M. officinalis* in central Europe.
- Uromyces briardi* Har. Brown rust pustules on leaves of *V. sativa* in France.
- Uromyces clavatus* Diet. See *Lathyrus*.
- Uromyces corrugatus* Speg. Powdery brown rust pustules on leaves of *V. patagonica* in Patagonia.
- Uromyces fischeri-eduardi* P. Magn. Powdery brown rust pustules on leaves of *V. cracca* and *Euphorbia cyparissia* in Europe.
- Uromyces heimerlianus* P. Magn. Brown rust pustules on leaves of *V. hirsuta* and *V. pannonica* in Yugoslavia, Hungary, and Austria.
- Uromyces johowii* Diet. and Neg. Yellow to brown rust pustules on leaves of *V. nigricans* in Chile.
- Uromyces nordenskjöldii* Diet. Yellow-brown rust pustules on leaves of *Vicia* sp. in South America.
- Uromyces orobi* (Pers.) Lévy. See *Lathyrus*.
- Uromyces unijugae* Ito. Leaf rust on *V. unijuga* in Japan.
- Uromyces valesiacus* Ed. Fisch. Yellow and brown leaf rust on *V. onobrychioides* in Macedonia and Switzerland.
- Uromyces viciae-cracca* Const. Powdery brown rust pustules on leaves of *V. cracca*, *V. tenuifolia*, and *Lens esculenta* in Europe.
- Uromyces viciae-unijugae* S. Ito. Leaf rust on *V. unijuga* in Japan.

- VIGNA.** COWPEA. Herbs grown as vegetables, forage, and green-manure crops.
- Aecidium caulicola** P. Henn. Rust on stems of *Vigna* sp. in Central Africa.
- Aecidium nigro-cinctum** Pat. and Har. Leaf rust on *Vigna* sp. in Indo-China.
- Aecidium vignae** Cke. Leaf rust on *V. catjang*, *V. marginata*, and *V. sinensis* in Ceylon, Uganda, and Union of South Africa.
- Ascochyta phaseolorum** Sacc. See Phaseolus.
- Ascochytopsis vignae** P. Henn. Dull-brown leaf spots on *V. vexillata* in east Africa.
- Auerswaldia vignae** P. Henn. Clear brownish spots on leaves of *V. vexillata* in Tanganyika.
- Cercospora vanderysti** P. Henn. Effuse brown leaf spots on *V. venulosa* in the Congo.
- Hypochnus cucumeris** Frank. See Cucumis.
- Mycosphaerella phaseolorum** Jachw. On leaves of *V. rubra* in Russia.
- Phyllachora phaseolina** Syd. See Phaseolus.
- Phyllosticta phaseolorum** Sacc. and Speg. Ochraceous leaf spots on *V. sinensis* and *Phaseolus vulgaris* in Japan, Formosa, and Italy.
- Phyllosticta vignae** Speg. Circular pale dull-brown leaf spots on *V. luteola* in Argentina.
- Puccinia** sp. Leaf rust on *V. angustifolia* in the Union of South Africa.
- Rhizoctonia** sp. Diseased plants wilt and die, the stems at the ground level being blackened and covered with numerous tiny black sclerotia. Among the hosts are *V. catjang*, *V. unguiculata*, *Corchorus capsularis*, *C. olitorius*, *Dolichos lablab*, *D. biflorus*, *Cucurbita* sp., *Citrullus* sp., *Gossypium* sp., *Lycopersicon esculentum*, *Medicago sativa*, *Nicotiana tabacum*, *Phaseolus lunatus*, *P. radiatus*, *Sesamum indicum*, and *Solanum tuberosum* in India.
- Rhizoctonia dimorpha** Matz. See Phaseolus.
- Septoria vignae** P. Henn. Circular dull-brown leaf spots on *Vigna* sp. in the Congo.
- Uredo purpurascens** P. Henn. Rust on circular to effuse purple spots on leaves and stems of *Vigna* sp. in the Congo.
- Uredo vignae** Bres. Brown leaf rust on *V. lutea*, *Glycine hispida*, *Phaseolus mungo*, and *Phaseolus* sp. in the Philippines and St. Thomas.
- Uromyces pазschkeanus** P. Henn. Black rust sori on leaves of *V. membranacea* and *V. triloba* in Abyssinia and Arabia.
- Uromyces punctiformis** Syd. Powdery black rust pustules on leaves of *V. strobiliphora* in Mexico.
- Uromyces vignae** Barcl. Brown to black rust pustules on leaves of *V. vexillata* in India.
- Vermicularia capsici** Syd. See Capsicum.
- VIGUIERA.** Shrubby or herbaceous composites.
- Coleosporium viguieriae** Diet. and Holw. Golden rust pustules on leaves of *V. helianthoides* in Mexico.
- Puccinia ensenadensis** Speg. Powdery black rust pustules on leaves of *Viguiera* sp. in Argentina.
- Puccinia inopinata** Syd. Dark-brown rust pustules on leaves of *V. stenophylla* in Argentina.
- Puccinia nanomitra** Syd. Powdery black rust pustules on leaves of *V. dentata*, *V. eriophora*, and *V. helianthoides* in Mexico.
- Puccinia punctoidea** Syd. Black rust pustules on leaves of *V. pringlei* in Mexico.
- Puccinia subglobosa** Diet. and Holw. Brown leaf rust on *V. palmeri* in Mexico.
- VINCA.** PERIWINKLE. Erect and procumbent herbs or shrubs cultivated for their flowers.
- Colletotrichum vineae** Speg. Circular ash leaf spots on *V. major* in Argentina.
- Macrophoma cylindrospora** Desm. See Hedera.
- Metasphaeria vineae** (Fr.) Sacc. On leaves of *V. minor* in Italy, France, and Germany.
- Peronospora vineae** Schroet. Downy mildew on leaves of *V. major* and *V. minor* in France and Russia.
- Phytophthora parasitica** Dast. See Ricinus.
- Phyllosticta vineae** Thuem. and var. **pedrosensis** Gz. Frag. Large irregular dull-brown to gray leaf spots on *V. media* and *V. minor* in Spain, Portugal, and France.
- Phyllosticta vineae-majoris** Allesch. Subcircular ochraceous leaf spots on *V. major* in Germany.
- Phyllosticta vineae-minoris** Bres. and Krieg. Dull-brown leaf spots on *V. minor* in Albania, Yugoslavia, and Germany.
- Phyllosticta vincicola** Oud. Irregular black leaf spots on *V. major* in Holland.
- Puccinia vineae** (DC.) Berk. Brown to black rust pustules on leaves of *V. acutiflora*, *V. herbacea*, *V. major*, *V. media*, and *V. minor* in Europe and Algeria. Reported from Massachusetts and Michigan.
- Ramularia vineae** Sacc. and var. **vineae-mediae** Maire. Yellow indefinite leaf spots on *V. major* and *V. media* in Spain and the Balearic Islands.
- VIOLA.** VIOLET. Pansy. Perennial herbs, cultivated for their flowers.
- Aphelenchus olesistus** Ritz. Bos. var. **longicollis** Schwartz. This nematode attacks the bases of the leaves and flower stalks of cultivated *Viola*, causing hypertrophy, with consequent stunting and fasciation and finally production of galls. The disease occurs in Europe.
- Ascochyta violae-hirtae** Bub. White subcircular leaf spots with brown margins on *V. hirta* in Yugoslavia.
- Ascochyta violicola** McAlp. Yellow to gray-white circular to ovate leaf spots on *V. odorata* in Australia.
- Cercospora** ii Trail. Indefinite dull-brown leaf spots on *V. palustris* in Scotland.
- Cercospora lilacina** Bres. On leaves of *V. palustris* in Germany.
- Cercospora violae-sylvatica** Oud. Subcircular pale-brown leaf spots on *V. sylvatica* in Holland.
- Cercospora violae-tricoloris** Br. and Cav. Large subcircular ash leaf spots on *V. tricolor* in Italy and Spain.
- Cladochytrium violae** Berl. On leaves of *V. odorata* in Europe.
- Entyloma anizanum** Pass. Smut sori on subcircular yellow leaf spots on *V. biflora* in Italy.
- Fusisporium lacteum** Desm. On leaves of *V. odorata* in France.
- Hendersonia triseptata** Da Cav. Subcircular leaf spots on *V. alba* in Portugal.
- Laetadial violae** (Lib.) Sacc. On leaves of *V. odorata* in Belgium.
- Melampsora lapponum** Lindf. See Salix.
- Mycosphaerella violae** A. Pot. On leaves of *V. hirta* in Russia. Possibly the perfect stage of *Ramularia lactea* Sacc.
- Ovularia acutata** (Bon.) Sacc. Small ash leaf spots with dull-brown margins on *V. odorata* and *V. sylvatica* in Sweden and Germany.
- Peronospora violae** De By. A downy mildew destroying the leaves of *V. biflora*, *V. tricolor*, and other species in Europe. American collections referred to this species should be referred to *Bremiella megasperma* (A. Berl.) G. W. Wils.
- Phyllosticta libertianae** Sacc. and March. Circular black leaf spots on *V. biflora* and *V. odorata* in Great Britain, France, Belgium, and Italy.
- Phyllosticta tricoloris** Sacc. On leaves of *V. odorata* and *V. tricolor* in Russia.
- Puccinia cingens** Bonm. and Rouss. Yellow and black rust pustules on leaves of *Viola* in Patagonia,

VIOLA—Continued.

- Puccinia depauperans** (Vize.) Syd. Yellow and brown to black rust pustules on leaves and stems of *V. cornuta*, *V. lutea*, *V. tricolor*, and *V. willkommii* in Europe.
- Puccinia hederaceae** McAlp. Powdery black rust pustules on leaf blades, petioles, and stems of *V. betonicifolia* and *V. hederacea* in Australia and Tasmania.
- Puccinia violae-glabbellae** Miura. Powdery brown rust pustules on leaves of *V. glabella* in Japan.
- Ramularia acutata** Bon. Subcircular, then confluent, spots occupying entire surface of leaf blades of *V. canina*, *V. riviniana*, *V. silvatica*, and *V. sylvestris* in Denmark, Russia, Scotland, and Austria.
- Ramularia agrestis** Sacc. Small subcircular olivaceous leaf spots on *V. altaica* and *V. tricolor* in Denmark, Estonia, and Italy.
- Ramularia biflorae** P. Magn. Subcircular greenish brown, then gray-white, irregularly zoned leaf spots on *V. biflora* in Austria.
- Ramularia defletores** Bres. On leaves of *V. tricolor* in Russia, Denmark, and Germany.
- Ramularia lactea** (Desm.) Sacc. Circular gray-white leaf spots with brown margins on *V. canina*, *V. hirta*, *V. odorata*, *V. sylvestris*, and *V. tricolor* in Europe and Argentina. Reported from Montana and Colorado.
- Septoria australiae** McAlp. Circular to irregular red-brown leaf spots with dark-brown margins on *V. betonicifolia* in Australia.
- Septoria violae-palustris** Died. On leaves of *V. palustris* in Austria.
- Septoria violicola** Sacc. On leaves of *V. biflora* in Great Britain, Switzerland, and Germany.
- Synchytrium alpinum** Thomas. Small galls on leaves and stems of *V. biflora* in Italy and Switzerland.
- Synchytrium globosum** Schroet. See *Potentilla*.
- Uredo alpestris** Schroet. Leaf rust on *V. biflora* in Japan, Switzerland, and Austria.
- Urocystis violae** Fisch. Smut sori causing blistering and distortion of leaf blades and petioles of *V. hirta*, *V. odorata*, and *V. tricolor* in Great Britain, France, Italy, Denmark, and Germany. The pustules finally split irregularly, exposing black sooty spore masses. Also reported from Utah and Minnesota on wild species.

VITEX. CHASTE TREE. Ornamental shrubs and trees with white or blue flowers.

- Aecidium viticis** Jul. Leaf rust on *Vitex* sp. in Trinidad.
- Helminthosporium viticis** Syd. Gray-brown leaf spots on *V. flavens* in Brazil.
- Phyllachora taruma** Speg. Shiny black circular stromata on leaves of *V. montevicensis* and *Vitex* sp. in Paraguay and Brazil.
- Phyllachora viticicola** P. Henn. Shiny black stromata on clear brown leaf spots on *Vitex* sp. in Brazil.
- Ramularia viticis** Syd. Circular to irregular brown leaf spots on *V. negundo* in India.
- Uredo raciborskiana** Sacc. and Trott. Yellow leaf rust on *V. leucocylon* in Java.
- Uredo viticis** Rac. Leaf rust on *V. leucocylon* in Java.
- Uredo viticis** Juel. Leaf rust on *Vitex* sp. in Paraguay. Hennings gives the same name to a species on *V. polygama* in Brazil.
- Uredo viticis-polygamae** P. Henn. Dull-brown rust pustules on circular leaf spots on *V. polygama* in Brazil.

VITIS. GRAPE. Woody vines.

- Aecidium vitis** A. L. Sm. Rust pustules on dull-brown leaf spots on *V. vulpina* in Tanganyika.
- Alternaria viticola** Brun. Ashen leaf spots on *V. vinifera* and *V. vulpina* in Italy.
- Ascochyta baccicola** P. Brun. Spots on mature fruit of *V. vinifera* in France.
- Ascochyta chlorospora** Speg. On leaves of *V. vinifera* in Australia and Argentina.
- Aureobasidium vitis** Viala and Boyer. Causes a dying back and blackening of twig tips, irregular red patches on leaves which fall prematurely and a spotting and shrivelling of the fruit of *V. vinifera* in Australia, Japan, Russia, France, Denmark, and Italy.
- Bacillus ampelopsorae** Trev. Said to cause small spongy galls on the roots and at the base of stems of *Vitis* in Argentina, France, Italy, and Germany. Probably crown-gall (*Bacterium tumefaciens*).
- Bacillus uvae** Cug. and Macc. Bacteriosis of *V. vinifera* in Italy.
- Bacillus vitivorus** Bacc. Said to cause malformation of the leaves, stripes on young branches, and dark lines in the wood of *V. vinifera* in Argentina and Italy.
- Cercospora roesleri** (Catt.) Sacc. Indefinite irregular olivaceous leaf spots on *V. vinifera* in Europe and Argentina.
- Cercospora vitis-heterophyllae** P. Henn. On leaves of *V. heterophylla* in Japan.
- Chrysomyxa vitis** Butl. Numerous yellow to brown powdery rust pustules on leaves of *V. latifolia* in India, causing defoliation.
- Clasterosporium putrefaciens crucipes** N. Spes. See *Morus*.
- Coryneum vitiphyllum** N. Spes. Numerous subcircular dull-brown leaf spots with purple-brown surrounding zones on *V. vinifera* in Russia.
- Exobasidium clandestinum** N. Spes. On leaves of *V. vinifera* in Russia.
- Fusidium peronosporae** Fautr. and Lamb. On leaves of *V. vinifera* in France.
- Fusicoccum bulgaricum** Bub. On *V. vinifera* in Austria.
- Helicobasidium tanakae** Miy. See *Morus*.
- Hendersonia rubi** West. See *Rubus*.
- Hypochnus burnati** Lendn. On trunks of *V. vinifera* in Switzerland.
- Kuehneola vitis** (Butl.) Syd. Powdery yellow and brown rust pustules on leaves of *V. latifolia* in India.
- Leptosphaeria vinealis** Pass. (*L. appendiculata* Pir.) On leaves and stems of *V. vinifera* in Italy.
- Marsonia viticola** Miy. On leaves of *V. vinifera* in China.
- Mycosphaerella rathayi** Nyp. On *Vitis* sp. in Belgium.
- Napcladium pusillum** Cav. On fruit of *V. vinifera* in Italy.
- Ovularia vitis** Rich. On leaves of *V. vinifera* in France, Austria, and Germany.
- Phakopsora ampelopsidis** Diet. and Syd. See *Ampelopsis*.
- Phakopsora coronatiformis** Diet. Yellow and brown rust pustules on leaves of *V. himalayana* in India. Causes leaf-fall.
- Phyllohendersonia vitiphylla** (N. Spes.) Tassi. Brown leaf spots on *V. vinifera* in Transcaucasia.
- Phyllosticta badhami** Cke. Indefinite elongate brown leaf spots on *V. vinifera* in Great Britain.
- Phyllosticta bizzozzeriana** C. Mass. Small irregular brown leaf spots on *V. vinifera* in Europe.
- Phyllosticta dumajensis** Bub. On leaves of *V. vinifera* in Bulgaria.
- Phyllosticta frankiana** Sacc. and Syd. Brown leaf spots with red margins on *V. vinifera* in Transcaucasia.
- Phyllosticta microspila** Pass. Small irregular dark-brown leaf spots on *V. vinifera* in Italy.
- Phyllosticta negeriana** (Thuem.) Allesch. Gray leaf spots, dark-brown beneath, on *V. vinifera* in Italy and Central Europe.
- Phyllosticta neurospileae** Sacc. and Berl. Red to ochraceous leaf spots on *V. antarectica* in Australia.
- Phyllosticta pilispora** N. Spes. Irregular dull-brown leaf spots on *V. vinifera* in Transcaucasia.

VITIS—Continued.

Phyllosticta succedanea (Pass.) Allesch. Dark-brown leaf spots on *V. vinifera* in Italy.

Phyllosticta viticola Sacc. and Speg. Gray-white leaf spots on *Vitis* sp. in Italy.

Phyllosticta vitis Sacc. Irregular gray leaf spots with brown borders on *V. vinifera* in Italy and central Europe.

Pseudopeziza tracheiphila Müller-Thurg. Said to attack the leaves of *V. vinifera* in Europe, causing the disease known as "rotbrenner."

Pseudocercospora vitis (Lév.) Sacc. On leaves of *V. rupestris* in Argentina and Brazil.

Septocylindrium dissiliens (Duby.) Sacc. Causes a drying of the leaves of *Vitis* sp. in Italy.

Septoria melanopsis Pat. Small numerous angular blackish leaf spots on *V. vinifera* in Tunis, Brazil, England, and Italy.

Septoria tassiana Syd. (*S. antarctica* Tess.) Red-ochraceous leaf spots on *V. antarctica* in Italy.

Valsaria ampelina Avena. A wound parasite causing cankers of stems and twig die-back of *V. rupestris* and *V. vinifera* in Brazil.

VITTADINIA. Composite herbs with white, yellow, or blue flower heads.

Puccinia vittadeniae MeAlp. Yellow and brown rust pustules on leaves of *V. australis* in Victoria.

VOCHYSIA. Resinous tropical trees.

Aecidium vochysiae P. Henn. Leaf rust on *Vochysia* sp. in Brazil.

Phylachora congregata Theiss. and Syd. Dull-black stromata on leaves of *V. elliptica* in Brazil.

Phylachora granulosa Lév. Black stromata on large leaf spots, yellow beneath, gray-brown above, on *Vochysia* sp. in Brazil.

Phylachora lehmanniana P. Henn. Black stromata on leaves of *V. lehmanni* in Colombia.

Phylachora vochysiae P. Henn. Black stromata on leaves of *V. haenkeana* in Brazil.

VOUACAPOUA. See Andira.**WARNERIA**. See Gardenia.**WATSONIA**. *MERIANA* Ag. BUGLE LILY. Summer-flowering bulbous herbs resembling *Gladiolus*.

Uromyces watsoniae Syd. Brown to dark-brown or black rust pustules on both leaf surfaces of *W. densiflora* in the Union of South Africa.

WEDELIA. Composite annual or perennial herbs or subshrubs.

Aecidium wedeliae-hispidae Diet. Leaf rust on *W. hispida* in Mexico.

Endophyllum wedeliae (Earle) W. and O. Yellow rust pustules on leaves of *W. trilobata* in Porto Rico, Haiti, and Trinidad.

Laestadia lorentzii Speg. Circular to angular pale-yellow leaf spots with ashen centers on *W. glauca* in Argentina.

Puccinia ecuadorensis Arth. Brown leaf rust on *Wedelia* sp. in Ecuador.

Puccinia wedeliae Mayor. Leaf rust on *W. trichostephia* in Colombia.

Septoria lorentzii Speg. Circular to angular leaf spots on *W. glauca* in Argentina.

Uredo nersiveda Syd. Leaf rust on *W. biflora* in the Philippines.

Uredo vicina Arth. Leaf rust on *W. jacquini* and *W. lanceolata* in Porto Rico and Trinidad.

Uredo wedeliae-biflorae Syd. Cinnamon-brown rust pustules on leaves of *W. biflora* in the Philippines.

Uromyces baccarinii Syd. Powdery dark-brown rust pustules on leaves of *Wedelia* sp. in Abyssinia.

Uromyces piauhensis P. Henn. Brown rust pustules on leaves of *W. reticulata*, *W. rugosa*, and *Wedelia* sp. in Brazil, Porto Rico, and Cuba.

Uromyces wedeliae P. Henn. Brown rust pustules on leaves of *W. prostrata* and *Wedelia* sp. in Japan, the Philippines, and Abyssinia.

WELWITSCHIA. *TUMBOA* Ag. Peculiar low, woody coniferous desert plants.

Ustilago welwitschiae Bres. Powdery dull-brown smut sori in the cone scales of *W. mirabilis* in south Africa.

WIGANDIA. Tall perennial herbs or subshrubs, used as bedding plants or grown in greenhouses.

Phyllosticta wigandiae Thum. On leaves of *W. imperialis* in south Europe.

WIKSTROMERIA. Ornamental shrubs and trees.

Melampsora yoshinagai P. Henn. Powdery yellow, then dark-brown, rust pustules on leaves of *W. cuneata*, *W. indica*, *W. japonica*, and *W. sikokiana* in Japan, Formosa, China, and India.

WISTERIA. *KRAUNHIA* Ag. Attractive woody climbers with pea-shaped flowers.

Ascochyta wistariae Tassi. Leaf spots on *W. sinensis* in Italy.

Bacterium montemartini Pav. Causes large irregular hollow spots on branches of *Wisteria* sp. in Italy.

Mycosphaerella wistariicola Turc. Causes a leaf spot of *W. chinensis* in Italy. Probably the perfect stage of *Ascochyta wistariae* Fl. Tass.

Oehropora kraunhaiae Diet. Yellow and brown rust pustules on leaves of *W. sinensis* and *W. floribunda* in Japan.

Phyllosticta affinis Tass. On leaves of *W. sinensis* in Italy.

Phyllosticta wistariae Sacc. Ochraceous leaf spots on *W. floribunda* (*W. brachybotrys*) and *W. sinensis* in Italy and France. Reported from Texas.

Septoria wistariae Brun. Irregular brown leaf spots on *W. sinensis* in France. Tharp has given the same name to a form on *W. sinensis* in Texas.

XANTHORHOEAE. GRASS TREE. Perennials with a thick woody caudex.

Coniothyrium xanthorrhoeae P. Henn. Pale-brown leaf spots on *X. gracilis* in Australia.

XANTHOSOMA. MALANGA. Yautia. Ornamental-leaved aroids, corms of some species edible.

Phyllosticta xanthosomatis Sacc. Subcircular dull-yellow leaf spots on *Xanthosoma* sp. in Mexico.

Phytophthora sp. A downy mildew is considered the cause of a wilt and soft white rot of the corms of *Xanthosoma* sp. in Hawaii and the Philippines. A similar parasite may be concerned in the wilt disease of *Xanthosoma* in Porto Rico. These diseases have not been studied in detail and the fungus present may be *P. colocasiae* Rac. See Colocasia.

Vasculomyces xanthosomae Ashby. This fungus was found in Jamaica in connection with a wilting and tuber rot of *X. sagittifolium*. The fungus discolors the vascular bundles and sets up a brown rot, cavities forming in the tissues. The leaves turn yellow.

Vermicularia xanthosomatis Sacc. Leaf spot on *X. sagittifolium* in China and the Philippines.

XANTHOXYLUM. See Xanthoxylum.**XERANTHEMUM**. Annual erect tomentose herbs.

Puccinia xeranthemi Mori. Brown rust pustules on leaves and stems of *X. annuum* in Italy.

XYLOPIA. Shrubs and trees, grown for their fruit and other products.

Aecidium xylopieae P. Henn. Leaf rust on *Zylopia* sp. in Brazil.

Diplochorella fertilissima Syd. Black stromata on leaves of *X. aethiopica* in Tanzania.

Phylachora zylopieae P. Henn. Irregular black stromata on large brown leaf spots on *X. grandiflora* in Peru.

Phyllosticta xylopieae Sacc. Subcircular white leaf spots with black raised margins on *Zylopia* sp. in Mexico.

Puccinia foveolata (B. and C.) P. Henn. (*Puccinia gregaria* Kze.) Dark-brown rust pustules on leaves of *X. grandiflora* and *Xylopia* sp. in Guatemala, Peru, Surinam, and Brazil.

XYLOSMA. Tropical trees and shrubs.

Phyllachora pittieri Speg. See Myrsine.

Phyllachora xylosmatis Speg. Smooth black stromata on leaves of *X. pubescens* in Argentina.

Phyllosticta tonduzi Speg. On leaves of *X. salzmanni* in Costa Rica.

Uredo recondita Speg. Leaf rust on *X. salzmanni* in Costa Rica.

YUCCA. Ornamental stiff-leaved plants.

Colletotrichum yuccae Pöhl. Anthracnose on leaves of *Y. filamentosa* in Italy.

Gloeosporium victoriosum Sacc. Leaf anthracnose of *Y. gloriosa* in Spain and Italy.

Sclerotium yuccae Roum. On *Yucca* sp. in France.

Septoria macrospora Alm. and Cam. On leaves of *Y. aloifolia* in Portugal.

ZANTEDESCHIA. CALLA. Perennial herbs grown for their ornamental corolla-like spathes.

Bacillus sp. A bacillus said to be distinct from *B. aroidae* Town. attacks *Zantedeschia* in Great Britain. Diseased plants stop developing, the tops turning yellow and wither, while roots and corms are attacked by a soft watery brown rot.

Gloeosporium callae Oud. Large yellow, then brown, leaf spots on *Z. aethiopica* in Holland.

Ramularia richardiae Kalchbr. On leaves of *Z. albomaculata* in the Union of South Africa.

Septoria callae (Lasch.) Sacc. Small, olivaceous leaf spots on *Z. palustris* in Germany.

ZANTHOXYLUM. PRICKLY ASH. Trees and shrubs grown for their handsome foliage and ornamental fruit.

Acanthothecium mirabile Speg. On leaves of *Zanthoxylum* sp. in Brazil.

Aecidium spissum Syd. Leaf rust on *Zanthoxylum* sp. in India.

Aecidium zanthoxyli-schinifolii Diet. Rust on yellow or brown spots on leaf blades and petioles of *Z. schinifolium* in Japan.

Bacterium citri Hassé. See Citrus.

Coleosporium zanthoxyli Diet. and Syd. Yellow rust pustules on leaves of *Z. ailanthoides*, *Z. bungei*, *Z. piperitum*, and *Z. schinifolium* in Japan, Formosa, and China.

Helicobasidium tanakae Miy. See Morus.

Phyllachora applanata Wint. Subcircular to elliptic, black stromata on leaves of *Zanthoxylum* sp. in Brazil.

Phyllachora tijucensis (Rehm.) Theiss. and Syd. Circular black stromata on dark-brown leaf spots on *Zanthoxylum* sp. in Brazil.

Phyllachora winteri Sacc. and Syd. Black stromata on leaves and leaf-sheaths of *Z. hiemale*, *Z. rhoifolium*, and *Zanthoxylum* sp. in Brazil.

Phyllachora zanthoxyli (Lév.) Cke. Black stromata on leaves of *Zanthoxylum* sp. in Java and Brazil.

Sorokina uleana Rehm. On leaves of *Zanthoxylum* sp. in Brazil.

Uredo asperata B. and C. Brown rust pustules on leaves of *Z. piperitum* in Japan.

Uredo fagaræ Syd. Leaf rust on *Z. (Fagara) nitidum* in Japan and Formosa.

ZEa. MAIZE. INDIAN CORN.

Dothiorella zeæ Foex and Berth. Grain, cobs, and glumes overgrown and rotted by mycelium of the fungus. On *Z. mays* in Indo-China.

Fusarium maydiperdum Bub. Causes a rot of cobs of *Z. mays* in Bulgaria.

Nigrospora panicl Zimm. See Triticum.

Phyllachora maydis Maubl. Circular to elongate slightly raised black stromata on leaves of *Z. mays* in Porto Rico, Cuba, and Mexico.

Physoderma maydis Miy. Numerous circular to elliptical or linear, then confluent, brown spots on culms, mid ribs, and base of husks of *Z. mays* in Japan. Possibly not distinct from *P. zeæ-maydis* Shaw, which occurs in India and the southern United States.

Puccinia pallescens Arth. See Tripsacum.

Rhizoctonia palida Matz. See Saccharum.

Sclerospora sp. A downy mildew of *Z. mays*, characterized by broad, pale-green stripes on the leaves, is reported from the Union of South Africa.

Sclerospora javanica Palm. Young plants remain stunted, with narrow, yellow or greenish-yellow leaves, with poorly developed root systems. Older plants show the yellow-striped leaves and in more advanced cases narrow, sharply defined yellow or brown stripes. The host is *Z. mays* in Java.

Sclerospora macrospora Sacc. See Triticum.

Sclerospora maydis (Rac.) Butl. The upper parts of infected plants turn white due to the disappearance of the chlorophyll in long streaks along the leaves. Growth of the plants is checked and the internodes are frequently shortened so as to give a bunched appearance. No grain is produced, the plants withering and dying before harvest time. In addition to *Zeæ*, *Coix lacrym-jobi*, and *Euchlaena mexicana* are attacked. This species is known only from India, although it has been confused with *S. javanica*, the Javan species, and reported from that country.

Sclerospora philippinensis Weston. This downy mildew is a very serious enemy of *Zeæ mays* in the Philippines, and together with *S. spontanea* would appear to be the limiting factor in corn production. Additional hosts are *Euchlaena luxurians* and *Holcus sorghum* and by inoculation *Saccharum spontaneum* and *Miscanthus japonicus*.

Plants of all ages are attacked. Young plants, when attacked, show at the base of the upper leaves two or three narrow linear stripes pale-yellow to white in color. Leaves developing above these are almost completely yellow or whitish, remain narrow, and become rigid instead of bending in normal manner. The leaf sheaths continue to develop and may inclose or grow beyond the stunted tassel. The plants remain dwarfed. The roots also become stunted. Such plants are either destroyed by secondary organisms (*Pythium*) or turn brown, wither, and die. Older plants when attacked show the same stripes, generally on the fourth or fifth leaves, but more extensive and broader. Subsequent leaves show the striping to an even greater degree, the last leaves being striped throughout their length. The leaves themselves are not altered in size but at times the midribs become brittle, break at the union with the sheaths, and hang straight down along the stems. The root systems are not affected. The tassels may appear normal, but there is a decreased pollen production. Ears are only partly produced, and then always abnormal. Malformations of many kinds occur, including fasciations and reduplications, torsions of the stem, and other abnormalities.

Sclerospora sacchari Miy. See Saccharum.

Sclerospora spontanea Weston. This species of downy mildew causes a disease not distinguishable macroscopically from that due to *S. philippinensis*. It appears to be limited to the southern part of the Philippine Islands, where it occurs naturally on *Z. mays*, *Saccharum officinarum*, and *S. spontaneum*. By artificial inoculations the fungus has been found capable of attacking in addition *Euchlaena luxurians* and *Miscanthus japonicus*.

Septoria maydis Schulz. and Sacc. On leaf sheaths of *Z. mays* in Spain and Bohemia.

Sphaerulina maydis P. Henn. Circular to angular brown leaf spots on *Z. mays* in Brazil.

Ustilago abortifera Speg. A smut involving ovaries and often rachides in dark-brown powdery spore masses. On *Z. mays* in Argentina. Probably not distinct from *U. maydis*.

[Undet.] An undetermined fungus is reported from South Africa as destroying grain of *Z. mays* during germination.

ZENOBIÆ. Ornamental evergreen shrubs.

Phyllosticta andromedæ West. See *Pieris*.

ZEPHYRANTHES. ATAMOSCO Ag. ZEPHYR LILY. Tunicate bulbous herbs.

Aecidium zephyranthis Shear. Yellow leaf rust on *Zephyranthes* sp. in Mexico.

Septoria psittacina F. Tassi. Circular to irregular yellow leaf spots, which often become confluent, on both leaf surfaces of *Z. lindleyana* in Italy.

ZINGIBER. GINGER. Perennial rhizomatous herbs.

Bacillus zingiberi Uyeda. Reported as the cause of a soft rot of the rhizomes of *Z. officinale* in Japan. Infected plants turn yellow and wither.

Mycosphaerella zingiberi Shir. and Hari. On leaves of *Z. mioga* in Japan.

Phyllosticta zingiberi Shir. and Hari. On leaves of *Z. officinale* in Japan.

Piricularia zingiberi Nish. Leaf spot on *Z. mioga* and *Z. officinale* in Japan.

Pythium butleri Subra. The leaves of infected plants turn yellow and die, hanging down parallel to the stem, due to a soft rot which works into the bases of the shoots and down into the rhizomes. The latter are completely destroyed. On *Z. officinale* in India.

Taphrina maculans Butl. See *Curcuma*.

Vermicularia zingiberæ. Small circular to oval yellow spots on both leaf surfaces of *Z. officinale* in India. The spots increase in size and become coalescent, forming large discolored patches. The centers often dry up and drop out. The leaf sheaths and scaly portions of the stems are similarly attacked. The disease causes reductions in yield.

ZINNIA. Annuals grown in gardens for their flowers.

Ascochyta zinniae Allesch. Irregular, often confluent, dark-brown leaf spots on *Z. elegans* in Germany.

Corticium sp. See *Thea*.

Phyllosticta zinniae P. Brun. Angular to irregular brown, then olivaceous, leaf spots on *Z. elegans* in France.

Puccinia melampodii Diet. and Holw. Powdery gray-black rust sori on irregular brown sunken leaf spots on *Z. tenuiflora*, *Eleutheranthera ruderalis*, *Synedrella nodiflora*, *Tridax procumbens*, and *Spilanthes oleracea* in Mexico, West Indies, and Central America.

ZIZANIA. WILD RICE. Aquatic grasses.

Uromyces coronatus Yosp. Brown rust pustules on leaves of *Z. aquatica* in Japan.

Ustilago esculenta P. Henn. Smut sori in the heads of *Z. aquatica* and *Z. latifolia* in China, Japan, Formosa, and Russia. These smut masses in the young condition are sold for food in China.

ZIZYPHUS. JUJUBE. Shrubs and small trees grown for their handsome foliage and some species for edible fruit.

Ascochyta ziziphi Putt. Circular, then irregular dark-brown leaf spots on *Z. spina-christi* in Brazil.

Cercospora ziziphi Petch. Yellow, then brown, leaf spots on *Z. oenoplia* in Ceylon.

Cladosporium ziziphi Karst. and Roum. Leaf spot on *Z. jujuba* in India and Amboina.

Cronartium ziziphi Syd. and Butl. Dull-brown rust pustules on leaves of *Z. oenoplia* and *Z. rugosa* in Ceylon, India, and Indo-China.

Hyalodema evansi P. Magn. Galls on stems and petioles of *Z. baclei* and *Z. mucronata* in the Union of South Africa. The fruits are also deformed.

Phakospora ziziphi-vulgaris (P. Henn.) Diet. Black rust pustules on leaves of *Z. jujuba*, *Z. rotundifolia*, and *Z. sativa* (*Z. vulgaris*) in India, Formosa, and Japan.

Phyllosticta ziziphi Thuem. Irregular gray-ochraceous leaf spots on *Z. vulgaris* in Portugal.

Septoria capensis Wint. Gray to grayish-white leaf spots on *Z. mucronata* in the Union of South Africa.

Septoria jujubæ Trav. and Spessa. Small irregular ochraceous to dull-brown leaf spots on *Z. jujuba* in Portugal.

Septoria ziziphi Sacc. Pale ochraceous to gray-white leaf spots on *Z. vulgaris* in Spain and Italy.

Uredo belamcandæ P. Henn. Brown leaf rust on *Z. jujuba* in India.

Uredo ziziphi Pat. Brown leaf rust on *Z. spina-christi* and *Zizyphus* sp. in Indo-China, Abyssinia, and Brazil.

ZOYSIA. Low perennial grasses used for lawns and putting greens.

Puccinia zoysiae Diet. Powdery black rust pustules on leaves of *Z. pungens* in Japan.

ZYGOPHYLLUM. Shrubs, sometimes cultivated.

Aecidium kuntzii Kaernb. and Wint. Leaf rust on *Z. atriplicoides* in Turkestan.

Aecidium lagena Sorok. A rust deforming leaves of *Zygophyllum* sp. in Central Asia.

Cercospora zygophylli S. Szembel. On leaves of *Z. fabago* in Russia.

Erysiphe taurica Lévl. See *Althæa*.

Septoria zygophylli Syd. Subcircular dull-brown, then gray-white, leaf spots with red-brown margins on *Z. fabago* in Germany.

Uredo zygophylli P. Henn. Brown rust pustules on leaves and stems of *Z. cornutum* and *Z. decumbens* in Algeria, Tunis, and Egypt.

Uromyces trollipii Kalchbr. and McO. Cinnamon-brown to black rust pustules on leaves of *Z. foetidum* in the Union of South Africa.

Uromyces vesiculosus Wint. Cinnamon-brown to black rust pustules on leaves, stems, and fruit of *Z. billardieri*, *Z. ammophilum*, and *Z. glaucescens* in Australia.

Anderson et al. [R.A.M. V. 563]

UNITED STATES DEPARTMENT OF AGRICULTURE



DEPARTMENT BULLETIN No. 1366



Washington, D. C.

April, 1926

CHECK LIST OF DISEASES OF ECONOMIC PLANTS IN THE UNITED STATES

By PAUL J. ANDERSON, *Pathologist in Charge, Connecticut Tobacco Experiment Station, and temporarily Associate Pathologist, Plant-Disease Survey*; ROYAL J. HASKELL, *Associate Pathologist in Charge of Plant-Disease Survey*; WALTER C. MUENSCHER, *Assistant Professor of Economic Botany, New York State College of Agriculture, and temporarily Associate Botanist, Plant-Disease Survey*; and CLARA J. WELD, *Assistant Pathologist*, JESSIE I. WOOD, *Junior Pathologist*, and G. HAMILTON MARTIN, *Assistant Pathologist, Office of Plant-Disease Survey and Pathological Collections, Bureau of Plant Industry*

CONTENTS

	Page		Page
Basis and scope of the list.....	2	Geographic areas specified in this list, with	
Host names.....	2	abbreviations.....	6
Names of pathogens.....	3	List of authorities.....	7
Common names of diseases.....	5	List of diseases.....	11
Distribution.....	5	Index to common names of hosts.....	109

The Office of Plant-Disease Survey and Pathological Collections has been collecting specimens and data regarding the occurrence and distribution of plant diseases in the United States for a number of years, but no attempt has heretofore been made to prepare a list of all the known diseases of economic importance in this country with their distribution. It is believed that such a list will be of great value to pathologists and others interested in plant diseases and their control, as it will provide a means of determining quickly and easily what diseases have been reported on our important economic plants and their present known distribution, with the name of the causal organism in the case of parasitic diseases and the common name of the disease. It is hoped also that the list will be helpful in bringing about greater uniformity and certainty in the use of both common and scientific names.

The list has been compiled from the office records, and much information and assistance have been given by members of the various pathological offices of the Bureau of Plant Industry and by some of the Plant-Disease Survey collaborators, to all of whom thanks are tendered.

BASIS AND SCOPE OF THE LIST

The data on which this list is based have come chiefly from four sources: (1) Reports of the Plant-Disease Survey collaborators in the different States, (2) specimens preserved in the Pathological Collections of the Bureau of Plant Industry, (3) the literature on plant pathology and mycology, and (4) the investigations of many specialists who have supplied information. The reports of collaborators have been used principally in the work on the geographic range of the various diseases. It will be seen at a glance that much more information is needed on this subject, and it is hoped that all who are in a position to add to our records will send reports with specimens. The records on specimens in the Pathological Collections have been used to obtain data regarding distribution. It has not been possible to verify the determinations of the fungi found in the herbarium, but all concerning which there was obvious doubt have been omitted. The material, however, is available for examination at any time in case question should arise regarding it. Records from the literature on the subject chiefly relate to additional information regarding the occurrence and geographic range of diseases.

The list is confined to the diseases of the principal economic plants of the United States, including most timber trees and many of the ornamentals. It has been the intention to include all economically important parasitic and nonparasitic diseases known to occur within the continental United States. The parasitic diseases are listed under each host alphabetically by causal organism. These are followed by the nonparasitic diseases and those for which no pathogen is known. These are indicated respectively by "nonpar." and "undet." Such troubles as winter injury, frost injury, and sun scald, which are liable to occur on any plant, have been omitted. There are no doubt other diseases in the literature which have been overlooked, and it is hoped that when such are observed they will be promptly reported.

It will be noted that the information concerning the geographic distribution of many diseases is obviously incomplete. It is hoped, however, that this preliminary list may stimulate more interest in the subject and lead all who have an opportunity to add to our information to do so. It is particularly important to have plant-disease material and specimens to verify the occurrence of new and little-known diseases from different localities.

HOST NAMES

The scientific and common names of the hosts included in this list are those used in Standardized Plant Names,¹ and no authority is cited in connection with these names. Occasionally, necessary exceptions which occur in the list are shown by the use of an asterisk.

The diseases of each host are listed separately whenever practicable, but no attempt has been made to separate the diseases of the varieties of the host species, except in a few cases where such varieties include rather distinct crop plants; for example, cabbage, cauliflower, kale, kohlrabi, etc., under *Brassica oleracea*.

¹ American Joint Committee on Horticultural Nomenclature. Standardized Plant Names. 546 pp Salem, Mass. 1923.

In some of the host genera, as, for example, *Quercus*, it has been impracticable to list the diseases for each species separately. In many genera, such as *Prunus* and *Rubus*, in which many of the records of our collaborators simply report a disease as occurring "on *Prunus*" or "on raspberry," it is not possible to give the particular species of the host; and in some genera the nomenclature of the species is so confused that it is almost impossible to determine what host the collaborator had in mind unless the authority for the species is given or a specimen supplied. For example, the name *Populus balsamifera* is sometimes applied to any one of a number of species of *Populus*. In some of the records the generic name of the host species only is given. In this connection it is desirable to urge upon collaborators the great importance of giving specific names of the host plants when sending reports and specimens, or accompanying the reports by specimens of the host as well as the fungus. This is essential in order that our records may be as complete and accurate as possible. The abbreviation "spp." after the generic name of a host means that all diseases reported on any species of the genus are listed and considered in recording the distribution of the diseases—for example, *Salix* spp. "Sp." after the generic name indicates that the diseases listed were reported largely on an undetermined or doubtfully determined species of the genus—for example, *Begonia* sp.

In certain host genera the diseases are listed separately for one or more of the common or important species, following which the diseases of all other species of the genus are grouped under "sp." or "spp." In such cases the diseases and their distribution do not necessarily include the diseases or distribution as listed under specific host species—for example, *Juniperus virginiana* and *Juniperus* spp.

NAMES OF PATHOGENS

As there is no general agreement at present among the mycologists and phytopathologists of the world in regard to the names of fungous parasites, it has been found necessary to adopt for the purposes of this list, as a rule, the names which are at present in most general use. For some large groups of parasitic fungi, of which more or less recent monographs are available and are largely used for reference by pathologists, it has seemed best to follow rather closely the nomenclature of such works.

Rusts.—The specific names of the rusts which appear in Volume VII of North American Flora have been adopted, except in a few cases where long usage has established a name so firmly that there seems to be no good reason for attempting to supplant it by another; for example, *Puccinia graminis* is used instead of *P. poculiformis* for the stem-rust of wheat and *P. coronata* instead of *P. rhamni* for the crown-rust of oats. Generic names of long standing and in general use, such as *Puccinia* and *Uromyces*, are also retained.

Smuts.—The names in this group are those of Clinton's monograph on the Ustilaginales in Volume VII of North American Flora.

Powdery mildews.—Salmon's monograph on the Erysiphaceæ in Volume 9 of the Memoirs of the Torrey Botanical Club has been followed in this group.

Downy mildews.—The papers of G. W. Wilson on the Peronosporales have been followed with the exception of the generic name

Rhysotheca, for which the common and well-established name *Plasmopara* is retained.

Polypores.—Of the wood-inhabiting polypores and other Hymenomyces, only those are included which are known to cause definite diseases or rots of living trees or timber and for which the symptoms have been described in pathological literature. Saprophytic species, or those whose economic importance has not been demonstrated, have been omitted. In the choice of names, the more common usage has been followed.

Bacteria.—The names used for the bacterial pathogens are those in most common use, as the bacteriologists apparently are not yet entirely agreed in regard to some recently proposed changes.




FIG. 1.—Geographic areas used in the list: New England, Great Plains States, upper Mississippi Valley, lower Mississippi Valley, Ohio Valley States, Middle Atlantic States, and South Atlantic States

Botrytis.—All pathogens of this genus, with a few exceptions, are here listed under the names *Botrytis* sp. or *Botrytis cinerea* Auct. It is probable that several distinct species are included under these names, but since there is no satisfactory basis now available for specific distinction, it seems best for present purposes to include all under these names, pending the appearance of a satisfactory revision of the genus.

Synonyms.—Synonyms are given in parentheses in certain pleomorphic species in which the perfect stage is known and in some other cases in which it seems desirable. In general, the name which has been applied to the perfect stage has been adopted.

COMMON NAMES OF DISEASES

The following principles, some of which have been adopted by the American Phytopathological Society,² have been used in selecting the common names in the present list. Many of these names have already been passed upon and approved by the society.

(1) First consideration has been given to names already in wide and general use.

(2) Names descriptive of symptoms have been selected where practicable.

(3) In general, the various diseases caused by a single organism on different hosts have been given the same common name.

(4) Many diseases which are not sufficiently well known or important to receive a descriptive name have been classed according to their major symptoms; for example, leaf-blight, root-rot, rust, and wilt.

(5) It is assumed that the generic name of the causal organism may be prefixed to such common names whenever it is deemed desirable. Such generic names have not, however, been repeated in this list.



FIG. 2.—Geographic areas used in the list: Western States, Missouri Valley States, Southern States, and Great Lakes States

It is hoped that as many of these common names as prove convenient and satisfactory for practical purposes will be generally adopted, in order that they may become fixed and definite in their application and thus secure uniformity and stability.

DISTRIBUTION

An endeavor has been made to give in a brief form the geographic range of each species as correctly as the available information will permit. Diseases reported from a few or widely scattered States are followed by lists of these States. Diseases which have been reported from a number of States in which the host crop is grown or grows naturally are indicated as "general." Diseases which have been reported from a large number of States and in different parts of the country are indicated as "widespread." If a disease occurs

² Shear, C. L. Report of the seventh annual meeting of the American Phytopathological Society. *In* *Phytopathology*, vol. 6, pp. 211-212. 1916.

in one or more restricted regions, the range is indicated by listing the region. The following list gives the States included in each of the regions commonly mentioned. The accompanying maps (figs. 1 to 4) show the outlines of the various regions.

GEOGRAPHIC AREAS SPECIFIED IN THIS LIST, WITH ABBREVIATIONS

New England: Maine (Me.), New Hampshire (N. H.), Vermont (Vt.), Massachusetts (Mass.), Connecticut (Conn.), Rhode Island (R. I.).

Middle Atlantic States: New Jersey (N. J.), Pennsylvania (Pa.), Delaware (Del.), Maryland (Md.), Virginia (Va.).

South Atlantic States: North Carolina (N. C.), South Carolina (S. C.), Georgia (Ga.), Florida (Fla.).

Southern States: Virginia (Va.), North Carolina (N. C.), South Carolina (S. C.), West Virginia (W. Va.), Tennessee (Tenn.), Georgia (Ga.), Florida (Fla.), Alabama (Ala.), Mississippi (Miss.), Louisiana (La.), Arkansas (Ark.), Oklahoma (Okla.), Texas (Tex.).




FIG. 3.—Geographic areas used in the list: Pacific Northwest, Cotton Belt, spring-wheat area, and the Southwest

Great Lakes States: Michigan (Mich.), Wisconsin (Wis.), Minnesota (Minn.).

Upper Mississippi Valley: Wisconsin (Wis.), Illinois (Ill.), Minnesota (Minn.), Iowa (Iowa), Missouri (Mo.).

Lower Mississippi Valley: Arkansas (Ark.), Mississippi (Miss.), Louisiana (La.).

Missouri Valley States: North Dakota (N. Dak.), South Dakota (S. Dak.), Nebraska (Nebr.), Iowa, Kansas (Kans.), Missouri (Mo.).

Ohio Valley States: Ohio, Indiana (Ind.), Kentucky (Ky.), West Virginia (W. Va.).

Middle West: Ohio, Indiana (Ind.), Illinois (Ill.), Michigan (Mich.), Wisconsin (Wis.), Minnesota (Minn.), Iowa, Missouri (Mo.), North Dakota (N. Dak.), South Dakota (S. Dak.), Nebraska (Nebr.), Kansas (Kans.).

Gulf States: Florida (Fla.), Alabama (Ala.), Mississippi (Miss.), Louisiana (La.), Texas (Tex.).

Great Plains States: North Dakota (N. Dak.), South Dakota (S. Dak.), Nebraska (Nebr.), Kansas (Kans.), Oklahoma (Okla.), Texas (Tex.), Montana (Mont.), Wyoming (Wyo.), Colorado (Colo.), New Mexico (N. Mex.).

Western States: Montana (Mont.), Wyoming (Wyo.), Colorado (Colo.), New Mexico (N. Mex.), Idaho, Utah, Nevada (Nev.), Arizona (Ariz.), Washington (Wash.), Oregon (Oreg.), California (Calif.).

Rocky Mountain States:

Northern—Montana (Mont.), Wyoming (Wyo.), Idaho.

Southern—Colorado (Colo.), New Mexico (N. Mex.), Utah, Nevada (Nev.), Arizona (Ariz.).

Pacific coast: Washington (Wash.), Oregon (Oreg.), California (Calif.).

Pacific Northwest: Washington (Wash.), Oregon (Oreg.), Idaho.

Southwest: Arizona (Ariz.), New Mexico (N. Mex.), California (Calif.).

Cotton Belt: North Carolina (N. C.), South Carolina (S. C.), Georgia (Ga.), Alabama (Ala.), Tennessee (Tenn.), Mississippi (Miss.), Louisiana (La.), Oklahoma (Okla.), Arkansas (Ark.), Texas (Tex.).

Spring-wheat area: North Dakota (N. Dak.), South Dakota (S. Dak.), Minnesota (Minn.).

East of Great Plains: All States east of and including the eastern part of North Dakota (N. Dak.), South Dakota (S. Dak.), Kansas (Kans.), Oklahoma (Okla.), Texas (Tex.).




FIG. 4.—Geographic areas used in the list: Pacific coast, northern Rocky Mountain States, southern Rocky Mountain States, Middle Western States, and Gulf States

It is hoped that all who use this list will report any errors or omissions observed and will send specimens and records of any diseases not reported here or from regions not specified in this list, in order that later editions may be made more complete and accurate than this one.

C. L. SHEAR,
*Senior Pathologist in Charge of Plant-Disease
Survey and Pathological Collections.*

LIST OF AUTHORITIES

The following is a complete list of the authorities used in the index, with the abbreviations of the names occurring most frequently:

Aderh. = R. Aderhold.
Alb. = Johannes Baptiste von Alber-
tini.
Allesch. = Andreas Allescher.
Alm. = J. V. d'Almeida.
Am. Auct. = of American authors.

Anderson, H. W.
Anderson, P. J.
Appel = O. Appel.
Arcangeli, G.
Arn. = G. Arnaud.
Arth. = J. C. Arthur.

- Arthur, John M.
 Atk. = George F. Atkinson.
 Auct. = of authors.
 Auer. = B. Auerswald.
 Baumler, J. A.
 Bain, S. M.
 Bak. = A. L. Bakke.
 Barclay, A.
 Barss = H. P. Barss.
 Barth. = E. T. Bartholomew.
 Bast. = H. C. Bastian.
 Batsch = August Johann Batsch.
 Beckm. = Paul Beckmann.
 Bellynck, A.
 Bereng. = A. de Berenger.
 Berk. = M. J. Berkeley.
 Berl. = A. N. Berlese.
 Bertero, C. G.
 Berth. = A. Berthold.
 Bessey, E. A.
 Bethel = E. G. Bethel.
 Biv. = A. de Bivona-Bernhardi.
 Bizz. = G. Bizzozero.
 Bjerk. = C. Bjerkander.
 Blox. = A. Bloxam.
 Bon. = H. F. Bonorden.
 Bonar, L.
 Bonquet, P. A.
 Bosc = Louis A. Guillaume Bosc.
 Br. = C. E. Broome.
 Braun = A. Braun.
 Braun, H.
 Brenckle = J. F. Brenckle.
 Bres. = Giacomo Bresadola.
 Briard, H.
 Briosi = G. Briosi.
 Brot. = Felix de Avellar Brotero.
 Brown, J. G.
 Brown, N. = Nellie A. Brown.
 Brun. = P. Brunaud.
 Bryan, M. K. = Mary K. Bryan.
 Bub. = F. Bubak.
 Bull. = Pierre Bulliard.
 Burk. = W. H. Burkholder.
 Burr. = T. J. Burrill.
 Butl. = E. J. Butler.
 Cam. = M. d'Souza da Camara.
 Carpenter, C. W.
 Carter, C. N.
 Casp. = R. Caspary.
 Cast. = L. Castagne.
 Catt. = A. Cattaneo.
 Cav. = Fridiano Cava.
 Cda. = A. C. J. Corda.
 Ces. = Vincenzo Cesati.
 Charles, V. K.
 Chester = F. D. Chester.
 Chev. = F. Chevallier.
 Childs = Le Roy Childs.
 Chupp, C.
 Cke. = M. C. Cooke.
 Clapp, G. L.
 Clausen, R. E.
 Clements, F. E.
 Clint. = G. P. Clinton.
 Cobb = N. A. Cobb.
 Coerper = Florence M. Coerper.
 Cohn = Ferdinand Julius Cohn.
 Colby, A. S.
 Com. S. A. B. = Committee of the Society of American Bacteriologists.
 Coons, G. H.
 Cordley, A. B.
 Coville = F. V. Coville.
 Crouan, P. L.
 Currey, F.
 Curt. = M. A. Curtis.
 Dan. = P. A. Dangeard.
 Dana, B. F.
 Davis = J. J. Davis.
 D By. = Anton de Bary.
 DC. = Augustin Pyramus de Candolle.
 Dearn. = John Dearness.
 Del. = G. Delacroix.
 DeLac. = l'Abbé S. de Lacroix.
 DeNot. = G. de Notaris.
 DeSeyn. = J. de Seynes.
 Desm. = J. B. H. J. Desmazieres.
 DeToni = G. B. de Toni.
 Died. = H. Diedicke.
 Diet. = P. Dietel.
 Dodge, B. O.
 Doidge = E. M. Doidge.
 Dorsett, P. H.
 Drechs. = Charles Drechsler.
 Duby = J. E. Duby.
 Dug. = B. M. Duggar.
 Dur. = Durieu de Maisonneuve.
 Durand, E. J.
 Earle = F. S. Earle.
 Eddy, E. D.
 Edg. = C. W. Edgerton.
 Edson, H. A.
 EFS. = Erwin F. Smith.
 Ehr. = C. G. Ehrenberg.
 Eidam = E. Eidam.
 Ell. = J. B. Ellis.
 Elliott, Charlotte.
 Elliott, J. A.
 Engelm. = George Engelmann.
 Enlows, = Ella M. A.
 Eriks. = J. Eriksson.
 Ev. = B. M. Everhart.
 Ezekiel, Walter N.
 Fairman = C. E. Fairman.
 Farl. = W. G. Farlow.
 Fautr. = F. Fautrey.
 Fawc. = H. S. Fawcett.
 Fekl. = L. Fuckel.
 Field, Ethel.
 Fisch. = A. Fischer von Waldheim.
 Fischer, Ed. = Eduard Fischer.
 Fitz. = H. M. Fitzpatrick.
 Foster, A. C.
 Fr. = Elias M. Fries.
 Frank, A. B.
 Fresen. = George Fresenius.
 Fromme, F. D.
 Frost, C. C.
 Gall. = B. T. Galloway.
 Gardner, M. W.
 Garovaglio, Santo.
 Gerard, W. R.
 Gidd. = N. J. Giddings.
 Gies. = Karl Giesenhagen.

- Gill. = Claude Casimir Gillet.
 Gloyer, W. O.
 Godfrey, G. H.
 Godkin, James.
 Goodding, Leslie N.
 Graves, A. H.
 Gray, S. F.
 Greef = R. Greef.
 Gregory, C. T.
 Griff. = David Griffiths.
 Griffin, F. L.
 Gross. = J. G. Grossenbacher.
 Grout, A. J.
 Gues. = H. T. Guessow.
 Hahn, G.
 Hall, J. G.
 Hall, van = Constant Johan Jacob van Hall.
 Hals. = Byron D. Halsted.
 Hanzawa, J.
 Hara, K.
 Hark. = H. W. Harkness.
 Harter = L. L. Harter.
 Hartig = Robert Hartig.
 Harz, C. O.
 Hasse = Clara Hasse.
 Heald = F. D. Heald.
 Hedge. = G. G. Hedgecock.
 Hedges = Florence Hedges.
 Hedw. f. = R. A. Hedwig.
 Hemmi, Takewo.
 Henn. = Ernst Henning.
 Henn., P. = Paul Hennings.
 Hesse = Rudolf Hesse.
 Hig. = B. B. Higgins.
 Hoehn. = F. von Hoehnel.
 Holw. = E. W. D. Holway.
 Hopkins, E. F.
 Hudson, William.
 Humph. = J. E. Humphrey.
 Hunt = N. Rex Hunt.
 Jaap, Otto.
 Jack. = H. S. Jackson.
 Jacq. = Nicholas Joseph Jacquin.
 Jacz. = Arthur de Jaczewski.
 Jagger, I. C.
 James, G. F.
 Jamieson, C.
 Jehle = R. A. Jehle.
 Jenkins, A.
 Jens. = J. L. Jensen.
 Johans. = C. J. Johanson.
 Johnson, A. G.
 Johnson, James.
 Johnson, T.
 Jones, F. R.
 Jones, L. R.
 Juel = H. O. Juel.
 Kab. = J. E. Kabat.
 Karst. = P. A. Karsten.
 Kauffman, C. H.
 Kell. = W. A. Kellerman.
 Kendrick, J. B.
 Kern = F. D. Kern.
 Khazanoff, A.
 King = Charlotte King.
 Kirchner, L.
 Kleb. = H. Klebahn.
 Klotzsch, J. F.
 Koern. = F. Koernicke.
 Krueger, W.
 Kuehn = J. Kuehn.
 Kuntze, O.
 Kze. = G. Kunze.
 Laf. = H. A. Lafferty.
 Lagh. = G. von Lagerheim.
 Lambert, C. R.
 Langlois, A. B.
 Lasch = W. G. Lasch.
 Leb. = Hermann Lebert.
 Lee, H. A.
 Leonian, L. H.
 Lév. = J. H. Léveillé.
 Lewis, I. M.
 Lib. = Marie Anne Libert.
 Lindau = G. Lindau.
 Lindr. = J. I. Lindroth.
 Liro, J. I.
 Lk. = Heinrich Friedrich Link.
 Long = W. H. Long.
 Ludwig, C. A.
 Ludwig, F.
 Luedi, R.
 Lyon, H. L.
 Mackie, W. W.
 Magn. = P. Magnus.
 Mains, E. B.
 Manns, T. F.
 Marignoni, G. B.
 Mart. = George Martin.
 Martius, C. = C. F. P. von Martius.
 Mass. = George Massee.
 Matz, J.
 Maub. = A. Maublanc.
 McAlp. = D. McAlpine.
 McC. = Lucia McCulloch.
 McClain, R. L.
 Meier, F.
 Meinecke, E. P.
 Melhus, I. E.
 Metcalf, H. A.
 Miles, L. E.
 Miyabe = Kingo Miyabe.
 Miyake = Tehiro Miyake.
 Molliard, M.
 Mont. = J. F. C. Montagne.
 Morgan, Andrew Price.
 Moug. = Jean Baptiste Mougeot.
 Muell. = C. Mueller.
 Munn, M. T.
 Murr. = W. A. Murrill.
 Murray, T. J.
 Nees = Christian Gottfried Nees von Esenbeck.
 Neg. = Friedrich Wilhelm Neger.
 Niessl = G. von Niessl.
 Nishida, T.
 Noack, F.
 Norton, J. B. S.
 Nutt. = Thomas Nuttall.
 O'Gara, P. J.
 Osner, George.
 Otth = G. H. Otth.
 Oud. = C. A. J. A. Oudemans.
 Overh. = L. O. Overholts.
 Pam. = L. H. Pammel.

- Pass. = Giovanni Passerini.
 Pat. = Narcisse Patouillard.
 Patterson, F. = Flora Wambaugh
 Patterson.
 Paxton, G. E.
 Pegl. = V. Peglion.
 Penz. = O. Penzig.
 Perc. = C. Percival.
 Pers. = Christian Hendrik Persoon.
 Peterson, A.
 Peth. = G. H. Pethybridge.
 Petr. = Franz Petrak.
 Phill. = William Phillips.
 Pierce, Newton B.
 Piper = C. V. Piper.
 Pk. = C. H. Peck.
 Plow. = C. B. Plowright.
 Poit. = A. Poiteau.
 Poll. = Gino Pollacci.
 Porte, W. S.
 Posey = G. B. Posey.
 Preuss = C. G. T. Preuss.
 Prill. = Ed. Prilleaux.
 Pritchard, F. J.
 Quel. = Lucien Quélet.
 Rabh. = G. L. Rabenhorst.
 Raciborski, M.
 Ramsey, G. B.
 Rand, F. V.
 Rankin, W. H.
 Rav. = H. W. Ravenel.
 Ravaz = L. Ravaz.
 Reade, J. M.
 Reb. = J. F. Rebentisch.
 Reddick, D.
 Reddy, C. S.
 Regel, E.
 Rehm = H. Rehm.
 Reinke = Johannes Reinke.
 Ricker, P. L.
 Riess, H.
 Ritzema Bos, J.
 Rob. = M. Roberge.
 Roberts, J. W.
 Rose, D. H.
 Rosen, H. R.
 Rosenbaum, J.
 Rosendahl, C. O.
 Rostew. = S. I. Rostewzew.
 Rostr. = E. Rostrup.
 Roum. = C. Roumeguère.
 Roze = E. Roze.
 Rudolph, A. B.
 S. A. B. (Com.) = Committee of
 Society of American Bacteriologists.
 Sacc. = P. A. Saccardo.
 Sacc., D. = D. Saccardo.
 Sackett, W. G.
 Sadeb. = R. E. B. Sadebeck.
 Salm. = E. S. Salmon.
 Scalia, G.
 Schaeff. = J. C. Schaeffer.
 Schilb. = K. Schilbersky.
 Schl. = Friedrich Franz Leonhard von
 Schlechtendahl.
 Schm. = J. K. Schmidt.
 Schmidt, A.
 Schneider, A.
 Schrad. = Heinrich Adolph Schrader.
 Schrank = E. P. von Schrank.
 Schrenk = Hermann von Schrenk.
 Schroet. = J. Schroeter.
 Schultz, E. S.
 Schulz. = S. Schulzer von Muggen-
 burg.
 Schum. = Christian Friedrich Schu-
 macher.
 Schw. = Lewis David von Schweinitz.
 Scrib. = F. L. Scribner.
 Seaver = F. J. Seaver.
 Seym. = A. B. Seymour.
 Shaw, F. J. F.
 Shear = C. L. Shear.
 Sheld. = J. L. Sheldon.
 Sherb. = C. D. Sherbakoff.
 Sideris, C. P.
 Siegler, E. A.
 Slagg, C. M.
 Smith, C. O.
 Smith, E. H.
 Smith, R. E.
 Smith, W. G.
 Southworth, E. A.
 Sow. = J. Sowerby.
 Spauld. = Perley Spaulding.
 Speg. = C. Spegazzini.
 Stein. = J. G. Steinbuch.
 Stevens, F. L.
 Stewart, F. C.
 Stewart, V. B.
 Ston. = Bertha M. Stoneman.
 Stone, R. E.
 Sw. = Walter T. Swingle.
 Syd. = H. and P. Sydow.
 Tak. = Yoshiro Takahashi.
 Tassi, F.
 Taub. = J. J. Taubenhaus.
 Tehon, L. R.
 Tharp = B. C. Tharp.
 Thax. = Roland Thaxter.
 Thom, Ch.
 Thomas, F.
 Thomas, R. C.
 Thore, Jean.
 Thuem. = F. von Thuemen.
 Tiegh. = P. E. L. van Tieghem.
 Tisdale, W. B.
 Tode = Heinrich J. Tode.
 Town. = C. O. Townsend.
 Tr. = S. M. Tracy.
 Trel. = W. Trelease.
 Trev. = V. Trevisan.
 Tub. = Karl Freiherr von Tubeuf.
 Tul. = L. R. Tulasne or C. and L. R.
 Tulasne.
 Turconi, M.
 Underw. = Lucien Marcus Under-
 wood.
 Ung. = F. Unger.
 Vahl = Martin Vahl.
 Van Breda de Haan = J. van Breda de
 Haan.
 Van Hall = Constant Johan Jacob van
 Hall.
 Vester. = T. Vestergren.
 Viala = P. Viala.

Vogl. = P. Voglino.	Wils. = G. W. Wilson.
Waite, M. B.	Wilson, M.
Wakker, J. H.	Wingard, S. A.
Wallr. = Karl Friedrich Wilhelm Wallroth.	Wint. = G. Winter.
Weber, G. F.	Wolf = F. A. Wolf.
Wehmer, C.	Woll. = H. W. Wollenweber.
Weir = J. R. Weir.	Wor. = M. S. Woronin.
Welch, D. C.	Wormald, H.
Went, F. A.	Wrob. = A. Wroblewski.
West. = G. D. Westendorp.	Wulf. = Franz Xavier von Wulfen.
Wetts. = Richard von Wettstein.	Young, E.
Whet. = H. H. Whetzel.	Young, P. A.
Wilcox, E. M.	Zeller, S. M.
Williamson, M. M.	Zinssmeister, C. L.
	Zopf = Friederich Wilhelm Zopf.

LIST OF DISEASES

[An asterisk (*) shows that the name of the host given differs from that appearing in Standardized Plant Names; nonpar. = nonparasitic; undet. = undetermined.]

ABIES.

A. amabilis, cascade fir.

- Calyptospora columnaris* (Alb. and Schw.) Kuehn, leaf blister rust. Wash.
Dimerosporium sp., sooty mold. Wash.
Echinodontium tinctorium Ell. and Ev., brown stringy rot. Wash., Oreg.
Fomes annosus Fr., white spongy rot. Wash.
F. pinicola Fr., brown crumbly rot. General with host.
Hypoderma robustum Tub., needle cast. Oreg.
Hypodermella abietis-concoloris Dearn., needle cast. Oreg.
Melampsorella elatina (Alb. and Schw.) Arth.
Polyporus schweinitzii Fr., red-brown butt-rot. Oreg.
Pucciniastrum pustulatum (Pers.) Diet., leaf blister rust. Wash.
Razoumofskyia douglasii (Engelm.) Piper, mistletoe. Wash.
R. occidentalis abietina (Engelm.) Coville, mistletoe witches'-broom. Oreg.
Uredinopsis copelandii Syd., leaf blister rust. Wash., Oreg.
U. pteridis Diet. and Holw., rust. Wash.

A. balsamea, balsam fir.

- Calyptospora columnaris* (Alb. and Schw.) Kuehn, leaf blister rust. Northern States.
Fomes pinicola Fr., brown crumbly rot. N. Y.
F. roseus (Alb. and Schw.) Fr., brown pocket rot. General within range of host.
Lophodermium nervisequum (DC.) Rehm, needle cast. Adirondack region.
Melampsora americana Arth., rust. Vt., Mich., Wis.
Melampsorella elatina (Alb. and Schw.) Arth., rust witches'-broom. New England, N. Y., Lake States.
Nectria cucurbitula Fr., bark canker. N. Y., Pa.
Polyporus schweinitzii Fr., red-brown butt-rot. General with host.
Pucciniastrum pustulatum (Pers.) Diet., leaf blister rust. Wis., Mich.
Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Common.
Uredinopsis sp. (*Peridermium balsameum* Pk.), leaf blister rust. N. H., N. Y., Pa., Wis., Minn., Wash.

A. concolor, white fir.

- Calyptospora columnaris* (Alb. and Schw.) Kuehn, leaf blister rust. Oreg., Calif.
Echinodontium tinctorium Ell. and Ev., brown stringy rot. Common in Western States.
Fomes officinalis Fr., reddish brown-rot. Western States.
F. pinicola Fr., brown crumbly rot. General in range of host.
Herpotrichia nigra Hartig, matting. Calif.
Hypodermella abietis-concoloris Dearn., needle cast. Colo., Oreg.
Lophodermium nervisequum (DC.) Rehm, needle cast. Oreg., Calif.
Melampsora americana Arth., rust. Colo.
Peridermium ornamentale Arth., rust. Oreg., Calif.

ABIES—Continued.

A. concolor, white fir—Continued.

- Phacidium infestans* Karst., needle blight. Idaho.
Polyporus schweinitzii Fr., red-brown butt-rot. General with host.
Pucciniastrum pustulatum (Pers.) Diet., rust. Colo.
Razoumofskyia douglasii abietina (Engelm.) Piper, mistletoe. Ariz. to Oreg.
R. occidentalis abietina (Engelm.) Coville, mistletoe witches'-broom. Utah, Oreg., Calif.
Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Calif.

A. grandis, great silver fir.

- Botrytis* sp., seedling twig-blight. Idaho.
Calyptospora columnaris (Alb. and Schw.) Kuehn, leaf blister rust. Mont.
Dimerosporium sp., sooty mold. Wash.
Echinodontium tinctorium Ell. and Ev., brown stringy rot. Mont., Idaho, Pacific coast.
Fomes officinalis Fr., reddish brown-rot. Idaho.
F. pinicola Fr., brown crumbly rot. Idaho, Wash.
Herpotrichia nigra Hartig, matting. Idaho.
Hydnum sp., hollow pocket rot. Idaho, Wash.
Lophodermium abietis Rostr., needle cast. Pacific Northwest.
L. nervisequium (DC.) Rehm, needle cast. Idaho.
Melampsora americana Arth., rust. Idaho, Wash., Oreg.
M. arctica Rostr., rust. Idaho, Mont.
Melampsorella elatina (Alb. and Schw.) Arth., rust witches'-broom. Pacific Northwest.
Peridermium rugosum Jack., rust. Wash., Oreg.
Phacidium infestans Karst., needle blight. Idaho.
Polyporus dryadeus (Pers.) Fr., heart-rot. Pacific Northwest.
P. schweinitzii Fr., red-brown butt-rot. General with host.
P. sulphureus (Bull.) Fr., brown-checked wood-rot.
Pucciniastrum pustulatum (Pers.) Diet., leaf blister rust. Idaho, Mont., Oreg., Wash.
Razoumofskyia douglasii (Engelm.) Piper, mistletoe. Pacific coast, Idaho.
R. occidentalis abietina (Engelm.) Coville, mistletoe witches'-broom. Idaho, Oreg., Calif.
Rhizina inflata (Schaeff.) Sacc., seedling blight. Mont.
Uredinopsis copelandii Syd., leaf blister rust. Pacific Northwest.
U. pteridis Diet. and Holw., rust. Mont., Idaho, Wash., Oreg., Calif.

A. lasiocarpa, alpine fir.

- Calyptospora columnaris* (Alb. and Schw.) Kuehn, leaf blister rust. Idaho, Mont., Wyo., Colo., Oreg.
Echinodontium tinctorium Ell. and Ev., brown stringy rot. Mont.
Fomes pinicola Fr., brown crumbly rot. General within range of host.
Herpotrichia nigra Hartig, matting. Colo., Wyo., Idaho.
Melampsora americana Arth., rust. Wash.
M. arctica Rostr., rust. Idaho, Mont.
Melampsorella elatina (Alb. and Schw.) Arth., rust witches'-broom. Mont., Wyo., Utah.
Peridermium ornamentale Arth., rust. Oreg.
Phacidium infestans Karst., needle blight. Idaho.
Polyporus schweinitzii Fr., red-brown butt-rot.
Pucciniastrum pustulatum (Pers.) Diet., leaf blister rust. Mont., Oreg., Colo., Idaho, Alaska.
Razoumofskyia douglasii (Engelm.) Piper, mistletoe. Colo., Mont.
R. laricis Piper, mistletoe witches'-broom. Mont.
Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Common with host.
Uredinopsis copelandii Syd., leaf blister rust. Colo.

A. magnifica, red fir.

- Calyptospora columnaris* (Alb. and Schw.) Kuehn, leaf blister rust. Calif., Oreg.
Echinodontium tinctorium Ell. and Ev., brown stringy rot. Calif.
Fomes officinalis Fr., reddish brown-rot. Calif.
F. pinicola Fr., brown crumbly rot. General within range of host.

ABIES—Continued.**A. magnifica**, red fir—Continued.

- Herpotrichia nigra* Hartig, matting. Calif.
Lophodermium nervisequium (DC.) Rehm, needle cast. Calif.
Polyporus schweinitzii Fr., red-brown butt-rot. General with host.
P. sulphureus (Bull.) Fr., brown-checked wood-rot.
Razoumofskya douglasii (Engelm.) Piper, mistletoe. Calif.
R. occidentalis abietina (Engelm.) Coville, mistletoe witches'-broom. Oreg., Calif.
Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Calif.

A. nobilis, noble fir.

- Echinodontium tinctorium* Ell. and Ev., brown stringy rot. Common in Western States.
Fomes pinicola Fr., brown crumbly rot. Common with host.
Hypoderma robustum Tub., needle cast. Oreg.
Hypodermella abietis-concoloris Dearn., needle cast. Oreg.
Polyporus schweinitzii Fr., red-brown butt-rot. General.
Pucciniastrum pustulatum (Pers.) Diet., rust. Oreg.
Razoumofskya douglasii (Engelm.) Piper, mistletoe. Wash.
R. occidentalis abietina (Engelm.) Coville, mistletoe witches'-broom. Oreg.
Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Common within range of host.
Uredinopsis pteridis Diet. and Holw., rust. Oreg.

A. veitchii, Veitch fir.

- Fusicoccum abietinum* (Hartig) Prill. and Del. Ohio.

ACACIA.**A. farnesiana**, sweet acacia.

- Ravenelia australis* Diet. and Neg., rust. Tex.
R. siliquae Long, rust. Tex., N. Mex.

ACER.**A. circinatum**, vine maple.

- Cytospora chrysosperma* (Pers.) Fr., canker. Mont.
Nectria galligena Bres., European canker. Oreg.
Septoria aceris-macrophylli Pk., leaf-spot. Wash.
S. circinata Ell. and Ev., leaf-spot. Wash., Oreg.

A. dasycarpum, silver maple.

- Fomes fomentarius* (L.) Fr., white sapwood rot.
F. igniarius (L.) Fr., white heart-rot. Vt., N. Y.
Gloeosporium acerinum West., leaf-blight. Kans., Mo.
G. apocryptum Ell. and Ev., leaf-blight. N. J.
Pholiotia adiposa Fr., wood-rot. Del.
Phoradendron flavescens (Pursh) Nutt., mistletoe. Ind.
Phyllactinia corylea (Pers.) Karst., powdery mildew. Wis.
Phyllosticta minima (Berk. and Curt.) Ell. and Ev., leaf-spot. General east of Great Plains.
P. minutissima Ell. and Ev., leaf-spot. Wis.
Polyporus delectans Pk. D. C.
Rhytisma acerinum (Pers.) Fr., tar-spot. General east of Great Plains.
Septogloeum acerinum (Pass.) Sacc., leaf-spot. Ala.
Septoria acerina Pk., leaf-spot. N. Y.
S. saccharina Ell. and Ev., leaf-spot. New England, N. Y., Wis.
S. salliae W. R. Gerard, leaf-spot. N. Y.
Taphrina sp., leaf blister. Mich.
Uncinula circinata Cke. and Pk., powdery mildew. General east of Great Plains.
Verticillium sp., wilt. Va.

A. glabrum, Rocky Mountain maple.

- Cylindrosporium consociatum* Dearn., leaf-spot. Pacific Northwest.
Cytospora chrysosperma (Pers.) Fr., canker. Mont., Pacific Northwest.
Fomes igniarius (L.) Fr., white heart-rot. Central States.
Phyllosticta minima (Berk. and Curt.) Ell. and Ev., leaf-spot. Rocky Mts. and Pacific Northwest.

ACER—Continued.

A. glabrum, Rocky Mountain maple—Continued.

- Septoria acerina* Pk., leaf-spot. Oreg.
S. circinata Ell. and Ev., leaf-spot. Colo., Calif.
S. saccharina Ell. and Ev., leaf-spot. Wash.

A. macrophyllum, Oregon maple.

- Cercospora aceris* Dearn. and Barth., leaf-spot. Wash., Oreg.
C. albopunctata Ell. and Ev., leaf-spot. Wash.
Fomes igniarius (L.) Fr., white heart-rot. Western States.
Nectria cinnabarina Fr., canker. Wash.
N. galligena Bres., European canker. Oreg.
Rhytisma acerinum (Pers.) Fr., tar-spot. Wash.
R. punctatum (Pers.) Fr., black-specked leaf-spot. Pacific coast.
Septoria aceris-macrophylli Pk., leaf-spot. Wash.
S. circinata Ell. and Ev., leaf-spot. Wash., Oreg.

A. negundo, boxelder.

- Cercospora negundinis* Ell. and Ev., leaf-spot. Nebr.
Cylindrosporium negundinis Ell. and Ev., leaf-spot. Nebr., Kans.
Pusarium negundi Sherb., red-stain. General with host.
Gloeosporium apocryptum Ell. and Ev., anthracnose. Wis., Ill., Kans., Okla.
G. negundinis Ell. and Ev., leaf-blight. Del., Tex.
Nectria cinnabarina Fr., canker. Ohio, Minn., N. Dak.
Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.
Phleospora aceris (Lib.) Sacc., leaf-spot. Calif.
Phyllosticta arida Earle, leaf-spot. Ala.
P. minima (Berk. and Curt.) Ell. and Ev., leaf-spot. Conn.
P. negundinis Sacc. and Speg., leaf-spot. Me., R. I., W. Va., Ala.
Pleurotus ostreatus Jacq., white line rot. N. Dak., S. Dak., Nebr.
P. ulmarius Bull., white sap-rot. Iowa, Kans.
Polyporus squamosus (Huds.) Fr., white strand-rot. Minn.
Rhytisma punctatum (Pers.) Fr., black-specked leaf-spot. N. Y.
Septogloeum acerinum (Pass.) Sacc., leaf-spot. Calif.
Septoria acerella Sacc., leaf-spot. Mo., Mont.
S. marginata Heald and Wolf, leaf-spot. Tex.
S. negundinis Ell. and Ev., leaf-spot. Mass., Utah.
S. samaræ Pk., leaf-spot. Colo.
Sphaeropsis albescens Ell. and Ev., twig-blight. S. Dak., Kans.
Stereum purpureum Pers., silver leaf. S. Dak.

A. negundo californicum, California boxelder.

- Phleospora aceris* (Lib.) Sacc., leaf-spot. Calif.
P. californica Ell. and Ev., leaf-spot. Calif.
Septoria marginata Heald and Wolf, leaf-spot. Tex.

A. nigrum, black maple.

- Gloeosporium saccharinum* Ell. and Ev., leaf-blight. Iowa.
Rhytisma punctatum (Pers.) Fr., black-specked leaf-spot. W. Va.
Taphrina sp., leaf blister. Ohio.

A. palmatum, Japanese maple.

- Nectria cinnabarina* Fr., canker. Conn., Pa.
Pseudographium sp., twig canker. Wash.

A. pennsylvanicum, striped maple, *moosewood.

- Fomes igniarius* (L.) Fr., white heart-rot. Vt., N. Y.
Gloeosporium acerinum West., leaf-blight. Vt.
G. apocryptum Ell. and Ev., anthracnose. Pa.
Phleospora aceris (Lib.) Sacc., leaf-spot. Pa.
Phyllosticta minutissima Ell. and Ev., leaf-spot. N. Y., Pa.
Rhytisma punctatum (Pers.) Fr., tar-spot. W. Va.
Septoria acerina Pk., leaf-spot. New England, N. Y., Pa.
Uncinula circinata Cke. and Pk., powdery mildew. N. Y.

A. platanoides, Norway maple.

- Gloeosporium apocryptum* Ell. and Ev., anthracnose. Me., N. Y.
Microstroma sp., leaf mildew. Ohio, Ill.

ACER—Continued.**A. platanoides**, Norway maple—Continued.*Nectria cinnabarina* Fr., canker. N. Y., N. J., Pa.*Phyllosticta minima* (Berk. and Curt.) Ell. and Ev., leaf-spot. Va., Mo.*P. platanoides* Sacc., leaf-spot. N. J.*Rhytisma acerinum* (Pers.) Fr., tar-spot. N. Y.*Verticillium* sp., wilt. General east of Miss. River.**A. pseudoplatanus**, sycamore maple.*Phyllosticta minima* (Berk. and Curt.) Ell. and Ev., leaf-spot. Conn.*Verticillium* sp., wilt. Va.**A. rubrum**, red maple.*Cytospora chrysosperma* Fr., canker. Wash.*Fomes igniarius* (L.) Fr., white heart-rot. Mich.*F. roseus* (Alb. and Schw.) Fr., brown pocket rot. Md.*Gloeosporium aceris* Cke., leaf-blight. S. C.*G. decolorans* Ell. and Ev., leaf-blight. N. Y.*Nectria cinnabarina* Fr., canker. Va.*Phleospora aceris* (Lib.) Sacc., leaf-spot. Ind.*Phoradendron flavescens* (Pursh) Nutt., mistletoe. Ind.*Phyllosticta minima* (Berk. and Curt.) Ell. and Ev., leaf-spot. General east of Miss. River; Minn.*Rhytisma acerinum* (Pers.) Fr., tar-spot. General east of Miss. River.*Taphrina* sp., leaf blister. N. Y., N. C.*Uncinula circinata* Cke. and Pk., powdery mildew. General east of Miss. River.*Verticillium* sp., wilt. N. Y., Md., Va.**A. saccharum**, sugar maple.*Armillaria mellea* (Vahl) Quel., root-rot. N. Y.*Fomes applanatus* (Pers.) Wallr. (*Ganoderma applanatum* (Pers.) Pat.), white streaked rot. N. Y.*F. fomentarius* (L.) Fr., white sapwood rot. New England, N. Y.*F. igniarius* (L.) Fr., white heart-rot. N. Y.*F. pinicola* Fr., brown crumbly rot. N. Y.*Gloeosporium* sp., leaf-blight. N. Y.*G. apocryptum* Ell. and Ev., anthracnose. Pa.*G. hysterioides* Dearn. and Barth., leaf-blight. N. Y.*Hydnum septentrionale* Fr., white sapwood rot. Mass.*Ozonium omnivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.*Phoradendron flavescens* (Pursh) Nutt., mistletoe. Ind.*Phyllactinia corylea* (Pers.) Karst., powdery mildew. Iowa.*Phyllosticta minima* (Berk. and Curt.) Ell. and Ev., leaf-spot. Mass., Pa., Ind., Mo.*Polyporus sulphureus* (Bull.) Fr., brown-checked wood-rot. N. Y.*Rhytisma punctatum* (Pers.) Fr., tar-spot. Pa.*Taphrina* sp., leaf blister. Me., N. H., N. Y., Pa., Ga., Ind., Mich., Mo.*Uncinula circinata* Cke. and Pk., powdery mildew. N. Y., Ind., Iowa., Mo.*Verticillium* sp., wilt. Conn. south to Va. and Tenn.**A. spicatum**, mountain maple.*Fomes applanatus* (Pers.) Wallr. (*Ganoderma applanatum* (Pers.) Pat.), white-streaked rot. N. Y.*F. igniarius* (L.) Fr., white heart-rot. N. Y.*Phyllosticta minima* (Berk. and Curt.) Ell. and Ev., leaf-spot. N. Y., Wis.*P. platanoides* Sacc., leaf-spot. N. Y.*Rhytisma punctatum* (Pers.) Fr., black-specked leaf-spot. New England, N. Y., Pa.*Septoria acerina* Pk., leaf-spot. W. Va.*Uncinula circinata* Cke. and Pk., powdery mildew. N. Y.**ACHRAS.****A. sapota**, *sapote, sapodilla.*Pestalozzia scirrofaciens* N. Brown, hard-gall. Fla.

ACONITUM.

Aconitum sp., monkshood

Erysiphe polygoni DC., powdery mildew. W. Va.

Urocystis sorosporioides Koern., smut. Utah.

Uromyces aconiti Fekl., rust. Colo., Utah, Calif.

AESCULUS.

A. californica, California buckeye.

Exoascus aesculi (Ell. and Ev.) F. Patterson, witches'-broom. Calif.

Gloeosporium carpogenum Cke., leaf-blight. Calif.

Phoradendron villosum Nutt., mistletoe. Oreg., Calif.

Phyllactinia corylea (Pers.) Karst., powdery mildew. Calif.

Septoria aesculi (Lib.) West., leaf-spot. Calif.

A. glabra, Ohio buckeye.

Aecidium aesculi Ell. and Kell., rust. Nebr., Kans.

Guignardia aesculi (Pk.) V. B. Stewart, leaf-blotch. Miss. Valley and Ohio Valley.

Phoradendron flavescens (Pursh) Nutt., mistletoe. Eastern States.

Phyllosticta aesculicola Sacc., leaf-spot. Va., Tex., Mo.

Septoria glabrae Ell. and Ev., leaf-spot. Ind.

Uncinula flexuosa Pk., powdery mildew. Miss. Valley and Ohio Valley.

A. hippocastanum, horsechestnut.

Botryosphaeria ribis Gross. and Dug., blight. N. Y., D. C.

Collybia velutipes Curt., white sap-rot. R. I.

Glomerella cingulata (Ston.) Spauld. and Schrenk, anthracnose. Conn., N. Y.

Guignardia aesculi (Pk.) V. B. Stewart, leaf-blotch. General, especially east of Rocky Mts.

Nectria cinnabarina Fr., coral spot. New England, N. Y.

Septoria sp., leaf-spot. Pa.

Uncinula flexuosa Pk., powdery mildew. Mass., N. Y., N. J., Del., Ind.

A. octandra, yellow buckeye.

Guignardia aesculi (Pk.) V. B. Stewart, leaf-blotch. Tex., Iowa.

Phyllosticta aesculicola Sacc., leaf-spot. Tex.

Uncinula flexuosa Pk., powdery mildew. Ala.

A. pavia, red buckeye.

Guignardia aesculi (Pk.) V. B. Stewart, leaf-blotch. Ala., Miss.

Phyllosticta aesculicola Sacc., leaf-spot. Miss.

Uncinula flexuosa Pk., powdery mildew. Ala.

AGROPYRON.

**A. repens*, couchgrass, quackgrass.

Bacterium coronafaciens atropurpureum C. S. Reddy and James Godkin, bacterial blight. Wis., N. Dak.

Claviceps purpurea (Fr.) Tul., ergot. Occurs scatteringly throughout the range of host.

Colletotrichum graminicolum (Ces.) Wils., anthracnose. N. Y., Pa., Minn.; probably other States.

Erysiphe graminis DC., powdery mildew. N. Y., Pa., Mich., Minn., Mont.

Helminthosporium giganteum Heald and Wolf, leaf-spot. D. C., Md.

H. sativum Pam., King, and Bak., spot-blotch. Wis., N. Dak.

Ophiobolus graminis Sacc., take-all. N. Y.

Phyllachora graminis (Pers.) Fekl., tar-spot. General throughout range of host.

Phyllosticta sp., leaf-spot. Pa.

Puccinia clematidis (DC.) Lagh., leaf-rust. General.

P. coronata Cda., rust. Vt.

P. graminis Pers., stem-rust. General.

Pyrenophora tritici-repentis (Died.) Drechs., leaf-blight. Wis.

Septoria agropyri Ell. and Ev., leaf-spot. Wis.

Tilletia earlei Griff., covered smut. N. Dak.

Urocystis agropyri (Preuss) Schroet., smut. Vt., Mass., Conn., N. Y., Pa.

Ustilago hypodytes (Schl.) Fr., smut. N. Y.

U. macrospora Desm., smut. Mass., Wis., Iowa.

U. striaeformis (West.) Niessl, stripe-smut. N. Y.

AGROSTIS.**A. capillaris**, bentgrass.*Puccinia graminis* Pers., stem-rust. General east of Great Plains.*Scolecotrichum graminis* Fekl., leaf-spot. Ind.*Ustilago striaeformis* (West.) Niessl, stripe-smut. Conn., Miss. Valley.**A. palustris**, redtop.*Claviceps purpurea* (Fr.) Tul., ergot. Pa., Mich., Minn., N. Dak.*Colletotrichum graminicolum* (Ces.) Wils., anthracnose. Pa.*Corticium vagum* Berk. and Curt., root-rot. General.*Erysiphe graminis* DC., powdery mildew. Minn.*Helminthosporium dematoides* Bub. and Wrob., leaf-spot. Conn., N. Y.*Ophiobolus graminis* Sacc., take-all. N. Y.*Puccinia graminis* Pers., stem-rust. General east of Great Plains.*Pythium* sp., damping-off. Conn.*Sclerotium rhizodes* Auer., Sclerotium blight. Conn.*Scolecotrichum graminis* Fekl., leaf-spot. Pa., Wis.*Urocystis agropyri* (Preuss) Schroet., smut. Pa., Wis.*Ustilago striaeformis* (West.) Niessl, stripe-smut. General east of Great Plains.**A. stolonifera**, carpet bentgrass.*Corticium vagum* Berk. and Curt., root-rot. General.*Helminthosporium giganteum* Heald and Wolf, leaf-spot. D. C.*H. stenacrum* Drechs., leaf-spot. Conn.**AILANTHUS.****A. glandulosa**, tree-of-heaven.*Cercospora glandulosa* Ell. and Kell., leaf-spot. Ala., Ill., Kans., Nebr., Tex.*Nectria cinnabarina* Fr., canker. N. J., Kans.**ALLIUM.****A. ascalonicum**, shallot.*Colletotrichum circinans* (Berk.) Vogl., smudge. Ill., Wis.*Heterodera radicola* (Greef) Muell., root-knot.*Peronospora schleideni* Ung., downy mildew. La.**A. cepa**, onion.*Aspergillus niger* Tiegh., black-mold. General.*Bacillus carotovorus* L. R. Jones (*Erwinia carotovora* (Jones) Com. S. A. B.), soft-rot. General.*B. cepivorus* Del., rot. W. Va.*Botrytis* sp., gray-mold. General.*B. allii* M. T. Munn, neck-rot. N. Y., Ohio, Ill., Mich., Wis., Pacific Northwest.*B. byssoidea* Walker, mycelial neck-rot. Conn., Ill., Wis.*B. squamosa* Walker, small sclerotial neck-rot. Ill., Wis.*Colletotrichum circinans* (Berk.) Vogl., smudge. General east of Great Plains.*Corticium vagum* Berk. and Curt., stem-rot. Ohio, Wash.*Cuscuta* sp., dodder. Ill., Wis., Calif.*Fusarium* sp., root-rot. Frequent occurrence.*F. cepae* J. Hanzawa, bulb rot. Ill.*F. mali* Taub., pink-root. Occurs in widely scattered States.*Heterodera radicola* (Greef) Muell., root-knot. Tex., Ind.*Macrosporium porri* Ell., mold. New England, Miss., Ind., Utah.*M. sarcinula parasiticum* Thuem., black stalk-rot. General.*Peronospora schleideni* Ung., downy mildew. General.*Puccinia porri* (Sow.) Wint., rust. Conn.*Pythium debaryanum* Hesse, damping-off. Mass.*Sclerotium cepivorum* Berk., white-rot. Va., Ky., Oreg.*S. rolfii* Sacc., stem-rot. Ala.*Urocystis cepulae* C. C. Frost, smut. General.

ALLIUM—Continued.*A. sativum*, garlic.*Macrosporium sarcinula parasiticum* Thuem., black stalk-rot. La.*Sclerotium cepivorum* Berk., white-rot. Oreg.*S. rolfii* Sacc., stem-rot. La.*Tylenchus dipsaci* (Kuehn) Bast., bulb nematode. Calif.*A. schoenoprasum*, chives.*Puccinia porri* (Sow.) Wint., rust. Conn.**ALNUS.***A. incana*, speckled alder.*Cylindrosporium vermiforme* Davis, leaf-spot. Wis.*Erysiphe aggregata* (Pk.) Farl., powdery mildew. Me., N. Y., Wis.*Exoascus tosquinetii* Sacc., catkin deformation. Minn.*Hymenochaete agglutinans* Ell., stem girdling. Pa.*Phyllactinia corylea* (Pers.) Karst., powdery mildew. New England, N. Y., Wis.*Taphrina robinsoniana* Gies., catkin deformation. N. Y., Md., Va.**A. rhombifolia*, white alder.*Fomes igniarius* (L.) Fr., white heart-rot. Common in host range.*Melampsoridium alni* (Thuem.) Diet., rust. Calif.**A. rubra*, alder.*Cercospora alni* Dearn. and Barth., leaf-spot. Wash., Oreg.*Exoascus* sp., catkin deformation. Wash., Oreg.*Fomes applanatus* (Pers.) Wallr. (*Ganoderma applanatum* (Pers.) Pat.), white-streaked rot. Wash.*F. igniarius* (L.) Fr., white heart-rot. Common in host range.*Gnomonia alni* Plow., leaf-spot. Calif.*Gnomoniella tubiformis* (Tode) Sacc., black-spot. Oreg.*Melampsoridium alni* (Thuem.) Diet., rust. Calif.*Microsphaera alni* (Wallr.) Wint., powdery mildew. Wash., Oreg.*Phyllactinia corylea* (Pers.) Karst., powdery mildew. Wash.*Polystictus hirsutus* (Wulf.) Fr., white-rot.*Septoria alnifolia* Ell. and Ev., leaf-spot. Wash., Oreg.*A. rugosa*, hazel alder.*Erysiphe aggregata* (Pk.) Farl., powdery mildew. N. Y., W. Va.*Exoascus tosquinetii* Sacc., catkin deformation. Me., Mass., Ala.*Gnomoniella tubiformis* (Tode) Sacc., black-spot. Pa.*Microsphaera alni* (Wallr.) Wint., powdery mildew. Ala.*Mycosphaerella hippocastani* (Otto Jaap) Kleb., leaf-spot. N. Y.*Phyllactinia corylea* (Pers.) Karst., powdery mildew. Ala., Ohio.*Taphrina robinsoniana* Gies., catkin deformation. Md., Va.**A. sinuata*, alder.*Microsphaera alni* (Wallr.) Wint., powdery mildew. Alaska.*Phyllachora* sp., leaf-spot. Wash.*A. tenuifolia*, mountain alder.*Cylindrosporium* sp., leaf-spot. Mont.*Exoascus* spp., catkin deformation. Mont., Idaho, Oreg.*E. tosquinetii* Sacc., catkin deformation. Idaho.*Fomes igniarius* (L.) Fr., white heart-rot. Mont.*Gnomoniella tubiformis* (Tode) Sacc., black-spot. Oreg.*Phyllactinia corylea* (Pers.) Karst., powdery mildew. Colo., N. Mex., Idaho, Oreg.*Polystictus hirsutus* (Wulf.) Fr., white-rot.**ALTHAEA.***A. rosea*, hollyhock.*Alternaria* sp., leaf-spot. Del., Mich., Nebr.*Ascochyta althaeina* Sacc. and Bizz., leaf-spot. N. Y.*A. parastitica* Fautr., leaf-spot. Conn., Del., W. Va.*Cercospora althaeina* Sacc., leaf-spot. General east of Great Plains.*C. kellermanii* Bub., leaf-spot. Ind., Ohio.*C. nebulosa* Sacc., leaf-spot. Nebr.

ALTHAEA—Continued.*A. rosea*, hollyhock—Continued.

- Colletotrichum malvarum* (Braun and Casp.) E. A. Southworth, anthracnose. N. Y., Ohio, Ind., Iowa.
Heterodera radicum (Greef) Muell., root-knot.
Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.
Phyllosticta althaeina Sacc., leaf-spot. N. Y., W. Va., Ohio.
Puccinia hibisciata (Schw.) Kell., rust. Nebr.
P. malvacearum C. G. Bertero, rust. General.
P. sherardiana Koern., rust. Calif.
Septoria fairmani Ell. and Ev., leaf-spot. N. Y.

AMARYLLIS.*A. belladonna*, belladonna-lily.

- Botrytis* sp., rot. Calif.
Cercospora amaryllidis Ell. and Ev., leaf-spot. La.

AMELANCHIER.*Amelanchier* sp., shadblow (*serviceberry, *juneberry, *shadbush, *saskatoon)

- Bacillus amylovorus* (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. Wash.
Dimerosporium collinsii (Schw.) Thuem. General.
Erysiphe polygoni DC., powdery mildew. Mont.
Fabraea maculata (Lév.) Atk., leaf-blight. Ala., Ind., N. Dak., Kans., Mont., Idaho.
Gymnosporangium blasdaleanum (Diet. and Holw.) Kern, rust. Oreg., Calif.
G. botryapites (Schw.) Kern, rust. Mass., Conn., N. Y., N. J., Pa.
G. clavariaeforme (Jacq.) DC., rust. N. Y. to Del., west to Utah.
G. corniculans Kern, rust. Vt. to Iowa, south to W. Va.
G. germinale (Schw.) Kern, rust. New England; N. Y., Mich., Wis.
G. inconspicuum Kern, rust. Colo., N. Mex., Utah.
G. juvenescens Kern, rust. Northern Great Plains States; Wis., Idaho.
G. kernianum Bethel, rust. Oreg.
G. nelsoni Arth., rust. Rocky Mt. States; Nebr.
G. nidus-avis Thax., rust. Mass. to Wis.; south to N. C.
Nummularia discreta (Schw.) Tul., blister canker. Mass., Ohio, Kans.
Phyllactinia corylea (Pers.) Karst., powdery mildew. Wash.
Phyllosticta innumerabilis Pk., leaf-spot. Nebr.
P. paupercula Pk., leaf-spot. Kans.
P. virginiana (Ell. and Hals.) Seaver, leaf-spot. N. Y., Pa., Wis.
Plowrightia morbosa (Schw.) Sacc., black-knot. Minn., Mont.
Podosphaera oxycanthae (DC.) D By., powdery mildew. Mass., Minn., Mo., Nebr.
Sclerotinia amelanchieris J. M. Reade, fruit-rot, blight. N. Y.
S. gregaria B. F. Dana, blight. Wash.

AMMOPHILA.*A. arenaria*, beachgrass.

- Claviceps purpurea* (Fr.) Tul., ergot. Mich.
Ustilago striaeformis (West.) Niessl, stripe smut. New England Coast States.

AMORPHA.*A. fruticosa*, indigobush.

- Cercospora passaloroides* Wint., leaf-spot. Ala., Ill., Nebr., Kans.
Uropyxis amorphae (Curt.) Schroet., rust. General east of Rocky Mts.

AMPELOPSIS.*A. arborea*, peppervine.

- Cercospora arboriae* Tharp, leaf-spot. Tex.
C. viticola (Ces.) Sacc., leaf-spot. La.
Guignardia bidwellii (Ell.) Viala and Ravaz, black-rot. Miss.

A. cordata, heartleaf ampelopsis.

- Cercospora truncata* Ell. and Ev., leaf-spot. La.
C. viticola (Ces.) Sacc., leaf-spot. Ala., La.

AMPELOPSIS—Continued.

A. quinquefolia, Virginia creeper.

Cercospora ampelopsidis Pk., leaf-spot. General east of Great Plains.

C. psedericola L. R. Tehon, leaf-spot. Ill.

C. pustula Cke., leaf-spot. Occurs in widely scattered States east of Great Plains.

Cladosporium herbarum Lk., die-back. N. J.

Guignardia bidwellii (Ell.) Viala and Ravaz, black-rot. General east of Great Plains.

Phyllosticta viticola (Berk. and Curt.) Thuem., leaf-spot. General.

Plasmopara viticola (Berk. and Curt.) Berl. and DeToni, downy mildew. Me., Ala., Wis., Iowa.

Septogloeum ampelopsidis (Ell. and Ev.) Sacc., leaf-spot. Ill., Wis., Iowa, Nebr.

Septoria ampelopsidis Ell., leaf-spot. Nebr.

Ucinula necator (Schw.) Burr., powdery mildew. General east of Great Plains.

A. tricuspidata, *Boston ivy, Japanese creeper.

Cladosporium sp., die-back. N. J.

Guignardia bidwellii (Ell.) Viala and Ravaz, black-rot. Atlantic Coast States; Tex.

AMYGDALUS.

A. communis, almond.

Armillaria mellea (Vahl) Quel., root-rot. Calif.

Bacillus amylovorus (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. Mass.

Bacterium tumefaciens EFS. and Town. (*Phytoplasma tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Ariz., Calif.

Cercospora circumscissa Sacc., shot-hole. Oreg., Calif.

Cladosporium carpophilum Thuem., scab. Calif.

Coryneum beijerinckii Oud., blight. Oreg., Calif.

Cyphella marginata McAlp., die-back. Oreg.

Gloeosporium sp., fruit-spot and twig-spot. Calif.

Heterodera radicola (Greef) Muell., root-knot. Ariz.

Pythiactis citrophthora E. H. and R. E. Smith, brown-rot, die-back. Calif.

Sclerotinia fructicola (Wint.) Rehm (*S. americana* (Wormald) Norton and Ezekiel), brown-rot. Calif.

Tranzschelia punctata (Pers.) Arth., rust. Calif.

A. persica, peach.

Armillaria mellea (Vahl) Quel., root-rot. Miss., Tex., Ark., Ind., Oreg., Calif., Alaska.

Bacterium cerasi F. L. Griffin, gummosis. Oreg.

B. pruni EFS. (*Phytoplasma pruni* (EFS.) Com. S. A. B.), bacterial spot. General east of Rocky Mts.

B. tumefaciens EFS. and Town. (*Phytoplasma tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. General.

Botrytis cinerea Auct., fruit-rot. Calif.

Cephalothecium roseum Cda., pink-rot. Md., Ill., Calif.

Cercospora circumscissa Sacc., shot-hole. General.

C. consobrina Ell. and Ev., leaf-spot. La., Ill.

Cercospora persicae Sacc., frosty mildew. General.

Choanephora persicaria E. D. Eddy, fruit-rot. In storage.

Cladosporium carpophilum Thuem., scab. General.

Clitocybe monadelpha (Morgan) Sacc., root-rot. Fla., Ark., Mo.

Comandra pallida DC., bastard toadflax. Wash.

Coryneum beijerinckii Oud., blight. General, especially in Western States.

Cyphella marginata McAlp., die-back. Oreg.

Diplodia natalensis Ev., gumming. Fla.

Ezoascus deformans (Berk.) Fekl., leaf-curl. General.

Fomes pinicola Fr., brown crumbly rot. Oreg.

F. roseus (Alb. and Schw.) Fr., brown pocket rot. Va.

Fusarium spp., fruit-rot. Calif.

F. gemmiperda Aderh., bud-rot. Ga.

Glomerella cingulata (Ston.) Spauld. and Schrenk, bitter-rot. Fla., Ark., Ohio.

AMYGDALUS—Continued.*A. persica*, peach—Continued.

Helminthosporium carpophilum Lév., pustular spot. N. Y., Ohio, Mich., Utah.

Heterodera radicola (Greef) Muell., root-knot. South Atlantic and Gulf States; Calif.

Lenzites saepiaria Fr., heart-rot. Oreg.

Monilia sp., western brown-rot. Oreg.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex., N. Mex., Ariz.

Phoma persicae Sacc., stem canker. Mass., Conn., Pa., Del., Md., Va., Ohio, Mich.

Phyllosticta circumscissa Cke., leaf-spot. N. Y., N. J., Wash.

Physalospora malorum (Pk.) Shear (*P. cydoniae* Am. Auct.), black-rot. Del., Ind.

Podosphaera oxycanthae (DC.) D By., powdery mildew. General.

Polyporus lacteus Fr., sap-rot and heart-rot.

Pythiactystis citrophthora E. H. and R. E. Smith, die-back. Calif.

Rhizopus nigricans Ehr., black-mold rot. Conn., Del., Ga., Fla., Ark., Ill., Ind., Kans.

Sclerotinia cinerea (Bon.) Schroet. (*Monilia oregonensis* Barss and Posey), blossom blight. Pacific coast.

S. fructicola (Wint.) Rehm (*S. americana* (Wormald) Norton and Ezekiel), brown-rot. General.

Sclerotium rolfsii Sacc., stem-rot (of seedlings). La.

Sphaerotheca pannosa (Wallr.) Lév., powdery mildew. General.

Stereum purpureum Pers., silver leaf. N. Y., Md.

Trametes carnea (Nees) Cke., wood-rot. Oreg.

Tranzschelia punctata (Pers.) Arth., rust. General.

Valsa leucostoma (Pers.) Fr., die-back. General.

Verticillium alboatrum Reinke and Berth., wilt. N. Y., N. J.

Chlorosis, nonpar. Southwest.

Gummosis, various causes. General.

Little peach, undet. East of Miss. River, especially Middle Atlantic States.

Phoney peach, undet. Central and southern Ga.

Rosette, undet. Tenn., Fla., S. C., Ga., Ala., Miss., La., Okla., Ark., Mo., Kans.

Yellows, nonpar. East of Miss. River and north of Gulf States.

A. persica nectarina, nectarine.

Cladosporium carpophilum Thuem., scab. Conn., N. Y., Ill.

Eoascus deformans (Berk.) Fckl., leaf-curl. Wash.

Sclerotinia fructicola (Wint.) Rehm (*S. americana* Norton and Ezekiel), brown-rot. Conn., N. Y.

ANANAS.*A. sativus*, pineapple.

Spegazzinia ornata Sacc. Fla.

Thielaviopsis paradoxa (DeSeyn.) Hoehn., soft-rot. Fla.

Red wilt, undet. Fla.

ANCHUSA.*A. capensis*, Cape bugloss.

Puccinia dispersa Eriks., rust. Ind.

A. officinalis, common bugloss.

Puccinia dispersa Eriks., rust. Ind.

ANETHUM.*A. graveolens*, dill.

Cercospora anethi Sacc., leaf-spot. N. Dak.

ANNONA.*A. cherimola*, cherimoya.

Gloeosporium sp., anthracnose. Fla.

Uredo cherimoliae Lagh., rust. Fla.

A. squamosa, sugar-apple.

Uredo cherimoliae Lagh., rust. Fla.

ANT RHINUM.

A. majus, snapdragon.

Colletotrichum antirrhini F. C. Stewart, anthracnose. General east of Miss. River.

Corticium vagum Berk. and Curt., rot (of cuttings). N. Y.

Fusarium sp., root and crown rot. Wash.

Heterodera radicolica (Greef) Muell., root-knot. D. C., Miss., Nebr.

Phyllosticta antirrhini Syd., leaf-spot, damping-off. Mass., N. Y., Va., Ind., Minn.

Puccinia antirrhini Diet. and Holw., rust. General.

Pythium sp., stem-rot, damping-off. N. Y.

Thielavia basicola (Berk. and Curt.) Zopf, black root-rot. N. J.

Verticillium alboatrum Reinke and Berth., wilt. Mass.

APIUM.

A. graveolens, celery.

Bacillus carotovorus L. R. Jones (*Erwinia carotovora* (Jones) Com. S. A. B.), soft-rot. General.

Bacterium apii I. C. Jagger (*Phytomonas apii* (Jagger) Com. S. A. B.), bacterial leaf-spot. N. Y., Ind., Mich., Minn.

Botrytis sp., gray-mold rot. N. Y., Idaho, Calif.

Cercospora apii Fresen., early-blight. General.

Corticium vagum Berk. and Curt., root-rot. General.

Fusarium sp., pink-root. Fla.

F. sp., root-rot. N. J., Fla., Ohio, Calif.

F. sp., yellows. Conn., Mass., N. Y., N. J., Pa., Ohio, Mich., Ind., Minn.

Heterodera radicolica (Greef) Muell., root-knot. N. J., Va., N. C., S. C., Fla., Ark., Ind., Kans., Calif.

Phoma apiicola Speg., root-rot. Ohio, Mich.

Phyllosticta apii Hals., leaf-spot. N. J., Del.

Pythium debaryanum Hesse, damping-off. Conn.

Sclerotinia minor I. C. Jagger, drop. N. Y., Pa., Fla.

S. sclerotiorum (Lib.) Mass., foot-rot. General.

Septoria apii Rostr., late-blight. General.

Bacterial heart-rot. East of Great Plains.

Black-heart, undet. N. Y., Pa., Fla., La., Calif.

Crinkle, nonpar. Pa.

Hollow-stem, nonpar. Pa., Calif.

Mosaic, undet. Mass., N. Y., N. J., Ind., Minn.

**A. graveolens rapaceum*, celeriac.

Cercospora apii Fresen., early-blight. Conn.

Phoma apiicola Speg., root-rot. N. Y.

Septoria apii Rostr., late-blight. Conn., N. Y.

AQUILEGIA.

Aquilegia spp., columbine.

Erysiphe polygoni DC., powdery mildew. Conn., N. Y., Ind., Ohio, Ill., Wis.

Gloeosporium aquilegiae Thuem., leaf-spot. Tex.

Puccinia clematidis (DC.) Lagh., rust. Ind., Mont., Colo., Utah, Idaho, Oreg., Wash., Calif.

Sclerotinia sclerotiorum (Lib.) Mass., crown-rot. Del.

Urocystis sorosporioides Koern., rust. Utah.

ARACHIS.

A. hypogaea, peanut.

Bacterium solanacearum EFS. (*Phytomonas solanaceara* (EFS.) Com. S. A. B.), bacterial wilt. N. C.

Botrytis sp., stem-rot. Tenn.

Cercospora personata (Berk. and Curt.) Ell. and Ev., leaf-spot. General where crop is grown.

Corticium vagum Berk. and Curt., stem-rot. N. C., Fla.

Fusarium vasinfectum Atk., wilt. Va., Ala.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Phoma sp., stem-rot. Ala.

ARACHIS—Continued.

A. hypogaea, peanut—Continued.

Puccinia arachidis Speg., rust. Fla.

Sclerotium rolfsii Sacc., stem-rot. General in South.

Thielavia basicola (Berk. and Br.) Zopf, black root-rot

ARALIA.

A. spinosa, devils-walkingstick.

Cercospora atromaculans Ell. and Ev., leaf-spot. La.

Phyllosticta everhartii Sacc. and Syd., leaf-spot. W. Va.

ARBUTUS.

A. menziesii, madrone.

Exobasidium vaccinii (Fckl.) Wor., gall. Calif.

Mycosphaerella arbuticola Pk., leaf-spot. Pacific coast.

Phyllosticta fimbriata Ell. and Ev., leaf-spot. Oreg.

Pucciniastrum sparsum (Wint.) Ed. Fischer, rust. Pacific coast.

Rhytisma arbuti Phill., tar-spot. Pacific coast.

ARCTOSTAPHYLOS.

Arctostaphylos spp., bearberry, manzanita.

Exobasidium vaccinii (Fckl.) Wor., red leaf-spot. Colo., Oreg., Wash., Calif.

Phoradendron villosum Nutt., mistletoe. Oreg., Calif.

Pucciniastrum sparsum (Wint.) Ed. Fischer, rust. Pacific coast.

ARISTOLOCHIA.

A. serpentaria, Virginia snakeroot.

Cercospora serpentaria Ell. and Ev., leaf-spot. Del., Ala.

ARONIA.

Aronia spp., chokeberry.

Bacillus amylovorus (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. W. Va.

Cercospora mali Ell. and Ev., leaf-spot. Ala.

C. piri Farl., leaf-spot. N. H., Wis.

Gymnosporangium clavariaeforme (Jacq.) DC., rust. Mass.

G. davisii Kern, rust. Me., Mich., Wis.

G. germinale (Schw.) Kern, rust. Me., Mass.

G. transfmans (Ell.) Kern, rust. Mass., Pa., N. J.

Phyllosticta arbutifolia Ell. and Mart., leaf-spot. N. J.

ARUNDINARIA.

Arundinaria sp., bamboo, cane.

Puccinia arundinariae Schw., rust. Tex.

A. macrosperma, southern cane.

Puccinia arundinariae Schw., rust. S. C.

A. tecta, small cane.

Puccinia arundinariae Schw., rust. S. C., Miss., La.

ASCLEPIAS.

A. tuberosa, butterflyweed, *pleurisy root.

Cercospora asclepiadorae Ell. and Kell., leaf-spot. Ala.

Phyllosticta tuberosa Ell. and Mart., leaf-spot. N. J.

Puccinia jamesiana (Pk.) Arth., rust. Okla., Iowa, Kans.

Uromyces howei Pk., rust. Del., Fla., Ill., Iowa, Kans.

ASIMINA.

A. triloba, papaw.

Cercospora asiminae Ell. and Kell., leaf-spot. Ala., Miss., Kans.

Phleospora asiminae Ell. and Morgan, leaf-blotch. Ohio, Ill., Mo., Kans.

Phyllosticta asiminae Ell. and Kell., leaf-spot. General.

ASPARAGUS.*A. officinalis*, asparagus.*Armillaria mellea* (Vahl) Quel., root-rot. Oreg.*Botrytis* sp., blight. N. Y., Calif.*Cercospora asparagi* Sacc., leaf-spot. Del., Va., Ala., Tex., Nebr., Calif.*C. caulicola* Wint., leaf-spot. Md., S. C., Ga., Ala., Mo.*Cladosporium fasciculare* (Pers.) Fr. S. C.*Colletotrichum* sp., anthracnose. Conn., Ill.*Corticium vagum* Berk. and Curt., root-rot. Mass., N. Y., Pa., N. J., Mich., N. Dak., Wash.*Fusarium* sp., dwarfing, stem-rot. Mass., Conn., N. Y., N. J., Ill., Minn. Calif.*Puccinia asparagi* DC., rust. General.

Dwarfing, undet. N. J.

Fasciation, undet. Conn., Calif.

Leopard spot, undet. Md., Nebr.

AVENA.*A. sativa*, oats.*Bacterium coronafaciens* Charlotte Elliott (*Phytomonas coronafaciens* (Elliott) Com. S. A. B.), halo leaf-blight. General.*Claviceps purpurea* (Fr.) Tul., ergot. Iowa, Minn., N. Dak., Kans., Colo., Wash.*Colletotrichum graminicolum* (Ces.) Wils., anthracnose. General east of Great Plains.*Erysiphe graminis* DC., powdery mildew. Scattered, especially in Pacific States.*Fusarium culmorum leteius* Sherb., seedling blight. Oreg.*Gibberella saubinetii* (Mont.) Sacc., scab. Scatteringly east of Great Plains.*Helminthosporium avenae* Eidam, leaf-spot. Widely distributed.*Leptosphaeria avenaria* G. F. Weber, speckled blotch. Ky., Minn., Wis.*Phyllachora graminis* (Pers.) Fekl., black-spot. Nebr.*Puccinia coronata* Cda., crown-rust. General.*P. graminis* Pers., stem-rust. General.*Scolecotrichum graminis* Fekl., leaf-spot. N. Y., Ill.*Ustilago avenae* (Pers.) Jens., loose smut. General.*U. levis* (Kell. and Sw.) Magn., covered smut. General.

Blast, undet. General.

AZALEA.*Azalea* spp., azalea.*Corticium vagum* Berk. and Curt., stem-rot. Ga.*Exobasidium vaccinii* (Fekl.) Wor., gall. Atlantic Coast and Gulf States, Ill.*Microsphaera alni* (Wallr.) Wint., powdery mildew. Conn., N. Y., Pa.*Pucciniastrum myrtilli* (Schum.) Arth., rust. Conn., Ga., Atlantic Coast States.*Ramularia angustata* Pk., leaf-spot. N. Y.*Septoria solitaria* Ell. and Ev., leaf-spot. Calif.*Synchytrium vaccinii* F. Thomas, red-gall. N. J.**BAMBUSA.***Bambusa* sp., bamboo.*Cladosporium graminum* (Pers.) Cda. S. C.*Coniosporium shiraianum* (Syd.) Bub. Fla.*Helminthosporium* sp., leaf-spot. Fla.*Puccinia melanocephala* Syd., rust. Fla.**BEGONIA.***Begonia* sp., begonia.*Aphelenchus olesistis* Ritzema Bos, nematode (nursery). N. Y., Pa.*Bacterium tumefaciens* EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Miss.*Botrytis* sp., blight. Ind., Ill.*Cercospora* sp., leaf-spot. Fla., Tex.*Gloeosporium* sp., anthracnose. Fla., La.

BEGONIA—Continued.

Begonia sp., begonia—Continued.

Heterodera radiculicola (Greef) Muell., root-knot. Ohio, Ind., Nebr., Calif. (greenhouse).

Thielavia basicola (Berk. and Br.) Zopf, black root-rot. Mass., Ohio.

BENZOIN.

B. aestivale, spicebush.

Hymenochaete agglutinans Ell. Appalachian Mts.

Phoradendron flavescens (Pursh) Nutt., mistletoe. Eastern States.

Phyllosticta linderae Ell. and Ev., leaf-spot. Del., W. Va.

P. lindericola Ell. and Ev., leaf-spot. W. Va.

BERBERIS.

B. canadensis, Allegheny barberry.

Puccinia graminis Pers., rust. Va., W. Va.

B. thunbergii, Japanese barberry.

Verticillium alboatrum Reinke and Berth., wilt. N. Y.

Bacterial leaf-spot, undet. N. Y., Minn.

B. vulgaris, common barberry.

Puccinia graminis Pers., rust. General wherever host occurs.

Bacterial leaf-spot, undet. N. Y., Wis., Iowa, Nebr., Wash.

BETA.

B. vulgaris, beet.

Actinomyces scabies (Thax.) Gues., scab. Frequent where beets are grown.

Bacterium teutlium H. A. Metcalf, soft-rot. Nebr.

B. tumefaciens EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.)

Com. S. A. B.), crown-gall. Conn., Pa., Wis., Idaho.

Cercospora beticola Sacc., leaf-spot. General.

Corticium vagum Berk. and Curt., root-rot. Occurs scatteringly, especially throughout the Northern States.

Fusarium culmorum letesii Sherb., seedling blight. Pacific coast.

Heterodera radiculicola (Greef) Muell., root-knot. Southern States, Ark., Ind., Kans., Colo., Calif.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Peronospora schachtii Fckl., downy mildew. N. Y., Calif.

Phoma betae (Oud.) Frank, root-rot and leaf-spot. N. J., N. C., Mich., Wis., N. Dak., S. Dak., Kans., Colo., N. Mex., Wash., Oreg.

Puccinia sarcobati (Pk.) Bethel, rust. Utah.

Pythium aphanidermatum (Edson) Fitz., damping-off. Wis.

P. debaryanum Hesse, damping-off. Conn.

Sclerotinia sclerotiorum (Lib.) Mass., watery soft-rot. Conn., Ill.

Sclerotium rolfsii Sacc., stem-rot. S. C., Ga., La.

Septoria betae West., leaf-spot. Mass., Del., Ind.

Tylenchus dipsaci (Kuehn) Bast., stem nematode. Kans.

Uromyces betae (Pers.) Lév., rust. Colo., Oreg., Calif.

Curly-top, undet. Tex., Colo., Utah, Calif.

B. vulgaris, sugar beet.

Actinomyces scabies (Thax.) Gues., scab. Conn., Ohio, Idaho.

Bacterium aptatum N. Brown and C. Jamieson (*Phytomonas aptatum* (Brown and Jamieson) Com. S. A. B.), leaf-spot. Utah.

B. beticolum EFS., N. Brown, and Town. (*Phytomonas beticolum* (EFS., Brown, and Town.) Com. S. A. B.), tuberculosis. Va., Kans., Colo., Utah.

B. tumefaciens EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Ohio, Utah, Colo., Idaho, Calif.

Cercospora beticola Sacc., leaf-spot. General.

Corticium vagum Berk. and Curt., root-rot. General.

Fusarium sp., root-rot. Ind., Mich., Kans., Western States.

Heterodera radiculicola (Greef) Muell., root-knot. Ohio, Texas, Western States.

H. schachtii A. Schmidt, sugar-beet nematode. Wyo., Colo., Utah, Idaho, Calif.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

BETA—Continued.*B. vulgaris*, sugar beet—Continued.

- Peronospora schachtii* Fekl., downy mildew. Calif.
Phoma betae (Oud.) Frank, root-rot and leaf-spot. General.
Puccinia sarcobati (Pk.) Bethel, rust. Colo., Utah.
Pythium sp., damping-off. Ohio, Wis., Utah, Idaho, Calif.
P. aphanidermatum (Edson) Fitz., damping-off. Wis., Utah.
Sclerotium rolfsii Sacc., stem-rot. Ga., Tex., Calif.
Tylenchus dipsaci (Kuehn) Bast., stem nematode. Kans.
T. penetrans Cobb, nematode. Utah.
Uromyces betae (Pers.) Lév., rust. Calif.
 Curly-top, undet. Western States.
 Mosaic, undet. Ind., Kans., Nebr., Colo., Utah, Calif.

B. vulgaris cicla, Swiss chard.

- Actinomyces scabies* (Thax.) Gues., scab. Pa., Wis.
Cercospora beticola Sacc., leaf-spot. General.
Corticium vagum Berk. and Curt., root-rot. N. Y.
Heterodera radicola (Greef) Muell., root-knot. Va., La.
Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.
Sclerotium rolfsii Sacc., stem-rot. S. C., La.
Uromyces betae (Pers.) Lév., rust. Calif.

B. vulgaris macrorrhiza, mangel-wurzel.

- Bacterium tumefaciens* EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Pa.
Cercospora beticola Sacc., leaf-spot. Conn., Minn., Iowa.
Corticium vagum Berk. and Curt., root-rot. Wash.
Phoma betae (Oud.) Frank, root-rot and leaf-spot. N. Y., Wis., Minn.
Pythium debaryanum Hesse, damping-off. Conn., Mich., Wis., Kans., Colo., Utah.
Sclerotium rolfsii Sacc., stem-rot. La.

BETULA.*B. alba*, white birch.

- Magnusiella flava* (Farl.) Sadeb., yellow leaf blister. Northeastern States.
Melanconium bicolor Nees, canker. N. Y., Ind.
Polyporus betulinus (Bull.) Fr., powdery sapwood rot. Vt., N. Y.

B. lenta, sweet birch.

- Fomes igniarius* (L.) Fr., white heart-rot.
Gloeosporium betularum Ell. and Mart., leaf-spot. Pa., N. J., Ill.
Microsphaera alni (Wallr.) Wint., powdery mildew. N. H.
Phyllactinia corylea (Pers.) Karst., powdery mildew. N. Y.
Phyllosticta betulae Ell. and Ev., leaf-spot. N. Y.
Septoria microsperma Pk., leaf-spot. Pa., W. Va.

B. lutea, yellow birch.

- Fomes applanatus* (Pers.) Wallr. (*Ganoderma applanatum* (Pers.) Pat.), white-streaked rot. Me., N. Y.
F. fomentarius (L.) Fr., white sapwood rot. New England, N. Y., Pa.
F. igniarius (L.) Fr., white heart-rot. Vt., N. Y.
Melampsoridium betulae (Schum.) Arth., rust. N. H., Ind.
Melanconium bicolor Nees, canker. N. Y.
Microsphaera alni (Wallr.) Wint., powdery mildew. Mass.
Nectria coccinea (Pers.) Fr., associated with canker. Mich.
Polyporus betulinus (Bull.) Fr., powdery sapwood rot. N. Y.

B. nigra, river birch.

- Fomes igniarius* (L.) Fr., white heart-rot. Conn.
Gloeosporium betularum Ell. and Mart., leaf-spot. Mass., Pa., Ind., Ill., Wis.
Nectria ditissima Tul., European canker. Pa.
Phoradendron flavescens (Pursh) Nutt., mistletoe. Ind.
Phyllactinia corylea (Pers.) Karst., powdery mildew. Ill., Mo.

B. occidentalis, western birch.

- Fomes fomentarius* (L.) Fr., white sapwood rot.
F. igniarius (L.) Fr., white heart-rot. Mont.

BETULA—Continued.*B. occidentalis*, western birch—Continued.

- Gnomonia campylostyla* Auer. Colo.
Melampsoridium betulae (Schum.) Arth., rust. Mont., Idaho.
Phyllactinia corylea (Pers.) Karst., powdery mildew. Mont., Wyo.
Septoria betulicola Pk., leaf-spot. Wash.
Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Idaho.

B. papyrifera, canoe birch.

- Exoascus betulinus* (Rostr.) Sadeb., witches'-broom. Wis.
Fomes applanatus (Pers.) Wallr. (*Ganoderma applanatum* (Pers.) Pat.), white-streaked rot. Me., Minn.
F. everhartii (Ell. and Gall.) Schrenk, white-rot. Minn.
F. fomentarius (L.) Fr., white sapwood rot. Mass., Minn.
F. igniarius (L.) Fr., white heart-rot. Mich., Minn.
Gloeosporium betulae-papyriferae Dearn. and Overh., anthracnose. Pa.
Magnusiella flava (Farl.) Sadeb., yellow leaf blister. Northeastern States.
Melampsoridium betulae (Schum.) Arth., rust. Me.
Phyllactinia corylea (Pers.) Karst., powdery mildew. Iowa.
Polyporus betulinus (Bull.) Fr., powdery sapwood rot. Mass., Conn., Minn.

B. populifolia, gray birch.

- Fomes fomentarius* (L.) Fr., white sapwood rot. Me.
Magnusiella flava (Farl.) Sadeb., yellow leaf blister. Northeastern States.
Melampsoridium betulae (Schum.) Arth., leaf-rust. Northeastern States.
Melanconium bicolor Nees, canker. N. Y.
Polyporus betulinus (Bull.) Fr., powdery sapwood rot. Northeastern States.
Septoria betulina Pk., leaf-spot. Wis.
Venturia ditricha (Fr.) Aderh. Me., N. Y.

BIGNONIA.*B. capreolata*, crossvine.

- Cercospora capreolata* Ell. and Ev., leaf-spot. Miss.
Dimerosporium tropicale Speg., sooty mold. Miss.

B. radicans, trumpetcreeper.

- Cercospora pallida* Ell. and Ev., leaf-spot. La.
C. sordida Sacc., leaf-blight. General east of Great Plains, especially in the South.
Erysiphe cichoracearum DC., powdery mildew. Ill.
Hendersonia peckii Clint. N. Y.
Microsphaera alni (Wallr.) Wint., powdery mildew. Va., Ala., Ohio, Ind., Miss.
Phyllosticta tecomae Sacc., leaf-spot. Miss.
Septoria tecomae Ell. and Ev., leaf-spot. W. Va., Tex.

BRASSICA.*B. campestris*, rutabaga.

- Albugo candida* (Pers.) O. Kuntze, white-rust. Mass.
Alternaria brassicae (Berk.) Sacc., black leaf-spot. Va.
Bacterium campestre (Pam.) EFS. (*Phytomonas campestre* (Pam.) Com. S. A. B.), black-rot. Va., Wis., Minn., Iowa.
Erysiphe polygoni DC., powdery mildew. Conn., Fla., Wash.
Heterodera radiculicola (Greef) Muell., root-knot. Ala.
Mycosphaerella brassicicola (Fr.) Lindau, ring-spot. Calif.
Peronospora parasitica (Pers.) D By., downy mildew. Fla., Iowa.
Phoma lingam (Tode) Desm., black-leg. Ga.
Plasmodiophora brassicae Wor., clubroot. Minn., N. Dak.

B. oleracea, cabbage.

- Albugo candida* (Pers.) O. Kuntze, white-rust. Ky., Idaho.
Alternaria brassicae (Berk.) Sacc., black leaf-spot. General.
A. herculea (Ell. and Mart.) J. A. Elliott, gray leaf-spot. Calif.
Bacillus carotovorus L. R. Jones (*Erwinia carotovora* (Jones) Com. S. A. B.), soft-rot. General.

BRASSICA—Continued.

B. oleracea, cabbage—Continued.

Bacterium campestre (Pam.) EFS. (*Phytomonas campestre* (Pam.) Com. S. A. B.), black-rot. General.

B. maculicolum McC. (*Phytomonas maculicola* (McC.) Com. S. A. B.), peppery leaf-spot. Conn., N. Y., Ala., Miss., La.

Botrytis sp., gray-mold. N. Y., Calif., Alaska.

Cercospora bloxami Berk. and Br., leaf-spot. Fla., Tex., Ill.

C. cruciferarum Ell. and Ev., leaf-spot. Ala.

Corticium vagum Berk. and Curt., wire-stem. General.

Erysiphe polygoni DC., powdery mildew. Md., Nebr., Tex., Oreg.

Fusarium conglutinans Woll., yellows. General except New England.

Heterodera radicola (Greef) Muell., root-knot. Southern States and Calif.

Mycosphaerella brassicicola (Fr.) Lindau, ring-spot. Oreg., Wash., Calif.

Olpidium brassicae (Wor.) Dan., seedling blight. N. Y., Miss., Wis.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Peronospora parasitica (Pers.) D By., downy mildew. General.

Phoma lingam (Tode) Desm., black-leg. General, especially in Eastern States.

Plasmidiophora brassicae Wor., clubroot. General.

Pythium debaryanum Hesse, damping-off. Conn., Pa.

Sclerotinia sclerotiorum (Lib.) Mass., drop, watery soft-rot. General.

Sclerotium rolfsii Sacc., stem-rot. Ala., Tex.

Malnutrition, nonpar. Southern States, Long Island, Mass.

Oedema, nonpar. N. Y.

Speck, undet. Frequent in stored cabbage.

Tipburn, nonpar. Ind.

B. oleracea acephala, kale, collards.

Albugo candida (Pers.) O. Kuntze, white-rust. Fla.

Alternaria brassicae (Berk.) Sacc., black leaf-spot. Southern States.

Bacterium campestre (Pam.) EFS. (*Phytomonas campestre* (Pam.) Com. S. A. B.), black-rot. N. C., Ala.

Corticium vagum Berk. and Curt., stem-rot. Wash.

Fusarium conglutinans Woll., yellows. N. Y., N. C., S. C., Ind., Calif.

Heterodera radicola (Greef) Muell., root-knot. N. Y., S. C., Ala.

Mycosphaerella brassicicola (Fr.) Lindau, ring-spot. Wash., Calif.

Peronospora parasitica (Pers.) D By., downy mildew. South Atlantic States; Oreg.

Phoma lingam (Tode) Desm., black-leg. N. C., Ga.

Plasmidiophora brassicae Wor., clubroot. Conn., Va., N. C., Wash.

Sclerotium rolfsii Sacc., stem-rot. Miss.

B. oleracea botrytis, cauliflower.

Alternaria brassicae (Berk.) Sacc., black leaf-spot. N. Y., Fla., La., Calif.

A. herculea (Ell. and Mart.) J. A. Elliott, gray leaf-spot. Calif.

Bacillus carotovorus L. R. Jones (*Erwinia carotovora* (Jones) Com. S. A. B.), soft-rot. Me., Mass., Va., Minn., Mo.

Bacterium campestre (Pam.) EFS. (*Phytomonas campestre* (Pam.) Com. S. A. B.), black-rot. General, especially in States east of Miss. River.

B. maculicolum McC. (*Phytomonas maculicola* (McC.) Com. S. A. B.), peppery leaf-spot. Conn., N. Y., Va., Ky., Fla.

Corticium vagum Berk. and Curt., stem-rot. La., Wash.

Fusarium conglutinans Woll., yellows. Del., Ind., Calif.

Mycosphaerella brassicicola (Fr.) Lindau, ring-spot. Mass., Oreg., Calif.

Peronospora parasitica (Pers.) D By., downy mildew. N. Y., Va., Mo., Calif.

Phoma lingam (Tode) Desm., black-leg. N. Y.

Plasmidiophora brassicae Wor., clubroot. East of Miss. River and north of Va. and Ky.

Pythium sp., damping-off. Ohio, Oreg.

Sclerotinia sclerotiorum (Lib.) Mass., drop, watery soft-rot. Va., Miss., La., Calif.

BRASSICA—Continued.

B. oleracea caulorapa, kohlrabi.

Heterodera radicum (Greef) Muell., root-knot. Ohio.

Peronospora parasitica (Pers.) D By., downy mildew. Va., Fla.

Plasmodiophora brassicae Wor., clubroot. Conn., Mass., N. Y., Ohio, Miss.

B. oleracea gemmifera, Brussels sprouts.

Bacillus carotovorus L. R. Jones (*Erwinia carotovora* (Jones) Com. S. A. B.), soft-rot. Minn.

Corticium vagum Berk. and Curt., stem-rot. La.

Mycosphaerella brassicicola (Fr.) Lindau, ring-spot. Calif.

Phoma lingam (Tode) Desm., black-leg. N. Y.

Plasmodiophora brassicae Wor., clubroot. Alaska.

Pythium debaryanum Hesse, damping-off. Minn.

Sclerotinia sclerotiorum (Lib.) Mass., drop, watery soft-rot. La.

B. pekinensis, petsai cabbage, Chinese cabbage.

Alternaria brassicae (Berk.) Sacc., black leaf-spot. Conn., Fla.

A. herculea (Ell. and Mart.) J. A. Elliott, gray leaf-spot. Fla., Calif.

Bacillus carotovorus L. R. Jones (*Erwinia carotovora* (Jones) Com. S. A. B.), soft-rot. Conn.

Bacterium campestre (Pam.) EFS. (*Phytomonas campestre* (Pam.) Com. S. A. B.), black-rot. Md.

Cercospora bloxami Berk. and Br., leaf-spot. Fla.

Cercospora albomaculans (Ell. and Ev.) Sacc., leaf-spot. Conn.

Cylindrosporium brassicae Fautr. and Roum., leaf-spot. Md., Mo.

Sclerotinia sclerotiorum (Lib.) Mass., drop. Mass.

Mosaic, undet.

B. rapa, turnip (white).

Albugo candida (Pers.) O. Kuntze, white-rust. Reported in seven scattered States.

Alternaria sp., black leaf-spot. Ind., Tex., Utah.

A. herculea (Ell. and Mart.) J. A. Elliott, gray leaf-spot. N. J., Ohio, Fla., Ala.

Bacillus carotovorus L. R. Jones (*Erwinia carotovora* (Jones) Com. S. A. B.), soft-rot. Reported from widely scattered States.

Bacterium campestre (Pam.) EFS. (*Phytomonas campestre* (Pam.) Com. S. A. B.), black-rot. Eastern States.

Cercospora albomaculans (Ell. and Ev.) Sacc., leaf-spot. General.

Colletotrichum brassicae Schult. and Sacc., leaf-spot. Ga.

C. higginsianum Sacc., leaf-spot. S. C., Ga., Fla., Ala., Miss.

Corticium vagum Berk. and Curt., stem-rot. Mass., Ohio, Wash.

Cylindrosporium brassicae Fautr. and Roum., leaf-spot. Ga.

Erysiphe polygoni DC., powdery mildew. Occurs in widely scattered States.

Fusarium conglutinans Woll., yellows. Ind.

Heterodera radicum (Greef) Muell., root-knot. S. C., Ga., Ala., Tex.

Peronospora parasitica (Pers.) D By., downy mildew. Frequent east of Great Plains.

Phoma napobrassicae Rostr., root-rot. Conn.

Plasmodiophora brassicae Wor., clubroot. East of Great Plains; Colo., Wash.; especially in Northeastern States.

Sclerotinia sclerotiorum (Lib.) Mass., drop, watery soft-rot. Conn., Tex.

Mosaic, undet. Ind.

BROMUS

B. inermis, *Hungarian brome-grass.

Bacterium coronafaciens atropurpureum C. S. Reddy and James Godkin, bacterial blight. Wis., N. Dak., S. Dak.

Claviceps purpurea (Fr.) Tul., ergot. Conn., Minn., Iowa, N. Dak., Nebr.

Erysiphe graminis DC., powdery mildew. Minn.

Helminthosporium giganteum Heald and Wolf, leaf-spot. D. C.

Phyllachora graminis (Pers.) Fekl., tar-spot. N. Dak.

Pyrenophora bromi (Died.) Drechs., leaf-blotch. Pa., Wis., Minn., N. Dak.

Rhynchosporium secalis (Oud.) Davis, scald. Wis.

Septoria bromi Sacc., leaf-blotch. Wis.

S. bromigena Sacc., leaf-spot. N. Dak.

Ustilago bromivora (Tul.) Fisch., smut. Iowa, Oreg.

BROMUS—Continued.**B. secalinus*, chess.

- Colletotrichum graminicolum* (Ces.) Wils., anthracnose. Pa., Ohio.
Ophiobolus graminis Sacc., take-all. N. Y.
Puccinia clematidis (DC.) Lagh., leaf-rust. N. C., Ind., Oreg., Calif.
P. graminis Pers., stem-rust. W. Va., Ind., Wis., Iowa, Kans.
Septoria bromi Sacc., leaf-blotch. Wis.
Ustilago bromivora (Tul.) Fisch., smut. Pacific coast.

BUDDLEIA.*Buddleia* sp., butterflybush.

- Heterodera radiculicola* (Greef) Muell., root-knot. Miss.

BULBILIS.B. dactyloides*, buffalo grass.

- Cercospora seminalis* Ell. and Ev., leaf-spot. S. Dak., Kans., Colo.
Helminthosporium turcicum Pass., leaf-blight. Kans.
Puccinia kansensis Ell. and Barth., rust. Nebr., Kans.
Tilletia buchloeana Kell. and Syd., covered smut. Tex., Kans.
Ustilago buchloes Ell. and Tr., loose smut. Nebr., N. Mex.

BUMELIA.*Bumelia* sp., bumelia.

- Cercospora lanuginosa* Heald and Wolf, leaf-spot. Texas.
Phyllosticta bumeliifolia Heald and Wolf, leaf-spot. Ala., Miss., Tex.
P. curtisii (Sacc.) Ell. and Ev., leaf-spot. Fla.
Septoria bumeliae Sacc., leaf-spot. Miss.

BUXUS.*B. sempervirens*, box.

- Macrophoma candollei* (Berk. and Br.) Berl. and Vogl., leaf-blight. East of Miss. River.
Nectria rousselliana Tul., canker. N. J., Pa., Md.
Phyllosticta sp., leaf-spot. N. Y.
P. auerswaldii Allesch., leaf-spot. N. J.

CAJANUS.*C. indicus*, pigeon-pea.

- Uromyces dolicholi* Arth., rust. Fla.

CALENDULA.*C. officinalis*, calendula.

- Corticium vagum* Berk. and Curt., stem-rot. Ind.
Heterodera radiculicola (Greef) Muell., root-knot. W. Va.
Puccinia emiliae P. Henn., rust. Ill., Mo., Nebr., Kans.

CALLICARPA.*C. americana*, American beautyberry.

- Cercospora callicarpae* Cke., leaf-spot. Gulf States.
Meliola cookeana Speg., sooty mold. Fla.

CALLISTEPHUS.*C. chinensis*, China-aster.

- Botrytis* sp., stem-blight. Conn., Pa.
Coleosporium solidaginis (Schw.) Thuem., rust. Frequent east of Great Plains.
Corticium vagum Berk. and Curt., stem-rot. Me., Mass., N. Y., Mich., Wash.
Erysiphe cichoracearum DC., powdery mildew. Minn., Wash.
Fusarium conglutinans callistephi Beach, wilt. General in Northern States.
Heterodera radiculicola (Greef) Muell., root-knot. Conn., W. Va.
Septoria callistephi W. O. Gloyer, leaf-spot. N. Y., Del., Ala., Ohio, Mich.
 Yellows, undet. General in Northern States.

CANNA.*C. indica*, canna.

Bacterium cannae M. K. Bryan (*Phytomonas cannae* (Bryan) Com. S. A. B.), bacterial bud-rot. Conn., Ind.

Fusarium sp., rot. Pa., Minn.

Puccinia cannae (Wint.) P. Henn., rust. Ohio, Fla.

CANNABIS.*C. sativa*, common hemp.

Botryosphaeria marconi (Cav.) V. K. Charles and A. Jenkins, wilt. Va.

Botrytis cinerea Auct., rot. Va., Oreg.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Sclerotium rolfsii Sacc., stem-rot. S. C.

Septoria cannabina Pk., leaf-spot. Iowa.

S. cannabis (Lasch) Sacc., leaf-spot. Ind., Ky., Minn.

CAPRIOLA.*C. dactylon*, Bermuda grass.

Helminthosporium cynodontis G. B. Marignoni, leaf-spot. Fla.

H. giganteum Heald and Wolf, leaf-spot. Fla., Tex.

Puccinia cynodontis DeLac., rust. Frequent in Southern States.

Ustilago cynodontis P. Henn., smut. Tex., Calif.

CAPSICUM.*C. annuum*, redpepper.

Alternaria sp., fruit-rot. Frequent, especially in the South.

Bacillus carotovorus L. R. Jones (*Erwinia carotovora* (Jones) Com. S. A. B.), soft-rot. Fla., Mich.

Bacterium solanacearum EFS. (*Phytomonas solanaceara* (EFS.) Com. S. A. B.), bacterial wilt. Pa., Va., N. C., La., Tex.

B. vesicatorium E. M. Doidge (*Phytomonas exitiosa* (Gardner and Kendrick) Com. S. A. B.), bacterial spot. Tenn., Ga., Fla., Ind.

Cercospora capsici Heald and Wolf, leaf and fruit spot. Ga., Fla., Tex., Calif.

Corticium vagum Berk. and Curt., stem-rot. Conn., N. J., Pa., Ga., La.

Fusarium sp., wilt. Del., Tenn., Ark., Calif.

F. annuum L. H. Leonian, wilt. N. Mex.

Glomerella cingulata (Ston.) Spauld. and Schrenk, anthracnose. General east of Miss. River, especially in the South; Tex.

Heterodera radiculicola (Greef) Muell., root-knot. Fla., N. Mex.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Phoma destructiva Plow., rot. Ga.

Phyllosticta sp., leaf-spot. N. J.

Phytophthora capsici L. H. Leonian, blight. N. Mex.

P. terrestris Sherb., fruit-rot. Ind.

Pythium debaryanum Hesse, fruit-rot. N. C.

Sclerotinia sclerotiorum (Lib.) Mass., damping-off. Conn.

Sclerotium bataticola Taub., fruit-rot. N. J.

S. rolfsii Sacc., stem-rot. South Atlantic and Gulf States.

Blossom-end rot, nonpar. Ga.

Mosaic, undet. Frequent in widely scattered States.

Sun scald, undet. N. J., Ind., Calif.

CARICA.*C. papaya*, papaya.

Colletotrichum sp., fruit-rot. Fla.

Heterodera radiculicola (Greef) Muell., root-knot.

Oidium caricae F. Noack, powdery mildew. Fla.

CARPINUS.*Carpinus* sp., hornbeam.

Fomes ignarius (L.) Fr., white heart-rot. Vt.

Fracchiacea calista (Berk. and Curt.) Sacc. Vt., Ala., Tex.

Gnomoniella fimbriata (Pers.) Sacc., leaf-spot. N. Y., W. Va.

CARPINUS—Continued.

C. caroliniana, American hornbeam, blue beech.

Dermatea carpinea (Pers.) Rehm, canker. N. Y., Pa.

Fusarium carpineum Davis. Wis.

Gloeosporium robergei Desm., leaf-spot. Pa.

Gnomoniella fimbriata (Pers.) Sacc., leaf-spot. N. Y.

Microsphaera alni (Wallr.) Wint., powdery mildew. Ind., Iowa.

Phyllactinia corylea (Pers.) Karst., powdery mildew. Ohio, Ind., Tex.

Phyllosticta hesperidearum (Catt.) Penz., leaf-spot. Colo. (greenhouse).

Septoria carpinea Davis, leaf-spot. Wis.

CASTANEA.

C. dentata, chestnut.

Endothia parasitica (Murr.) P. J. and H. W. Anderson, blight. New England, N. Y., Atlantic Coast States except Florida; also W. Va., Ky., Tenn., Ala., Ohio, Ind., Mich., Iowa, Mo., Nebr.

Fusicoccum sp., canker. Calif. (*Castanea* sp.).

Marssonina ochroleuca (Berk. and Curt.) Humph., leaf-spot. Me. to N. C., Ohio.

Melanconis modonia Tul., black canker. Ohio.

Microsphaera alni (Wallr.) Wint., powdery mildew. Conn., N. Y., Pa.

Monochaetia desmazieri Sacc., large leaf-spot. Va., N. C., Ga.

Phoradendron flavescens (Pursh) Nutt., mistletoe. Eastern States.

Phyllactinia corylea (Pers.) Karst., powdery mildew. Mass., N. Y., Ind.

Phyllosticta castanea Ell. and Ev., leaf-spot. Md., W. Va.

P. fusispora Ell. and Ev., leaf-spot. Ala.

Phyalospora malorum (Pk.) Shear (*P. cydoniae* Am. Auct.), twig-blight. Central Eastern States.

Polyporus croceus Fr., white-piped rot. Eastern States.

P. frondosus Fr., straw-colored rot.

P. spraguei Berk., heart-rot. Ind.

P. sulphureus (Bull.) Fr., brown-checked wood-rot. N. Y.

Septoria ochroleuca Berk. and Curt., leaf-spot. Mass., Kans.

Strumella coryneoides Sacc. and Wint., canker. Northeastern States.

C. japonica, Japanese chestnut.

Endothia parasitica (Murr.) P. J. and H. W. Anderson, blight. Middle Atlantic States, Mass., Conn., N. Y., W. Va., N. C.

Septoria gillettiana Sacc., leaf-spot. La.

C. pumila, chinquapin.

Endothia parasitica (Murr.) P. J. and H. W. Anderson, blight. N. H., Conn., Pa., Md., Va., W. Va., Tenn., S. C., Ga.

Microsphaera alni (Wallr.) Wint., powdery mildew. Va., N. C.

Polyporus croceus Fr., white-piped rot. Ark.

C. sativa, Spanish chestnut.

Endothia parasitica (Murr.) P. J. and H. W. Anderson, blight. Middle Atlantic States, Conn., N. Y., Ohio.

Marssonina ochroleuca (Berk. and Curt.) Humph., leaf-spot. Conn., N. J.

Microsphaera alni (Wallr.) Wint., powdery mildew. Mass.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Phyllactinia corylea (Pers.) Karst., powdery mildew. Ohio.

CASTANOPSIS.

C. chrysophylla, giant chinquapin.

Dothidella castanopsidis Dearn., leaf-spot. Oreg.

CATALPA.

Catalpa sp., catalpa.

Cercospora catalpae Wint., leaf-spot. Mass., N. J., Fla., Ala., La., Mo., Tex.

Cladosporium sp., leaf-spot. Ind.

Collybia velutipes Fr., heart-rot. Ind.

Gloeosporium catalpae Ell. and Ev., leaf-spot. N. Y., Pa.

Heterodera radicola (Greef) Muell., root-knot. Ala., Fla.

Macrosporium catalpae Ell. and Mart., leaf-spot. Frequent east of Miss. River; Mo.

CATALPA—Continued.*Catalpa* sp., catalpa—Continued.*Microsphaera alni vaccinii* (Schw.) Salm., powdery mildew. General east of Great Plains.*Ozonium omnivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.*Phyllactinia corylea* (Pers.) Karst., powdery mildew. S. C., Ohio, Ind., Ill.*Phyllosticta catalpae* Ell. and Mart., leaf-spot. General east of Great Plains.*Polyporus catalpae* Schrenk, brown butt-rot. Mass.*Polystictus versicolor* (L.) Fr., yellowish wood-rot. Mass., Ohio, Ind., Nebr.*Thielavia basicola* (Berk. and Br.) Zopf, black root-rot. Ohio.

Mosaic, undet. Md.

CELASTRUS.*C. scandens*, American bittersweet.*Cercospora melanochaeta* Ell. and Ev., leaf-spot. N. Dak., Nebr., Kans.*Microsphaera alni* (Wallr.) Wint., powdery mildew. Wis.*Phyllactinia corylea* (Pers.) Karst., powdery mildew. General.*Phyllosticta celastri* Ell. and Ev., leaf-spot. W. Va.*Ramularia celastri* Ell. and Mart., leaf-spot. General.**CELTIS.***Celtis* sp., hackberry, sugarberry.*Cylindrosporium celtidis* Earle, leaf-spot. Ala.*C. defoliatum* Heald and Wolf, leaf-blight. Tex.*Ozonium omnivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.*Phleospora celtidis* Ell. and Mart., leaf-spot. Mass., Mo., Kans.*Phoradendron flavescens* (Pursh) Nutt., mistletoe. Tex., Ind., Ariz.*Phyllosticta celtidis* Ell. and Kell., leaf-spot. East of Great Plains; N. Mex.*Pseudoperonospora celtidis* (M. B. Waite) Wils., downy mildew. D. C.*Ramularia celtidis* Ell. and Kell., leaf-spot. Frequent east of Great Plains.*Septogloeum celtidis* Dearn., leaf-spot. N. Y.*Septoria celti-gallae* W. R. Gerard, leaf-spot. N. Y.*S. gigaspora* Ell. and Ev., leaf-spot. Mass., Kans.*Sphaerotheca phytophila* Kell. and Sw., powdery mildew. East of Great Plains.*Uncinula confusa* Mass., powdery mildew. N. C.*U. parvula* Cke. and Pk., powdery mildew. General.*U. polychaeta* (Berk. and Curt.) Ell., powdery mildew. Mass., Ala., Miss., Tex.**CEPHALANTHUS.***C. occidentalis*, buttonbush.*Ascochyta cephalanthi* Ell. and Ev., leaf-spot. La.*Cercospora perniciosa* Heald and Wolf, leaf-blight. Tex.*Coniothyrium cephalanthi* Ell. and Ev., leaf-spot. La.*Microsphaera alni* (Wallr.) Wint., powdery mildew. N. H., Ala., Ind., Ill.*Phyllactinia corylea* (Pers.) Karst., powdery mildew. Ind.*Phyllosticta cephalanthi* Tharp, leaf-spot. Tex.*Puccinia cephalanthi* (Seym.) Barth., rust. N. Y., Ind., Ill., Kans.*Ramularia cephalanthi* (Ell. and Kell.) Heald, leaf-spot. Southern States.*Septoria cephalanthi* Ell. and Kell., leaf-spot. Kans.**CERCIS.***C. canadensis*, redbud, *Judas-tree.*Cercospora cercidicola* Ell., leaf-spot. General throughout range of host.*C. chionea* Ell. and Kell., leaf-spot. Ind., Ill., Kans.*Pestalozzia* sp., leaf-spot. Fla.*Phyllosticta cercidicola* Ell. and Ev., leaf-spot. W. Va., Ind.*P. siliquastri* Sacc. and Speg., leaf-spot. Ala.**C. occidentalis*, California redbud.*Cercospora cercidicola* Ell., leaf-spot. Tex.

CHAETOCHLOA.

C. italica, millet.

Helminthosporium sp., leaf-spot. Minn.

Piricularia grisea (Cke.) Sacc., leaf-spot. Pa., Va., Ala., Tex., Ind., Ill., Iowa, Mo.

Sclerospora graminicola (Sacc.) Schroet., downy mildew. North-Central States.

Ustilago crameri Koern., smut. General east of Great Plains.

CHAMAECYPARIS.

C. nootkatensis, *Alaska cedar, Nootka cypress.

Gymnosporangium nootkatensis (Trel.) Arth., rust. Wash., Oreg.

C. thyoides, whitecedar.

Gymnosporangium botryapites (Schw.) Kern, rust. Atlantic Coast States from Mass. to Ala.

G. ellisii (Berk.) Farl., rust. Atlantic Coast States from Mass. to Del.; also Fla. and Ala.

G. transformans (Ell.) Kern, rust. Mass., N. J.

CHAYOTA.

C. edulis, chayote.

Cercospora cucurbitae Ell. and Ev., leaf-spot. Fla.

Colletotrichum lagenarium (Pass.) Ell. and Hals., anthracnose. Fla.

CHENOPODIUM.

**C. ambrosioides anthelminticum*, wormseed.

Cercospora anthelmintica Atk., leaf-spot. Ala., Miss., Tex.

C. chenopodii Fr., leaf-spot. Tex.

Heterodera radiculicola (Greef) Muell., root-knot. Ala.

Phyllosticta ambrosioides Thuem. and Syd., leaf-spot. Del., N. J., La., Ind.

CHIONANTHUS.

C. virginica, white fringetree.

Phyllactinia corylea (Pers.) Karst., powdery mildew. Md.

Phyllosticta chionanthi Thuem., leaf-spot. N. J., W. Va.

Septoria chionanthi Cke., leaf-spot. S. C.

Sclerospora Sacc., leaf-spot. Tex.

CHRYSANTHEMUM.

Chrysanthemum sp., chrysanthemum.

Aphelenchus sp., bud nema. Md.

Ascochyta chrysanthemi F. L. Stevens, ray-blight. N. C., S. C.

Botrytis sp., bud-rot. Mass., Ky., Minn., Mo.

B. cinerea Auct., flower deformation. N. Y.

Cercospora chrysanthemi Heald and Wolf., leaf-spot. Tex.

Corticium vagum Berk. and Curt., stem-rot. Md.

Cylindrosporium chrysanthemi Ell. and Dearn., leaf-spot. Mass., Conn., Pa., N. C., Ga., Miss.

Erysiphe cichoracearum DC., powdery mildew. General.

Fusarium sp., wilt. Mass., N. J., Ala., Fla., Ohio, Kans.

Heterodera radiculicola (Greef) Muell., root-knot. Miss., Tex., Wash.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Phyllosticta chrysanthemi Ell. and Dearn., leaf-spot. Mass., Fla.

Puccinia chrysanthemi Roze, rust. General.

Septoria chrysanthemella Cav., leaf-spot. General east of Great Plains.

S. leucanthemi Sacc. and Speg., leaf-spot. Conn.

Verticillium alboatrum Reinke and Berth., wilt. N. J.

Chlorosis, nonpar. Pa., Md.

Mosaic, undet. Md., Ohio.

Yellows, undet. Mich.

CICHORIUM.

C. endivia, endive.

Corticium vagum Berk. and Curt., damping-off. Conn.

Pythium debaryanum Hesse, root-rot. Conn.

Sclerotinia sclerotiorum (Lib.) Mass., watery soft-rot. La.

C. intybus, chicory.

Puccinia hieracii (Schum.) Mart., rust. Vt., N. Y., Wis., Wash., Oreg.

CINNAMOMUM.

C. camphora, camphor-tree.

Diplodia sp., canker. Tex.

Gloeosporium camphorae Sacc., leaf-spot. Fla., Ala., Miss., Tex.

Glomerella cingulata (Ston.) Spauld. and Schrenk, anthracnose. Ga., Fla.

Tylenchus penetrans Cobb, nematode. Fla.

CITRULLUS.

C. vulgaris, watermelon.

Bacillus tracheiphilus EFS. (*Erwinia tracheiphila* (EFS.) Com. S. A. B.), bacterial wilt. General.

Cercospora citrullina Cke., leaf-spot. Ind. to Tex. and eastward.

Colletotrichum lagenarium (Pass.) Ell. and Huls., anthracnose. General east of Rocky Mts.; also in Calif.

Corticium vagum Berk. and Curt., fruit-rot. La., Wash., Oreg.

Diplodia sp., *Diplodia* rot. Southern States.

Erysiphe cichoracearum DC., powdery mildew. Va.

Fusarium sp., *Fusarium* rot. Va., Kans.

F. niveum EFS., wilt. Southern two-thirds of United States.

Heterodera radicolica (Greef) Muell., root-knot. S. C., Ala., Tex., Calif.

Macrosporium cucumerinum Ell. and Ev., leaf-blight. Occurs scatteringly, mostly east of Rocky Mts.

M. sarcinula parasiticum Thuem., leaf-spot. Kans.

Mycosphaerella citrullina (C. O. Smith) Gross., gummy stem-blight. N. Y., Va., S. C., Tenn., Fla., La., Mo.

Pseudoperonospora cubensis (Berk. and Curt.) Rostew., downy mildew. Atlantic Coast and Gulf States; W. Va., Ohio, Mich., Ark., Calif.

Pythium sp., blossom-end rot. Va.

P. aphanidermatum (Edson) Fitz., rot.

P. debaryanum Hesse, rot. Va., Mich.

Sclerotinia sclerotiorum (Lib.) Mass., watery soft-rot. N. J.

Sclerotium rolfsii Sacc., stem-rot. Southern States.

Septoria citrulli Ell. and Ev., leaf-spot. N. Y.

Thielavia basicola (Berk. and Br.) Zopf, black root-rot. Utah, Oreg.

Mosaic, undet. Del., Md., Mich.

CITRUS.

Citrus spp., citrus.

Tylenchus semipenetrans Cobb, nematode. Fla., Calif.

C. aurantifolia, lime.

Bacterium citri (Hasse) Jehle (*Phytomonas citri* (Hasse) Com. S. A. B.), citrus canker. Fla.

Colletotrichum gloeosporioides Penz. (*Glomerella cingulata* (Ston.) Spauld. and Schrenk), associated with citrus wither-tip. Fla.

Gloeosporium limeticolum R. F. Clausen, lime wither-tip. Fla.

Phytophthora terrestris Sherb., foot-rot. Fla.

Sphaeropsis tumefaciens Hedges, knot. Fla.

Sporotrichum citri Butl. (*Sphaceloma fawcettii* A. Jenkins), scab. Fla.

C. aurantium, sour, or Seville, orange.

Armillaria mellea (Vahl) Quel., root-rot. Calif.

Bacterium citrarefaciens H. A. Lee (*Phytomonas citrarefaciens* (Lee) Com. S. A. B.), citrus blast. Calif.

B. citri (Hasse) Jehle (*Phytomonas citri* (Hasse) Com. S. A. B.), citrus canker. Fla., Miss.

Colletotrichum gloeosporioides Penz. (*Glomerella cingulata* (Ston.) Spauld. and Schrenk), associated with citrus wither-tip. Fla., Tex., Calif.

Gloeosporium intermedium Sacc., anthracnose. Fla.

CITRUS—Continued.

C. aurantium, sour, or Seville, orange—Continued.

- Penicillium glaucum* Lk., blue-mold rot. La.
Phomopsis citri Fawc., stem-end rot, melanose. Fla., Ala.
Phytophthora terrestris Sherb., foot-rot. Fla.
Sporotrichum citri Butl. (*Sphaceloma fawcettii* A. Jenkins), scab. Gulf States.
 Chlorosis, undet. Calif.
 Die-back, undet. Fla.

C. grandis, grapefruit.

- Bacterium citrarefaciens* H. A. Lee (*Phytophthora citrarefaciens* (Lee) Com. S. A. B.), citrus blast. Calif.
B. citri (Hasse) Jehle (*Phytophthora citri* (Hasse) Com. S. A. B.), citrus canker. Fla., Ala., La., Miss.
Cephaleuros virescens Kze., algal leaf-spot. Fla.
Colletotrichum gloeosporioides Penz. (*Glomerella cingulata* (Ston.) Spauld. and Schrenk), associated with citrus wither-tip. Fla., Ala., Tex., Miss.
Diplodia natalensis Ev., stem-end rot. Fla.
Phomopsis citri Fawc., stem-end rot, melanose. Fla., Tex., Ariz.
Pythiacystis citrophthora E. H. and R. E. Smith, brown-rot. Ariz.
Sclerotium rolfsii Sacc., seedling blight (greenhouse). Ala.
Sporotrichum citri Butl. (*Sphaceloma fawcettii* A. Jenkins), scab. Fla., Ala.
Tylenchus semipenetrans Cobb, root nematode. Fla., Calif.
 Die-back, undet. Fla.
 Gummosis, undet. Fla.

C. limonia, lemon.

- Alternaria* sp., fruit-rot. Calif.
Armillaria mellea (Vahl) Quel., root-rot. Calif.
Bacterium citrarefaciens H. A. Lee (*Phytophthora citrarefaciens* (Lee) Com. S. A. B.), citrus blast. Calif.
B. citri (Hasse) Jehle (*Phytophthora citri* (Hasse) Com. S. A. B.), citrus canker. Fla., Ala., La., Tex.
Botrytis cinerea Auct., gray-mold rot. Calif.
Colletotrichum gloeosporioides Penz. (*Glomerella cingulata* (Ston.) Spauld. and Schrenk), associated with citrus wither-tip. Fla., Tex.
Gloeosporium spegazzinii Sacc., leaf-scorch. Fla.
Oospora citri-aurantii C. O. Smith, sour-rot. Calif.
Penicillium sp., blue-mold rot. Calif.
Phomopsis californica Fawc., rot. Calif.
P. citri Fawc., stem-end rot, melanose. Fla.
Phyllosticta hesperidearum (Catt.) Penz., leaf-spot. Colo. (greenhouse).
Phytophthora terrestris Sherb., foot-rot. Fla.
Pythiacystis citrophthora E. H. and R. E. Smith, brown-rot. N. J. (greenhouse), Calif.
Sclerotinia sclerotiorum (Lib.) Mass., cottony rot. Calif.
Sporotrichum citri Butl. (*Sphaceloma fawcettii* A. Jenkins), scab. Fla.
 Die-back, undet. Fla.
 Gummosis, undet. Calif.
 Internal decline, nonpar. Calif.

C. nobilis, king orange, tangerine.

- Bacterium citri* (Hasse) Jehle (*Phytophthora citri* (Hasse) Com. S. A. B.), citrus canker. Tex.
Colletotrichum gloeosporioides Penz. (*Glomerella cingulata* (Ston.) Spauld. and Schrenk), associated with citrus wither-tip. Miss., Calif.
Oidium tingitaninum C. N. Carter, powdery mildew. Calif.
Phomopsis citri Fawc., stem-end rot, melanose. Fla.
Sporotrichum citri Butl. (*Sphaceloma fawcettii* A. Jenkins), scab. Fla., Miss.
 Die-back, undet. Fla.

C. nobilis unshiu, Satsuma orange.

- Alternaria citri* N. B. Pierce, black-rot. Ala., Calif.
Bacterium citri (Hasse) Jehle (*Phytophthora citri* (Hasse) Com. S. A. B.), citrus canker. Fla.
Colletotrichum gloeosporioides Penz. (*Glomerella cingulata* (Ston.) Spauld. and Schrenk), associated with citrus wither-tip. Ala.
Diplodia natalensis Ev., stem-end rot. Ala., Miss.
Oospora sp., soft-rot. Ala., Miss.

CITRUS—Continued.

C. nobilis unshiu, Satsuma orange—Continued.*Penicillium digitatum* (Fr.) Sacc., green-mold rot. Ala.*P. italicum* C. Wehmer, blue-mold rot. Ala.*Phomopsis citri* Fawc., melanose, stem-end rot. Fla., Ala., Miss.

Die-back, undet. Fla.

Gummosis, undet. Fla.

C. sinensis, orange.*Alternaria citri* N. B. Pierce, black-rot. Fla., Ariz., Calif.*Bacterium citrarefaciens* H. A. Lee (*Phytomonas citrarefaciens* (Lee) Com. S. A. B.), citrus blast. Calif.*B. citri* (Hasse) Jehle (*Phytomonas citri* (Hasse) Com. S. A. B.), citrus canker. Gulf States.*Cerospora aurantia* Heald and Wolf, leaf-spot. Tex.*Colletotrichum gloeosporioides* Penz. (*Glomerella cingulata* (Ston.) Spauld. and Schrenk), die-back, associated with citrus wither-tip. Gulf States.*Diplodia natalensis* Ev., stem-end rot. Fla.*Fusarium limonis* Briosi, foot-rot. Ala.*Oospora* sp., soft-rot. La.*Penicillium* sp., blue-mold. General.*Phomopsis citri* Fawc., melanose, stem-end rot. Fla., Ala., Miss., La.*Phyllosticta adusta* Ell. and Mart., leaf-spot. Fla.*Phytophthora terrestris* Sherb., foot-rot. Fla.*Pythiacystis citrophthora* E. H. and R. E. Smith, brown-rot. Ariz.*Sclerotinia* sp., Calif.*Septobasidium* sp., girdle. Fla., La.*Sporotrichum citri* Butl. (*Sphaceloma fawcettii* A. Jenkins), scab. Fla., Miss., Tex.*Stemphylium citri* F. Patterson and V. K. Charles, end-rot. Ariz.*Tylenchus semipenetrans* Cobb, root nematode. Fla., Calif.

Chlorosis, undet. Fla., Calif.

Die-back, undet. Miss., Calif.

Greasy spot, undet. Fla.

Gummosis, undet. Fla., Calif.

Mottled-leaf, frenching, undet. Fla., Miss., Calif.

Psorosis, California scaly bark, undet. Fla., Ariz., Calif.

Scaly bark, attributed to *Cladosporium herbarum citricolum* Fawc. Fla.*C. trifoliata*, hardy orange.*Bacterium citri* (Hasse) Jehle (*Phytomonas citri* (Hasse) Com. S. A. B.) citrus canker. Ala., Fla.*Colletotrichum gloeosporioides* Penz. (*Glomerella cingulata* (Ston.) Spauld. and Schrenk), associated with citrus wither-tip. Fla., Tex.*Diplodia aurantii* Catt., twig-blight. Tex.*Phoma* sp., damping-off. Tex.*Phomopsis citri* Fawc., melanose. Fla., Miss.

Die-back, undet. Fla.

CLEMATIS.

Clematis sp., clematis, virgins-bower.*Ascochyta clematidina* (Thuem.) emend. W. O. Gloyer, stem-rot and leaf-spot. N. Y., Wis.*Bacterium tumefaciens* EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Minn.*Cercospora rubigo* Cke. and Hark., leaf-spot. Wash., Calif.*C. squalidula* Pk., leaf-spot. Frequent in widely scattered States.*Cylindrosporium clematidis* Ell. and Ev., leaf-spot. Conn., N. Y., Del., Miss., Wis.*Erysiphe polygoni* DC., powdery mildew. Frequent in widely scattered States.*Heterodera radiculicola* (Greef) Muell., root-knot. Ala., Ohio, Wash., Oreg.*Phleospora adusta* Heald and Wolf, leaf-blight. Tex.*Puccinia clematidis* (DC.) Lagh., rust. General.*P. stromatica* Berk. and Curt., rust. Ala.*Ramularia clematidis* Dearn. and Barth., leaf-spot. Mont.*Septoria jackmani* Ell. and Ev., leaf-spot. N. Y.

COCOS.

C. nucifera, coconut.

Alternaria sp., leaf-spot. Fla.

Colletotrichum sp., anthracnose. Fla.

Cytospora palmarum Cke., leaf-spot. Fla.

Diplodia epicocos Cke., stem-rot. Fla.

Fusarium sp., dry-rot. Fla.

Gloeosporium nanoti Prill. and Del., anthracnose. Fla.

Pestalozzia palmarum Cke., leaf-spot. Fla.

Phytophthora faberi Maub., bud-rot. Fla.

Pythium sp., wilt. Fla.

Ramularia sp., leaf-spot. Fla.

Thielaviopsis paradoxa (DeSeyn.) Hoehn., root-rot. Fla.

COFFEA.

C. arabica, coffee.

Glomerella cingulata (Ston.) Spauld. and Schrenk, anthracnose. Greenhouse, D. C.

Stilbum flavidum Cke., leaf-spot. Fla.

COLOCASIA.

C. esculenta, dasheen.

Bacillus carotovorus L. R. Jones (*Erwinia carotovora* (Jones) Com. S. A. B.), soft-rot. Storage.

Diplodia spp., Java black-rot. Storage.

Fusarium solani (Mart.) Sacc., associated with various rots. Fla.

Heterodera radicola (Greef) Muell., root-knot. Fla.

Sclerotium rolfsii Sacc., storage rot.

COMPTONIA.

C. asplenifolia, sweetfern.

Cronartium comptoniae Arth., rust. N. H., Vt., N. Y. to N. C., Wis.

CORNUS.

Cornus spp., dogwood.

Ascochyta cornicola Sacc., leaf-spot. Ohio, Wis.

Botrytis sp., twig-blight. Pa.

Cercospora cornicola Tr. and Earle, leaf-spot. Ga., Fla., Miss., Tex.

Diaporthe albocarnis Ell. and Ev., canker. Colo.

Dimarosporium pulchrum Sacc., sooty mold. Miss., Tex., Ill., Iowa.

Microsphaera alni (Wallr.) Wint., powdery mildew. Frequent in North-eastern States.

Myxosporium nitidum Berk. and Curt., twig-blight. Frequent in North-eastern States.

Phoma corni Fekl., canker. Minn.

Phyllactinia corylea (Pers.) Karst., powdery mildew. General.

Phyllosticta corni West., leaf-spot. W. Va.

P. globifera Ell. and Ev., leaf-spot. Kans.

Physalospora corni Ell. and Ev. Colo.

Septoria cornicola Desm., leaf-spot. General.

C. florida, flowering dogwood.

Cercospora cornicola Tr. and Earle, leaf-spot. Miss.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Phyllactinia corylea (Pers.) Karst., powdery mildew. S. C., Ala., Ind., Ill., Iowa.

Phyllosticta globifera Ell. and Ev., leaf-spot. Del., W. Va.

P. starbaeckii Sacc. and Syd., leaf-spot. Pa.

Septoria cornicola Desm., leaf-spot. Pa., S. C., Ind., Ky.

C. nuttalli, Pacific dogwood.

Fomes ignarius Fr., white wood-rot.

Nectria galligena Bres., European canker. Wash., Oreg.

Phyllactinia corylea (Pers.) Karst., powdery mildew. Wash., Oreg.

CORYLUS.

Corylus sp., filbert, hazelnut.

Bacterium sp., blight. Oreg.

Cryptosporella anomala (Pk.) Sacc., blight. Conn. to N. C.

Cylindrosporium vermiforme Davis, leaf-spot. Wis.

Gloeosporium coryli Ell. and Ev., leaf-spot. Me., N. J., Pa., Wash., Oreg.

Gnomoniella coryli (Batsch) Sacc., leaf-spot. General in Northern States.

Microsphaera alni (Wallr.) Wint., powdery mildew. Conn., Ill., Iowa, N. Dak.

Phyllactinia corylea (Pers.) Karst., powdery mildew. Northeastern States.

Phyllosticta coryli West., leaf-spot. N. J., Ind., Iowa, Mo.

Septogloeum profusum (Ell. and Ev.) Sacc., leaf-spot. Miss.

Septoria corylina Pk., leaf-spot. Mass., N. Y., Mich., Pacific coast.

Taphrina coryli T. Nishida, leaf-curl. Pa., Wis.

COSMOS.

Cosmos sp., cosmos.

Bacterium solanacearum EFS. (*Phytomonas solanaceara* (EFS.) Com. S. A. B.), bacterial wilt. N. C.

Erysiphe cichoracearum DC., powdery mildew. Nebr.

Phomopsis stewartii Pk., stem canker. Ohio, Mich., S. Dak.

Mosaic, undet. Fla.

CRATAEGUS.

Crataegus spp., hawthorn.

Bacillus amylovorus (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. Conn., N. Y., Ill., Utah, Wash., Oreg.

Cercospora crataegi Heald and Wolf, leaf-spot. Tex.

Cercosporella mirabilis Pk., leaf-spot. Colo.

Cladosporium carpophilum Thuem., scab. Iowa.

Cylindrosporium brevispina Dearn., leaf-spot. Mont., Idaho, Calif.

C. crataegi Ell. and Ev., leaf-spot. W. Va.

Fabraea maculata (Lév.) Atk., leaf-spot. General.

Fomes applanatus (Pers.) Wallr. (*Ganoderma applanatum* (Pers.) Pat.), white-streaked rot. Calif.

Gymnosporangium betheli Kern, rust. N. Mex., Northern Rocky Mt. States; Pacific Northwest.

G. blasdaleanum (Diet. and Holw.) Kern, rust. Oreg., Calif.

G. exiguum Kern, rust. Tex.

G. floriforme Thax., rust. Ala., Ark.

G. germinale (Schw.) Kern, rust. General east of Great Plains.

G. globosum Farl., rust. General east of Great Plains.

G. hyalinum (Cke.) Kern, rust. N. C. to Fla.

G. trachysorum Kern, rust. N. C., Ala., La., Miss., Tex.

G. tubulatum Kern, rust. Mont., Idaho.

Hendersonia crataegi Brenckle, leaf-spot. N. Dak.

H. foliorum Fckl., leaf-spot. Tex.

Phoradendron flavescens (Pursh) Nutt., mistletoe. Tex.

Phyllactinia corylea (Pers.) Karst., powdery mildew. N. Y., Fla., Ala., Ill., Iowa, Wash.

Phyllosticta crataegi (Cke.) Sacc., leaf-spot. N. Y., S. C., N. Dak., N. Mex.

P. grisea Pk., leaf-spot. N. Y.

P. solitaria Ell. and Ev., blotch. N. J.

Podosphaera oxyacanthae (DC.) D By., powdery mildew. East of Great Plains.

Venturia inaequalis (Cke.) Aderh., scab. N. Y.

CROTON.

Croton sp., croton.

Cercospora capitati Tharp, leaf-spot. S. C., Ala., Tex.

C. crotonis Ell. and Ev., leaf-spot. Kans.

C. maritima Tr. and Earle, leaf-spot. Miss.

Gloeosporium sp., leaf-spot. N. J.

Phakopsora crotonis (Burr.) Arth., rust. General in southern half of the United States.

Uromyces graminicola Burr., rust. Ill., Kans.

CUCUMIS.

C. melo, cantaloupe, muskmelon.

Bacillus melonis Gidd. (*Erwinia melonis* (Gidd.) Com. S. A. B.), soft-rot. Vt., Mass.

B. tracheiphilus EFS. (*Erwinia tracheiphila* (EFS.) Com. S. A. B.), bacterial wilt. General.

Cercospora cucurbitae Ell. and Ev., leaf-spot. Del.

Cladosporium cucumerinum Ell. and Arth., scab. Mass., Conn., N. J., Del., Ohio, Mich., Calif.

Colletotrichum lagenarium (Pass.) Ell. and Hals., anthracnose. General east of Great Plains.

Erysiphe cichoracearum DC., powdery mildew. Va., Fla.

Fusarium niveum EFS., wilt. General in southern half of the United States.

Heterodera radiculicola (Greef) Muell., root-knot. General in sandy soils in Southern States.

Macrosporium cucumerinum Ell. and Ev., leaf-blight. General.

Mycosphaerella citrullina (C. O. Smith) Gross., gummy stem-blight. N. Y.

Phyllosticta cucurbitacearum Sacc., leaf-spot. Ohio.

Pseudoperonospora cubensis (Berk. and Curt.) Rostew., downy mildew. General east of Great Plains, especially in Gulf and Atlantic Coast States.

Sclerotinia sclerotiorum (Lib.) Mass., timber rot. Mass.

Sclerotium rolfsii Sacc., stem-rot. South Atlantic and Gulf States.

Septoria cucurbitacearum Sacc., leaf-spot. N. Y., Wis.

Mosaic, undet. Frequent in widely scattered States.

C. sativus, cucumber.

Acremonium sp., leaf-blight. Mass., Ohio.

Bacillus sp., soft-rot. Fla.

B. melonis Gidd. (*Erwinia melonis* (Gidd.) Com. S. A. B.), soft-rot. Ala.

B. tracheiphilus EFS. (*Erwinia tracheiphila* (EFS.) Com. S. A. B.), bacterial wilt. General.

Bacterium lachrymans EFS. and Mary K. Bryan, angular leaf-spot. East of Great Plains; Colo., Calif. (in greenhouse).

Cercospora sp., leaf-spot. Ind.

Cladosporium cucumerinum Ell. and Arth., scab. General.

Colletotrichum lagenarium (Pass.) Ell. and Hals., anthracnose. East of Great Plains, Colo., Ariz., Calif.

Corticium vagum Berk. and Curt., stem-rot. Ohio, Wash.

Erysiphe cichoracearum DC., powdery mildew. Widespread, especially in Northern States.

Fusarium niveum EFS., wilt. Frequent east of Great Plains, especially in the South; N. Mex., Calif.

Heterodera radiculicola (Greef) Muell., root-knot. In sandy soils in Southern States.

Macrosporium cucumerinum Ell. and Ev., leaf-blight. General.

Mycosphaerella citrullina (C. O. Smith) Gross., gummy stem-blight. Fla.

Phyllosticta cucurbitacearum Sacc., leaf-spot. Ohio.

Pseudoperonospora cubensis (Berk. and Curt.) Rostew., downy mildew.

General east of Great Plains, especially in Gulf and Atlantic Coast States; Calif.

Pythium aphanidermatum (Edson) Fitz., cottony leak. N. C., S. C.

P. debaryanum Hesse, damping-off. Mass., N. Y., Ohio, Mich., Minn.

Sclerotinia sclerotiorum (Lib.) Mass., timber rot. Widespread.

Sclerotium rolfsii Sacc., stem-rot. S. C.

Stemphylium cucurbitacearum Geo. Osner, leaf-spot. Ohio, Ind.

Mosaic, undet. General.

CUCURBITA.

Cucurbita sp., gourd.

Bacterium lachrymans EFS. and Mary K. Bryan, angular leaf-spot.

Cercospora cucurbitacea Ell. and Gall., leaf-spot. Ala.

Colletotrichum lagenarium (Pass.) Ell. and Hals., anthracnose. Ill.

Pseudoperonospora cubensis (Berk. and Curt.) Rostew., downy mildew. Ohio, Ind.

C. maxima, squash; *C. pepo condensa*, summer squash.

Bacillus tracheiphilus EFS. (*Erwinia tracheiphila* (EFS.) Com. S. A. B.), bacterial wilt. East of Great Plains; Colo., Calif.

CUCURBITA—Continued.

C. maxima, squash; *C. pepo condensa*, summer squash—Continued.

Cercospora cucurbitae Ell. and Ev., leaf-spot. N. J., Ohio.

Choanephora cucurbitarum (Berk. and Rav.) Thax., brown-rot, blossom blight. Mass., Conn., N. Y., N. C., S. C., Fla., Ohio, Tex.

Cladosporium cucumerinum Ell. and Arth., scab. Me.

Colletotrichum lagenarium (Pass.) Ell. and Hals., anthracnose. Mass., Conn., Pa., Md., Ohio, Minn.

Corticium vagum Berk. and Curt., stem-rot. Mass.

Erysiphe cichoracearum DC., powdery mildew. Conn., Del., N. Mex., Utah.

Fusarium nivium EFS., wilt. N. C., Tex., Ohio, Nebr., N. Mex., Ariz., Calif.

Heterodera radiculicola (Greef) Muell., root-knot. Tex.

Macrosporium cucumerinum Ell. and Ev., leaf-blight. N. Y., Minn.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Pseudoperonospora cubensis (Berk. and Curt.) Rostew., downy mildew. Conn., Del., Va., S. C., Tex.

Rhizopus sp., black-mold rot. N. Y., Tex.

Septoria cucurbitacearum Sacc., leaf-spot. Ind.

Mosaic, undet. Vt., Conn., N. J., Pa., La., Tex., Minn.

C. pepo, pumpkin.

Bacillus tracheiphilus EFS. (*Erwinia tracheiphila* (EFS.) Com. S. A. B.), bacterial wilt. Ohio, Calif.

Botrytis cinerea Auct., gray-mold rot. Tex.

Cercospora cucurbitae Ell. and Ev., leaf-spot. Del.

Corticium vagum Berk. and Curt., stem-rot. Tex.

Erysiphe cichoracearum DC., powdery mildew. Conn., Del., N. Mex., Calif.

Heterodera radiculicola (Greef) Muell., root-knot. La., Tex.

Macrosporium cucumerinum Ell. and Ev., leaf-blight. W. Va., Minn.

Phyllosticta orbicularis Ell. and Ev., leaf-spot. N. Y., Del.

Pseudoperonospora cubensis (Berk. and Curt.) Rostew., downy mildew. Conn., N. J., Va., Ohio.

Septoria cucurbitacearum Sacc., leaf-spot.

CUPRESSUS.

C. arizonica, Arizona cypress.

Gymnosporangium cupressi Long and L. N. Goodding, rust. Ariz.

C. glabra, smooth cypress.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Phoma sp., nursery blight. Pa., Ill., Iowa, Nebr., Kans.

CYCAS.

C. revoluta, sago cycas.

Ascochyta cycadina G. Scalia, leaf-spot. Tex.

CYCLAMEN.

Cyclamen sp., cyclamen.

Botrytis cinerea Auct. Pa., Ohio, Calif.

Glomerella cingulata (Ston.) Spauld. and Schrenk, anthracnose. Mass., Va.

Heterodera radiculicola (Greef) Muell., root-knot. W. Va., Ohio, Ind., Nebr.

Phoma cyclamenae Hals., leaf-spot. Ohio.

Phyllosticta cyclaminicola Trel., leaf-spot. Ill.

Ramularia cyclaminicola Trel., leaf-spot. Ill., Minn.

Thielavia basicola (Berk. and Br.) Zopf, black root-rot. Conn.

CYDONIA.

C. japonica, flowering quince.

Bacillus amylovorus (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. Mich.

Gymnosporangium blasdaleanum (Diet. and Holw.) Kern, rust. Oreg.

G. germinale (Schw.) Kern, rust. Me., Conn., N. J.

CYDONIA—Continued.

C. oblonga, common quince.

- Bacillus amylovorus* (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. General.
- Bacterium tumefaciens* EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. N. J., Pa., Ohio, N. C., Tex., Wash., Oreg.
- Corticium stevensii* Burt (*Hypochnus ochroleucus* F. Noack), leaf-blight. N. C.
- Fabraea maculata* (Lév.) Atk., leaf-blight. General.
- Glomerella cingulata* (Ston.) Spauld. and Schrenk, bitter-rot. Mass., R. I., Pa., Va., W. Va., Mich., Ind., Kans.
- Gymnosporangium blasdaleanum* (Diet. and Holw.) Kern, rust. Oreg.
- G. clavariaeforme* (Jacq.) DC., rust. Me., N. H., Conn.
- G. germinale* (Schw.) Kern, rust. General east of Great Plains.
- G. nelsoni* Arth., rust. Colo.
- G. nidus-avis* Thax., rust. Conn., N. Y.
- Hendersonia cydoniae* Cke. and Ell., leaf-spot. Conn.
- Neofabraea malicorticis* (A. B. Cordley) Jack., anthracnose. Oreg.
- Ozonium omnivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.
- Phoma cydoniae* Sacc., pale-rot. Ill., Mich.
- P. pomi* Pass., fruit-spot. N. H., N. J., Pa., Md.
- Phyllactinia corylea* (Pers.) Karst., powdery mildew. Va.
- Phyalospora malorum* (Pk.) Shear (*P. cydoniae* Am. Auct.), black-rot. East of Great Plains.
- Podosphaera leucotricha* (Ell. and Ev.) Salm., powdery mildew. Wash.
- P. oxycanthae* (DC.) D By., powdery mildew. N. Y., Ill., Ind.
- Sclerotinia cinerea* (Bon.) Schroet. (*Monilia oregonensis* Barss and Posey), blossom blight. Wash., Oreg.
- S. fruticola* (Wint.) Rehm (*S. americana* (Wormald) Norton and Ezekiel), brown-rot. N. Y., W. Va., Oreg.

CYNARA.

C. scolymus, globe artichoke.

- Botrytis cinerea* Auct., rot. N. Y., Calif. General with host.
- Cercospora obscura* Heald and Wolf, leaf-spot. Tex.
- Cladosporium* sp., leaf-blight. S. C.
- Corticium vagum* Berk. and Curt., stem-rot. Miss.
- Ozonium omnivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.
- Sclerotium rolfsii* Sacc., stem-rot. Ga.

DACTYLIS.

D. glomerata, orchard grass.

- Claviceps microcephala* (Wallr.) Tul., ergot. Ind., Alaska.
- C. purpurea* (Fr.) Tul., ergot. N. Y., Va., W. Va., Minn.
- Colletotrichum graminicolum* (Ces.) Wils., anthracnose. Pa., Miss., Ohio, Minn.
- Erysiphe graminis* DC., powdery mildew. N. Y., Pa., Minn., Idaho, Utah.
- Gibberella saubinetii* (Mont.) Sacc., scab. W. Va.
- Helminthosporium* sp., leaf-spot. Minn.
- Puccinia graminis* Pers., stem-rust. General.
- Rhynchosporium secalis* (Oud.) Davis, scald. Wis.
- Sclerotium graminis* Fekl., leaf-spot. Widespread.
- Uromyces dactylidis* Otth, leaf-rust. Va., N. C.
- Ustilago striaeformis* (West.) Niessl, stripe-smut. General.

DAHLIA.

Dahlia sp., dahlia.

- Bacterium solanacearum* EFS. (*Phytomonas solanaceara* (EFS.) Com. S. A. B.), bacterial wilt. N. C.
- Botrytis* sp., bud-rot. N. J., Pa., Mich.
- B. sp.*, storage root-rot. N. J.
- Corticium vagum* Berk. and Curt., stem-rot. Md.
- Erysiphe cichoracearum* DC., powdery mildew. Conn., Md., N. C., W. Va., Ohio, Wash., Calif.

DAHLIA—Continued.*Dahlia* sp., dahlia—Continued.*E. polygoni* DC., powdery mildew. Del., Ga., Iowa, Mo., Calif.*Fusarium* sp., wilt. N. J.*Heterodera radicola* (Greef) Muell., root-knot. Miss., Calif.*Macrosporium* sp., leaf-blight. Pa., Mo.*Phoma dahliae* Berk., blight. Ohio.*Pythium debaryanum* Hesse, damping-off. Conn., Wash.*Sclerotinia sclerotiorum* (Lib.) Mass., stem-rot. Calif.*Sclerotium rolfsii* Sacc., stem-rot. Fla.*Sporotrichum* sp., root-rot. N. J.*Verticillium albo-atrum* Reinke and Berth., wilt.

Mosaic or yellows, undet. Conn., Del.

DAUCUS.*D. carota*, carrot.*Alternaria radicina* F. Meier, Drechs., and E. D. Eddy, black-rot. Mass., N. Y.*Bacillus carotororus* L. R. Jones (*Erwinia carotorora* (Jones) Com. S. A. B.), soft-rot. General, especially east of Rocky Mts.*Botrytis* sp., gray-mold rot. Pa., Wash.*Cercospora* sp., leaf-blight. Md.*C. apii carotae* Pass., leaf-spot. N. Y., Pa., N. J., Md., Ohio, Ind., Mich., Mo.,*Corticium vagum* Berk. and Curt., stem-rot. Mass., N. Y., Mich., Ill., Tex. Wash., Oreg.*Heterodera radicola* (Greef) Muell., root-knot. Occurs in widely scattered States.*Macrosporium carotae* Ell. and A. B. Langlois, leaf-blight. Occurs scatteringly in Eastern States.*Ozonium omnivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.*Rhizoctonia crocorum* (Pers.) DC., violet root-rot. Wash.*Sclerotinia intermedia* G. B. Ramsey, root-rot. Ill.*S. sclerotiorum* (Lib.) Mass., watery soft-rot. Conn., N. Y., N. J., La., Miss., Ind., Ill.*Sclerotium rolfsii* Sacc., stem-rot. S. C.**DELPHINIUM.***Delphinium* spp., larkspur.*Bacterium delphinii* (EFS.) Bryan, black-spot. Me., N. H., Mass., Conn., N. Y., Pa., Ill., Oreg.*Botrytis* sp., gray-mold. Conn.*Cercospora delphinii* Thuem., leaf-spot. Colo.*Corticium vagum* Berk. and Curt., stem-rot. N. Y., Mich.*Erysiphe polygoni* DC., powdery mildew. Me., Conn., N. Y., Pa., Minn.*Fusarium* sp., wilt. Fla.*Phyllosticta delphinii* F. E. Clements, leaf-spot. Colo.*Puccinia clematidis* (DC.) Lagh., rust. Nebr., Colo., Mont., Idaho, Oreg.*Sclerotium delphinii* D. S. Welch, crown-rot. N. Y., Pa., N. J., Ind.*S. rolfsii* Sacc., stem-rot. N. J., Fla., La., Ind., Ill.*Septoria delphinella* Sacc., leaf-spot. Kans.*Urocystis sorosporioides* Koern., smut. Calif.**DEUTZIA.***Deutzia* sp., deutzia.*Cercospora deutziae* Ell. and Ev., leaf-spot. Del.*Heterodera radicola* (Greef) Muell., root-knot. Miss.*Phyllosticta deutziae* Ell. and Ev., leaf-spot. N. J.**DIANTHUS.***D. barbatus*, sweet-william.*Corticium vagum* Berk. and Curt., stem-rot. Conn., N. Y.*Puccinia arenariae* (Schum.) Wint., rust. Conn., Mass., N. Y., Pa., Miss.*Septoria dianthi* Desm., leaf-spot. Ala.*Uromyces caryophyllinus* (Schränk) Wint., rust. Nebr.

DIANTHUS—Continued.

D. caryophyllus, carnation.

Alternaria dianthi F. L. Stevens and J. G. Hall, leaf-spot. Occurs scatteringly in Eastern States.

Bacterium woodsii EFS., bacterial spot. Okla.

Botrytis sp., bud-blight. Conn., Pa., Ohio, Ill.

Corticium vagum Berk. and Curt., stem-rot. General.

Fusarium sp., wilt. Widespread.

Heterodera radiculicola (Greef) Muell., root-knot. Mass., N. C., S. C., Ohio, Tex., Oreg., Calif.

Heterosporium echinulatum (Berk.) Cke., leaf-spot. Conn., N. Y., Oreg., Calif.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Septoria dianthi Desm., leaf-spot. Mass., Conn., N. Y., Del., S. C., Miss.

Sporotrichum anthophilum Pk., bud-rot. N. Y., N. J., Ind., Ill., Nebr.

S. poae Pk., bud-rot. Mass., Pa., N. J., Nebr.

Uromyces caryophyllinus (Schrank) Wint., rust. General.

Volutella dianthi (Hals.) Atk., anthracnose. Pa.

DIERVILLA.

D. trifida, bush-honeysuckle.

Phyllosticta diervillae Davis, leaf-spot. Wis.

Ramularia diervillae Pk., leaf-spot. N. H., N. Y., Wis.

Septoria diervillae Ell. and Ev., leaf-spot. Mass., Iowa, Minn., Wis.

DIGITALIS.

D. purpurea, foxglove.

Colletotrichum sp., anthracnose. Mass.

Phyllosticta digitalis A. Bellyneck, leaf-spot. N. Y., Alaska.

DIOSPYROS.

D. kaki, Japanese persimmon.

Cercospora diospyri Thuem., leaf-spot. S. C.

C. fuliginosa Ell. and Kell., leaf-spot. Fla., Tex., Ill.

C. kaki Ell. and Ev., leaf-spot. Ga., Ala., La., Tex.

Heterodera radiculicola (Greef) Muell., root-knot. Tex.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Phyllosticta biformis Heald and Wolf, leaf-spot. Tex.

Physalospora cydoniae Arn. (?), fruit-rot. Ala.

D. virginiana, common persimmon.

Cercospora atra Ell. and Ev., leaf-spot. Del.

C. diospyri Thuem., leaf-spot. General throughout range of host.

C. flexuosa Tr. and Earle, leaf-spot. Miss.

C. fuliginosa Ell. and Kell., leaf-spot. Ala., Kans., Tex.

C. virginiana Thuem., leaf-spot. Ala.

Gloeosporium diospyri Ell. and Ev., anthracnose. Del., W. Va., Ala., Miss.

Macrophoma diospyri Earle, fruit-spot. Ala.

Phoradendron flavescens (Pursh) Nutt., mistletoe. Tex.

Podosphaera oxycanthae (DC.) D By., powdery mildew. Ind.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

ELAEAGNUS.

E. angustifolia, Russian-olive.

Aecidium arctoum Arth., rust. N. Dak.

Cercospora elaeagni Heald and Wolf, leaf-spot. Tex.

Septoria argyraea Sacc., leaf-spot. Iowa, N. Dak.

E. argentea, silverberry.

Puccinia coronata Cda., crown-rust. N. Dak., Mont.

ERIOBOTRYA.

E. japonica, loquat.

Bacillus amylovorus (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. Fla., La., Calif.

Fusicladium eriobotryae Cav., scab. Fla., Calif.

ERYTHRINA.

E. crista-galli, coraltree.

Cercospora erythrinae Ell. and Ev., leaf-spot. La.

ESCHSCHOLTZIA.

E. californica, California poppy.

Entyloma eschscholtziae Hark., smut. Calif.

Heterodera radiculicola (Greef) Muell., root-knot.

EUCALYPTUS.

Eucalyptus spp., eucalyptus.

Bacterium tumefaciens EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Calif.

Botrytis sp., gray-mold. Calif.

Diaporthe eucalyptica Hark. Calif.

Fomes applanatus (Pers.) Wallr. (*Ganoderma applanatum* (Pers.) Pat.), white-streaked rot. Calif.

Hendersonia coryneoides Cke. and Hark., leaf-spot. Calif.

H. eucalypti Cke. and Hark., leaf-spot. Calif.

Macrophoma molleriana (Thuem.) Berl. Calif.

Pezizella lythri (Desm.) Shear and B. O. Dodge, leaf-spot. Calif.

Phyllosticta eucalypti Ell. and Ev., leaf-spot. Calif.

P. extensa Sacc. and Syd., leaf-spot. Calif.

Physalospora latitans Sacc. Calif.

P. suberumpens Ell. and Ev. Calif.

Polyporus schweinitzii Fr., red-brown butt-rot. Calif.

Septoria euthosporoides (Cke.) Sacc., leaf-spot. Calif.

S. mortolensis Penz. and Sacc., leaf-spot. Calif.

Damping-off (seedlings), undet. Calif.

Powdery mildew (seedlings), undet. Calif.

EUCHLAENA.

E. mexicana, teosinte.

Helminthosporium turcicum Pass., leaf-blight. Tex.

Physoderma zeae-maydis Shaw, brown-spot. Fla., La.

Puccinia sorghi Schw., rust. Nebr.

Ustilago zeae (Beckm.) Ung., smut. Conn., Ala., Ill., Iowa, Kans.

EUONYMUS.

Euonymus spp., burningbush, wahoo.

Bacterium tumefaciens EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Conn.

Cercospora destructiva Rav., leaf-spot. S. C., Miss.

C. evonymi Ell., leaf-spot. Ky., Ill., Nebr., Kans.

Cladosporium fasciculatum Cda. S. C.

Colletotrichum griseum Heald and Wolf, anthracnose. La., Tex.

Exosporium concentricum Heald and Wolf, leaf-spot. Tex.

Gloeosporium frigidum Sacc., anthracnose. Miss.

Marssonina thomasiana Sacc., leaf-spot. Mich., Wis., Ill.

Microsphaera alni (Wallr.) Wint., powdery mildew. Ind., Ill., Iowa, Nebr., Miss.

M. evonymi (DC.) Sacc., powdery mildew. N. J., Miss., Ohio, Minn.

Oidium evonymi-japonici (G. Arcangeli) Sacc., powdery mildew. La.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Phyllosticta evonymi Sacc., leaf-spot. Del.

P. pallens Ell. and Ev., leaf-spot. Ala., Tex., Ind.

Ramularia evonymi Ell. and Kell., leaf-spot. Mo., Kans.

Septoria evonymi Rabh., leaf-spot. Va.

EUPHORBIA.

E. marginata, snow-on-the-mountain.

Alternaria tenuis Nees, leaf-spot. Kans.

Microsphaera euphorbiae (Pk.) Berk. and Curt., powdery mildew. Ind., Mo., Kans.

Puccinia pammelii (Trel.) Arth., rust. Miss. Valley and Mo. Valley States.

Uromyces proeminens (DC.) Pass., rust. Great Plains, Upper Miss. Valley.

FAGOPYRUM.

F. esculentum, buckwheat.

Corticium vagum Berk. and Curt., stem-rot. N. C., Minn., Wash.

Phyllosticta polygonorum Sacc., leaf-spot. N. J.

Ramularia anomala Pk., leaf-spot. Ind.

R. rufo-maculans Pk., leaf-spot. Conn., Mich.

FAGUS.

F. americana, American beech.

Fomes applanatus (Pers.) Wallr. (*Ganoderma applanatum* (Pers.) Pat.), white-streaked rot. General throughout range of host.

F. fomentarius (L.) Fr., white sapwood rot. New England, N. Y., Pa.

F. igniarius (L.) Fr., white heart-rot. Mass.

Gloeosporium fagi (Desm.) Ell. and Ev., leaf-spot. Conn., N. Y., Pa.

Hydnum septentrionale Fr., white sapwood rot. Ind.

Microsphaera alni (Wallr.) Wint., powdery mildew. East of Miss. River.

Phyllactinia corylea (Pers.) Karst., powdery mildew. N. Y., Ohio, Ill.

Phyllosticta faginea Pk., leaf-spot. N. Y., Ohio, Ind., Wis.

FESTUCA.

Festuca spp., fescue.

Claviceps purpurea (Fr.) Tul., ergot. Me., N. Y., Pa., Va., Ill., Minn.

Colletotrichum graminicolum (Ces.) Wils., anthracnose. Pa., Minn.

Corticium vagum Berk. and Curt., stem-rot. General.

Helminthosporium dictyoides Drechs., net-blotch. New England, N. Y., Pa., Md., D. C., Va.

Ophiobolus graminis Sacc., take-all. N. Y., Ark.

Pleomeris piperi (Ricker) Syd., rust. Utah, Oreg., Calif.

Puccinia abundans (Pk.) Jack., rust. N. Dak., Colo., Utah, Calif.

P. coronata Cda., crown-rust. Conn., N. Y., Pa., Ky., Ind., Mo., Nebr., Kans., Colo., Oreg.

P. ephylla (L.) Wetts., rust. Oreg.

P. graminis Pers., stem-rust. N. Y., Pa., Tenn., Ind., Mich., Minn., Iowa, Colo., Wyo., Wash., Oreg., Calif.

Septoria tenella Cke. and Ell., leaf-spot. N. J., Kans., Colo.

Tilletia fusca Ell. and Ev., covered smut. Western States.

Uromyces hordei Tr., rust. Miss., Tex.

Ustilago mulfordiana Ell. and Ev., smut. Western States.

U. striaeformis (West.) Niessl, stripe-smut. Md., N. Dak.

FICUS.

F. aurea, Florida strangler fig.

Bacterium tumefaciens EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Fla.

Cerotelium fici (Cast.) Arth., rust. Fla.

Colletotrichum elasticae F. Tassi, anthracnose. Fla.

Ophiodothella fici E. Bessey, leaf-spot. Fla.

F. carica, fig.

Aspergillus niger Tiegh., fruit-rot (smut). Calif.

Botrytis cinerea Auct., twig-blight, canker. Wash., Calif.

Cercospora fici Heald and Wolf, leaf-spot. General in South.

Cerotelium fici (Cast.) Arth., rust. General in South.

Colletotrichum elasticae F. Tassi, anthracnose. Miss.

Corticium laetum Karst., limb-blight. Ga., Ala., La., Miss.

Diplodia sycina syconophila Sacc., die-back. Tex.

Fusarium moniliforme Sheld., fruit-rot. Calif.

Glomerella cingulata (Ston.) Spauld. and Schrenk, anthracnose. Gulf States.

Heterodera radicola (Greef) Muell., root-knot. Gulf States, N. Mex., Ariz.

Macrophoma fici Alm. and Cam., canker, fruit-rot. Tex.

Mycosphaerella bolleana Hig., leaf-spot. General in South.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Rhizoctonia microsclerotia J. Matz, leaf-blight. Fla., Ala., La., Tex.

Rhizopus nigricans Ehr., rot. Ala., La.

Sclerotinia sclerotiorum (Lib.) Mass., die-back.

Tubercularia fici Edg., canker. Ala., La.

FICUS—Continued.

F. elastica, India rubber-tree.

Glomerella cingulata (Ston.) Spauld. and Schrenk, anthracnose. Miss.

Phyllosticta roberti L. Boyer and Jacz., leaf-spot. Fla.

FILIPENDULA.

Filipendula sp., meadowsweet.

Cylindrosporium sp. Conn.

Septoria ulmariae Oud., leaf-spot. Conn.

Sphaerotheca humuli (DC.) Burr., powdery mildew. Vt.

Triphragmium ulmariae (Schum.) Lk., rust. Ind.

FORSYTHIA.

Forsythia sp., forsythia, *goldenbell.

Bacterium tumefaciens EFS. and Town. (*Phytoplasma tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Miss.

Heterodera radiculicola (Greef) Muell., root-knot. Miss.

Phyllosticta discincola Ell. and Ev., leaf-spot. Md.

P. terminalis Ell. and Mart., leaf-spot. Fla.

FRAGARIA.

Fragaria sp., strawberry.

Armillaria mellea (Vahl) Quel., root-rot. Oreg.

Botrytis cinerea Auct., gray-mold rot. General.

Corticium vagum Berk. and Curt., crown-rot. Wash.

Dendrophoma obscurans (Ell. and Ev.) H. W. Anderson, angular spot. N. J., Md., Va., W. Va., S. C., Ill., Mich., Minn., Mo.

Diplocarpon earliana (Ell. and Ev.) Wolf (*Marssonina fragariae* Sacc.), leaf-scorch. East of Miss. River; Wis., Mont.

Heterodera radiculicola (Greef) Muell., root-knot. Ark., Miss., Wash.

Mycosphaerella fragariae (Tul.) Lindau, leaf-spot. General.

Pezizella lythri (Desm.) Shear and B. O. Dodge, tan brown-rot. Me., Md., Va., Fla., La., Ark., Ohio, Wis.

Phyllosticta fragaricola Desm. and Rob., leaf-spot. N. J., Fla.

Phytophthora cactorum (Leh. and Cohn) Schroet., leather rot. Ky., Tenn., Ark., Ill., Mo., Miss., La.

Pythium sp., lilac soft-rot. Tenn.

Rhabdospora fragaria Atk., leaf-spot. Ill.

Rhizoctonia sp., hard brown-rot (fruit). Tenn., N. C., Fla.

Rhizopus nigricans Ehr., leak. General east of Great Plains.

Sclerotinia sclerotiorum (Lib.) Mass., crown-rot. La., Miss., Tenn., Ark., Mo., Ohio.

Sclerotium rolfsii Sacc., stem-rot. Fla.

Sphaeronemella fragariae F. L. Stevens and A. Peterson, fruit-rot. Fla., La.

Sphaeropsis sp., black-rot. Ill.

Sphaerotheca humuli (DC.) Burr., powdery mildew. General.

Tylenchus dipsaci (Kuehn) Bast., stem nematode. Wash., Oreg.

Black-root, undet. Tenn., Fla., Ala., Mich., Wis., Colo.

FRAXINUS.

Fraxinus spp., ash.

Cercospora fraxinea Ell. and Ev., leaf-spot. Kans.

C. fraxinites Ell. and Ev., leaf-spot. Fla., La., Tex.

C. superflua Ell. and Holw., leaf-spot. Iowa.

C. texensis Ell. and Gall., leaf-spot. Tex.

Cercospora trichophila Davis, leaf-spot. Wis.

Cladosporium epiphyllum (Pers.) C. Martius, leaf-spot. N. Y.

Cylindrosporium californicum Earle, leaf-spot. Oreg.

C. fraxini Ell. and Kell., leaf-spot. Md., Mich., Kans., Pacific coast.

C. minor Ell. and Kell., leaf-spot. Kans.

C. viridis Ell. and Kell., leaf-spot. La., Tex., Wash.

Fomes fraxinophilus Pk., white heartwood rot. Widespread.

Gloeosporium aridum Ell. and Holw., anthracnose. Conn., Mass., Pa., Ohio, Mich., Wis.

G. decipiens Ell. and Ev., spot. Iowa, Kans., Tex.

G. fraxini Hark., anthracnose. N. Y., Mich., Calif.

Leptosphaeria fraxini Ell. and Ev. Mo.

Marssonina fraxini Ell. and Davis, leaf-spot. Wis.

FRAXINUS—Continued.

Fraxinus spp., ash—Continued.

- Ozonium omnivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.
Phoradendron flavescens (Pursh) Nutt., mistletoe. Tex., Ind., Ariz.
Phyllactinia corylea (Pers.) Karst., powdery mildew. Northeastern States.
Phyllosticta fraxinicola (F. Currey) Ell. and Ev., leaf-spot. East of Great Plains
P. viridis Ell. and Kell., leaf-spot. Widespread.
Piggotia fraxini Berk. and Curt., leaf-spot. General.
Puccinia fraxinata (Lk.) Arth., rust. General east of Great Plains.
Ramularia fraxinea Davis, leaf-spot. Wis.
Septoria besseyi Pk., leaf-spot. Wis., Iowa.
S. fraxini Desm., leaf-spot. Fla., La., Mich., Minn.
S. leucostroma Ell. and Ev., leaf-spot. Mass., Ind.
S. submaculata Wint., leaf-spot. Mo., Tex.

GAILLARDIA.

G. aristata, gaillardia.

Bullaria gaillardiae (Diet. and Holw.) Arth. and E. B. Mains (*Uredo gaillardiae* Diet. and Holw.), rust. Calif.

Erysiphe cichoracearum DC., powdery mildew. Mont.

G. pulchella, rose ring gaillardia.

Entyloma polysporum (Pk.) Farl., smut. Nebr., Kans.

Septoria gaillardiae Ell. and Ev., leaf-spot. Kans.

GARDENIA.

G. florida, Cape-jasmine.

Heterodera radicularis (Greef) Muell., root-knot. Tex., Calif. (greenhouse).

GARRYA.

G. elliptica, silktassel-bush.

Cercospora garryae Hark., leaf-spot. Calif.

C. glomerata Hark., leaf-spot. Calif.

Phyllosticta garryae Cke. and Hark., leaf-spot. Calif.

GAULTHERIA.

G. shallon, salal.

Phyllosticta gaultheriae Ell. and Ev., leaf-spot. Pacific coast.

GAYLUSSACIA.

Gaylussacia sp., huckleberry.

Exobasidium vaccinii (Fekl.) Wor., hypertrophy. Atlantic Coast States; Mich., Wis.

Guignardia vaccinii Shear, scald. Md.

Microsphaera alni vaccinii (Schw.) Salm., powdery mildew. Mass., N. J., Md., Ill., Wis.

Pucciniastrum myrtilli (Schum.) Arth., rust. Mass., Conn., N. Y., Pa., Va.

Rhytisma vaccinii Fr., tar-spot. Mass.

Synchytrium vaccinii F. Thomas, red-gall. N. J.

GELSEMIUM.

G. sempervirens, Carolina-jessamine.

Phyllosticta gelsemii Ell. and Ev., leaf-spot. N. J.

GLADIOLUS.

Gladiolus sp., gladiolus.

Alternaria fasciculata (Cke. and Ell.) L. R. Jones and A. J. Grout. Pa.

Bacterium gummisudans McC., bacterial blight. Mich., Minn., N. Dak.

B. marginatum McC. (*Phytophthora marginata* (McC.) Com. S. A. B.), stem-rot, scab. Widely distributed.

Corticium vagum Berk. and Curt., root-rot. N. Dak.

Didymellina iridis (Desm.) Hoehn., leaf-spot. Md.

Fusarium sp., rot. Widely distributed.

Phyllosticta gladioli Ell. and Ev., leaf-spot. Ala.

Sclerotium sp., dry-rot. Widespread.

Septoria gladioli Pass., hard-rot, leaf-spot. General wherever host is grown.

GLEDITSIA.

G. triacanthos, honeylocust.

Cercospora condensata Ell. and Kell., leaf-spot. Ill., Kans.

C. olivacea (Berk. and Rav.) Ell., leaf-spot. Ala., Miss., Ill., Iowa.

Cladosporium gleditschiae Cke. S. C.

Fomes applanatus (Pers.) Wallr. (*Ganoderma applanatum* (Pers.) Pat.), white-streaked rot. N. Y.

Melasmia hypophylla (Berk. and Rav.) Sacc., leaf-spot. Widespread east of Great Plains.

Microsphaera alni (Wallr.) Wint., powdery mildew. General east of Great Plains.

Phoradendron flavescens (Pursh) Nutt., mistletoe. Ind., Tex.

Polyporus sulphureus (Bull.) Fr., brown-checked wood-rot.

Ravenelia opaca Diet., rust. Ill.

GORDONIA.

G. lasianthus, loblolly-bay.

Phyllosticta gordoniae Ell. and Mart., leaf-spot. Fla.

GOSSYPIUM.

G. hirsutum, Upland cotton, *G. barbadense*, Sea Island cotton.

Alternaria sp., leaf-spot. S. C.

Ascochyta gossypii Syd., blight. Va., N. C., S. C., Ark.

Bacterium malvacearum EFS. (*Phytomonas malvacearum* (EFS.) Com. S. A. B.), angular leaf-spot. General in Cotton Belt States; Ariz.

Cercospora althaeina Sacc., leaf-spot. Ill.

Cerotelium gossypii (Lagh.) Arth. (*Uredo gossypii* Lagh.), rust. Fla.

Corticium vagum Berk. and Curt., stem-rot, sore-shin. Cotton Belt States; Ariz.

Diplodia gossypina Cke., boll-rot. General in Cotton Belt States.

Fusarium spp., boll-rot. General in Cotton Belt States during wet seasons.

F. vasinfectum Atk., wilt. General in sandy soils of Cotton Belt States.

Glomerella gossypii (E. A. Southworth) Edg., anthracnose. Cotton Belt States.

Heterodera radiculicola (Greef) Muell., root-knot. General in sandy soils in Cotton Belt States.

Macrosporium nigricantium Atk., black leaf-spot. Cotton Belt States; Ariz.

Mycosphaerella gossypina (Cke.) Earle, leaf-spot. General in Cotton Belt States.

Olpitrichum carpophilum Atk., boll-rot. N. C., S. C., Ala., La.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex., Okla., N. Mex., Ariz., Calif.

Phlyctaena gossypii Sacc., leaf-spot. S. C., Ala.

Phyllosticta gossypina Ell. and Mart., leaf-spot. N. C., S. C., Ala., Miss., Ill.

Puccinia hibisciata (Schw.) Kell. (*Aecidium gossypii* Ell. and Ev.), rust. Tex., Ariz., Calif.

Pythium debaryanum Hesse, damping-off. Ala.

Ramularia areola Atk., frosty mildew. Cotton Belt States; Fla.

Septoria gossypina Cke., leaf-spot. S. C., Miss.

Thielavia basicola (Berk. and Br.) Zopf, black root-rot. Ala.

Tylenchus penetrans Cobb, nematode. N. C., Ga.

Verticillium alboatrum Reinke and Berth., wilt. Va.

Malnutrition, "rust," nonpar. Poorer soils, Cotton Belt States.

GYMNOCLADUS.

G. dioicus, Kentucky coffeetree.

Cercospora gymnocladi Ell. and Kell., leaf-spot. Ill., Nebr., Kans., Mo.

Marssonina sp., leaf-spot. Nebr.

HAMAMELIS.

H. virginiana, witch-hazel.

Cercospora hamamelidis Ell. and Ev., leaf-spot. Fla.

Gonatobotryum maculicolum (Wint.) Sacc., leaf-spot. N. Y.

Phyllactinia corylea (Pers.) Karst., powdery mildew. Wis.

HAMAMELIS—Continued. *

H. virginiana, witch-hazel—Continued.

Phyllosticta hamamelidis Cke., leaf-spot. General east of Miss. River.

Podosphaera biuncinata Cke. and Pk., powdery mildew. General east of Miss. River.

Ramularia hamamelidis Pk., leaf-spot. N. Y., W. Va., Mich., Ill., Wis.

HEDERA.

H. helix, English ivy.

Glomerella cingulata (Ston.) Spauld. and Schrenk, anthracnose. Conn., Md., N. C., Tex.

Phyllosticta concentrica Sacc., leaf-spot. N. Y., N. J., S. C., Ala., Tex.

P. hederacola Dur. and Mont., leaf-spot. N. Y., N. J.

Ramularia hedericola Heald and Wolf, leaf-spot. Tex.

Vermicularia trichella Fr., leaf-spot. Mass., S. C., Md., Ohio.

HELIANTHUS.

H. annuus, sunflower.

Armillaria mellea (Vahl) Quel., root-rot. Oreg.

Botrytis sp., stem-rot. Oreg., Calif.

Cercospora pachypus Ell. and Kell., leaf-spot. Kans., Tex.

Entyloma polysporum (Pk.) Farl., smut. Mont.

Erysiphe cichoracearum DC., powdery mildew. General.

Fusarium sp., damping-off. N. J.

Heterodera radicola (Greef) Muell., root-knot. W. Va., Ala.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Plasmopara halstedii (Farl.) Berl. and DeToni, downy mildew. Ohio, W. Va., Ind., Iowa, Minn., N. Dak., Kans.

Puccinia helianthi-mollis (Schw.) Jack., rust. General.

Sclerotinia sclerotiorum (Lib.) Mass., stem-rot. Mich., Wis., Minn., Iowa

N. Dak., Mont., Idaho, Wash., Oreg.

Septoria helianthi Ell. and Kell., leaf-spot. Widespread.

H. tuberosus, Jerusalem-artichoke.

Cercospora helianthi Ell. and Ev., leaf-spot. Kans.

Coleosporium helianthi (Schw.) Arth., rust. Va., S. C., Ala., Ill.

Erysiphe cichoracearum DC., powdery mildew. Northeastern and North-Central States.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Plasmopara halstedii (Farl.) Berl. and DeToni, downy mildew. Vt., Del., Wis., Iowa, Kans.

Puccinia helianthi-mollis (Schw.) Jack., rust. East of Great Plains.

Sclerotinia sp., wilt. Wash.

Sclerotium rolfsii Sacc., stem-rot. S. C., La.

HELLEBORUS.

H. niger, Christmas-rose.

Coniothyrium hellebori Cke. and Mass. N. Y., Oreg.

HEMEROCALLIS.

H. fulva, daylily.

Cercospora hemerocallis L. R. Tehon, leaf-spot. Ill.

Didymellina iridis (Desm.) Hoehn., leaf-spot. N. J.

HIBISCUS.

H. esculentus, okra, gumbo.

Ascochyta abelmoschi Harter, pod-spot. N. Y., Md., S. C.

Botrytis sp., rot. N. Y.

Cercospora althaeina Sacc., leaf-spot. La.

C. brachypoda Speg., leaf-spot. La.

C. hibisci Tr. and Earle, leaf-spot. Fla., Ala., La., Miss.

Corticium vagum Berk. and Curt., stem-rot. Ala., Tex.

Erysiphe cichoracearum DC., powdery mildew. Conn.

Fusarium vasinfectum Atk., wilt. Southern States; Conn., Pa., Ariz.

Heterodera radicola (Greef) Muell., root-knot. Southern States.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex., Ariz.

HIBISCUS—Continued.

H. esculentus, okra, gumbo—Continued.*Phyllosticta hibiscina* Ell. and Ev., leaf-spot. Ill.*Verticillium alboatrum* Reinke and Berth., wilt. N. J., S. C., Ala., Oreg., Calif.*H. syriacus*, *rose-of-Sharon.*Heterodera radicola* (Greef) Muell., root-knot. Tex.*Kuehneola malvicola* (Speg.) Arth., rust. La., Miss.*Nectria* sp., canker. Ohio.*Ozonium omnivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.*Phyllosticta syriaca* Sacc., leaf-spot. N. Y.

HICORIA.

Hicoria spp., hickory.*Armillaria mellea* (Vahl) Quel., root-rot. Ind.*Bacterium tumefaciens* EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Md., Kans.*Fomes applanatus* (Pers.) Wallr. (*Ganoderma applanatum* (Pers.) Pat.), white-streaked rot. Conn.*Fusicladium effusum* Wint., scab. Ill., Kans.*Gnomonia caryae* Wolf, anthracnose. East of Miss. River; Mo.*Hendersonia davisii* Ell. and Ev., leaf-spot. Wis.*Marssonina juglandis* (Lib.) Sacc., leaf-spot. N. J., W. Va., Iowa.*Microsphaera alni* (Wallr.) Wint., powdery mildew. Ala., Miss., Ind., Ill., Mo.*Microstroma juglandis* (Bereng.) Sacc., witches'-broom. General east of Great Plains.*Mycosphaerella convexula* (Schw.) F. V. Rand, leaf-spot. S. C., Ga., Fla., Ala., Ohio.*Phoradendron flavescens* (Pursh) Nutt., mistletoe. Tex., Ind.*Phyllactinia corylea* (Pers.) Karst., powdery mildew. Wis.*Phyllosticta caryae* Pk., leaf-spot. Northeastern and North-Central States; Gulf States.*P. convexula* Bub., leaf-spot. Mo.*P. subtilis* Pk., leaf-spot. N. Y.*Rosellinia caryae* L. Bonar, canker. Mich.*Septoria caryae* Ell. and Ev., leaf-spot. Del.*S. hickoriae* Tharp, leaf-spot. Tex.*Strumella coryneoides* Sacc. and Wint., canker. Pa.*H. pecan*, pecan.*Bacterium tumefaciens* EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. S. C., Gulf States, Calif.*Botryosphaeria berengeriana* DeNot., die-back. S. C., Ala., Fla.*Cercospora fusca* (Heald and Wolf) emend. F. V. Rand, brown leaf-spot. S. C., Gulf States, N. Mex.*Fusicladium effusum* Wint., scab. N. C. to Okla. and Tex.*Glomerella cingulata* (Ston.) Spauld. and Schrenk, anthracnose. S. C., Gulf States.*Heterodera radicola* (Greef) Muell., root-knot.*Microsphaera alni* (Wallr.) Wint., powdery mildew. S. C., Gulf States.*Microstroma juglandis* (Bereng.) Sacc., leaf-spot. Ga., Ala., Tex.*M. juglandis robustum* Hig., flower-blight. Ga.*Mycosphaerella convexula* (Schw.) F. V. Rand, leaf-blotch. S. C., Ga., Ala., Fla., Ohio.*Oidium* sp., powdery mildew. Tex.*Ozonium omnivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.*Pestalozzia* sp., leaf-spot. Fla.*Phoma* sp., twig-blight. Fla.*Phoradendron flavescens* (Pursh) Nutt., mistletoe. Tex.*Phyllosticta caryae* Pk., leaf-spot. Va., S. C., Gulf States.*Septoria caryae* Ell. and Ev., leaf-spot. Tex.Kernel-spot, southern stinkbug (*Nezara viridula* L.). N. C., S. C., Ga., Fla., Ala., Miss., La., Tex.

Rosette, undet. Del., Va., N. C., S. C., Gulf States; Ark., Calif.

HOLCUS.

H. halepensis, Johnson grass.

Cercospora sorghi Ell. and Ev., leaf-spot. Ala., La., Tex., Nebr.

Colletotrichum lineola Cda., anthracnose. Ala., La., Tex.

Helminthosporium turcicum Pass., leaf-blight. Ala., Tex.

Phyllosticta sorghina Sacc., leaf-spot. Tex.

Puccinia purpurea Cke., rust. S. C., Fla., Ala., La., Okla., Tex., Calif.

Septoria pertusa Heald and Wolf, leaf-spot. Tex.

Sphacelotheca sorghi (Lk.) Clint., smut. Utah, Wash.

H. sorghum, sorghum.

Bacterium andropogoni EFS., stripe. General.

Cercospora sorghi Ell. and Ev., leaf-spot. Ga., Ala., La., Miss.

Colletotrichum lineola Cda., anthracnose. Conn., W. Va., N. C., Ga., Fla., Ala., La., Tex., Minn.

Helminthosporium turcicum Pass., leaf-blight. Kans., Tex.

Puccinia purpurea Cke., rust. Tex., Okla., La., Fla., Calif.

Sclerotium rolfsii Sacc., stem-rot. La.

Sorosporium reilianum (Kuehn) McAlp., head-smut. General.

Sphacelotheca cruenta (Kuehn) Potter, loose kernel smut. Ala., Tex., Ark., N. Mex.

S. sorghi (Lk.) Clint., covered kernel smut. General.

**H. sorghum saccharatus*, sorgo, sugar sorghum.

Bacterium andropogoni EFS., stripe.

Colletotrichum lineola Cda., anthracnose. N. J., Va.

Phoma insidiosa F. Tassi, leaf-spot. Tex., Miss., Ark., Kans.

Puccinia purpurea Cke., rust. Tex.

Sphacelotheca sorghi (Lk.) Clint., covered kernel smut. Colo.

H. sorghum sudanensis, Sudan grass.

Colletotrichum graminicolum (Ces.) Wils., anthracnose. N. C., Ala., Tex., N. Dak.

C. lineola Cda., anthracnose. Ala.

Helminthosporium turcicum Pass., leaf-blight. Tex.

Phoma insidiosa F. Tassi, leaf-spot. Va.

Sorosporium reilianum (Kuehn) McAlp., head-smut. Wash.

Sphacelotheca sorghi (Lk.) Clint., covered kernel smut. N. C., Nebr., Kans.

**H. sorghum technicus*, broomcorn.

Bacterium andropogoni EFS., stripe.

Colletotrichum lineola Cda., anthracnose. Occurs in widely scattered States.

Sorosporium reilianum (Kuehn) McAlp., head-smut. Ohio.

Sphacelotheca cruenta (Kuehn) Potter, loose kernel smut. Tex.

S. sorghi (Lk.) Clint., covered kernel smut. Ohio.

HOLODISCUS.

H. discolor, *oceanspray.

Cylindrosporium filipendulae Thuem., leaf-spot. Wash.

Phyllactinia corylea (Pers.) Karst., powdery mildew. Wash.

Podosphaera oxyacanthae (DC.) D By., powdery mildew. Idaho.

HORDEUM.

H. vulgare, barley.

Bacterium translucens L. R. Jones, A. G. Johnson, and C. S. Reddy, bacterial blight. S. C., Tex., Ark., Ind., Wis., Minn., Iowa, N. Dak., S. Dak., Mont., Colo., Idaho, Oreg.

Claviceps purpurea (Fr.) Tul., ergot. Infrequent but widely distributed.

Colletotrichum graminicolum (Ces.) Wils., anthracnose. General east of Great Plains.

Erysiphe graminis DC., powdery mildew. General.

Fusarium culmorum leteius Sherb., seedling blight. Oreg., Calif.

Gibberella saubinetii (Mont.) Sacc., scab. General east of Great Plains.

Helminthosporium californicum W. W. Mackie and G. E. Paxton, rusty blotch. Calif.

H. gramineum Rabh., stripe. General.

H. sativum Pam., King, and Bak., spot-blotch. General.

Marasmius tritici P. A. Young. Ill.

Ophiobolus graminis Sacc., take-all, foot-rot. Calif.

HORDEUM—Continued.*H. vulgare*, barley—Continued.

- Puccinia anomala* Rostr., leaf-rust. General.
P. glumarum (Schm.) Eriks. and Henn., stripe-rust. West of Great Plains.
P. graminis Pers., stem-rust. General.
Pyrenophora teres (Died.) Drechs., net-blotch. General.
Rhizoctonia sp., foot-rot. Wash.
Rhynchosporium secalis (Oud.) Davis, scald. Upper Miss. Valley; Pacific Northwest; Calif.
Septoria passerinii Sacc., leaf-spot. Mich., Minn., Iowa, N. Dak., S. Dak., Idaho, Oreg.
Ustilago hordei (Pers.) Kell. and Sw., covered smut. General.
U. nuda (Jens.) Kell. and Sw., loose smut. General.

HUMULUS.*H. lupulus*, hop.

- Armillaria mellea* (Vahl) Quel., root-rot. Oreg.
Colletotrichum sp., anthracnose. N. Y.
Cylindrosporium humuli Ell. and Ev., leaf-spot. N. Y., N. J., Del., Mich., Wis., Iowa.
Erysiphe cichoracearum DC., powdery mildew. Utah.
Phyllosticta decidua Ell. and Kell., leaf-spot. Wis.
P. humuli Sacc. and Speg., leaf-spot. Mass., Iowa.
Plasmopara humuli Miyabe and Tak., downy mildew. Wis.
Septoria lupulina Ell. and Kell., leaf-spot. Kans.
Sphaerotheca humuli (DC.) Burr., powdery mildew. Me., N. Y., Ala., Wyo., Pacific coast.

HYACINTHUS.*H. orientalis*, common hyacinth.

- Bacterium hyacinthi* J. H. Wakker (*Phytomonas hyacinthi* (Wakker) Com. S. A. B.), yellow disease. Mass.
Fusarium sp., bulb-rot (dry). Mo., Wash.
F. bulbigenum Cke. and Mass., bulb-rot. R. I.
Tylenchus dipsaci (Kuehn) Bast., stem and bulb nematode. Wash.
Mosaic, gray disease, undet. Wash.

HYDRANGEA.*Hydrangea* sp., hydrangea.

- Ascochyta* sp., leaf-spot. Pa.
Botrytis cinerea Auct., blight. Wash.
Cercospora hydrangeana Tharp, leaf-spot. Ala., Tex.
Heterodera radiculicola (Greef) Muell., root-knot. Miss.
Phyllosticta hydrangeae Ell. and Ev., leaf-spot. N. J., Ala., Ind.
Pucciniastrum hydrangeae (Berk. and Curt.) Arth., rust. Frequent east of Miss. River.
Septoria hydrangeae Bizz., leaf-spot. Miss.

HYDRASTIS.*H. canadensis*, goldenseal.

- Botrytis* sp., stem-rot. Conn., N. Y., Ind., Mich., Wis., Wash.
Fusarium sp., wilt. N. Y., Ohio, Ill.
Heterodera radiculicola (Greef) Muell., root-knot. Ohio.

ILEX.*Ilex* spp., holly, winterberry.

- Cercospora ilicis* Ell., leaf-spot. N. J., Ala., Miss.
Microsphaera alni (Wallr.) Wint., powdery mildew. Ala., Ill., Wis.
Phyllosticta concomitans Ell. and Ev., leaf-spot. La.
P. haynaldi Roum. and Sacc., leaf-spot. W. Va.
P. ilicicola (Cke. and Ell.) Ell. and Ev., leaf-spot. N. Y.
P. opaca Ell. and Ev., leaf-spot. W. Va.
P. terminalis Ell. and Mart., leaf-spot. Fla.
Ramularia alaterni Thuem., leaf-spot. Mich.
R. prinus Pk., leaf-spot. N. Y., Wis.
Rhytisma concavum Ell. and Kell., tar-spot. Ohio.
R. ilicicolum Schw., tar-spot. Va., Miss.

ILEX—Continued.

Ilex spp., holly, winterberry—Continued.

R. ilicis-canadensis Schw., tar-spot. Me., Mich., Ind.

R. velatum (Schw.) Fr., tar-spot. Miss.

Septoria ilicifolia Cke. and Ell., leaf-spot. N. J.

I. opaca, American holly.

Cercospora ilicicola Maub., leaf-spot. Tex.

C. pulvinula Cke. and Ell., leaf-spot. N. J., Tex.

Diplodia sp., foot-rot. Fla.

Melampsoropsis ilicinum (Ell. and Ev.) Jack., rust.

Phyllactinia corylea (Pers.) Karst., powdery mildew. Ill.

Phyllosticta ilicicola (Cke. and Ell.) Ell. and Ev., leaf-spot. N. Y., N. J.

P. opaca Ell. and Ev., leaf-spot. W. Va., S. C.

P. terminalis Ell. and Mart., leaf-spot. Miss.

Rhytisma curtisii Berk. and Rav., tar-spot. Md., S. C., Ala., Tex.

R. ilicis-canadensis Schw., tar-spot. W. Va.

IPOMOEA.

I. batatas, sweetpotato.

Albugo ipomoeae-panduranae (Schw.) Sw., white-rust. General where sweetpotatoes are grown.

Botrytis cinerea Auct., gray-mold rot. N. J.

Coleosporium ipomoeae (Schw.) Burr., rust. Miss.

Corticium vagum Berk. and Curt., stem-rot. Frequent in sweetpotato States.

Cystospora batata (Ell. and Hals.) J. A. Elliott, soil rot, pox. Middle Atlantic and Gulf States, Ohio, Kans., Ark., Okla.

Diaporthe batatatis (Ell. and Hals.) Harter and E. Field, dry-rot. N. J. to Kans. and Tex.

Diplodia tubericola (Ell. and Ev.) Taub., Java black-rot. General, Del. to Kans., Ga., Tex.

Fusarium batatatis Woll. and *F. hyperoxysporum* Woll., stem-rot. General in sweetpotato States.

F. oxysporum Schl., surface rot. N. J., Ala., Miss.

Heterodera radicola (Greef) Muell., root-knot. Southern States, Calif.

Monilochaetes infuscans Ell. and Hals., scurf. General.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Phyllosticta batatas (Thuem.) Cke., leaf-spot. N. J. and Ohio to Kans. and Tex.

Plenodomus destruens Harter, foot-rot. N. J. to Iowa and southward, Calif.

Pythium sp., rootlet rot. N. Y., Del., Md., Va. and in States south and west.

Rhizopus nigricans Ehr., soft-rot, ring-rot. General as a storage rot.

Sclerotinia sp., damping-off. N. J.

Sclerotium bataticola Taub., charcoal rot. Frequent throughout country as a storage rot.

S. rolfsii Sacc., stem-rot. S. C. to Tex.

Septoria bataticola Taub., leaf-spot. Iowa to Tex. and eastward.

Sphaeronema fimbriatum (Ell. and Hals.) Sacc., black-rot. General in all sweetpotato States.

Trichoderma koningi Oud., storage rot. N. J., Del.

Mosaic, undet. Fla., Miss., Ark., Tex., Kans.

Mottle-necrosis (associated with *Pythium* sp.). N. J., D. C.

I. purpurea, morning-glory.

Albugo ipomoeae-panduranae (Schw.) Sw., white-rust. Del., Va., W. Va., Ga., Ala., Miss., Kans.

Cercospora alabamensis Atk., leaf-spot. Ala.

C. ipomoeae Wint., leaf-spot. Ala.

C. viridula Ell. and Ev., leaf-spot. N. J., N. C., Nebr.

Coleosporium ipomoeae (Schw.) Burr., rust. Miss. Valley States, Ala., Tex.

Heterodera radiciola (Greef) Muell., root-knot.

Puccinia crassipes Berk. and Curt., rust. Okla.

IRIS.

Iris sp., iris.

- Bacillus carotovorus* L. R. Jones (*Erwinia carotovora* (Jones) Com. S. A. B.), soft-rot. Conn., N. Y., N. J., Ohio, Ind., Mich., Minn.
Cylindrosporium iridis Ell. and Hals., leaf-spot. Ind., Iowa.
Didymellina iridis (Desm.) Hoehn., leaf-spot. Widespread in northern two-thirds of United States.
Macrosporium iridicolum Ell. and Ev., leaf-spot. Idaho.
M. iridis Cke. and Ell., leaf-spot. Ala., Ohio, Mont.
Pseudomonas iridis Van Hall, rhizome rot. S. Dak.
Puccinia iridis (DC.) Rabh., rust. Conn., N. Y., Mich., Colo., Pacific coast.
Sclerotinia sclerotiorum (Lib.) Mass., stem-rot. Wash.
Sclerotium rolfii Sacc., crown-rot. Md., Ind., Miss., Calif.
Scolecotrichum iridis Fautr. and Roum., leaf-spot. Miss., Minn.
S. punctulatum Tr. and Earle, leaf-spot. Miss.
Septoria sp., leaf-spot. N. Dak.

I. versicolor, blueflag iris.

- Cylindrosporium iridis* Ell. and Hals., leaf-spot. Ind.
Didymellina iridis (Desm.) Hoehn., leaf-spot. Ind.
Phyllosticta iridis Ell. and Mart., leaf-spot. N. Y., Fla., Ohio, Mich.
Puccinia iridis (DC.) Rabh., rust. Northeastern States.
P. majanthae (Schum.) Arth. and Holw., rust. Mass., N. Y., Ind., Mich., Wis., Minn., Iowa, Nebr.

JUGLANS.

J. cinerea, butternut.

- Cercospora juglandis* Kell. and Sw., leaf-spot. Mass.
Fomes igniarius (L.) Fr., white heart-rot. Vt., N. J.
Gnomonia leptostyla (Fr.) Ces. and DeNot., anthracnose. General throughout range of host.
Melanconis juglandis (Ell. and Ev.) A. H. Graves, canker. East of Miss. River; Wis., Iowa.
Microsphaera alni (Wallr.) Wint., powdery mildew. Ill.
Microstroma juglandis (Bereng.) Sacc., leaf-spot. General over range of host.
Phyllactinia coryleae (Pers.) Karst., powdery mildew. Ind.
Polyporus sulphureus (Bull.) Fr., brown-checked wood-rot.

J. hindsii, Hinds walnut.

- Cylindrosporium juglandis* Wolf, leaf-spot. Calif.
Microstroma juglandis (Bereng.) Sacc., leaf-spot. Calif.

J. nigra, black walnut.

- Armillaria mellea* (Vahl) Quel., root-rot. Oreg.
Bacterium tumefaciens EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Kans., Calif.
Cercospora juglandis Kell. and Sw., leaf-spot. Kans.
Fomes everhartii (Ell. and Gall.) Schrenk, heart-rot.
F. igniarius (L.) Fr., white heart-rot. Ohio.
Gnomonia leptostyla (Fr.) Ces. and DeNot., anthracnose. General throughout range of host.
Melanconis juglandis (Ell. and Ev.) A. H. Graves, canker. N. Y.
Microsphaera alni (Wallr.) Wint., powdery mildew. Ind., Ill., Kans., Miss.
Microstroma juglandis (Bereng.) Sacc., leaf-spot. Del., Md., N. C., Ohio, Ill., Mo., Kans.
Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.
Phoradendron flavescens (Pursh) Nutt., mistletoe. Ind.
Phleospora multimaculans Heald and Wolf, leaf-spot. Tex.
Polyporus sulphureus (Bull.) Fr., brown-checked wood-rot.
Septoria juglandis Berk. and Curt., leaf-spot. Pa., N. C.

J. regia, Persian walnut (English walnut).

- Armillaria mellea* (Vahl) Quel., root-rot. Oreg., Calif.
Bacterium juglandis (N. B. Pierce) EFS., bacterial blight. N. Y., Del., Va., Pacific coast.
B. tumefaciens EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Pacific coast.

JUGLANS—Continued.

J. regia, Persian walnut (English walnut)—Continued.

Cylindrosporium sp., leaf-spot. Ala.

C. juglandis Wolf, leaf-spot. N. C.

Dothiorella gregaria Sacc., melaxuma. Calif.

Glomerella cingulata (Ston.) Spauld. and Schrenk, anthracnose. Md.

Melanconis juglandis (Ell. and Ev.) A. H. Graves, canker. Conn., N. J.

Microstroma juglandis (Bereng.) Sacc., leaf-spot. Conn., Calif.

Sphaeropsis sp., canker. Conn.

J. sieboldiana, Japanese walnut.

Bacterium tumefaciens EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Pa.

Melanconis juglandis (Ell. and Ev.) A. H. Graves, canker. Conn., N. Y., N. J.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

JUNIPERUS.

Juniperus spp., juniper, redcedar.

Cercospora sequoiae juniperi Ell. and Ev., leaf-spot. Wis.

Cyanospora albicedrae Heald and Wolf, whitening. Tex. (*J. montana*).

Fomes earlei (Murr.) Sacc. and D. Sacc., yellow pocket rot. Tex., Colo., N. Mex., Ariz.

F. juniperinus Schrenk, white pocket rot. Central States.

F. roseus (Alb. and Schw.) Fr., brown pocket rot. Fla., Ala.

F. texanus (Murr.) Hedge. and Long, brown fibrous rot. Tex., N. Mex., Ariz.

Gymnosporangium bermudianum (Farl.) Earle, rust. Ala., Miss.

G. betheli Kern, rust. N. Dak., Mont., Colo., N. Mex.

G. clavariaeforme (Jacq.) DC., rust. East of Great Plains.

G. corniculans Kern, rust. Mich., Wis.

G. cornutum (Pers.) Arth., rust. Mich., Colo.

G. davisii Kern, rust. Me., Wis.

G. exiguum Kern, cedar rust. Tex.

G. germinale (Schw.) Kern., rust. Eastern States.

G. globosum Farl., rust. Eastern States.

G. gracilens (Pk.) Kern and Bethel, rust. Colo., Utah, N. Mex., Ariz., Nev.

G. harknessianum (Ell. and Ev.) Kern, rust. Oreg.

G. inconspicuum Kern, rust. Colo., Utah.

G. japonicum Syd., rust. Conn., Mass., Oreg.

G. juniperinum (L.) C. Martius, rust. Iowa, Colo., Utah, Wash.

G. juniperi-virginianae Schw., rust. East of Great Plains.

G. juvenescens Kern, rust. Mont., Colo., Utah, N. Mex., Idaho.

G. kernianum Bethel, rust. Colo., N. Mex., Utah, Idaho.

G. multiporum Kern, rust. Colo.

G. nelsoni Arth., rust. Northern Rocky Mt. States.

G. nidus-avis Thax., rust. Mass., N. Y., Nebr.

G. trachysorum Kern, rust. Fla.

G. tubulatum Kern, rust. Mont.

Herpotrichia nigra Hartig, matting. Colo. (*J. sibirica* and *J. communis*).

Lophodermium juniperinum (Fr.) DeNot., leaf-cast. Iowa, Colo.

Phomopsis juniperovora G. Hahn, nursery blight. Pa., Ohio, Ill., Mo., Iowa, Nebr., Kans.

Phoradendron juniperinum Engelm., mistletoe. Southwestern and Western States.

J. virginiana, redcedar.

Cercospora sequoiae juniperi Ell. and Ev., leaf-spot. Wis.

Fomes annosus Fr., white spongy rot. Del.

F. juniperinus Schrenk, white pocket rot. Md.

F. roseus (Alb. and Schw.) Fr., brown pocket rot. General.

Gymnosporangium bermudianum (Farl.) Earle, rust. Fla., La., Miss.

G. corniculans Kern, rust. N. Y.

G. effusum Kern, rust. N. Y., N. J., S. C., Va.

G. exiguum Kern, rust. Tex.

G. exterum Arth. and Kern, rust. Ky.

G. floriforme Thax., rust. Gulf States.

JUNIPERUS—Continued.*J. virginiana*, redcedar—Continued.

- G. germinale* (Schw.) Kern, rust. General east of Great Plains.
G. globosum Farl., rust. General east of Great Plains.
G. juniperi-virginianae Schw., rust. General east of Great Plains.
G. juvenescens Kern, rust. Wis., Nebr.
G. nidus-avis Thax., rust. General east of Miss. River.
G. trachysorum Kern, rust. S. C., Fla., Miss.
Phomopsis juniperovora G. Hahn, nursery blight. N. Y. to Kans.
Poria purpurea Fr., brown laminated rot. Iowa.

KALMIA.*K. latifolia*, mountain-laurel.

- Cercospora kalmiae* Ell. and Ev., leaf-spot. Conn., Pa., W. Va.
C. sparsa Cke., leaf-spot. S. C., Miss.
Mycosphaerella colorata (Pk.) Earle, leaf-spot. N. Y., Pa., Ala.
Pestalotzia kalmicola Ell. and Ev., leaf-spot. N. J., Del.
Phomopsis kalmiae Enlows, leaf-blight. D. C.
Phyllosticta kalmicola (Schw.) Ell. and Ev., leaf-spot. N. Y., N. J., Pa., Del., Md., Miss.
P. latifolia Ell. and Ev., leaf-spot. N. J., Del., Miss.
Septoria angustifolia Ell. and Ev., leaf-spot. Atlantic Coast States; Ohio.
Venturia kalmiae Pk., leaf-spot. N. Y.

KERRIA.*K. japonica*, kerria.

- Coccomyces kerriae* V. B. Stewart, twig and leaf blight. Conn., Del., Ala., Ky.
Cylindrosporium sp., leaf-spot. Md.
Septoria sp., leaf-spot. N. J., Md.

LACTUCA.*L. sativa*, lettuce.

- Alternaria* sp., leaf-spot. Ohio, Fla., La., Wash.
Aplanobacter rhizoctonia R. C. Thomas, bacterial rosette. Ohio.
Bacillus carotovorus L. R. Jones (*Erwinia carotovora* (Jones) Com. S. A. B.), soft-rot. General.
Bacterium marginale N. A. Brown (*Phytomonas marginale* (Brown) Com. S. A. B.), leaf-spot. N. J., Kans.
B. viridilividum N. A. Brown (*Phytomonas viridilivida* (Brown) Com. S. A. B.), lettuce rot. Va., La., Mich.
B. vitians N. A. Brown (*Phytomonas vitians* (Brown) Com. S. A. B.), bacterial wilt. N. J., Va., S. C., N. Mex.
Botrytis cinerea Auct., gray-mold rot. General.
Bremia lactucae E. Regel, downy mildew. General.
Corticium vagum Berk. and Curt., bottom rot. General.
Heterodera radiculicola (Greef) Muell., root-knot. Southern States.
Macrosporium cladosporioides Desm., leaf-spot. Fla.
Marssonina panattoniana (Berl.) Magn., anthracnose. N. Y., N. C., Ohio, Mich., Kans., Utah, Oreg., Wash.
Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex., Ariz.
Puccinia hieraciata (Schw.) Jack., rust. Minn., Ind.
Pythium debaryanum Hesse, damping-off. Conn., N. Y., Pa., Fla., Ohio, Mich.
Sclerotinia minor I. C. Jagger, drop, pink-rot. Mass., N. Y., Pa., Fla.
S. sclerotiorum (Lib.) Mass., drop. General.
Sclerotium rolfsii Sacc., stem-rot. Fla.
Septoria lactucae Pass., leaf-spot. East of Great Plains.
Mosaic, undet. Conn., N. Y., Pa., S. C., Fla., Minn.
Rio Grande disease (rabbit-ear, white-heart), undet. N. Y., Miss., La., Tex., Ill.
Tipburn, nonpar. Widespread.

LAGERSTROEMIA.*L. indica*, crapemyrtle.

- Cercospora lythracearum* Heald and Wolf, leaf-spot. Tex.
Phyllosticta lagerstroemiae Ell. and Ev., leaf-spot. La., Tex.
Powdery mildew, undet. S. C., Ga., Ala., La.

LARIX.

L. europaea, European larch.

Melampsora bigelowii Thuem., rust. N. Y., Ind.

M. medusae Thuem., rust. Ind.

L. laricina, American larch, or tamarack.

Fomes officinalis Fr., reddish brown-rot. Wis.

F. pinicola Fr., brown crumbly rot. General with host.

Melampsora bigelowii Thuem., rust. N. Y., Wis.

M. medusae Thuem., rust. N. Y., Mich., Ind.

Melamporidium betulae (Schum.) Arth., rust. Wis.

Polyporus schweinitzii Fr., red-brown butt-rot.

Razoumofskyia laricis Piper, mistletoe witches'-broom. Idaho.

R. pusilla (Pk.) O. Kuntze, mistletoe. Northern States.

Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Minn.

L. occidentalis, western larch.

Botrytis sp., seedling twig-blight. Idaho.

Fomes officinalis Fr., reddish brown-rot. Mont., Idaho.

F. pinicola Fr., brown crumbly rot. Idaho.

F. roseus (Alb. and Schw.) Fr., brown pocket rot. General with host.

Hypodermella laricis Tub., leaf-cast. Mont., Idaho, Oreg.

Lophodermium laricinum Duby, leaf-cast. Idaho, Wash.

Melampsora bigelowii Thuem., rust. Idaho.

Polyporus berkeleyi Fr., string and ray rot. Idaho.

P. schweinitzii Fr., red-brown butt-rot. Idaho.

P. sulphureus (Bull.) Fr., brown-checked wood-rot. Idaho.

Razoumofskyia laricis Piper, mistletoe witches'-broom. Mont., Pacific Northwest.

Rhizina inflata (Schaeff.) Sacc., seedling blight. Idaho.

Sparassis radicata Weir, yellow root-rot. Mont., Pacific Northwest.

Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Idaho.

LATHYRUS.

L. odoratus, sweet pea.

Bacillus lathyrus Manns and Taub., streak. Mass., N. Y., Pa., Del.

Colletotrichum pisi Pat., anthracnose. S. C., Ga.

Corticium vagum Berk. and Curt., stem-rot. Widespread.

Erysiphe polygona DC., powdery mildew. General.

Fusarium lathyrus Taub., wilt, stem-rot. Conn., Mass., N. Y., Va., Fla., Ind., N. Dak.

F. martii pisi F. R. Jones, root-rot. N. Y.

Glomerella cingulata (Ston.) Spauld. and Schrenk, anthracnose. Occurs scatteringly east of Great Plains.

Heterodera radiculicola (Greef) Muell., root-knot.

Microsphaera alni (Wallr.) Wint., powdery mildew. General.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Va.

Phytophthora cactorum (Leb. and Cohn) Schroet., root-rot. Conn.

Pythium debaryanum Hesse, root-rot. Conn.

Sclerotinia sclerotiorum (Lib.) Mass., stem-rot, collar rot. Pa.

Sclerotium rolfsii Sacc., stem-rot. Fla.

Thielavia basicola (Berk. and Br.) Zopf, black root-rot. Conn., N. Y., Pa., Del., Ohio, Calif.

Mosaic, undet. Atlantic coast; Ohio, Wis., N. Mex.

LAVATERA.

Lavatera sp., treemallow.

Colletotrichum malvarum A. Braun and Casp., anthracnose. Calif.

Puccinia malvacearum Bertero, rust. Calif.

LESPEDeza.

**L. striata*, Japan clover.

Heterodera radiculicola (Greef) Muell., root-knot. Ala.

Microsphaera diffusa Cke. and Pk., powdery mildew. S. C., Ala., Miss,

LIBOCEDRUS.

L. decurrens, incense-cedar.

Gymnosporangium blasdaleanum (Diet. and Holw.) Kern, rust. Oreg., Calif.
Herpotrichia nigra Hartig, matting. Calif.
Phoradendron juniperinum libocedri Engelm., mistletoe. Nev., Oreg., Calif.
Polyporus amarus Hedge, pecky heartwood rot. Oreg., Calif.

LIGUSTRUM.

Ligustrum sp., privet.

Armillaria mellea (Vahl) Quel., root-rot. Miss.
Bacterium tumefaciens EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Miss.
Cercospora adusta Heald and Wolf, leaf-spot. Tex.
C. ligustri Roum., leaf-spot. Miss.
Exosporium concentricum Heald and Wolf, leaf-spot. Tex.
Glomerella cingulata (Ston.) Spauld. and Schrenk, anthracnose. Widespread east of Great Plains.
Heterodera radicolica (Greef) Muell., root-knot. Miss.
Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.
Phyllosticta ovalifolia Brun., leaf-spot. Tex.

LILIUM.

Lilium spp., lily (mostly native species).

Corticium vagum Berk. and Curt., stem-rot. Mich.
Uromyces holwayi Lagh., rust. Widespread.

L. longiflorum, Easter lily.

Botrytis cinerea Auct., gray-mold. Fla.
Cercospora unicolor Sacc. and Penz., leaf-spot. Fla.
Fusarium sp., wilt. Fla.
Sclerotinia sclerotiorum (Lib.) Mass., wilt. Fla.

LINUM.

L. usitatissimum, flax.

Alternaria sp., boll disease. Minn., N. Dak.
Colletotrichum linicolum Peth. and Laf., anthracnose. Mich., Minn., N. Dak., Mont.
Fusarium lini H. L. Bolley, wilt. General in States where flax is grown.
Heterodera radicolica (Greef) Muell., root-knot.
Macrosporium sp., blight. N. Dak.
Melampsora lini (Schum.) Desm., rust. General.
Phlyctaena linicola Speg., pasmo. Mich., Wis., Minn., N. Dak., S. Dak.
Rhizoctonia sp., root-rot. Minn., N. Dak.
Heat canker, nonpar. Wis., Minn., N. Dak., S. Dak., Mont., Idaho.

LIQUIDAMBAR.

L. styraciflua, sweetgum.

Cercospora liquidambaris Cke. and Ell., leaf-spot. Md., Ga., Gulf States.
C. tuberculans Ell. and Ev., leaf-spot. La., Miss., Mo.
Hymenochaete agglutinans Ell., twig-blight.
Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.
Phoradendron flavescens (Pursh) Nutt., mistletoe. Ohio, Tex.
Septoria liquidambaris Cke. and Ell., leaf-spot. Mass., N. J., Ga.
Stereum subpileatum Berk. and Curt., honeycomb heart-rot. Fla.

LIRIODENDRON.

L. tulipifera, tuliptree.

Cercospora liriodendri Ell. and Hark., leaf-spot. Middle Atlantic States, Gulf States.
Cylindrosporium cercosporoides Ell. and Ev., leaf-spot. Md.
Erysiphe polygoni DC., powdery mildew. General over range of host.
Gloeosporium liriodendri Ell. and Ev., anthracnose. Conn., Del., Tex., Ohio.
Myxosporium longisporium Edg., twig canker. N. Y.
Phyllactinia corylea (Pers.) Karst., powdery mildew. N. Y., Ala., Ind., Ill.

LIRIODENDRON—Continued.*L. tulipifera*, tuliptree—Continued.

Phyllosticta liriodendrica Sacc., leaf-spot. General over range of host.

Polyporus sulphureus (Bull.) Fr., brown-checked wood-rot.

Rhytisma liriodendri Wallr., tar-spot. Ohio.

Verticillium sp., canker. Ohio.

LOLIUM.*L. multiflorum*, Italian ryegrass.

Claviceps purpurea (Fr.) Tul., ergot. Wash., Oreg.

Colletotrichum sp., anthracnose. Wash.

Helminthosporium siccans Drechs., leaf-spot. Md., Va.

L. perenne, perennial ryegrass.

Claviceps purpurea (Fr.) Tul., ergot. Va., W. Va., S. Dak., Nebr., Kans., Idaho, Calif.

Colletotrichum graminicolum (Ces.) Wils., anthracnose. Pa.

Gibberella sp., root-rot. Pa.

Helminthosporium siccans Drechs., leaf-spot. Md., Va.

Puccinia coronata Cda., rust. Pacific coast.

P. glumarum (Schm.) Eriks. and Henn., stripe-rust. Wash., Oreg.

P. graminis Pers., stem-rust. Minn.

LONICERA.*Lonicera* spp., honeysuckle.

Bacterium tumefaciens EFS. and Town., crown-gall. Conn.

Cercospora antipus Ell. and Holw., leaf-spot. Iowa, Nebr.

C. varia Pk., leaf-spot. Tex.

Fusarium sp., stem-blight. Ohio.

Gloeosporium lonicerae Hark., leaf-spot. Calif.

Leptothyrium periclymeni (Desm.) Sacc. Wis., Idaho.

Marssonina lonicerae Hark., leaf-spot. Colo.

Microsphaera alni (Wallr.) Wint., powdery mildew. Frequent in northern half of United States.

Puccinia festucae Plow., rust. Iowa.

LUPINUS.*L. perennis*, sun-dial lupine.

Ascochyta pisi Lib. (*Mycosphaerella pinodes* (Berk. and Blox.) R. E. Stone), leaf-spot.

Cercospora filispora Pk., leaf-spot. N. Y.

C. longispora Pk., leaf-spot. N. Y.

Erysiphe polygoni DC., powdery mildew. Mass., N. Y., N. J., Md., Ohio, Ill., Mich.

Sclerotinia sclerotiorum (Lib.) Mass., stem-rot. N. Y.

L. polyphyllus, Washington lupine.

Uromyces lupini Berk. and Curt., rust. Wash.

LYCOPERSICUM.*L. esculentum*, tomato.

Alternaria sp. (not *A. solani* nor *M. tomato*), leaf-spot. Calif.

A. solani (Ell. and Mart.) L. R. Jones and A. J. Grout, early-blight. General east of Great Plains; nailhead spot especially in Southern States.

Aplanobacter michiganense EFS. (*Phytomonas michiganense* (EFS.) Com. S. A. B.), bacterial canker (Grand Rapids disease). Mass., Conn., N. J., Pa., Ga., Ohio, Ind., Mich.

Ascochyta lycopersici Brun. (*Didymella lycopersici* Kleb., *Diplodina destructiva* (Plow.) Petr.), leaf-spot. N. J.

Bacillus aroideae Town. (*Erwinia aroidea* (Town.) Com. S. A. B.), soft-rot. Va.

B. carotovorus L. R. Jones (*Erwinia carotovora* (Jones) Com. S. A. B.), soft-rot. N. J., Fla.

Bacterium solanacearum EFS. (*Phytomonas solanaceara* (EFS.) Com. S. A. B.), bacterial wilt. For the most part east of Great Plains and south of Missouri and Ohio Rivers.

LYCOPERSICUM—Continued.*L. esculentum*, tomato—Continued.

- B. vesicatorium* Doidge (*Phytophthora blight*) (Gardner and Kendrick) Com. S. A. B.), bacterial spot. N. Y., Pa., Tenn., Ga., Fla., Ind., Ill., Mich., Iowa, Kans.
- Basisporium gallarum* M. Molliard, fruit-rot. Calif.
- Botrytis cinerea* Auct., gray-mold rot. N. J., Ohio, Ky., Ala.
- Cercospora cruenta* Sacc., leaf-spot. S. C., Ala., Iowa.
- Cladosporium fulvum* Cke., leaf-mold. General.
- Colletotrichum phomoides* (Sacc.) Chester, anthracnose. East of Great Plains; Wash., Calif.
- Corticium vagum* Berk. and Curt., stem-rot, soil-rot. General.
- Fusarium lycopersici* Sacc., Fusarium wilt. Conn. to Nebr. and southward, Calif.
- Glomerella cingulata* (Ston.) Spauld. and Schrenk, anthracnose. N. Y., Mich., Fla., La.
- Heterodera radicola* (Greef) Muell., root-knot. General in South.
- Isaria clonostachyodes* F. J. Pritchard and W. S. Porte, fruit-rot. Va., D. C.
- Macrosporium tomato* Cke., nailhead spot. Fla. and other Southern States.
- Melanconium* sp., fruit-spot. Ohio, Wis.
- Nematosporea lycopersici* A. Schneider, fruit-rot. Calif.
- Olpidium* sp., seedling blight. Wis.
- Oospora lactis parasitica* F. J. Pritchard and W. S. Porte, fruit-rot. N. J., Va., D. C.
- Ozonium omnivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex., Ariz.
- Phoma destructiva* Plow. (*Diplodina destructiva* (Plow.) Petr.), ripe-rot, leaf-spot. Widespread.
- Phyllosticta* sp., leaf-spot. Fla.
- Phytophthora* sp. (D. Reddick), stem girdle. N. Y.
- P. infestans* (Mont.) D By., late-blight, damping-off. Atlantic and Pacific Coast States.
- P. terrestris* Sherb., buckeye rot. South Atlantic and Gulf States; Va., Ark., Tenn., Ind.
- Pythium debaryanum* Hesse, damping-off. New England, N. Y., Ohio, Pa., Wash.
- Rhizopus nigricans* Ehr., black-mold rot. Frequent in harvested crop.
- Sclerotinia sclerotiorum* (Lib.) Mass., stem-rot, fruit-rot. General.
- Sclerotium rolfsii* Sacc., stem-rot. Southern States.
- Septoria lycopersici* Speg., leaf-spot. General, but infrequent in West.
- Synchytrium endobioticum* (Schilb.) Perc., wart. Pa.
- Verticillium albo-atrum* Reinke and Berth., wilt. Northern States.
- V. lycopersici* F. J. Pritchard and W. S. Porte, collar rot. N. J., Del., Md.
- Blossom-end rot, nonpar. General.
- Mosaic, undet. General.
- Oedema, nonpar. N. Y., Ind., Fla.
- Streak or winter blight, undet. Common in greenhouses, Mass., N. Y., N. J., Pa., Ohio, Ind., Mich.
- Yellow blight, undet. Western States.

LYONIA.*L. ligustrina*, *maleberry.

- Exobasidium vaccinii* Wor., hypertrophy. Mass., N. Y., N. J., Fla., Ala.
- Microsphaera alni vaccinii* Berk. and Curt., powdery mildew. Ala.
- Pucciniastrum myrtilli* (Schum.) Arth., rust. Del., Va., Ala.
- Rhytisma andromedae* Fr., tar-spot. N. J., Va.
- R. decolorans* Schw., tar-spot. Me., Mass., N. J., Fla., Ga., Ala.

MACLURA.*M. pomifera*, Osage-orange.

- Cerotelium fici* (Cast.) Arth., rust. S. C., Ga., La.
- Ovularia maculuræ* Ell. and A. B. Langlois, cottony leaf-spot. La., Tex.
- Phoradendron flavescens* (Pursh) Nutt., mistletoe. Tex., Ind.
- Phyllosticta maculuræ* Ell. and Ev., leaf-spot. N. J.
- Sporodesmium maculuræ* Thuem., leaf-spot. Mo.

MAGNOLIA.

Magnolia sp., magnolia.

Cladosporium fasciculatum Cda., Tex.

Colletotrichum sp., anthracnose. Mass., S. C., Fla.

Coniothyrium olivaceum grandiflorae Sacc., leaf-spot. Tex.

Phyllosticta cookei Sacc., leaf-spot. N. Y., S. C., Miss., Calif.

P. terminalis Ell. and Mart., leaf-spot. N. J.

Septoria magnoliae Cke., leaf-spot. Fla.

M. acuminata, cucumbertree.

Phyllactinia corylea (Pers.) Karst., powdery mildew. Ohio.

Phyllosticta cookei Sacc., leaf-spot. W. Va.

M. glauca, sweetbay.

Cercospora magnoliae Ell. and Hark., leaf-spot. N. J.

Cladosporium fasciculatum Cda. N. J.

Dimerosporium magnoliae Tr. and Earle, sooty mold. Miss.

Macrosporium martindalei Ell. and Mart., leaf-spot. N. J.

Phyllosticta cookei Sacc., leaf-spot. N. J., Fla.

P. glauca Cke., leaf-spot. S. C., Fla., Ala.

M. grandiflora, southern magnolia.

Coniothyrium olivaceum grandiflorae Sacc., leaf-spot. Tex.

Leptothyrella langloisii (Ell. and Ev.) Sacc. La.

Phyllosticta cookei Sacc., leaf-spot. Fla., Miss., La.

Septoria magnoliae Cke., leaf-spot. S. C., Ga., La., Tex.

S. niphostoma Berk. and Curt., leaf-spot. N. C., S. C., Ga.

MAHONIA.

Mahonia spp. (native western species), hollygrape, Oregon grape.

Puccinia graminis Pers., rust. Mich., Mont., Colo., N. Mex., Wash.

P. koeleriae Arth., rust. Mont., Colo., Idaho, Oreg.

P. oxalidis Diet. and Ell., rust. N. Mex.

Uropyxis sanguinea (Pk.) Arth., rust. Western States.

MALUS.

M. baccata, Siberian crab apple.

Bacillus amylovorus (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. Ohio, W. Va., Tex., Wash.

Gymnosporangium globosum Farl., rust. Kans.

G. juniperi-virginianae Schw., rust. Md.

Phyllosticta solitaria Ell. and Ev., blotch. Ohio.

Physalospora malorum (Pk.) Shear (*P. cydoniae* Am. Auct.), black-rot. Ohio.

Venturia inaequalis (Cke.) Aderh., scab. Ohio, Nebr.

M. coronaria, wild sweet crab apple.

Ascochyta coronaria Ell. and Davis, leaf-spot. Wis.

Bacillus amylovorus (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. W. Va.

Cercospora persicae Sacc., leaf-spot. Ill.

C. pyrina Ell. and Ev., leaf-spot. Wis.

Gymnosporangium juniperi-virginianae Schw., rust. General east of Great Plains.

Nummularia discreta (Schw.) Tul., blister canker. W. Va.

Phyllosticta solitaria Ell. and Ev., blotch. W. Va., Ind.

Physalospora malorum (Pk.) Shear (*P. cydoniae* Am. Auct.), black-rot. N. Y., Ill.

Podosphaera ozyacanthae (DC.) D By., powdery mildew. Wis.

M. ioensis, prairie crab apple.

Cercospora pyrina Ell. and Ev., leaf-spot. Wis.

Gymnosporangium juniperi-virginianae Schw., rust. East of Great Plains.

Marssonina coronariae Sacc. and Dearn., leaf-spot. Mo.

Phyllosticta zonata Ell. and Ev., leaf-spot. Iowa.

Venturia inaequalis (Cke.) Aderh., scab. Ill., Wis., Iowa, Nebr.

M. sylvestris, apple.

Alternaria sp., fruit-spot, fruit-rot. Reported from 11 scattered States.

A. mali J. W. Roberts, leaf-spot. Va., Md., Tenn., Ark., Mo.

MALUS—Continued.

M. sylvestris, apple—Continued.

- Armillaria mellea* (Vahl) Quel., root-rot. General, especially in Pacific Northwest.
- Bacillus amylovorus* (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. General.
- Bacterium tumefaciens* EFS. and Town. (*Phytoplasma tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall, hairy-root. General.
- Botryosphaeria ribis* Gross. and Dug., black-rot. Va., Ga., Ark.
- Botrytis cinerea* Auct., fruit-rot. Wash.
- Cephalothecium roseum* Cda., pink-rot. General, especially east of Rocky Mts.
- Cercospora mali* Ell. and Ev., leaf-spot. Ala., Miss., La., Tex.
- Clitocybe monadelphica* (Morgan) Sacc., root-rot. Va., W. Va., N. C., Ind., Mo., Ark., Okla., Tex., Colo., Ariz., Wash.
- Comandra pallida* DC., bastard toadflax. Wash.
- Corticium laetum* Karst., limb-blight. La.
- C. stevensii* Burt (*Hypochnus ochroleucus* F. Noack), leaf-blight. N. C., S. C., Ga., Ala., Miss., Tex., Ind.
- Cytospora* sp., canker. Oreg.
- Fomes applanatus* (Pers.) Wallr. (*Ganoderma applanatum* (Pers.) Pat.), white-streaked rot. Ind.
- F. fomentarius* (L.) Fr., white sapwood rot. Mont.
- F. ignarius* (L.) Fr., white heart-rot. Conn., N. Y.
- Fusicoccum pyrorum* C. Chupp and G. L. Clapp, canker. N. Y., Md.
- Gloeodes pomigena* (Schw.) A. S. Colby (*Phyllachora pomigena* (Schw.) Sacc.), sooty blotch. New England to N. C. and westward to Nebr.
- Gloeosporium perennans* Zeller and Childs, perennial canker. Wash., Oreg.
- Glomerella cingulata* (Ston.) Spauld. and Schrenk, bitter-rot. Mass. to Nebr. and southward; occasionally in Pacific Coast States.
- Gymnosporangium blasdaleanum* (Diet. and Holw.) Kern, rust. Oreg.
- G. germinale* (Schw.) Kern, rust. Mass., R. I., Va.
- G. globosum* Farl., rust. New England south to Ga.
- G. juniperi-virginianae* Schw., rust. General east of Great Plains.
- Leptosphaeria coniothyrium* (Fekl.) Sacc., orange pox canker. Middle Atlantic States; Ohio, Ind., Minn., Kans.
- Leptothyrium pomi* (Mont. and Fr.) Sacc., flyspeck. East of Great Plains; occasionally collected in Northwest.
- Myxosporium corticolum* Edg. (*Dermateia corticola* (Edg.) Arn.), surface bark canker. New England, N. Y., Pa., Md., Ohio, Mich., S. Dak., Oreg.
- Nectria galligena* Bres., European canker. Me. to Ill. and N. C., Minn., Pacific coast.
- Neofabraea malicorticis* (A. B. Cordley) Jack., anthracnose, black-spot canker. Pacific Northwest; Mont., Nev., Calif.
- Nummularia discreta* (Schw.) Tul., blister canker. East of Rocky Mts., most destructive in Miss. Valley States; N. Dak., S. Dak.
- Ozonium omnivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Ark., Tex., N. Mex., Ariz.
- Penicillium expansum* (Lk.) emend. Thom., blue-mold rot. General in stored fruit.
- Phoma mali* Schulz. and Sacc., twig-blight. Me., Mass., Md., W. Va., Okla., N. Mex.
- P. pomi* Pass., fruit-spot. Me. to N. C., and westward to Ark. and Iowa.
- Phomopsis mali* J. W. Roberts, rough-bark disease. Va., Ohio, Ill., Pacific Coast States.
- Phoradendron flavescens* (Pursh) Nutt., mistletoe. Tex.
- Phyllosticta clypeata* Ell. and Ev., leaf-spot. Oreg.
- P. limitata* Pk., leaf-spot. Mass., N. Y., W. Va., N. C., Ind.
- P. prunicola* Sacc., leaf-spot. Ohio.
- P. solitaria* Ell. and Ev., blotch. East of Great Plains except New England and Fla.
- Phyalospora malorum* (Pk.) Shear (*P. cydoniae* Am. Auct.), black-rot, frog-eye. General, especially east of Great Plains.
- Phytophthora cactorum* (Leb. and Cohn) Schroet., fruit-rot. Conn., N. Y., Pa., Va., Ind., Mich., Idaho, Wash.
- Plenodomus fuscomaculans* G. H. Coons, canker. Mich., Calif.

MALUS—Continued.*M. sylvestris*, apple—Continued.

- Podosphaera leucotricha* (Ell. and Ev.) Salm., powdery mildew. General, most destructive in Western States.
P. oxyacanthae (DC.) D By., powdery mildew. General.
Polyporus lacteus Fr., sap-rot and heart-rot. Va., Ind., Idaho, Wash., Oreg.
Polystictus versicolor (L.) Fr., yellowish wood-rot. N. Y., Mo., Kans., Pacific coast.
Pseudomonas papulans D. H. Rose, blister spot. Mo.
Schizophyllum commune Fr., heart-rot. North of Ohio and Mo. Rivers; Kans., Nebr., Pacific coast.
Sclerotinia cinerea (Bon.) Schroet. (*Monilia oregonensis* Barss and Posey), blossom blight. Wash., Oreg.
S. fructicola (Wint.) Rehm (*S. americana* (Wormald) Norton and Ezekiel) brown-rot. East of Great Plains; Pacific coast.
Septobasidium sp., canker. N. Y., Miss., Tex.
S. pseudopedicellatum Burt, canker. N. C.
S. retiforme (Berk. and Curt.) Pat., canker. Southern States.
Stereum purpureum Pers., associated with silver-leaf. Me., N. Y., Minn., Wash.
Valsa spp. (*Cytospora* spp.), die-back, canker. New England, N. Y., Pa., Ill., Minn., Western States.
Venturia inaequalis (Cke.) Aderh., scab. Reported from every State except Fla., La., Tex., Ariz., Nev., Utah, Wyo. Most serious in East.
Volutella fructi F. L. Stevens and J. G. Hall, spongy dry-rot. Mass., N. Y., Pa., W. Va., N. C., Ind.
Xylaria spp., black root-rot. N. Y., Pa., Va., W. Va., Tenn., N. C., S. C., Ga., Ark., Ind., Ill., Mich., Iowa, Kans.
Bitter-pit (stippen, Baldwin spot), undet. General.
Black-heart, nonpar. Md., Va., Nebr., Mont.
Chlorosis, nonpar. Me., N. Y., Pa., S. Dak., N. Mex., Wyo., Wash.
Jonathan spot, undet. Eastern apple sections to Nebr. and Kans.; Colo., Idaho, Pacific coast.
Measles, undet. Pa., Md., Va., W. Va., S. C., Ala., Ark., Tex., Ohio, Ill., Mich., Mo., Kans., Colo., N. Mex.
Mosaic, undet. N. Y.
Rosette, undet. Del., Md., Kans., Colo., Pacific Northwest.
Scald, nonpar. General on stored fruit.
Tan disease, undet. Md., Wash.
Water-core, nonpar. General.

MANGIFERA.*M. indica*, mango.

- Glomerella cingulata* (Ston.) Spauld. and Schrenk, anthracnose. Fla.
Phyllosticta mortonii Fairman, leaf-spot. Fla.

MANIHOT.*M. esculenta*, cassava.

- Cercospora cassavae* Ell. and Ev., leaf-spot. Fla.
Gloeosporium manihotis P. Henn., wither-tip. La.

MEDICAGO.*M. maculata*, southern bur clover.

- Cercospora medicaginis* Ell. and Ev., leaf-spot. Southern States.
Pseudoplea medicaginis Miles, leaf-spot. Ala.

M. sativa, alfalfa.

- Aplanobacter insidiosum* McC., bacterial root-rot. Ala., Ind., Ill., Wis., Colo.
Ascochyta imperfecta Pk., leaf-spot. N. Y., Ky., Ill., Wash.
Bacterium medicaginis (Sack.) EFS. (*Phytomonas medicaginis* (Sack.) Com. S. A. B.), bacterial blight. General, especially in Western States.
B. tumefaciens EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Md., Va., Ky., Ala.
Cercospora medicaginis Ell. and Ev., leaf-spot. East of Great Plains.
Colletotrichum trifolii S. M. Bain, anthracnose. General east of Great Plains.

MEDICAGO—Continued.*M. sativa*, alfalfa—Continued.

Cuscuta sp., dodder. Frequent where alfalfa is grown.

Fusarium sp., root-rot. S. C., Va., Ohio, N. Dak., Kans., N. Mex., Utah, Nev.

Gloeosporium medicaginis Ell. and Kell., anthracnose. Va., Kans., Okla.

Heterodera radicola (Greef) Muell., root-knot. N. Y., N. J., Tex.

Macrosporium sp., leaf-spot. Me., Pa., N. J., Md., Va., W. Va., La., Tex., Mich., Mo.

M. sarcinaeforme Cav., black-mold. Ill.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Ark., Okla., Tex., Southwest.

Peronospora trifoliorum D By., downy mildew. General.

Pleosphaerulina briosiana Poll., leaf-spot. Va., Ky., Ala., Ga., Ind., Ill., Wis., Minn., Iowa, Mo., S. Dak., Kans.

Pseudopeziza medicaginis (Lib.) Sacc., leaf-spot. General.

Pyrenopeziza medicaginis Fekl., yellow leaf-blotch. General.

Rhizoctonia crocorum (Pers.) DC., violet root-rot. Middle Atlantic States; Miss. Valley; Pacific Northwest.

Sclerotinia trifoliorum Eriks., stem-rot. Mostly east of Miss. River; Pacific coast.

Sclerotium rolfsii Sacc., stem-rot. Va., La.

Stagonospora sp., leaf-spot. N. Y., N. J., Del., Ill.

Thielavia basicola (Berk. and Br.) Zopf, black root-rot. Mass., N. J., Tex.

Tylenchus dipsaci (Kuehn) Bast., stem nematode. Wyo., Colo., N. Mex., Idaho, Oreg., Calif.

Uromyces medicaginis Pass., rust. General.

Urophlyctis alfalfae (Lagh.) Magn., crown-wart. West of Rocky Mts.

Girdle, undet. Ariz.

Yellowing, undet. General, especially in the eastern United States.

MELIA.*M. azedarach*, chinaberry.

Cercospora leucosticta Ell. and Ev., leaf-spot. Ala., Miss., La.

C. meliae Ell. and Ev., leaf-spot. La.

Nectria cinnabarina (Tode) Fr., canker. S. C.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex., Ariz.

Phoradendron flavescens (Pursh) Nutt., mistletoe. Tex.

Phyllosticta meliae Ell. and Ev., leaf-spot. La.

M. azedarach umbraculiformis, Texas umbrella-tree.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

MELILOTUS.*M. alba*, white sweetclover.

Ascochyta meliloti (Trel.) Davis, leaf-spot. Occurs in widely scattered States.

Cercospora davisii Ell. and Ev., leaf-spot. Ala., Md., Ind., Wis., Minn., Iowa, Kans.

Colletotrichum trifolii S. M. Bain, anthracnose. Ill.

Corticium vagum Berk. and Curt., stem-rot. Ala., Mich., Mont., Wash.

Fusarium sp., root-rot. Tenn., N. Dak., N. Mex.

Gloeosporium sp., anthracnose. Ill., Wis., Kans.

Heterodera radicola (Greef) Muell., root-knot. S. C., Ala.

Mycosphaerella lethalis R. E. Stone, stem-spot. N. Y.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Peronospora trifoliorum D By., downy mildew. Wis.

Sclerotinia trifoliorum Eriks., root-rot. Ky., Wash.

Stagonospora meliloti (Lasch) Petr., leaf-spot. Wis.

Mosaic, undet. Ark., N. Dak.

MENISPERMUM.

M. canadense, moonseed.

Cercospora menispermii Ell. and Holw., leaf-spot. W. Va., Iowa, Kans.

Entyloma menispermii Farl. and Trel., smut. North-Central States.

Phyllosticta abortiva Ell. and Kell., leaf-spot. Kans.

Ramularia contexta Ell. and Ev., leaf-spot. N. Dak.

MENTHA.

M. piperita, peppermint.

Puccinia menthae Pers., rust. Mass., Calif.

M. spicata, spearmint.

Phyllactinia corylea (Pers.) Karst., powdery mildew. Wash.

Phyllosticta decidua Ell. and Kell., leaf-spot. Ohio.

Puccinia menthae Pers., rust. Mass., N. Y., Ohio, Mich., Ind., Ky., Wis., Wash.

MIRABILIS.

M. jalapa, common four-o'clock.

Albugo platensis (Speg.) Sw., white-rust. Tex.

Cercospora mirabilis Tharp, leaf-spot. Tex.

MORUS.

Morus sp., mulberry.

Armillaria mellea (Vahl) Quel., root-rot. Del.

Bacterium mori (Boyer and Lambert) emend. EFS. (*Phytomonas mori* (Boyer and Lambert) Com. S. A. B.), mulberry blight. Widespread.

Cercospora missouriensis Wint., leaf-spot. Tex.

C. moricola Cke., leaf-spot. Widespread east of Great Plains.

C. pulvinulata Sacc. and Wint., leaf-spot. Nebr., Kans.

Cercospora mori Pk., leaf-spot. Nebr., Tex.

Exosporium sp., leaf-spot. Fla.

Gibberella moricola Ces. and DeNot., twig-blight. Pa., Wis.

Hymenochaete agglutinans Ell., heart-rot. Md.

Myxosporium diedickei Syd., twig-blight. Tex., Wash.

Mycosphaerella morifolia (Fekl.) Lindau, leaf-spot. Ind.

Nectria sp., bark disease, canker. N. J., Ky., Ala., Ark., Ind., Kans.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex., Ariz.

Phleospora maculans (Bereng.) Allesch., leaf-spot. Widespread east of Great Plains.

P. moricola (Pass.) Sacc., leaf-spot. Pa., Md., Kans.

Phyllactinia corylea (Pers.) Karst., powdery mildew. Ohio.

Sclerotinia carunculoides E. A. Siegler and A. Jenkins, pop-corn disease. S. C., Ga., Ala., Miss., La., Tex.

Uncinula geniculata W. R. Gerard, powdery mildew. Widespread east of Great Plains.

MUSA.

Musa sp., banana.

Fusarium sp., Panama wilt. Fla.

Gloeosporium musarum Cke. and Mass., stem-rot, fruit-rot. Fla.

Heterodera radiculicola (Greef) Muell., root-knot. Fla.

MYRICA.

Myrica spp. (*M. cerifera* and *M. carolinensis*), bayberry, waxmyrtle.

Cercospora diffusa Ell. and Ev., leaf-spot. N. J.

C. myricae Tr. and Earle, leaf-spot. Miss.

C. penicillus Ell. and Ev., leaf-spot. N. J., Miss.

Gymnosporangium myricatum (Schw.) F. D. Fromme, rust. Mass., N. J., Pa.

Phyllosticta myricae Cke., leaf-spot. N. J., S. C., Fla.

Ramularia moniloides Ell. and Mart., leaf-spot. Mass., N. Y.

Septoria myricae Ell. and Ev., leaf-spot. N. J.

NARCISSUS.

Narcissus sp., narcissus, jonquil.

Botrytis sp., blight. Wash.

Tylenchus dipsaci (Kuehn) Bast., bulb nematode. Ill., Oreg., Calif.

Mosaic (gray disease), undet. Wash.

NERIUM.

N. oleander, oleander.

Bacterium savastanoi EFS. (*Phytomonas savastanoi* (EFS.) Com. S. A. B.), gall. Ariz.

Cercospora neriella Sacc., leaf-spot. La., Tex.

Macrosporium nerii Cke., leaf-spot. Ga., Fla., Miss., Calif.

Phyllosticta nerii West., leaf-spot. Fla., La., Mich.

Septoria sp., specked leaf-spot. Fla.

S. oleandrina Sacc., leaf-spot. Fla., La.

NICOTIANA.

N. tabacum, tobacco.

Bacillus carotovorus L. R. Jones (*Erwinia carotovora* (Jones) Com. S. A. B.), hollow-stalk. Conn.

Bacterium angulatum F. D. Fromme and T. J. Murray, angular leaf-spot, black-fire. Mass., Conn., N. Y., Md., Va., N. C., S. C., Tenn., Ky., Ind., Wis.

B. melleum James Johnson, bacterial leaf-spot. Ky., Wis.

B. solanacearum EFS. (*Phytomonas solanaceara* (EFS.) Com. S. A. B.), bacterial wilt, Granville wilt. Southern States; Ohio.

B. tabacum Wolf and A. C. Foster (*Phytomonas tabaca* (Wolf and Foster) Com. S. A. B.), wild-fire. General.

Cercospora nicotianae Ell. and Ev., frog-eye. Atlantic Coast States; Wis., Tex.

Corticium vagum Berk. and Curt., sore-shin. General.

Fusarium affine C. M. Slagg, leaf-spot. Conn., Ky., Wis.

F. oxysporum nicotianae James Johnson, wilt. Md., Ohio, Ky., Tenn.

Heterodera radiculicola (Greef) Muell., root-knot. Southern States.

Macrosporium longipes Ell. and Ev., brown-spot. Pa., N. C., Mo.

M. tabacinum Ell. and Ev., white-speck. Mass., Conn., N. C., Ohio, Mo.

Olpidium sp., seedling blight. Wis.

Orobanche ramosa L., broom-rape. Ohio, Ky.

Peronospora hyoscyami D By., downy mildew. Ga., Fla.

Phyllosticta nicotiana Ell. and Ev., leaf-spot. Mass., Va., N. C.

Phytophthora nicotianae (Speg.) Van Breda de Haan, black-shank. Ga., Fla.

Pythium debaryanum Hesse, damping-off. General.

Sclerotinia sclerotiorum (Lib.) Mass., stem-rot. Conn.

Sclerotium rolfsii Sacc., stem-rot. N. C.

Thielavia basicola (Berk. and Br.) Zopf, black root-rot. General.

Brown root-rot, undet. Mass., Conn.

Frenching, undet. Va., N. C., Ky., Ohio, Wis.

Leaf-spots including "rust," firing, ring-spot, etc., nonpar. Widespread.

Mosaic, undet. General.

Sand-drown, nonpar. N. C.

NYSSA.

N. aquatica, big tupelo.

Aplospora nyssae (Ell.) E. B. Mains, rust. Miss.

Phyllosticta nyssae Cke., leaf-spot. Ga., Ala.

N. sylvatica, *sour gum, tupelo, *pepperidge.

Leptothyrium dryinum Sacc., leaf-spot. Ala.

Phoradendron flavescens (Pursh) Nutt., mistletoe. Tex.

Phyllosticta nyssae Cke., leaf-spot. N. Y.

Pirostoma nyssae L. R. Tehon, leaf-spot. Ill.

OLEA.

O. europaea, olive.

Armillaria mellea (Vahl) Quel., root-rot. Calif.

Bacterium savastanoi EFS. (*Phytomonas savastanoi* (EFS.) Com. S. A. B.), knot. Calif.

OLEA—Continued*O. europaea*, olive—Continued.*Cycoconium oleaginum* Cast., leaf-spot. Calif.*Ozonium omnivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.**ORYZA.***O. sativa*, rice.*Cercospora oryzae* Miyake, leaf-spot. La.*Corticium vagum* Berk. and Curt., foot-rot. Ark.*Fusarium minimum* Fekl., wilt. La.*Helminthosporium* sp., leaf-spot. S. C., Ark.*H. oryzae* Van Breda de Haan, blight. Fla., La.*Ophiobolus* sp., foot-rot. Ark.*Phoma glumarum* Ell. and Tr., glume-blight. Miss.*Piricularia grisea* (Cke.) Sacc., blast. Occurs in all States where rice is grown.*Sclerotium oryzae* Catt., stem-rot. La., Ark.*S. rolfsii* Sacc., seedling blight. La.*Sphaerotheca* sp., powdery mildew. N. C.*Tilletia horrida* Tak., black-smut. S. C., Ga., La., Ark.*Ustilaginoides virens* (Cke.) Tak., green-smut. Ga., La.

Straighthead, nonpar. General in States in which rice is grown.

OSMANTHUS.*Osmanthus* sp., osmanthus.*Phyllosticta oleae* Ell. and Mart., leaf-spot. Fla.*P. sinuosa* Ell. and Mart., leaf-spot. Fla., Miss.*P. terminalis* Ell. and Mart., leaf-spot. Fla.**OSTRYA.***O. virginiana*, hophornbeam.*Fomes igniarius* (L.) Fr., white heart-rot. N. Dak., S. Dak.*Gloeosporium robergiae* Desm., leaf-spot. Wis.*G. robergiae dendriticum* Davis, leaf-spot. Pa.*Microsphaera alni* (Wallr.) Wint., powdery mildew. N. Y., Ill., Iowa.*Phyllactinia corylea* (Pers.) Karst., powdery mildew. General over range of host.*Septoria ostryae* Pk., leaf-spot. N. Y.*Taphrina virginica* Seym. and Sadeb., leaf-spot. N. H., Mass., Ala., Wis. Iowa.**OXYDENDRUM.***O. arboreum*, sourwood.*Cercospora oxydendri* Tr. and Earle, leaf-spot. Ala.*Phyllosticta oxydendri* Ell. and Ev., leaf-spot. W. Va.**PAEONIA.***Paeonia* sp., peony.*Armillaria mellea* (Vahl) Quel., root-rot. Mich., Oreg.*Bacterium tumefaciens* EFS. and Town. (*Phytoplasma tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Mich.*Botrytis* sp., seed-rot. Pa.*B. paeoniae* Oud., blight. Widespread.*Cerospora varicolor* Wint., leaf-spot. Ind., Ky., Mo., Nebr.*Cladosporium paeoniae* Pass., leaf-mold. General.*Corticium vagum* Berk. and Curt., stem-rot. Minn.*Fusarium* sp., root-rot, stem-rot. Ind., Mo., Nebr.*Heterodera radicola* (Greef) Muell., root-knot. N. Y., Pa., S. C., Miss., Ohio, Ind., N. Dak., Nebr., Wash.*Phyllosticta commonsii* Ell. and Ev., leaf-spot. Del., Ind.*Phytophthora* sp., blight. Conn., Pa., Ind., Kans.*Septoria paeoniae berolinensis* Allesch., leaf-spot. N. Y.*Verticillium alboatrum* Reinke and Berth., wilt. Kans.

Mosaic, undet. Mass., Conn., Va., Ohio, Kans., Wash.

PANAX.

P. quinquefolium, American ginseng.

Alternaria panax Whet., blight. General; N. Y. to Minn. southward to Kans. and N. C.

Botrytis sp., gray-mold. Conn., N. Y., Mich., Wash.

Corticium vagum Berk. and Curt., stem-rot. N. Y., Wash.

Fusarium sp., wilt. N. Y., Ala., Ohio, Mo.

Heterodera radicolica (Greef) Muell., root-knot. N. Y., Pa., Ohio, Mich., Wis.

Pestalozzia funerea Desm., leaf anthracnose. Mo.

Phytophthora cactorum (Leb. and Cohn) Schroet., blight, downy mildew. Conn., N. Y., Pa., Ohio, Ind., Mich., Iowa, Wash.

Ramularia destructans C. L. Zinssmeister, root-rot. Wis.

R. panacicola C. L. Zinssmeister, root-rot. N. Y., Wis.

Sclerotinia panacis W. H. Rankin, black-rot. N. Y., Mich., Wis.

S. sclerotiorum (Lib.) Mass., white-rot. N. Y., Pa., Ohio, Mich., Wis.

S. smilacina E. J. Durand, black-rot. N. Y., Mich., Wis.

Thielavia basicola (Berk. and Br.) Zopf, black root-rot. General where ginseng is grown.

Vermicularia dematium (Pers.) Fr., stem anthracnose. N. Y., Va., N. C., Tenn., Ohio, Ill., Minn., Mo.

Verticillium alboatrum Reinke and Berth., wilt. N. Y., Pa., N. J., Tenn., Ky., Ohio, Mich., Ind.

Damping-off, associated with various fungi. General.

Papery leaf-spot, nonpar. N. Y., Mich.

Soft-rot, undet. N. Y., Pa., Ohio, Ind.

PANICUM.

P. miliaceum, proso millet, broomcorn millet.

Bacterium panici C. Elliott, bacterial stripe. Wis., S. Dak.

Fusarium rhizochromatistes C. P. Sideris, root-rot. Calif.

Phyllachora graminis (Pers.) Fekl., black-blotch. Tex.

Piricularia grisea (Cke.) Sacc., leaf-spot. Minn., Ohio.

Puccinia emaculata Schw., rust. Ind.

Sclerospora graminicola (Sacc.) Schroet., downy mildew. Wis., Minn., Iowa, N. Dak., S. Dak.

Sorosporium syntherismae (Pk.) Farl., smut. N. J., Wis., Minn., Mo., N. Dak., Nebr.

Ustilago panici-miliacei (Pers.) Wint., smut. N. Dak., Colo., Wash.

PARKINSONIA.

P. aculeata, parkinsonia.

Cylindrosporium parkinsoniae Heald, leaf-spot. Tex.

Dimerosporium parkinsoniae Heald and Wolf, sooty mold. Tex.

Phyllosticta parkinsoniae Ell. and Ev., leaf-spot. Tex.

PASTINACA.

P. sativa, parsnip.

Bacillus carotovorus L. R. Jones (*Erwinia carotovora* (Jones) Com. S. A. B.), soft-rot. Conn., Mass., Minn.

Bacterium tumefaciens EFS. and Town. (*Phytoplasma tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Va.

Cercospora aptii pastinacae Farl., leaf-spot. North Atlantic and North-Central States.

Corticium vagum Berk. and Curt., stem-rot. Tex., Wash.

Heterodera radicolica (Greef) Muell., root-knot. Va., Ala., La., Ohio, Ind., Kans.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Ramularia pastinacae (Karst.) Lindr. and Vester., leaf-spot. N. Y., Ind., N. Dak.

Sclerotinia sclerotiorum (Lib.) Mass., watery soft-rot. Ind.

Septoria pastinacina Sacc., leaf-spot, blight. N. Y., Utah.

PAULOWNIA.

P. tomentosa, *princess tree.

Ascochyta paulowniae Sacc. and Brun., leaf-spot. D. C.

Phyllosticta paulowniae Sacc., leaf-spot. N. Y., D. C.

PELARGONIUM.

Pelargonium sp., geranium.

Alternaria sp., leaf-spot. Mass., La., Mich., Calif.

Bacterium erodii I. M. Lewis, bacterial leaf-spot. Conn., Pa., N. Y., Miss., Ohio, Ind., Ill., Wash.

B. pelargoni N. A. Brown, bacterial leaf-spot. Mass., N. J., Md., Va., D. C.

Botrytis cinerea Auct., gray-mold. Conn., Md., La., Ohio, Mich., Minn., Kans.

Cercospora brunckii Ell. and Gall., leaf-spot. Ala., Miss., Tex.

Pythium complectens H. Braun, stem-rot. D. C., Okla.

PERSEA.

P. americana, avocado.

Cercospora sp., blotch. Fla.

Glomerella cingulata (Ston.) Spauld. and Schrenk, anthracnose, leaf-spot, fruit-spot. Fla., Ga., Tex.

Phyllosticta sp., leaf-spot. Fla.

Pythiacystis citrophthora E. H. and R. E. Smith, bark disease. Calif.

Sporotrichum citri Butl. (*Sphaceloma fawcettii* A. Jenkins), scab. Fla.

P. borbonia, redbay.

Helminthosporium fumosum Ell. and Mart., leaf-blotch. Fla.

Phyllosticta micropuncta Cke., leaf-spot. D. C., Fla., Miss., La., Tex.

PETROSELINUM.

P. hortense, parsley.

Cercospora apii Fresen., leaf-spot. N. J.

Corticium vagum Berk. and Curt., root canker. N. Y.

Heterodera radiculicola (Greef) Muell., root-knot. Tex.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Sclerotinia sclerotiorum (Lib.) Mass., stem-rot. Conn., N. Y., Pa., Va., La., Tex.

Septoria petroselini Desm., leaf-spot. Conn., N. J.

PETUNIA.

P. hybrida, petunia.

Ascochyta petuniae Speg., leaf-spot.

Cercospora sp., leaf-blotch. Fla.

Corticium vagum Berk. and Curt., stem-rot. Pa.

Heterodera radiculicola (Greef) Muell., root-knot.

Macrosporium sarcinula parasiticum Thuem., black stalk-rot. Tex.

Mosaic, undet. Conn., Pa., S. C., La., Iowa.

PHASEOLUS.

P. acutifolius latifolius, tepary bean.

Fusarium sp., wilt. Calif.

F. martii phaseoli Burk., dry root-rot. N. Y.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Uromyces appendiculatus (Pers.) Lk., rust. Tex., Calif.

P. angularis, adzuki bean.

Amerosporium oeconomicum Ell. and Tr., leaf-spot. Va.

Bacterium phaseoli EFS. (*Phytomonas phaseoli* (EFS.) Com. S. A. B.), bacterial blight. Ind.

Fusarium martii phaseoli Burk., dry root-rot. N. Y.

F. vasinfectum Atk., wilt. Ala.

Mosaic, undet. Ind.

P. aureus, mung bean.

Bacterium phaseoli EFS. (*Phytomonas phaseoli* (EFS.) Com. S. A. B.), bacterial blight. Ind.

Sclerotinia sclerotiorum (Lib.) Mass., stem-rot. Va.

Mosaic, undet. Ind.

PHASEOLUS—Continued.

P. coccineus, Scarlet Runner bean.

Colletotrichum lindemuthianum (Sacc. and Magn.) Briosi and Cav. (*Glomerella lindemuthiana* Shear), anthracnose.

Fusarium martii phaseoli Burk., root-rot. N. Y.

P. lunatus, civet bean, and *P. lunatus macrocarpus*, Lima bean.

Bacterium phaseoli EFS. (*Phytomonas phaseoli* (EFS.) Com. S. A. B.), leaf-spot. Occurs scatteringly throughout Eastern States.

B. viridifaciens W. B. Tisdale and M. M. Williamson, bacterial leaf-spot. N. Y., Ind., Wis.

Cercospora cruenta Sacc., leaf-spot. N. Y., Miss., Mo., Tex.

C. phaseolorum Cke., leaf-spot. La.

Colletotrichum lindemuthianum (Sacc. and Magn.) Briosi and Cav. (*Glomerella lindemuthiana* Shear), anthracnose. Conn., N. Y., Pa., W. Va., La., Kans.

Corticium vagum Berk. and Curt., stem-rot. Pa., Calif.

Diaporthe phaseolorum (Cke. and Ell.) Sacc., pod-blight. Widespread in eastern United States.

Erysiphe polygoni DC., powdery mildew. Conn., Calif.

Fusarium sp., root-rot. Ohio, Calif.

F. martii phaseoli Burk., dry root-rot. N. Y.

Heterodera radiculicola (Greef) Muell., root-knot. Calif.

Nematospora phaseoli S. A. Wingard, yeast-spot. Va., Tenn., Ala., Ill.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Phyllosticta phaseolina Sacc., leaf-spot. N. J., W. Va., Fla., Ind., Mich.

Phytophthora phaseoli Thax., downy mildew. Atlantic Coast States; W. Va., Ky., Ohio, Calif.

Sclerotinia sp., stem-rot. Ind.

Uromyces appendiculatus (Pers.) Lk., rust. Pa., Md., Va., Ohio, S. C., La., Calif.

Mosaic, undet. W. Va., Ga., Ind., Mo.

P. vulgaris, bean.

Alternaria fasciculata (Cke. and Ell.) L. R. Jones and A. J. Grout, leaf-spot. La., Mich., Ill., N. Dak., Utah.

Bacillus lathyri Manns and Taub., streak. Del.

Bacterium flaccumfaciens Hedges (*Phytomonas flaccumfaciens* (Hedges) Com. S. A. B.), bacterial wilt. Md., Va., Mich., S. Dak., Mont., Oreg.

B. phaseoli EFS. (*Phytomonas phaseoli* (EFS.) Com. S. A. B.), blight. General.

B. solanacearum EFS. (*Phytomonas solanaceara* (EFS.) Com. S. A. B.), bacterial wilt. Fla.

Botrytis cinerea Auct., gray-mold rot. N. Y., Miss., Tex.

Cercospora cruenta Sacc., leaf-blotch. Fla., Ga.

Colletotrichum caulicolum Heald and Wolf, stem canker. Tex.

C. lindemuthianum (Sacc. and Magn.) Briosi and Cav. (*Glomerella lindemuthiana* Shear), anthracnose. General.

Corticium vagum Berk. and Curt., stem-rot. General.

Erysiphe polygoni DC., powdery mildew. Widespread.

Fusarium martii phaseoli Burk., dry root-rot. General.

Heterodera radiculicola (Greef) Muell., root-knot. Southern States; Calif.

Isariopsis griseola Sacc., angular leaf-spot. Atlantic Coast States; Fla., Miss.

Microsphaera diffusa Cke. and Pk., powdery mildew. Ga.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Okla., Tex.

Phoma sp., stem-rot. Ohio, Ark.

Phyllosticta phaseolina Sacc., leaf-spot. Conn., N. Y., N. J., Va., W. Va., Ind., Miss., La., Tex.

Pythium debaryanum Hesse, damping-off. Del., N. J., Ohio.

Sclerotinia sclerotiorum (Lib.) Mass., stem-rot. Middle Atlantic and Gulf States; W. Va., Mich., Ind.

Sclerotium rolfsii Sacc., stem-rot. Southern States.

Thielavia basicola (Berk. and Br.) Zopf, black root-rot. N. H., N. Y.

Uromyces appendiculatus (Pers.) Lk., rust. General.

Mosaic, undet. General.

PHILADELPHUS.

Philadelphus spp., *syringa, mockorange.

Cercospora angulata Wint., leaf-spot. Mo.

Gymnosporangium gracilens (Pk.) Kern and Bethel, rust. Colo., Utah, N. Mex.

Phyllactinia corylea (Pers.) Karst., powdery mildew. Mont.

Ramularia philadelphi Sacc., leaf-spot. Wash.

Septoria philadelphi Ell. and Ev., leaf-spot. Wash.

PHLEUM.

P. pratense, timothy.

Cercospora graminicola Tr. and Earle, leaf-spot. Miss.

Claviceps microcephala (Wallr.) Tul., ergot. Ind., Ky., Alaska.

C. purpurea (Fr.) Tul., ergot. Vt., Pa., Ind., Minn.

Colletotrichum graminicolum (Ces.) Wils., anthracnose. Pa., Ohio, Ind., Minn.

Entyloma crastophilum Sacc., smut. Ill., Wis., Iowa.

Erysiphe graminis DC., powdery mildew. Minn.

Heterosporium phlei C. T. Gregory, leaf-spot. N. Y., Pa.

Ophiobolus graminis Sacc., take-all. N. Y.

Puccinia coronata Cda., crown-rust. Iowa.

P. graminis Pers., stem-rust. General.

Scolecotrichum graminis Fckl., leaf-spot. Widespread in Northern States.

Septoria sp., leaf-spot. Pa., Ill.

Sporotrichum poae Pk., silver top. N. Y., Ind., Iowa.

Ustilago striaeformis (West.) Niessl, stripe-smut. General.

PHLOX.

Phlox spp., phlox.

Ascochyta phlogis phlogina Fairman, leaf-spot. Mass.

Bacterium tumefaciens EFS. and Town. (*Phytoplasma tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. N. J.

Cercospora amphakodes Ell. and Hals., leaf-spot. Ind., Wis., Iowa.

C. phlogina Pk., leaf-spot. Va., Minn.

Erysiphe cichoracearum DC., powdery mildew. General.

Heterodera radiculicola (Greef) Muell., root-knot. Mass.

Leptosphaeria phlogis Oud., leaf-spot. Md., Ind., Minn., Iowa, Nebr.

Phyllactinia corylea (Pers.) Karst., powdery mildew. Ohio.

Puccinia douglasii Ell. and Ev., rust. N. J., Pa., Nebr., Western States.

P. plumbaria Pk., rust. Ill., Iowa, Nebr., Western States.

Septoria divaricata Ell. and Ev., leaf-spot. N. Y., Va., Ill.

Sclerotium rolfsii Sacc., stem-rot. Ill.

Sphaerotheca humuli (DC.) Burr., powdery mildew. Ohio, Kans.

Tylenchus dipsaci (Kuehn) Bast., stem nematode. N. J.

Uromyces polemonii (Pk.) Arth., rust. S. C., Miss., Ind., Ill., Wis., Iowa.

PHOENIX.

P. canariensis, Canary Island date palm.

Exosporium palmivorum Sacc., leaf-spot. La., Tex.

Graphiola phoenicis (Moug.) Poit., false smut. Fla., La., Miss., Tex., Calif.; in hothouses in North.

P. dactylifera, date palm.

Exosporium palmivorum Sacc., leaf-spot. La.

Graphiola phoenicis (Moug.) Poit., false smut. Fla., Tex., Ariz., Calif.

P. reclinata, Senegal date palm.

Graphiola phoenicis (Moug.) Poit., false smut. S. C.

PHOTINIA.

P. arbutifolia, Christmasberry.

Cercospora heteromeles Hark., leaf-spot. Calif.

Fabraea maculata (Lév.) Atk., leaf-spot. Calif.

Fusicladium photinicola R. L. McClain, scab. Calif.

Phyllosticta heteromeles Cke. and Hark., leaf-spot. Calif.

Septoria photiniae Berk. and Curt., leaf-spot. Calif.

PHYLLOSTACHYS.

Phyllostachys sp., bamboo.

Cylindrosporium bambusae Miyake and Hara. Ga.

Melanconium bambusae Turconi. Ga.

Schirrhia bambusae Turconi. Ga.

P. bambusoides, Japanese timber bamboo.

Coniosporium shiraianum (Syd.) Bub. Fla.

Puccinia arundinariae Schw., rust. Fla.

P. melanocephala Syd., rust. Ga., Fla.

Ustilago shiraiana P. Henn., smut. Fla., La., Calif.

P. nigra, blackjoint bamboo.

Ustilago shiraiana P. Henn., smut. Fla., La., Calif.

PHYSOCARPUS.

P. opulifolius, ninebark.

Gloeosporium neilliae Hark., anthracnose. Calif.

Marssonia lonicerae Hark., leaf-spot. Wis.

M. neilliae Hark., leaf-spot. Calif.

Ramularia ulmariae Cke., leaf-spot. N. Y.

Sphaerotheca humuli (DC.) Burr., powdery mildew. Wis.

PICEA.

P. canadensis, white spruce.

Fomes pinicola Fr., brown crumbly rot. N. Y.

F. roseus (Alb. and Schw.) Fr., brown pocket rot. General.

Melampsoropsis pyrolae (DC.) Arth., rust. Me.

Pythium debaryanum Hesse, damping-off.

Razoumofskyia pusilla (Pk.) O. Kuntze, mistletoe. Northeastern States.

Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Vt., N. Y.

P. engelmanni, Engelmann spruce.

Chrysomyxa weirii Jack., rust. Mont., Idaho, Oreg.

Corticium vagum Berk. and Curt., damping-off. D. C., Calif.

Echinodontium tinctorium Ell. and Ev., brown stringy rot. Mont., Idaho, N. Mex.

Fomes officinalis (L.) Fr., reddish brown rot.

F. pinicola Fr., brown crumbly rot. Mont.

F. roseus (Alb. and Schw.) Fr., brown pocket rot.

Herpotrichia nigra Hartig, matting. Colo., Wyo., Idaho, Oreg.

H. quinqueseptata Weir, brown felt-blight. Idaho.

Hydnum sp., honeycomb rot.

Melampsorella elatina (Alb. and Schw.) Arth., rust. Idaho.

Melampsoropsis ledicola (Pk.) Arth., rust. Colo.

M. pyrolae (DC.) Arth., rust. Mont.

Peridermium coloradense (Diet.) Arth. and Kern, rust witches'-broom. Northern Rocky Mts.; Pacific Northwest.

Polyporus dryadeus (Pers.) Fr., heart-rot. Pacific Northwest.

P. schweinitzii Fr., red-brown butt-rot.

P. sulphureus (Bull.) Fr., brown-checked wood-rot. Me.

Pythium debaryanum Hesse, damping-off.

Razoumofskyia douglasii microcarpa Engelm., mistletoe. Idaho, Colo., Mont., Ariz., N. Mex.

R. laricis Piper, mistletoe. Mont.

Sparassia radicata Weir, yellow root-rot. Mont., Pacific Northwest.

Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Mont., Idaho.

P. excelsa, Norway spruce.

Melampsorella elatina (Alb. and Schw.) Arth., rust. Mont., Wyo.

P. mariana, black spruce, and *P. rubra*, red spruce.

Fomes annosus Fr., white spongy rot. Me. -

F. pinicola Fr., brown crumbly rot. Me.

Melampsoropsis abietina (Alb. and Schw.) Arth., leaf blister rust. N. H., Mich.

M. cassandrae (Pk. and Clint.) Arth., leaf blister rust. Mich., Minn.

M. ledicola (Pk.) Arth., leaf blister rust. Me., Minn.

Polyporus borealis Fr., cuboidal wood-rot. Minn.

PICEA—Continued.

P. mariana, black spruce, and *P. rubra*, red spruce—Continued.

P. schweinitzii Fr., red-brown butt-rot. Minn.

Razoumofskyia pusilla (Pk.) O. Kuntze, witches'-broom. New England, N. Y., Mich.

Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Me., Minn.

P. sitchensis, Sitka spruce.

Fomes officinalis Fr., reddish brown-rot. Wash.

F. pinicola Fr., brown crumbly rot. General within range of host.

F. roseus (Alb. and Schw.) Fr., brown pocket rot. General within range of host.

Lophodermium pinastri (Schrad.) Chev., needle cast. Alaska.

Melampsoropsis ledicola (Pk.) Arth., leaf blister rust. Wash., Alaska.

Polyporus berkeleyi Fr., string and ray rot. Calif.

P. schweinitzii Fr., red-brown butt-rot. Oreg.

Pythium debaryanum Hesse, damping-off.

Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Oreg.

PINUS.

P. albicaulis, whitebark pine.

Fomes pinicola Fr., brown crumbly rot. General within range of host.

Neopeckia coulteri (Pk.) Sacc., brown felt-blight. Idaho, Calif.

Polyporus schweinitzii Fr., red-brown butt-rot.

Razoumofskyia cyanocarpa (A. Nelson) P. A. Rydberg, mistletoe. Idaho, Mont.

R. laricis Piper, mistletoe. Mont.

P. attenuata, knobcone pine.

Cronartium cerebrum (Pk.) Hedge. and Long, oak-rust. Calif.

C. coleosporioides (Diet. and Holw.) Arth., rust. Calif.

Fomes pinicola Fr., brown crumbly rot. General within range of host.

Razoumofskyia americana (Nutt.) O. Kuntze, mistletoe. Oreg.

R. campylopoda (Engelm.) Piper, mistletoe. Pacific coast.

P. banksiana, jack pine.

Coleosporium helianthi (Schw.) Arth., rust. Mich.

Corticium vagum Berk. and Curt., damping-off. Nebr.

Cronartium cerebrum (Pk.) Hedge. and Long, oak-rust. Conn., Mich., Minn.

C. comptoniae Arth., rust. Conn., Wis., Minn.

C. pyriforme (Pk.) Hedge. and Long, rust. Minn.

Lophodermium amplum Davis, needle cast. Wis.

L. pinastri (Schrad.) Chev., needle cast. Wis., Minn.

Polyporus schweinitzii Fr., red-brown butt-rot. Minn.

Pythium debaryanum Hesse, damping-off.

Razoumofskyia americana Nutt., mistletoe.

P. caribaea, slash pine.

Coleosporium apocynaceum Cke., rust. Fla.

C. carneum (Bosc) Jack., rust. Ga., Fla.

C. delicatulum (Arth. and Kern) Hedge. and Long, rust. Fla., La.

C. elephanthopodis (Schw.) Thuem., rust. Fla.

C. ipomoeae (Schw.) Burr., rust. Fla.

Cronartium strobilinum (Arth.) Hedge. and G. Hahn, rust. Fla.

P. contorta, shore pine, and *P. murrayana*, lodgepole pine.

Cenangium piniphilum Weir, canker. Idaho.

Coleosporium solidaginis (Schw.) Thuem., rust. Colo., Mont., Idaho, Oreg., Wash.

Cronartium coleosporioides (Diet. and Holw.) Arth., rust. Colo., S. Dak., Mont., Idaho, Wyo., Wash., Calif.

C. cerebrum (Pk.) Hedge. and Long, oak-rust. Wyo., Utah, Idaho.

C. comptoniae Arth., rust. Mich.

C. filamentosum (Pk.) Hedge., pine-gall rust. Pacific Northwest.

C. harknessii (Moore) E. P. Meinecke, western pine-gall rust. Pacific coast; Idaho, Mont.

C. pyriforme (Pk.) Hedge. and Long, rust. Mich., Colo., Mont., Wyo., Idaho.

PINUS—Continued.

P. contorta, shore pine, and *P. murrayana*, lodgepole pine—Continued.

Fomes pinicola Fr., brown crumbly rot. Mont., Idaho.

F. roseus (Alb. and Schw.) Fr., brown pocket rot. General in range of host.

Hypoderma deformans Weir, needle cast, witches'-broom. Oreg., Calif.

H. lineare (Pk.) Thuem., needle cast. Colo.

Hypodermella montivaga (Petr.) Dearn., needle cast. Idaho, Oreg.

Lophodermium pinastri (Schr.) Chev., needle cast. Colo., Alaska.

Neopeckia coulteri (Pk.) Sacc., brown felt-blight. Colo., Utah, Idaho, Wash., Oreg.

Peridermium betheli Hedge. and Long, rust. Colo.

Polyporus schweinitzii Fr., red-brown butt-rot. Idaho.

P. sulphureus (Bull.) Fr., brown-checked wood-rot.

Razoumofskyia americana Nutt., mistletoe. Rocky Mt. region; Pacific Northwest; eastern Calif.

Rhizina inflata (Schaeff.) Sacc., seedling blight. Mont.

Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Mont., Idaho.

P. echinata, shortleaf pine.

Coleosporium carneum (Bosc) Jack., rust. Southeastern States.

C. delicatulum (Arth. and Kern) Hedge. and Long, rust. Pa., Md., S. C., Ga.

C. elephantopodis (Schw.) Thuem., rust. Fla.

C. helianthi (Schw.) Arth., rust. Ga.

C. inconspicuum (Long) Hedge. and Long, rust. Md., N. C., Ga.

C. ipomoeae (Schw.) Burr., rust. Pa., Va., N. C., S. C., Tenn., Ga., Ala., Ark., Tex.

C. solidaginis (Schw.) Thuem., rust. Del., Miss.

C. terebinthinaceae (Schw.) Arth., rust. Southeastern States.

Cronartium cerebrum (Pk.) Hedge. and Long, oak-rust. Ark.

C. comptoniae Arth., sweet-fern rust. N. C.

C. strobilinum Arth., rust. Miss.

Fomes pinicola Fr., brown crumbly rot. General in host range.

Lophodermium pinastri (Schr.) Chev., needle cast. Ala.

Polyporus schweinitzii Fr., red-brown butt-rot.

Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Mo.

P. edulis, nut pine.

Coleosporium ribicola (Cke. and Ell.) Arth., rust. Colo., N. Mex.

Cronartium occidentale Hedge., Bethel, and Hunt; piñon blister rust. Colo., N. Mex., Utah.

Fomes pinicola Fr., brown crumbly rot. General within host range.

Polyporus schweinitzii Fr., red-brown butt-rot.

Razoumofskyia divaricata (Engelm.) Coville, mistletoe. Colo., Utah, Calif.

Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot.

P. jeffreyi, Jeffrey pine.

Cronartium comptoniae Arth., sweet-fern rust.

C. filamentosum (Pk.) Hedge. and Long, rust. Oreg.

C. harknessii (Moore) E. P. Meinecke, rust. Nev., Calif.

Fomes pinicola Fr., brown crumbly rot. General with host.

Polyporus schweinitzii Fr., red-brown butt-rot.

P. sulphureus (Bull.) Fr., brown-checked wood-rot.

Razoumofskyia americana Nutt., mistletoe. Calif.

R. campylopoda (Engelm.) Piper, mistletoe. Calif.

P. lambertiana, sugar pine.

Armillaria mellea (Vahl) Quel., root-rot. Calif.

Cronartium ribicola Fisch., blister rust. N. Y.

Fomes officinalis Fr., reddish brown-rot. Calif.

F. pinicola Fr., brown crumbly rot. Calif.

Lophodermium pinastri (Schr.) Chev., needle cast. Oreg.

Polyporus schweinitzii Fr., red-brown butt-rot.

Razoumofskyia cyanocarpa (A. Nelson) P. A. Rydberg, mistletoe. Oreg., Calif.

Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Calif.

PINUS—Continued.

P. monophylla, singleleaf pine.

- Cronartium occidentale* Hedge., Bethel, and Hunt, blister rust. Ariz., Calif.
Fomes pinicola Fr., brown crumbly rot. General within host range.
Polyporus schweinitzii Fr., red-brown butt-rot.
Razoumofskyia divaricata (Engelm.) Coville, mistletoe. Ariz., Calif.

P. montana, Swiss mountain pine.

- Cronartium comptoniae* Arth., sweet-fern rust. N. H., Conn.
Herpotrichia nigra Hartig, matting.

P. monticola, western white pine.

- Cronartium ribicola* Fisch., blister rust. Wash.
Dasyscypha fuscousanguinea Rehm, canker. Mont., Idaho, Wash.
Fomes officinalis Fr., reddish brown-rot. Idaho.
F. pinicola Fr., brown crumbly rot.
Hypoderma lineare (Pk.) Thuem., needle cast. Pacific Northwest.
Lophodermium pinastri (Schr.) Chev., needle cast. Wash., Idaho, Mont.
Neopeckia coulteri (Pk.) Sacc., brown felt-blight. Calif.
Polyporus schweinitzii Fr., red-brown butt-rot.
P. sulphureus (Bull.) Fr., brown-checked wood-rot.
Razoumofskyia cyanocarpa (A. Nelson) P. A. Rydberg, mistletoe. Calif.
Rhizina inflata (Schaeff.) Sacc., seedling blight. Mont., Idaho.
Scleroderma bacillifera (Karst.) Sacc., branch canker. Wash., Oreg.
Sparassis radicata Weir, yellow root-rot. Mont., Pacific Northwest.
Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Idaho.

P. nigra, Austrian pine and Corsican pine.

- Coleosporium carneum* (Bosc) Jack., rust. Ohio.
C. delicatulum (Arth. and Kern) Hedge. and Long, rust. Pa.
C. solidaginis (Schw.) Thuem., rust. Pa.
Corticium vagum Berk. and Curt., damping-off.
Cronartium comptoniae Arth., sweet-fern rust. Conn., Ohio.
C. pyriforme (Pk.) Hedge. and Long, rust. Conn.
Peridermium sp., leaf-rust. Ohio, Nebr.
Pythium debaryanum Hesse, damping-off.

P. palustris, longleaf pine.

- Coleosporium apocynaceum* Cke., rust. Fla.
C. carneum (Bosc) Jack., rust. S. C. to Miss.
C. delicatulum (Arth. and Kern) Hedge. and Long, rust. S. C., Ga., Fla., Miss.
C. elephantopodis (Schw.) Thuem., rust. N. C., S. C., Fla., Ala., Miss.
C. inconspicuum (Long) Hedge. and Long, rust. S. C.
C. ipomoeae (Schw.) Burr., rust. S. C., Fla., Miss.
C. laciniariae Arth., rust. Ga., Fla.
C. terebinthinaceae (Schw.) Thuem., rust. S. C.
Cronartium cerebrum (Pk.) Hedge. and Long, oak-rust. Ga.
C. fusiforme (Pk.) Hedge. and Hunt, rust. Tex.
C. strobilinum Arth., pine-cone rust. Fla.
Fomes pinicola Fr., brown crumbly rot.
Peridermium floridanum Hedge. and G. Hahn, rust. Fla.
Polyporus schweinitzii Fr., red-brown butt-rot.
Septoria acicola (Thuem.) Sacc., leaf-spot. La.
Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot.

P. ponderosa, western yellow pine.

- Botrytis cinerea* Auct., gray-mold. Utah.
Cenangium abietis (Pers.) Rehm, twig-blight. N. Mex., Ariz., Mont., Idaho, Wash., Oreg.
C. piniphilum Weir, canker. Mont., Idaho, Wash.
Ceratostomella pilifera (Fr.) Wint., bluing. S. Dak.
Coleosporium carneum (Bosc) Jack., rust. D. C., Ohio.
C. delicatulum Arth. and Kern, rust. D. C.
C. solidaginis (Schw.) Thuem., rust. Md.
Corticium vagum Berk. and Curt., damping-off. Nebr.
Cronartium cerebrum (Pk.) Hedge. and Long, oak-rust. Pa., Wis., Nebr., Idaho.

PINUS—Continued.

P. ponderosa, western yellow pine—Continued.

- C. comptoniae* Arth., sweet-fern rust. Mass., N. H., Conn., Mich.
C. filamentosum (Pk.) Hedge. and Long, rust. Colo., Mont., Ariz., Utah, Wash., Calif.
C. harknessii (Moore) E. P. Meinecke, rust. Idaho, Wash., Calif.
C. pyriforme (Pk.) Hedge. and Long, rust. Mont., Wyo., Colo., S. Dak., Ariz., Utah, Idaho; Pacific coast.
C. stalactiforme Arth., rust. Mont.
Fomes officinalis Fr., reddish brown-rot. S. Dak., Ariz., Wash.
F. pinicola Fr., brown crumbly rot. Western States.
F. roseus (Alb. and Schw.) Fr., brown pocket rot. General in host range.
Fusarium sp., damping-off. Vt., N. Y., N. J., Nebr.
Hypoderma deformans Weir, needle cast. Pacific Northwest and Mont.
Lentinus sp., red-brown heart-rot. Ariz.
Lophodermium pinastri (Schrad.) Chev., needle cast. Idaho, Oreg.
Neopeckia coulteri (Pk.) Sacc., brown felt-blight. Calif.
Pestalozzia sp., leaf-blight. N. Y., Nebr.
Polyporus ellisianus (Murr.) Long, red-rot. S. Dak., N. Mex., Ariz., Wash.
P. ponderosus Schrenk, red-rot. S. Dak.
P. schweinitzii Fr., red-brown butt-rot. Mont.
P. sulphureus (Bull.) Fr., brown-checked wood-rot.
Pythium debaryanum Hesse, damping-off.
Razoumofskyia americana Nutt., mistletoe. Mont., Idaho, Oreg.
R. campylopoda (Engelm.) Piper, mistletoe. Northern Rocky Mt. States; Pacific Northwest.
R. cryptopoda (Engelm.) Coville, mistletoe. Rocky Mt. States.
Rhizina inflata (Schaeff.) Sacc., seedling blight. Mont.
Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. S. Dak., Mont., Oreg., Calif.
P. pungens, Table Mountain pine.
Coleosporium solidaginis (Schw.) Thuem., rust. Md.
Cronartium pyriforme (Pk.) Hedge. and Long, rust. Pa., Northwestern States.
P. radiata, Monterey pine.
Cronartium cerebrum (Pk.) Hedge. and Long, rust. Calif.
Lophodermium pinastri (Schrad.) Chev., needle cast. Calif.
Razoumofskyia campylopoda (Engelm.) Piper, mistletoe. Calif.
P. resinosa, red pine.
Coleosporium delicatulum (Arth. and Kern) Hedge. and Long, rust. Vt.
C. solidaginis (Schw.) Thuem., needle rust. N. H., Vt., N. Y., Pa., S. C., Minn.
Corticium vagum Berk. and Curt., damping-off. Conn., Mich., Minn.
Cronartium comptoniae Arth., sweet-fern rust. N. H.
Fomes pinicola Fr., brown crumbly rot. General within range of host.
Fusoma parasiticum Tub., damping-off. Minn.
Polyporus schweinitzii Fr., red-brown butt-rot.
P. sulphureus Fr., brown-checked wood-rot. Minn.
Pythium debaryanum Hesse, damping-off. Minn.
Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot.
P. rigida, pitch pine.
Armillaria mellea (Vahl) Quel., root-rot. Pa.
Coleosporium campanulae (Pers.) Lévl., rust. Ohio.
C. carneum (Bosc) Jack., rust. Conn., Pa., D. C., Va., N. C., W. Va., Ohio, Ind.
C. delicatulum (Arth. and Kern) Hedge. and Long, rust. New England, south to N. C.
C. elephantopodis (Schw.) Thuem., rust. D. C., N. C.
C. ipomoeae (Schw.) Burr., rust. Pa., Md., S. C., Ga.
C. lacinariae Arth., rust. N. J.
C. solidaginis (Schw.) Thuem., rust. Conn. to Md.
C. terebinthinaceae (Schw.) Arth., rust. N. C.
Cronartium cerebrum (Pk.) Hedge. and Long, oak-rust. Mass., N. Y., N. J., Pa.

PINUS—Continued.

P. rigida, pitch pine—Continued.

- C. comploniae* Arth., sweet-fern rust. New England, N. Y., Pa.
C. pyriforme (Pk.) Hedgec. and Long, rust. Vt., Conn., N. J., North-western States.
Fomes pinicola Fr., brown crumbly rot. General within range of host.
Hypoderma brachysporum (Rostr.) Tub., needle cast. Mass.
Lophodermium pinastri (Schr.) Chev., needle cast. R. I., N. J.
Pestalozzia funerea Desm., leaf-blight. Mass.
Polyporus schweinitzii Fr., red-brown butt-rot.
Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Me., N. J., Ala., Ky.

P. sabiniana, digger pine.

- Cronartium cerebrum* (Pk.) Hedgec. and Long, oak-rust. Calif.
C. harknessii (Moore) E. P. Meinecke, rust. Calif.
Razoumofskyia campylopoda (Engelm.) Piper, mistletoe. Calif.

P. serotina, pond pine.

- Coleosporium carneum* (Bosc) Jack., rust. S. C., Fla., Ga.
C. delicatulum (Arth. and Kern) Hedgec. and Long, rust. N. J., S. C., Ga., Fla.
C. elephantopodis (Schw.) Thuem., rust. Fla.
C. ipomoeae (Schw.) Burr., rust. S. C.
C. solidaginis (Schw.) Thuem., rust. N. C., S. C.
C. terebinthinaceae (Schw.) Arth., rust. S. C.

P. strobus, white pine.

- Armillaria mellea* (Vahl) Quel., root-rot. Pa.
Cenangium abietis (Pers.) Rehm, canker. Ohio.
Coccomyces pini (Alb. and Schw.) Karst., bark-blight. Ga.
Corticium vagum Berk. and Curt., damping-off. Conn., Mass., N. Y., Minn.
Cronartium cerebrum (Pk.) Hedgec. and Long, oak-rust. Ga.
C. ribicola Fisch., blister rust. Me. to Minn., south to N. J. and Pa.
Cytospora sp. Pa.
Fomes roseus (Alb. and Schw.) Fr., brown pocket rot. General in host range.
Fusarium sp., damping-off. Vt., N. Y., Iowa.
Fusoma parasiticum Tub., damping-off. Minn.
Hendersonia foliicola (Berk.) Fckl., leaf-spot. Mass.
Hypoderma lineare (Pk.) Thuem., needle cast. N. Y., Pa.
H. strobicola Tub., needle cast. New England, Pa., N. C.
Lophodermium pinastri (Schr.) Chev., needle cast. New England, Pa., Wis.
Nectria cucurbitula Fr., canker. Vt., Pa.
Pestalozzia sp., leaf-blight. Vt., Mass., N. Y.
Phoma harknessii Sacc., canker. Conn., Mass., N. Y.
Polyporus berkeleyi Fr., string and ray rot. Minn.
P. ellisianus (Murr.) Long, red-rot. Vt.
P. schweinitzii Fr., red-brown butt-rot. Me.
Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. N. H., Pa., N. C., Minn.

P. sylvestris, Scotch pine.

- Coleosporium sonchi-arvensis* (Pers.) Lév., rust. Wis.
Corticium vagum Berk. and Curt., damping-off. Conn.
Cronartium cerebrum (Pk.) Hedgec. and Long, oak-rust. Pa.
C. comploniae Arth., rust. N. H., Mass., N. Y., Ohio.
C. pyriforme (Pk.) Hedgec. and Long, rust. Conn.
Cytospora pinastri Fr., needle blight. Pa.
Fusoma parasiticum Tub., damping-off. Minn.
Pestalozzia sp., leaf-blight. Vt., Nebr.
Phytophthora cactorum (Leb. and Cohn) Schroet., damping-off. Minn.
Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. N. J.

PINUS—Continued.

P. taeda, loblolly pine.*Coleosporium apocynaceum* Cke., rust. Fla.*C. carneum* (Bosc) Jack., rust. Southeastern States, N. J., Ark.*C. delicatulum* (Arth. and Kern) Hedge. and Long, rust. Va., N. C., S. C., Fla., Tex.*C. elephantopodis* (Schw.) Thuem., rust. South Atlantic and Gulf States.*C. ipomoeae* (Schw.) Burr., rust. Southeastern States.*C. laciniariae* Arth., rust. Fla.*C. pini* Gall., rust. Md.*C. solidaginis* (Schw.) Thuem., rust. Southeastern States.*C. terebinthinaceae* (Schw.) Arth., rust. S. C., Ala.*Cronartium cerebrum* (Pk.) Hedge. and Long, oak-rust. Fla., Ala., Miss., La.*C. comptoniae* Arth., sweet-fern rust. N. J., Ga.*C. pyriforme* (Pk.) Hedge. and Long, rust. N. J.*Fomes annosus* Fr., white spongy rot. New England.*F. pinicola* Fr., brown crumbly rot.*Lophodermium pinastri* (Schrad.) Chev., needle cast. La.*Peridermium minutum* Hedge. and Hunt, rust. Fla.*Polyporus schweinitzii* Fr., red-brown butt-rot.*Septoria acicola* (Thuem.) Sacc., leaf-spot. Fla.*Trametes pini* (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot.*P. virginiana*, scrub pine.*Coleosporium helianthi* (Schw.) Arth., rust. Pa. to Ga.*C. inconspicuum* (Long) Hedge. and Long, rust. Md. to Ga.*C. pini* Gall., rust. Middle Atlantic States, Tenn.*Cronartium cerebrum* (Pk.) Hedge. and Long, oak-rust. Pa. to Ala.*C. comptoniae* Arth., sweet-fern rust. Md.*Fomes pinicola* Fr., brown crumbly rot.*F. roseus* (Alb. and Schw.) Fr., brown pocket rot.*Lophodermium pinastri* (Schrad.) Chev., needle cast. Md.*Polyporus schweinitzii* Fr., red-brown butt-rot.*Trametes pini* (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Md., Ga.

PISUM.

P. sativum, pea.*Aphanomyces euteiches* Drechs., root-rot. Conn., N. Y., Pa., N. J., Del., Md., Ohio, Ind., Ill., Mich., Wis., Mont., Utah, Idaho, Calif.*Bacillus lathyri* Manns and Taub., bacterial streak. Del.*Bacterium pisi* (W. G. Sackett) EFS. (*Phytophthora pisi* (Sack.) Com. S. A. B.), bacterial blight. N. Y., Pa., Del., Md., S. C., Fla., Ill., Ind., Mich., Wis., N. Dak., Colo., Mont., Oreg.*Colletotrichum pisi* Pat., anthracnose. Wis.*Corticium vagum* Berk. and Curt., stem-rot. Widespread.*Erysiphe polygoni* DC., powdery mildew. General.*Fusarium martii* minus Sherb., stem-rot. Widespread.*F. martii pisi* F. R. Jones, root-rot. Widespread.*Heterodera radicum* (Greef) Muell., root-rot. Southern States, Calif.*Isariopsis griseola* Sacc., leaf-spot. R. I.*Mycosphaerella pinodes* (Berk. and Blox.) R. E. Stone (*Ascochyta pisi* Lib.), blight. General.*Ozonium omnivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.*Peronospora viciae* (Berk.) D By., downy mildew. N. Y., Pa., Del., Md., Tenn., Wis., Iowa, Kans., Oreg., Calif.*Phytophthora cactorum* (Leb. and Cohn) Schroet., blight. Conn.*Pythium* spp., root-rot. Conn., Del., Ill., Mich., Wis., Mont.*Sclerotinia sclerotiorum* (Lib.) Mass., stem-rot. Pa., Del., Fla.*Septoria flagellifera* Ell. and Ev., leaf-spot. Wis., S. Dak.*S. pisi* West., leaf-spot. North of Ohio and Missouri Rivers; Tex., Mont., Colo., Calif.*Thielavia basicola* (Berk. and Br.) Zopf, black root-rot. Conn., N. Y., Pa., Md., Ark.*Uromyces fabae* (Pers.) D By., rust. Minn., N. Dak., Nebr.

PLATANUS.

Platanus sp., *sycamore, planetree.

- Armillaria mellea* (Vahl) Quel., root-rot. Md., W. Va.
Cercospora platanicola Ell. and Ev., leaf-spot. Ga., La.
Gnomonia veneta (Sacc. and Speg.) Kleb., anthracnose. General.
Microsphaera alni (Wallr.) Wint., powdery mildew. East of Great Plains.
Phleospora multumaculans Heald and Wolf, leaf-blight. Tex.
Phoradendron flavescens (Pursh) Nutt., mistletoe. Calif.
Phyllosticta platani Sacc. and Speg., leaf-spot. Ala., Kans.
Septoria platanifolia Cke., leaf-spot. Md., S. C.
Stigmata platani (Fckl.) Sacc., leaf-spot. W. Va., Ga., Miss., Okla., Ark., Ind., Ill., Mo., Calif.

POA.

P. compressa, Canada bluegrass.

- Erysiphe graminis* DC., powdery mildew. Minn.
Ophiobolus graminis Sacc., take-all. N. Y.
Puccinia coronata Cda., crown-rust. Mo., Kans.
P. epiphylla Wetts., rust. Minn., Colo.
P. graminis Pers., stem-rust. Pa., Ind., Mich.
Scolecotrichum graminis Fckl., leaf-spot. Wis., Minn., Nebr.

P. pratensis, Kentucky bluegrass.

- Claviceps microcephala* (Wallr.) Karst., ergot. Ind.
C. purpurea (Fr.) Tul., ergot. Ky., Ill., Minn.
Colletotrichum graminicolum (Ces.) Wils., anthracnose. Ohio, Ind., Minn.
Corticium vagum Berk. and Curt., brown-patch. General.
Entyloma irregulare Johans., smut. Ill., Minn.
Epichloe typhina (Pers.) Tul. Del.
Erysiphe graminis DC., powdery mildew. General, especially east of Great Plains.
Helminthosporium vagans Drechs., leaf-spot. New England, N. Y., Pa., Md., Ill., Wis.
Phyllachora graminis (Pers.) Fckl., tar-spot. Nebr.
Puccinia glumarum (Schm.) Eriks. and Henn., stripe-rust. Idaho.
P. epiphylla Wetts., rust. General.
P. triticea Eriks., leaf-rust. Colo.
Sclerotium sp., damping-off.
Scolecotrichum graminis Fckl. Pa., Ind., Minn.
Septoria sp., leaf-blotch. Pa., Ind., Wis., Minn.
Uromyces poae Rabh., rust. Ind.
Ustilago striaeformis (West.) Niessl, stripe-smut. General.

POINCIANA.

P. regia, royal poinciana.

- Gloeosporium* sp., anthracnose. Fla.

POPULUS.

Populus spp., poplar, cottonwood, aspen.

- Bacterium tumefaciens* EFS. and Town. (*Phytoplasma tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Conn., N. J., Minn., Kans., Wyo., Utah.
Cercospora populina Ell. and Ev., leaf-spot. Ala., La.
Cladosporium subsessile Ell. and Barth., leaf-spot. Wis.
Cytospora chrysosperma (Pers.) Fr., canker. General.
Dimerosporium populi Ell. and Ev., sooty mold. Mont.
Dothichiza populea Sacc. and H. Briard, canker. General east of Great Plains.
Fomes applanatus (Pers.) Wallr. (*Ganoderma applanatum* (Pers.) Pat.), white-streaked rot. Conn., Minn., N. Dak., S. Dak., N. Mex.
F. everhartii (Ell. and Gall.) Schrenk, white-rot. Mont.
F. fomentarius (L.) Fr., white sapwood rot. Minn.
F. ignarius (L.) Fr., white heart-rot. General.
Hypoxylon pruinatum (Klotzsch) Cke., canker. Me., N. Y., Mich.
Marssonina brunnea (Ell. and Ev.) Sacc., anthracnose. N. J., Wis., Wash.
M. curvata Bub. and Kab., leaf-spot. Utah.
M. populi (Lib.) Magn., leaf-spot. Wis.

POPULUS—Continued.

Populus spp., poplar, cottonwood, aspen—Continued.

- M. rhabdospora* (Ell. and Ev.) Magn., leaf-spot. N. J., Wis.
Melampsora abietis-canadensis (Farl.) F. Ludwig, rust. General.
M. albertensis Arth., rust. Western States.
M. larici-populina Kleb., rust. General.
M. medusae Thuem., rust. General.
M. occidentalis Jack., rust. Western States; Wis.
Napicladium tremulae (Frank) Sacc., twig-blight. Minn.
Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.
Phoradendron flavescens (Pursh) Nutt., mistletoe. Southwest.
P. flavescens macrophyllum Engelm., mistletoe. N. Mex., Ariz.
Phyllosticta alcides Sacc., leaf-spot. Ohio, Idaho.
P. brunnea Dearn. and Barth., leaf-spot. Mont., Colo., Ariz.
P. intermixta Seaver, leaf-spot. Kans.
P. maculans Ell. and Ev., leaf-spot. N. Y., Kans.
Polyporus dryadeus (Pers.) Fr., white root-rot.
P. dryophilus Berk., white pocket heartwood rot. Me., N. H., Wis., Colo.
Rhytisma nervale (Alb. and Schw.) Rehm, tar-spot. N. Y.
Sclerotium bifrons Ell. and Ev., leaf-spot. New England and Adirondack region; Wis., Wyo., Colo., Utah, Oreg.
Septoria musiva Pk., leaf-spot. Widespread.
S. populicola Pk., leaf-spot. Nebr., Tex.
Stereum purpureum Fr., silver-leaf. Wash.
Taphrina aurea (Pers.) Fr., yellow leaf blister. Me., Mass., N. Y., N. J., Iowa; Pacific coast.
Trochila popularum Desm., leaf-spot. Widespread.
Uncinula salicis (DC.) Wint., powdery mildew. General.
Venturia tremulae Aderh., scab. Me., Wash.

PRIMULA.

Primula sp., primrose.

- Botrytis cinerea* Auct., gray-mold. Conn., S. C., Ohio.
Heterodera radicolica (Greef) Muell., root-knot. Conn., W. Va.
Ramularia primulae Thuem., leaf-spot. Conn., N. Y., Del.
 Mosaic, undet. Pa. (greenhouse).

PROSOPIS.

Prosopis sp., prairie mesquite.

- Cercospora prosopidis* Heald and Wolf, leaf-blight. Tex.
Erysiphe sp., powdery mildew. Tex.
Gloeosporium leguminum Cke., pod-spot, anthracnose. Tex.
Neoravenelia holwayi (Diet.) Long, rust. Tex., Calif.
Phoradendron californicum Nutt., mistletoe. Tex., Ariz., Calif.
P. flavescens (Pursh) Nutt., mistletoe. Tex.
P. flavescens macrophyllum Engelm., mistletoe. Ariz., N. Mex.
Phyllosticta juliflora Ell. and Barth., leaf-spot. Tex.
Ravenelia arizonica Ell. and Ev., rust. Tex., Ariz., Calif.

PRUNUS.

Prunus sp., cherries (sweet and sour).

- Armillaria mellea* (Vahl) Quel., root-rot. Ohio, Mich., Ark., Pacific coast.
Bacillus amylovorus (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. Ohio, Mich., Kans., N. Mex., Wash.
Bacterium cerasi F. L. Griffin, bacterial gummosis. Okla., Ind., Mich., Idaho; Pacific coast.
B. pruni EFS. (*Phytophthora pruni* (EFS.) Com. S. A. B.), bacterial spot. N. Y., Ala., Okla., Ill.
B. tumefaciens EFS. and Town. (*Phytophthora tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Mass., N. Y., Okla., Ohio, N. Mex., Colo.; Pacific Northwest.
Cercospora cerasella Sacc., leaf-spot. Ga., Fla., Ala., Miss.
C. circumscissa Sacc., shot-hole. Widespread.
C. rosicola Pass., leaf-spot. Oreg.

PRUNUS—Continued.

Prunus sp., cherries (sweet and sour)—Continued.

- Cladosporium carpophilum* Thuem., scab. Me., N. H., N. Y., Mich., Iowa, Oreg.
Clitocybe parasitica E. M. Wilcox, root-rot. Ark.
Coccomyces hiemalis Hig., leaf-spot. General.
Coryneum beijerinckii Oud., blight. Pacific coast, Idaho, Utah.
Eoascus cerasi (Fekl.) Sadeb., witches'-broom. North Atlantic Coast States; Tenn., Pacific coast.
E. pruni Fekl., pockets. Minn.
Fomes applanatus (Pers.) Wallr. (*Ganoderma applanatum* (Pers.) Pat.), white-streaked rot. N. Y.
Phyllosticta prunicola Sacc., leaf-spot. Wis., Minn.
Plowrightia morbosa (Schw.) Sacc., black-knot. East of Great Plains.
Podosphaera oxyacanthae (Fr.) D By., powdery mildew. General.
Polyporus sulphureus (Bull.) Fr., brown-checked wood-rot.
Schizophyllum commune Fr., sap-rot. Ohio, Minn., Nebr.
Sclerotinia fructicola (Wint.) Rehm (*S. americana* (Wormald) Norton and Ezekiel), brown-rot. General.
Stereum purpureum Fr., associated with silver-leaf. N. Y., Minn., Oreg.
Tranzschelia punctata (Pers.) Arth., rust. Miss., Mich., Nebr., Calif.
Valsa leucostoma (Pers.) Fr., die-back. Ohio, Mo.
 —*Venturia cerasi* Aderh., scab. Ind.

P. avium, sweet cherry.

- Alternaria* sp., fruit-rot. Mich., Idaho.
A. citri cerasi B. A. Rudolph, leaf-spot. Calif.
Armillaria mellea (Vahl) Quel., root-rot. Pacific coast.
Bacillus amylovorus (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. Wash.
Bacterium cerasi F. L. Griffin, bacterial gummosis. Oreg., Calif.
B. pruni EFS. (*Phytomonas pruni* (EFS.) Com. S. A. B.), bacterial spot. N. Y., Mich.
Cercospora cerasella Sacc., leaf-spot. Ga., Ala.
C. circumscissa Sacc., shot-hole. N. J., Pa., Ohio.
Coccomyces hiemalis Hig., leaf-spot. N. Y., Pa., Md., Wash., Oreg.
Coryneum beijerinckii Oud., blight. Pacific Northwest.
Eoascus cerasi (Fekl.) Sadeb., witches'-broom. N. Y., Wash., Oreg., Alaska.
E. communis Sadeb., pockets. Iowa.
E. pruni Fekl., pockets. Okla.
Physalospora malorum (Pk.) Shear (*P. cydoniae* Am. Auct.) black-rot. East of Rocky Mts.
Plowrightia morbosa (Schw.) Sacc., black-knot. N. Y., Mich.
Podosphaera oxyacanthae (DC.) D By., powdery mildew. N. Y., Md., Iowa, Wash.
Polyporus lacteus Fr., heart-rot. Oreg.
Sclerotinia cinerea (Bon.) Schroe¹. (*Monilia oregonensis* Barss and Posey), blossom blight. Pacific coast.
S. fructicola (Wint.) Rehm (*S. americana* (Wormald) Norton and Ezekiel), brown-rot. General.
Tranzschelia punctata (Pers.) Arth., rust. Okla.

P. cerasus, sour cherry.

- Alternaria* sp., fruit-rot. Mich.
Bacterium pruni EFS. (*Phytomonas pruni* (EFS.) Com. S. A. B.), bacterial spot. N. Y.
Cercospora cerasella Sacc., leaf-spot. Ala.
Coccomyces hiemalis Hig., leaf-spot. General east of Great Plains and north of Ky. and Va.
Eoascus cerasi (Fekl.) Sadeb., witches'-broom. Md., Wis.
E. communis Sadeb., pockets. Iowa.
Phyllosticta prunicola Sacc., leaf-spot.
Podosphaera oxyacanthae (DC.) D By., powdery mildew. General.
Plowrightia morbosa (Schw.) Sacc., black-knot. N. Y., Pa., Md., Va., W. Va., N. C.
Sclerotinia fructicola (Wint.) Rehm (*S. americana* (Wormald) Norton and Ezekiel), brown-rot. N. J., Ind.

PRUNUS—Continued.

P. besseyi, *P. cuneata*, *P. pumila*, sand cherry.

Bacterium tumefaciens EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Minn.

Eoascus communis Sadeb., pockets. N. Y., Wis., N. Dak., S. Dak., Kans.

Plowrightia morbosa (Schw.) Sacc., black-knot. N. Dak.

Podosphaera oxyacanthae (DC.) D By., powdery mildew. Iowa, N. Dak., S. Dak., Nebr.

Sclerotinia cinerea (Bon.) Schroet. (*Monilia oregonensis* Barss and Posey), blossom blight. Wash., Oreg.

S. fruticola (Wint.) Rehm (*S. americana* (Wormald) Norton and Ezekiel), brown-rot. Minn., Nebr.

Tranzschelia punctata (Pers.) Arth., rust. Nebr., Kans.

P. pennsylvanica, pin cherry.

Coccomyces hiemalis Hig., leaf-spot. N. Y., W. Va.

Eoascus cerasi (Fekl.) Sadeb., witches'-broom. N. H.

Phyllosticta congesta Heald and Wolf, leaf-spot, Wis.

Physalospora malorum (Pk.) Shear (*P. cydoniae* Am. Auct.), black-rot.

Plowrightia morbosa (Schw.) Sacc., black-knot. Me., W. Va., Minn.

Podosphaera oxyacanthae (DC.) D By., powdery mildew. Ill.

Tranzschelia punctata (Pers.) Arth., rust. Ill.

P. serotina, wild black cherry.

Coccomyces hiemalis Hig., leaf-spot. Widespread east of Great Plains.

C. lutescens Hig., leaf-blight. N. Y., Pa., Ga., Miss., Ill.

Eoascus cerasi (Fekl.) Sadeb., witches'-broom. Mass., Ala.

E. farlowii Sadeb., leaf-curl. Ala.

E. pruni Fekl., pockets. Va., Ala.

E. varius Atk., leaf deformation. Del., Fla., Ala.

Fomes fomentarius (L.) Fr., white sapwood rot.

Gloeosporium serotinum Ell. and Ev., leaf-spot. Del.

Phoradendron flavescens (Pursh) Nutt., mistletoe. Ind.

Phyllosticta prunicola Sacc., leaf-spot. N. Y., Wis.

P. serotina Cke., leaf-spot. Widespread east of Great Plains.

P. virginiana (Ell. and Hals.) Seaver, leaf-spot. Iowa, Kans.

Plowrightia morbosa (Schw.) Sacc., black-knot. Widespread east of Miss. River.

Podosphaera oxyacanthae (DC.) D By., powdery mildew. N. Y., Ill.

Sclerotinia fruticola (Wint.) Rehm (*S. americana* (Wormald) Norton and Ezekiel), brown-rot. S. C.

S. seaveri Rehm. N. Y., Pa., Del., Md., Va., W. Va., Ga., Wis., Iowa.

Septoria cerasina Pk., leaf-spot. Mass.

S. ravenelii Thuem., leaf-spot. Pa., N. J., S. C., Ill.

Tranzschelia punctata (Pers.) Arth., rust. S. C., Ala., Tex., Ind., Ill., Iowa, Mo.

Valsa leucostoma (Pers.) Fr., die-back. N. Y.

P. virginiana, *P. demissa*, chokecherry.

Bacterium pruni EFS. (*Phytomonas pruni* (EFS.) Com. S. A. B.), bacterial spot. N. Y.

Cercospora cerasella Sacc., leaf-spot. Kans.

C. circumscissa Sacc., shot-hole. N. Dak., Kans.

Coccomyces hiemalis Hig., leaf-spot. Widespread in northern United States.

C. lutescens Hig., leaf-blight. N. Y., Wash.

Cylindrosporium tubeufianum Allesch., leaf-spot. N. Y.

Cytospora chrysosperma (Pers.) Fr., canker. Mont.

Eoascus cerasi (Fekl.) Sadeb., witches'-broom. Nebr., Colo.

E. confusus Atk., fruit deformation. Mass.

E. farlowii (Sadeb.) Sacc., leaf-curl. Wash.

E. pruni Fekl., pockets. N. Dak., Wash.

Gloeosporium prunicolum Ell. and Ev., leaf-spot. Wis.

Phyllactinia corylea (Pers.) Karst., powdery mildew. Wash.

Phyllosticta circumscissa Cke., leaf-spot. Kans., Wash.

P. prunicola Sacc., leaf-spot. N. Y.

P. virginiana (Ell. and Hals.) Seaver, leaf-spot. Wis., Iowa, Kans., Mont.

Physalospora malorum (Pk.) Shear (*P. cydoniae* Am. Auct.), black-rot.

PRUNUS—Continued.

P. virginiana, *P. demissa*, chokecherry—Continued.

Plowrightia morbosa (Schw.) Sacc., black-knot. Widespread in Northern States.

Podosphaera oxyacanthae (DC.) D By., powdery mildew. Widespread in Northern States.

Sclerotinia angustior Rehm. N. Y., Pa., Iowa, N. Dak.

S. demissa B. F. Dana, blight. Idaho, Wash.

S. fructicola (Wint.) Rehm (*S. americana* (Wormald) Norton and Ezekiel), brown-rot. Conn., Fla., Kans.

Septogloeum nuttallii Hark., leaf-spot. Oreg.

Tranzschelia punctata (Pers.) Arth., rust. S. C., Mich., Ill.

P. armeniaca, apricot.

Armillaria mellea (Vahl) Quel., root-rot. Calif.

Bacillus amylovorus (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. Pa., Tex., Nebr., Colo.

Bacterium cerasi F. L. Griffin, bacterial gummosis. Calif.

B. pruni EFS. (*Phytophthora pruni* (EFS.) Com. S. A. B.), bacterial spot. Ill., Nebr.

B. tumefaciens EFS. and Town. (*Phytophthora tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Colo., Ariz., Idaho, Pacific coast.

Botrytis cinerea Auct., fruit-rot. Calif.

Cercospora circumscissa Sacc., shot-hole. Tex., Calif.

Cladosporium carpophilum Thuem., scab. Widespread.

Coccomyces sp., leaf-spot. Mass., Tex.

Coryneum beijerinckii Oud., blight. Utah, Idaho, Pacific coast.

Heterodera radicola (Greef) Muell., root-knot. Ariz.

Monilia sp., blossom blight. Calif.

Monochaetia rosenwaldia A. Khazanoff, tumor. Calif.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Ariz.

Phyllosticta circumscissa Cke., shot-hole. Calif.

Plowrightia morbosa (Schw.) Sacc., black-knot. N. Y., Colo.

Sclerotinia cinerea (Bon.) Schroet. (*Monilia oregonensis* Barss and Posey), blossom blight. Wash., Oreg.

S. fructicola (Wint.) Rehm (*S. americana* (Wormald) Norton and Ezekiel), brown-rot. Mass., Conn., N. Y., Md., Ohio, Ind., Ill., Tex., Okla., Colo., Pacific coast.

S. sclerotiorum (Lib.) Mass., green-rot. Calif.

Septoria cerasina Pk., leaf-spot. Calif.

Tranzschelia punctata (Pers.) Arth., rust. Miss., Tex., Calif.

Valsa leucostoma (Pers.) Fr., twig-blight. Tex.

Verticillium sp., wilt. Calif.

P. domestica (in part), prune.

Armillaria mellea (Vahl) Quel., root-rot. Pacific coast.

Bacillus amylovorus (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. Pacific Northwest.

Bacterium cerasi F. L. Griffin, bacterial gummosis. Pacific coast.

B. pruni EFS. (*Phytophthora pruni* (EFS.) Com. S. A. B.), bacterial spot. N. Y., Pa.

B. tumefaciens EFS. and Town. (*Phytophthora tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Pacific Northwest; Calif.

Cercospora circumscissa Sacc., shot-hole. Wash.

Cladosporium carpophilum Thuem., scab. Ohio, Nebr., Pacific Northwest.

Coccomyces prunophorae Hig., leaf-spot. N. Y., Del., Nebr., Pacific coast.

Coryneum beijerinckii Oud., blight. Wash., Oreg.

Exoascus pruni (Berk.) Fekl., pockets. Nebr.

Plowrightia morbosa (Schw.) Sacc., black-knot. Mich., Nebr.

Sclerotinia cinerea (Bon.) Schroet. (*Monilia oregonensis* Barss and Posey), blossom blight. Wash., Oreg.

S. fructicola (Wint.) Rehm (*S. americana* (Wormald) Norton and Ezekiel), brown-rot. Widespread.

Tranzschelia punctata (Pers.) Arth., rust. Nebr., Kans., Tex., Oreg., Calif.

PRUNUS—Continued.*Prunus* spp., plum.

- Armillaria mellea* (Vahl) Quel., root-rot. Minn., N. Dak., Tex., Calif.
Bacillus amylovorus (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. Vt., N. J., Pa., Miss., Tex., Ohio, Mich., S. Dak., Nebr., Colo., Ariz., Wash.
Bacterium cerasi F. L. Griffin, bacterial gummosis. Tex., Kans., Ariz., Calif.
B. pruni EFS. (*Phytophthora pruni* (EFS.) Com. S. A. B.), bacterial spot. General east of Great Plains.
B. tumefaciens EFS. and Town. (*Phytophthora tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Occurs in widely scattered States.
Botrytis cinerea Auct., fruit-rot. Wash.
Cercospora circumscissa Sacc., shot-hole. N. Y., Kans., Wash.
Cladosporium carpophilum Thuem., scab. Widespread.
Coccomyces prunophorae Hig., leaf-spot. General.
Coryneum beijerinckii Oud., blight. Pacific coast, Idaho.
Diplodia pruni Fekl., twig-blight. Fla.
Exoascus communis Sadeb., pockets. Iowa.
E. mirabilis Atk., hypertrophy. Widespread east of Great Plains.
E. pruni (Berk.) Fekl., pockets. General.
Fomes fulvus Fr., brown heartwood rot. Minn., Nebr.
F. pinicola Fr., brown crumbly rot. Oreg.
Heterodera radicola (Greef) Muell., root-knot. S. C.
Lasiodiplodia triflorae Hig., wilt. Md., Va., Ga.
Lenzites saepiaria Fr., dry-rot. Oreg.
Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Ariz.
Phoradendron flavescens (Pursh) Nutt., mistletoe. Calif.
Phyllosticta circumscissa Cke., shot-hole. N. J., Ga., Wash.
P. congesta Heald and Wolf, blotch. Ga., Tex.
P. prunicola Sacc., leaf-spot. Minn.
Ploerightia morbosa (Schw.) Sacc., black-knot. General except Pacific coast.
Podosphaera oxyacanthae (DC.) D By., powdery mildew. Widespread east of Great Plains.
Polyporus lacteus Fr., heart-rot. Va., Ind., Wash., Oreg.
Polystigma rubrum (Pers.) DC., red leaf-spot.
Pythiacystis citrophthora E. H. and R. E. Smith, brown-rot. Calif.
Schizosphyllum commune Fr., sap-rot. Minn., Nebr.
Sclerotinia fructicola (Wint.) Rehm (*S. americana* (Wormald) Norton and Ezekiel), brown-rot. General.
Septobasidium sp., canker. Fla.
S. retiforme (Berk. and Curt.) Pat., canker. Ala., Fla.
Septoria cerasina Pk., leaf-spot. Mass.
Trametes carnea (Nees) Cke., wood-rot. Oreg.
Tranzschelia punctata (Pers.) Arth., rust. Widespread.
Valsa leucostoma (Pers.) Fr., die-back. Pa., S. C., Tex., Ohio, Mo., Colo., Idaho.

P. americana, American wild plum.

- Cercospora circumscissa* Sacc., shot-hole. Nebr.
C. prunicola Ell. and Ev., leaf-spot. Ala., La.
Cladosporium carpophilum Thuem., scab. Wis., Iowa, Calif.
Coccomyces prunophorae Hig., shot-hole. Ind., S. Dak.
Exoascus cerasi (Fekl.) Sadeb., witches'-broom. Ill., Iowa.
E. communis Sadeb., pockets. General over range of host.
E. decipiens Atk., leaf-curl. Mont.
E. mirabilis Atk., hypertrophy. Miss.
E. pruni Fekl., pockets. N. Dak., Iowa, Miss.
Fomes fulvus Fr., brown heartwood rot. Ohio, Ind., N. Dak., S. Dak., Nebr., Kans.
Phyllosticta circumscissa Cke., leaf-spot. Ill.
P. virginiana (Ell. and Hals.) Seaver, leaf-spot. Kans.
Ploerightia morbosa (Schw.) Sacc., black-knot. General over range of host.
Podosphaera oxyacanthae (DC.) D By., powdery mildew. General over range of host.
Sclerotinia fructicola (Wint.) Rehm (*S. americana* (Wormald) Norton and Ezekiel), brown-rot. N. Y., Va., Miss., Iowa.

PRUNUS—Continued.

P. americana, American wild plum—Continued.

Septoria pruni Ell., leaf-spot. Ky., Kans.

S. purpureocincta Ell. and Ev., leaf-spot. Kans.

Tranzschelia punctata (Pers.) Arth., rust. General over range of host.

PSEUDOTSUGA.

P. douglasii, Douglas fir.

Botrytis cinerea Auct., gray-mold twig-blight. Conn.

Corticium vagum Berk. and Curt., damping-off. Colo.

Echinodontium tinctorium Ell. and Ev., brown stringy rot. Utah, Mont., Idaho.

Fomes officinalis Fr., reddish brown-rot. Mont.

F. pinicola Fr., brown crumbly rot. Western States.

F. roseus (Alb. and Schw.) Fr., brown pocket rot. Ariz., Wash.

Melampsora albertensis Arth., rust. Mont., Colo., Utah, Wash.

Phacidium infestans Karst., needle blight. Idaho.

Polyporus schweinitzii Fr., red-brown butt-rot. Mont., Idaho, Wash., Oreg.

P. sulphureus (Bull.) Fr., brown-checked wood-rot. Idaho, Wash.

Pythium debaryanum Hesse, damping-off.

Razoumofskyia douglasii (Engelm.) O. Kuntze, mistletoe. Colo., Pacific Northwest.

Rhabdocline pseudotsugae Syd., needle cast. Oreg.

Sparassis radicata Weir, yellow root-rot. Mont., Pacific Northwest.

Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Mont., Wyo., Idaho, Wash.

PSIDIUM.

P. guajava, guava.

Clitocybe monadelpha (Morgan) Sacc., root-rot. Fla.

Colletotrichum sp., die-back. Fla.

Glomerella cingulata (Ston.) Spauld. and Schrenk, ripe-rot. Fla.

PTELEA.

P. trifoliata, hoptree.

Cercospora afflata Wint., leaf-spot. Mo.

C. pteleae Wint., leaf-spot. Ind., Mo.

Phleospora pteleae Tharp. Utah.

Puccinia windsorise Schw., rust. Scattering throughout range of host.

Septoria pteleae Ell. and Ev., leaf-spot. Ind.

PUNICA.

P. granatum, pomegranate.

Cercospora lythracearum Heald and Wolf, leaf-spot. Tex.

Heterodera radicolica (Greef) Muell., root-knot.

Sterigmatocystis castanea F. Patterson, fruit-rot. Tex., Ariz., Calif.

PYRACANTHA.

P. coccinea, scarlet firethorn.

Bacillus amylovorus (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. Idaho, Wash.

PYRUS.

P. communis, pear.

Armillaria mellea (Vahl) Quel., root-rot. Pacific coast.

Bacillus amylovorus (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. General.

Bacterium tumefaciens EFS. and Town. (*Phytoplasma tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. General.

Botryosphaeria ribis Gross. and Dug., black-rot. Va.

Botrytis sp., spur-blight. Oreg.

B. cinerea Auct., fruit-rot. Wash.

Cephalothecium roseum Cda., pink-mold rot. Conn., R. I., Ill.

Cercospora minima Tr. and Earle, leaf-spot. Miss., Tex.

C. piri Farl., leaf-spot. Mich.

Corticium stevensii Burt (*Hypochnus ochroleucus* F. Noack), leaf-blight. W. Va., N. C., S. C., Ga., Fla., Ala., Tex.

PYRUS—Continued.

P. communis, pear—Continued.

- Coryneum foliicolum* Fekl., leaf-spot. Ind.
Fabraea maculata (Lév.) Atk., leaf-blight. General.
Gloeodes pomigena (Schw.) A. S. Colby (*Phyllachora pomigena* (Schw.) Sacc.), sooty blotch. Mass. to N. C., westward to Ill., Tex.
Gloeosporium perennans Zeller and Childs, perennial canker. Wash., Oreg.
Glomerella cingulata (Ston.) Spauld. and Schrenk, bitter-rot. Conn., Mass., N. J., S. C., Ga., Tex., Ark.
Gymnosporangium blasdaleanum (Diet. and Holw.) Kern, incense-cedar rust. Oreg.
G. globosum Farl., rust. Conn., N. Y., Ohio, Colo.
G. nelsoni Arth., rust. Colo.
Hendersonia foliorum Fekl., leaf-spot. Tex.
Leptothyrium carpophilum Pass., flyspeck. Tex.
Mycosphaerella sentina (Fr.) Schroet., leaf-spot. General east of Great Plains; Oreg.
Myxosporium corticolum Edg. (*Dermatea corticola* (Edg.) Arn.), surface bark canker. N. Y., Mich., Oreg.
Nectria galligena Bres., European canker. Oreg.
Neofabraea malicorticis (A. B. Cordley) Jack., anthracnose. Wash., Oreg.
Nummularia discreta (Schw.) Tul., blister canker. Del.
Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex., N. Mex., Ariz.
Phoradendron flavescens (Pursh) Nutt., mistletoe. Tex.
P. flavescens macrophyllum Engelm., mistletoe. Ariz., N. Mex.
Phyllosticta pyrorum Cke., leaf-spot. S. C., Ill.
Physalospora malorum (Pk.) Shear (*P. cydoniae* Am. Auct.), black-rot. Widespread east of Great Plains; Oreg.
Phytophthora cactorum (Leb. and Cohn) Schroet., rot. Conn., Ind., Mich.
Podosphaera leucotricha (Ell. and Ev.) Salm., powdery mildew. Colo., Wash.
P. oxycanthae (DC.) D By., powdery mildew. Colo.
Pythiacystis citrophthora E. H. and R. E. Smith, citrus brown-rot. Calif.
Sclerotinia cinerea (Bon.) Schroet. (*Monilia oregonensis* Barss and Posey), blossom blight. Pacific coast.
S. fructicola (Wint.) Rehm (*S. americana* (Wormald) Norton and Ezekiel), brown-rot. Conn., Pa., Del., S. C., Ga., Fla., Miss., Ind., Ill., Wash.
Septoria nigerina Fekl., leaf-spot. N. Y.
Venturia pyrina Aderh., scab. General.
Xylaria sp., root-rot. Ind.
 Bitter-pit, undet. N. Y., Wash., Oreg.
 Brown-blotch, undet. N. J.
 Chlorosis, nonpar. Tex., N. Mex., Wash.
 Measles, undet. Fla.

QUERCUS.

Quercus spp., oaks.

- Actinopelte japonica* Sacc., leaf-spot. N. J.
Armillaria mellea (Vahl) Quel., root-rot. Widespread.
Cercospora polytricha Cke., leaf-spot. S. C., Ga.
Cronartium cerebrum (Pk.) Hedge. and Long, rust. East of Great Plains; Calif.
C. fusiforme (Pk.) Hedge. and Hunt, rust. Ohio.
C. strobilinum (Arth.) Hedge. and Hahn, rust. Fla.
Cylindrosporium microspilum Sacc. and Wint., leaf-spot. Mo.
Cytospora chrysosperma (Pers.) Fr., canker. R. I., N. J., Pa.
Diplodia longispora Cke. and Ell., twig-blight. Conn., N. Y., N. J., Pa., Md., Va., W. Va.
Fomes applanatus Fr. (*Ganoderma applanatum* (Pers.) Pat.), white-streaked rot. Me., N. H., Pa., Ky., Minn., Calif.
F. everhartii (Ell. and Gall.) Schrenk, white-rot. Conn., N. J., Pa., N. C., Ohio, Ind., Minn., Mont.
F. fomentarius (L.) Fr., white sapwood rot.
F. ignarius (L.) Fr., white heart-rot. Conn., Mass., N. C., Minn., Okla.
F. lobatus Schw., spongy white-rot. Ark.

QUERCUS—Continued.

Quercus spp., oaks—Continued.

Gnomonia veneta (Sacc. and Speg.) Kleb., anthracnose. East of Great Plains.

Hydnium erinaceus Fr., wet heartwood rot. Ala., Ind., Ark.

Leptothyrium dryinum Sacc., leaf-spot. N. J., Va., Ga., Miss., Tex., Wis.

Marssonina martini (Sacc. and Ell.) Magn., leaf-spot. Miss. Valley; Pa., Va.

M. quercus Pk., leaf-spot. Ind., Mo., Tex.

Microsphaera alni (Wallr.) Wint., powdery mildew. General east of Great Plains; Wash., Calif.

Monochaetia desmazierii Sacc., large leaf-spot. Pa., Southeastern States,

Nectria cinnabarina Fr., canker. Mass., Pa., Ala.

N. galligena Bres., European canker. Oreg.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Phoradendron flavescens (Pursh) Nutt., mistletoe. Tex., Ind., Ariz.

P. villosum Nutt., mistletoe. Oreg.

Phyllactinia corylea (Pers.) Karst., powdery mildew. East of Miss. River.

Phyllosticta agrifolia Ell. and Ev., leaf-spot. Calif.

P. livida Ell. and Ev., leaf-spot. Fla., La., Wis., Calif.

P. ludoviciana Ell. and Mart., leaf-spot. Fla., La.

P. phomiformis Sacc., leaf-spot. Occurs scatteringly east of Great Plains.

P. quercus Sacc. and Speg., leaf-spot. N. Y., Ind., Wis.

P. quercus prini Ell. and Ev., leaf-spot. W. Va.

P. vesicatoria Thuem., leaf-spot. S. C.

P. wislizeni Ell. and Ev., leaf-spot. Calif.

Polyporus berkeleyi Fr., string-rot and ray-rot. N. Y., Pa., N. C., Ala., Ark., Ind., Mo.

P. croceus Fr., white-piped rot. N. Y., N. C., Ark., Ohio, Ind.

P. dryadeus Fr., white wood-rot. Pa., Va., Tex., Okla., Ark., Ohio, Ind., Minn.

P. frondosus Fr., straw-colored rot. Mass., Va., Ark., Ind.

P. obtusus Berk., soft heartwood rot. Tenn., Ga., Ala., Minn.

P. rheades Fr., white pocket rot. General.

P. spraguei Berk., heart-rot.

P. sulphureus (Bull.) Fr., brown-checked wood-rot. General.

Rhytisma erythrosporium Berk. and Curt., tar-spot. Tex.

Septoria dryina Cke., leaf-spot. Mass., S. C., Kans.

S. neglecta Earle, leaf-spot. Ala.

Sphaeropsis quercina Cke. and Ell., twig canker. Pa.

Sphaerotheeca lanestrus Hark., powdery mildew. Ala., Miss., Ind., Ill., Iowa, Calif.

Stereum subpileatum Berk. and Curt., honeycomb heart-rot. Southern States; Ohio, Ky., Mo.

Strumella coryneoidea Sacc. and Wint., canker. Conn., Mass., N. Y., Pa., Ill., Mo., Oreg.

Taphrina coerulescens (Mont. and Desm.) Tul., leaf blister. General east of Great Plains.

RADICULA.

R. armoracia, horseradish.

Albugo candida (Pers.) O. Kuntze, white-rust. Mass. to Mont., south to Kans.

Alternaria brassicae (Berk.) Sacc., leaf-spot. Nebr., Mo.

A. herculea (Ell. and Mart.) J. A. Elliott, gray leaf-spot. Conn., N. J., Ind.

Bacillus sp., bacterial root-rot. N. J.

Cercospora armoraciae Sacc., leaf-spot. Northern States.

Corticium vagum Berk. and Curt., stem-rot. N. J., Mich., Ill., Wash.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Phyllosticta decidua Ell. and Kell., leaf-spot. Wis.

P. orbicula Ell. and Ev., leaf-spot. N. Y.

Ramularia armoraciae Fekl., leaf-spot. General.

Thielavia basicola (Berk. and Br.) Zopf, black root-rot. N. J.

RAPHANUS.*R. sativus*, radish.*Actinomyces scabies* (Thax.) Gues., scab. Ohio, Ind.*Albugo candida* (Pers.) O. Kuntze, white-rust. General.*Alternaria brassicae macrospora* Sacc., leaf-spot. Conn.*Ascochyta* sp., leaf-spot. Mass.*Bacillus carotovorus* L. R. Jones (*Erwinia carotovora* (Jones) Com. S. A. B.), soft-rot. N. Y.*Bacterium campestre* (Pam.) EFS. (*Phytophthora campestre* (Pam.) Com. S. A. B.) black-rot. Ohio.*Cercospora cruciferarum* Ell. and Ev., leaf-spot. Ala.*Corticium vagum* Berk. and Curt., stem-rot. Mass., N. Y., Del., Ohio, Ill., Wash.*Erysiphe polygoni* DC., powdery mildew. Tex., Mo.*Peronospora parasitica* (Pers.) D By., downy mildew. Widespread.*Plasmodiophora brassicae* Wor., clubroot. Conn., N. Y., Pa., Minn.*Puccinia sarcobati* (Pk.) Bethel, rust. Southwestern States.*Pythium aphanidermatum* (Edson) Fitz., black-root. Widely scattered States east of Miss. River.*P. debaryanum* Hesse, damping-off. Va., Minn.*Sclerotinia sclerotiorum* (Lib.) Mass., stem-rot. Minn.**RESEDA.***R. odorata*, mignonette.*Cercospora resedae* Fekl., leaf-spot. Widespread east of Great Plains.*Corticium vagum* Berk. and Curt., stem-rot. Conn.*Fusarium* sp., root-rot, N. Y.**RHAMNUS.***R. cathartica*, buckthorn.*Cercospora rhamni* Fekl., leaf-spot. N. J.*Puccinia coronata* Cda., crown-rust. Vt., N. Y., Ill., Wis., Minn., Iowa, N. Dak., S. Dak.*R. purshiana*, cascara buckthorn.*Cylindrosporium rhamni* Ell. and Ev., leaf-spot. Idaho.*Marssonina rhamni* Ell. and Ev., leaf-spot. Wash.*Phyllosticta rhamnigena* Sacc., leaf-spot. Wash.*Puccinia coronata* Cda., crown-rust. Pacific Northwest; Mont.*Septoria blasdalei* Sacc. and Syd., leaf-spot. Calif.**RHEUM.***R. rhaponticum*, rhubarb.*Ascochyta rhei* Ell. and Ev., leaf-spot. General east of Rocky Mts.*Bacillus carotovorus* L. R. Jones (*Erwinia carotovora* (Jones) Com. S. A. B.), soft-rot. Ohio.*Colletotrichum erumpens* Sacc., anthracnose. Pa., W. Va., Ill.*Corticium vagum* Berk. and Curt., stem-rot. N. Y., Ill., Wash.*Ozonium omnivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.*Peronospora rumicis* Cda., downy mildew. Calif.*Phyllosticta straminella* Bres., leaf-spot. N. Y., Pa., Ind., Ill., Minn., Mo., N. Dak., Nebr.*Phytophthora* sp., crown-rot. N. J., Pa., Del., Tenn., Ill., Mich., Mo.*P. cactorum* (Leb. and Cohn) Schroet., crown-rot. Pa.*P. parasitica rhei* G. H. Godfrey, foot-rot. Md., D. C., Va., Kans.*Puccinia rubella* (Pers.) Arth., rust. Nebr.*Sclerotium rolfsii* Sacc., stem-rot. Va.**RHODODENDRON.***Rhododendron* sp., rhododendron.*Armillaria mellea* (Vahl) Quel., root-rot. Wash.*Coccomyces rhododendri* (Schw.) Sacc., leaf-spot. W. Va. (on *R. maximum*).*Coryneum rhododendri* Schw., leaf-spot. Wash., Oreg. (on *R. californicum*).*Cryptostictis* sp., leaf-spot. Pacific coast.*Exobasidium vaccinii* (Fekl.) Wor., hypertrophy. Pa., N. C., Fla., Ind., Iowa, Wash., Oreg.

RHODODENDRON—Continued.*Rhododendron* sp., rhododendron—Continued.

- Hendersonia concentrica* Ell. and Ev., leaf-spot. N. C.
Laestadia rhodora Berl. and Vogl., leaf-spot. N. Y.
Melampsoropsis piperiana Arth., rust. Oreg., Wash. (on *R. californicum*).
M. roanensis Arth., rust. Tenn.
Pestalozzia funerea Desm., leaf-spot. Va.
P. guepini Desm., leaf-spot. Conn., N. Y., N. J., Pa., Del., Ind., Wash.
Phyllosticta maxima Ell. and Ev., leaf-spot. Conn., N. Y., N. J.
P. saccardoi Thuem., leaf-spot. N. Y.
Septoria rhododendri Cke., leaf-spot. Me.
Sporocybe azaleae (Pk.) Sacc., bud-blight. Pacific coast.
 Witches'-broom and white-leaf, undet. Pacific coast (on *R. californicum*).

RHODORA.*R. canadense*, rhodora.

- Exobasidium vaccinii* (Fekl.) Wor., hypertrophy. N. H.
Pucciniastrum myrtilli (Schum.) Arth., rust. N. H.

RHODOTYPOS.*R. kerrioides*, jetbead.

- Ascochyta rhodotyphi* H. W. Anderson, leaf-spot. Ill.

RHUS.*R. copallina*, shining sumac.

- Cercospora copallina* Cke., leaf-spot. S. C., Fla., La.
C. rhoina Cke. and Ell., leaf-spot. Widespread east of Great Plains.
Cylindrosporium toxicodendri Ell. and Ev., leaf-spot. N. J.
Exoascus purpurascens (Ell. and Ev.) Sadeb., leaf-curl. Mass., N. J., Ga.
Sphaerotheca humuli (DC.) Burr., powdery mildew. Ill., Mo.

R. cotinus, smoketree.

- Cercospora rhoina* Cke. and Ell., leaf-spot. Ala.
Septoria rhoina Berk. and Curt., leaf-spot. New England, N. Y.

R. glabra, smooth sumac.

- Cercospora rhoina* Cke. and Ell., leaf-spot. Va. to Nebr. and southward.
Cladosporium aromaticum Ell. and Ev. Nebr.
Septoria rhoina Berk. and Curt., leaf-spot. Ill., Minn., Kans.
Sphaerotheca humuli (DC.) Burr., powdery mildew. Iowa, Miss.

R. typhina, staghorn sumac.

- Cercospora rhoina* Cke. and Ell., leaf-spot. Md.
Exoascus purpurascens (Ell. and Ev.) Sadeb., leaf-curl. Md.
Phyalospora malorum (Pk.) Shear (*P. cydoniae* Am. Auct.), canker. N. Y.
Septoria rhoina Berk. and Curt., leaf-spot. Miss., Iowa.
Sphaerotheca humuli (DC.) Burr., powdery mildew. Pa.

RIBES.*R. nigrum*, European black currant.

- Cronartium ribicola* Fisch., blister rust. New England, N. Y., Wash., Oreg.
Microsphaera grossulariae (Wallr.) Lév., powdery mildew. Mont.
Mycosphaerella grossulariae (Fr.) Lindau, leaf-spot. Conn., Iowa.
Pseudopeziza ribis Kleb., anthracnose. N. H.
Puccinia grossulariae (Schum.) Lagh., rust. Conn.

R. vulgare, red currant, white currant.

- Botryosphaeria ribis* Gross. and Dug., cane-blight. Mass., Conn., N. Y. to Va., west to Ill.
Cercospora angulata Wint., leaf-spot. N. Y. to Va., west to Iowa and Kans.
Cronartium ribicola Fisch., blister rust. North Atlantic States; Mich., Minn.
Fomes ribis (Schum.) Fr., root-rot. N. Y., N. J., Ind., Utah.
Glomerella cingulata (Ston.) Spauld. and Schrenk, bitter-rot. Conn.
Mycosphaerella grossulariae (Fr.) Lindau, leaf-spot. General in Northern States.
Nectria ditissima Tul., canker. N. Y.
Phyllosticta grossulariae Sacc., leaf-spot. N. J.
Plasmopara ribicola Schroet., downy mildew. Wis.

RIBES—Continued.

R. vulgare, red currant, white currant—Continued.

Pleonectria berolinensis Sacc., doubtfully parasitic on canes. Conn., Mass., Md., Ohio, Mich., Ind., N. Dak., Utah.

Pseudopeziza ribis Kleb., anthracnose. Widespread in Northern States.

Puccinia grossulariae (Schum.) Lagh., rust. Mass., Conn., N. Y., Mich., Ind., Minn., N. Dak., Iowa.

Sphaerotheca mors-uvae (Schw.) Berk. and Curt., powdery mildew. Widespread in Northern States.

**R. grossulariae* and other species, gooseberry.

Armillaria mellea (Vahl) Quel., root-rot. Oreg.

Botrytis cinerea Auct., die-back. Wash., Oreg.

Cercospora sp., leaf-spot. N. Y., W. Va., Ind., Mich., Mo., Nebr.

Coleosporium ribicola (Cke. and Ell.) Arth., rust. Wyo., Colo., N. Mex.

Cronartium occidentale Hedgec., Bethel, and Hunt, piñon blister rust. Colo., Ariz., Calif.

C. ribicola Fisch., blister rust. New England, Minn.

Leptosphaeria coniothyrium (Fckl.) Sacc., cane-blight. Ind., Mo.

Mycosphaerella grossulariae (Fr.) Lindau, leaf-spot. General in Northern States.

Phyllactinia corylea (Pers.) Karst., powdery mildew. N. Y.

Phyllosticta grossulariae Sacc., leaf-spot. N. J., Wis., Wash.

Plasmopara ribicola Schroet., downy mildew. W. Va., Minn., Oreg.

Pseudopeziza ribis Kleb., anthracnose. General in Northern States.

Puccinia grossulariae (Schum.) Lagh., rust. New England to N. Dak., south to Kans. and Md.

Ramularia sp., leaf-spot. Mich.

Septoria grossulariae (Lib.) West., leaf-spot. Me., Nebr., Kans.

Sphaerotheca mors-uvae (Schw.) Berk. and Curt., powdery mildew. General.

RICINUS.

R. communis, castor-bean.

Bacterium solanacearum EFS. (*Phytophthora solanacearum* (EFS.) Com. S. A. B.), bacterial wilt. Ga., Fla., Ala.

Botrytis cinerea Auct., gray-mold. Fla.

Cercospora albido-maculans Wint., leaf-spot. Mo.

C. ricinella Sacc. and Berl., leaf-spot. Gulf States, Calif.

Corticium vagum Berk. and Curt., damping-off. Fla., Kans.

Heterosporium sp. Calif.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Sclerotinia ricini G. H. Godfrey, gray-mold. Fla., Miss., La., Tex.

ROBINIA.

R. pseudacacia, black locust.

Cylindrosporium solitarium Heald and Wolf, leaf-spot. Tex.

Fomes rimosus Berk., yellow wood-rot. Widespread east of Great Plains.

F. robiniae Murr., spongy heart-rot. Miss. Valley and Eastern States.

Gloeosporium revolutum Ell. and Ev., leaf-spot. N. J.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Phoradendron flavesceus macrophyllum Engelm., mistletoe. Ariz., N. Mex.

Polyporus obtusus Berk., soft heartwood rot.

P. sulphureus (Bull.) Fr., brown-checked wood-rot. Mass.

Septoria curvata (Rabh. and A. Braun) Sacc., leaf-spot. Ohio.

Trametes robinophila Murr., white heart-rot. Mich.

ROSA.

Rosa spp., rose (cultivated and wild species).

Armillaria mellea (Vahl) Quel., root-rot. Oreg.

Bacterium tumefaciens EFS. and Town. (*Phytophthora tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Widespread.

Botryosphaeria ribis Gross. and Dug., cane-blight. Md., Va.

Botrytis cinerea Auct., bud-blight. General.

Cercospora rosicola Pass., leaf-spot. General east of Great Plains; Wash.

C. rosigena Tharp, leaf-spot. Tex.

ROSA—Continued.

Rosa spp., rose (cultivated and wild species)—Continued.

Cladosporium fuscum Lk., leaf-spot. Miss., Minn.

Coniothyrium rosarum Cke. and Hark., graft canker. N. J., Iowa.

Coryneum sp., stem canker. Wash.

C. microstictum Berk. and Br., canker. Pa.

Cylindrocladium scoparium Morgan, crown canker. Mass., N. Y., Pa., Ga., Ind., Mich., Mo.

Diaporthe umbrina A. Jenkins, canker. Atlantic coast, W. Va., Ky., Miss.

Dicoccum rosae Bon., blight, leaf-spot. Fla., Ind.

Diplocarpon rosae Wolf, black-spot. General.

Diplodia sp., twig-blight. Fla.

Discosia artocreas (Tode) Fr., blossom blight. La.

Dothiorella sp., blossom blight. La.

Gloeosporium rosae Hals., twig-blight. Widespread east of Miss. River; Tex., Okla.

Heterodera radicola (Greef) Muell., root-knot. N. J., W. Va., Ala., Ohio, Mich., Nebr.

Leptosphaeria coniothyrium (Fckl.) Sacc., cane-blight. General.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Peronospora sparsa Berk., downy mildew. N. Y., Pa., Ohio, Tex., Calif.

Phomopsis sp., canker. Ky.

Phragmidium americanum (Pk.) Diet., rust. Widespread in Northeastern States; upper Miss. Valley.

P. disciflorum (Tode) G. F. James, rust. General.

P. montivagum Arth., rust. Western States.

P. rosae-acicularis Liro, rust. Mich., Wis., Minn., N. Dak., Mont., Idaho, Alaska.

P. rosae-arkansanae Diet., rust. Widespread.

P. rosae-californicae Diet., rust. Pacific coast.

P. rosae-setigeræ Diet., rust. Upper Miss. Valley.

P. speciosum Cke., rust. General.

P. subcorticinum (Schränk) Wint., rust. Widespread.

Phyllosticta rosae Desm., leaf-spot. Ind., Fla.

Physalospora malorum (Pk.) Shear (*P. cydoniae* Am. Auct.), canker. Tex.

Sclerotinia sp., blight. Miss.

Septoria rosae Desm., leaf-spot. Miss.

Sphaerotheca humuli (DC.) Burr., powdery mildew. Conn., Pa., Tex., Ohio, Mich., Wis., Nebr., Calif.

S. pannosa (Wallr.) Lév., powdery mildew. General.

Uromyces rosicola Ell. and Ev., rust. Nebr.

RUBUS.

Rubus sp., blackberry.

Armillaria mellea (Vahl) Quel., root-rot. Wash.

Bacterium tumefaciens EFS. and Town. (*Phytoplasma tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. General.

Botrytis cinerea Auct., gray-mold. General on harvested fruit.

Cercospora rubi Sacc., blotch. Fla., La., Ill.

Corticium vagum Berk. and Curt., crown-rot. Colo.

Cylindrosporium rubi Ell. and Morgan, leaf-spot. N. C.

Fusarium sp., rosette. Md., Gulf States.

Fusisporium rubi Wint., double-blossom. N. Y., N. J., Del., Md., Va., Tenn., south to Fla., Ala., La., Ill.

Gloeodes pomigena (Schw.) A. S. Colby (*Phyllachora pomigena* (Schw.) Sacc.), sooty blotch. Ind.

Gymnoconia interstitialis (Schl.) Lagh., orange rust. General, especially in Eastern States.

Kuehneola uredinis (Lk.) Arth., yellow rust. General east of Miss. River.

Leptosphaeria coniothyrium (Fckl.) Sacc., cane-blight. East of Great Plains except Southern States; Pacific Northwest.

Leptothyrium pomi (Mont. and Fr.) Sacc., flyspeck. Ind.

Mycosphaerella rubi E. W. Roark, leaf-spot. General.

M. rubina (Pk.) Jacz., spur-blight. N. Y., Wis., Minn.

Peronospora rubi Rabh., downy mildew. Wis.

Pezizella lythri (Desm.) Shear and B. O. Dodge, fruit-rot.

RUBUS—Continued.

Rubus sp., blackberry—Continued.

Phragmidium imitans Arth., leaf-rust. Wash.
Plectodiscella veneta (Speg.) Burk., anthracnose. General.
Sphaerotheca humuli (DC.) Burr., powdery mildew. Ohio, Ill., Minn.
Verticillium alboatrum Reinke and Berth., bluestem, wilt. N. Y., Minn., Wash.

Dwarf, undet. Wash., Oreg.

Mosaic, undet. N. Y., Ind., Iowa, Oreg., Calif.

Rubus sp., dewberry (cultivated and wild).

Cercospora rubi Sacc., blotch. Fla., La., Tex., Ark.
Fusisporium rubi Wint., double-blossom. N. Y., N. J., Del., Md., Miss., La., Tex., Ohio.
Gymnoconia interstitialis (Schl.) Lagh. (not on *Lucretia* dewberry), orange rust. General, especially in Eastern States.
Kuehneola uredinis (Lk.) Arth., yellow rust. General east of Miss. River.
Leptosphaeria coniothyrium (Fekl.) Sacc., cane-blight. N. C., Miss., Idaho.
Mycosphaerella rubi E. W. Roark, leaf-spot. General.
Peronospora potentillae D By., downy mildew. Conn.
P. rubi Rabb., downy mildew. Fla., Wash.
Peizella lythri (Desm.) Shear and B. O. Dodge, fruit-rot.
Phyllosticta ruborum Sacc., leaf-spot. N. Y., Miss.
Plectodiscella veneta (Speg.) Burk., anthracnose. General.
Pucciniastrum americanum (Farl.) Arth., leaf-rust. Wis. (on **R. arcticus triflorus*).
Sphaerotheca humuli (DC.) Burr., powdery mildew. Pa.

Rubus sp., Logan blackberry.

Armillaria mellea (Vahl) Quel., root-rot. Oreg.
Bacterium tumefaciens EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Wash., Oreg.
Corticium vagum Berk. and Curt., crown-rot. Wash.
Mycosphaerella rubi E. W. Roark, leaf-spot. Mich., Wash., Oreg.
M. rubina (Pk.) Jacz., spur-blight. Wash.
Plectodiscella veneta (Speg.) Burk., anthracnose. N. Y., Wash., Oreg.
Dwarf, undet. Wash., Oreg.
Mosaic, undet. Wash., Oreg., Calif.

Rubus sp., raspberry.

Armillaria mellea (Vahl) Quel., root-rot. Wash., Oreg.
Ascospora ruborum (Oud.) S. M. Zeller, leaf-spot. Mass., Oreg.
Bacterium tumefaciens EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. General.
Botrytis cinerea Auct., gray-mold rot. General on harvested fruit.
Cercospora rubi Sacc., blotch. Fla., Ala.
Corticium vagum Berk. and Curt., crown-rot. Idaho.
Cuscuta sp., dodder. Wis.
Cylindrosporium rubi Ell. and Morgan, leaf-spot. Tex., Ohio, Wis., Mo.
Fusisporium rubi Wint., double-blossom. N. Y., Pa., Md.
Gymnoconia interstitialis (Schl.) Lagh., orange rust. General, especially in Eastern States.
Leptosphaeria coniothyrium (Fekl.) Sacc., cane-blight. East of Great Plains except Southern States; Pacific Northwest.
Leptothyrium pomi (Mont. and Fr.) Sacc., flyspeck. Mass., Ind.
Mycosphaerella rubi E. W. Roark, leaf-spot. General.
M. rubina (Pk.) Jacz., spur-blight. Widespread in Northern States.
Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.
Peronospora rubi Rabb., downy mildew. Wash.
Phragmidium imitans Arth., leaf-rust. Widespread in Northern States.
P. occidentale Arth., rust. Colo.
Phyllosticta bicolor Pk., leaf-spot. N. Y.
Plectodiscella veneta (Speg.) Burk., anthracnose. General.
Pucciniastrum americanum (Farl.) Arth., rust. Conn., Mass., Vt., Md., Va., Wis., Iowa, N. Dak.
Rhizopus nigricans Ehr., black-mold rot. General.

RUBUS—Continued.*Rubus* sp., raspberry—Continued.*Sphaerotheca humuli* (DC.) Wint., powdery mildew. Conn., N. Y., Ohio, Ill., Minn., Utah, Wash.*Verticillium alboatrum* Reinke and Berth., bluestem, wilt. N. Y., Ohio, Mich., Minn., Wash., Oreg., Calif.

Leaf-curl, undet. N. Y., Md., Pa., Ohio, Ind., Mich., Wis., Minn., Wash., Oreg., Calif.

Mosaic, undet. Northeastern States; Minn., Pacific Northwest.

Streak, undet. Conn., N. Y., Pa., Md., Ohio, Ind., Ill., Wis., Oreg.

RUDBECKIA.*R. laciniata*, *goldenglow, coneflower.*Cercospora tabacina* Ell. and Ev., leaf-spot. Wis.*Entyloma compositarum* Farl., smut. Ohio, Iowa, Mo.*Erysiphe cichoracearum* DC., powdery mildew. N. H., Conn., Md., W. Va., Ohio, Ind., Mich., Minn.*Phyllosticta rudbeckiae* Ell. and Ev., leaf-spot. Wis., Iowa.*Plasmopara halstedii* Farl., downy mildew. N. Dak., Iowa.*Ramularia rudbeckiae* Pk., leaf-spot. N. Y., Pa., Mich., Iowa, Mont., Colo.*Septoria rudbeckiae* Ell. and Hals., leaf-spot. Nebr., Kans.*Uromyces perigynius* Hals., rust. Pa., Wis., Iowa, Mo., N. Dak., S. Dak., Nebr., Mont.*U. rudbeckiae* Arth. and Holw., rust. Ind., Ill., Miss. River to Rocky Mts.; N. Mex.**SACCHARUM.***S. officinarum*, sugarcane.*Cercospora vaginiae* W. Krueger, red-spot. Fla.*Colletotrichum falcatum* F. A. Went, red-rot. Gulf States; Ark.*Gnomonia iliaui* H. L. Lyon, stem-rot, Iliau disease. La.*Helminthosporium sacchari* Butl., eye leaf-spot. Ga., Fla., Miss.*Heterodera radiculicola* (Greef) Muell., root-knot. Ga., Fla.*Leptosphaeria sacchari* Van Breda de Haan, ring-spot. Fla., Ala.*Marasmius sacchari* J. H. Wakker, root-rot. Gulf States.*M. stenophyllus* Mont., root-rot. La., Tex.*Melanconium sacchari* Mass., rind disease. Fla., La.*Phyllosticta* sp., leaf-spot. Ga.*Sclerotium rolfsii* Sacc., sheath-rot. Ga., Fla., Ala., La.*Thielaviopsis paradoxa* (DeSeyn.) Hoehn., pineapple disease. Miss., La.

Mosaic, undet. Gulf States; Ga., Ark.

SALIX.*Salix* spp., willow.*Bacterium tumefaciens* EFS. and Town. (*Phytoplasma tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Conn., Tex.*Cercospora salicina* Ell. and Ev., leaf-spot. Ill., La., Tex.*Coryneum salicinum* (Cda.) Sacc., leaf-spot. N. Dak., Alaska.*Cytospora chrysosperma* (Pers.) Fr., canker. Western States; Ind., Minn.*Gloeosporium maculans* Pk., leaf-spot. Idaho.*G. salicis* West., leaf-spot. N. Y., N. J., Wis., Oreg.*Marssonina* sp., twig-blight. Wash.*Marssonina populi* (Lib.) Magn., leaf-spot. Wis.*Melampsora americana* Arth., rust. General.*M. arctica* Rostr., rust. Colo.*M. bigelowii* Thuem., leaf-rust. General.*M. confluens* (Pers.) Jack., rust. General, especially in Western States.*Nectria galligena* Bres., European canker. Oreg.*Phoradendron flavescens* (Pursh) Nutt., mistletoe. Ind., Calif.*Phyllosticta apicalis* Davis, leaf-spot. Wis.*Ramularia rosea* (Fekl.) Sacc., leaf-spot. Wis., Colo.*Rhytisma salicinum* Fr., tar-spot. Widespread.*Septoria albaniensis* Thuem., leaf-spot. Mass., N. Y., Wis., Minn.*S. didyma* Fekl., leaf-spot. Wis.*S. salicina* Pk., leaf-spot. N. Y., Miss., Kans., Colo.*Trametes suaveolens* Fr., white wood-rot. N. Y., N. J., Wis., Minn., Mo.*Uncinula salicis* (DC.) Wint., powdery mildew. General.

SAMBUCUS.

Sambucus spp., elderberry.

- Ascochyta wisconsinia* Davis, leaf-spot. Wis.
Cercospora catenosporea Atk., leaf-spot. Ala., Miss., Tex., Kans.
C. depazeoides (Desm.) Sacc., leaf-spot. N. Y., Ala., Ga., Ind., Iowa, Kans.
C. lateritia Ell. and Hals., leaf-spot. Iowa, Kans.
C. sambucina Ell. and Kell., leaf-spot. Wash.
Cercosporella prolificans (Ell. and Holw.) Sacc., leaf-spot. Calif.
Cytospora chrysosperma (Pers.) Fr., canker. Kans., Wash.
Gloeosporium tineum Sacc., leaf-spot. Miss.
Microsphaera alni (Wallr.) Wint., powdery mildew. N. Y., Pa., Ohio, Ill.
M. grossulariae (Wallr.) Lév., powdery mildew. Widespread east of Great Plains.
Nectria cinnabarina Fr., canker. Ind., N. Dak., Mont., Wash.
Phyllosticta sp., leaf-spot. Pa.
Puccinia sambuci (Schw.) Arth., rust. East of Great Plains.
Ramularia glauca Ell. and Ev., leaf-spot. Colo.
R. sambucina Pk., leaf-spot. N. Y., Mo., Wash.
Septoria sambucina Pk., leaf-spot. Widespread.
Sphaeropsis sambucina (Cke.) Sacc., canker. N. Y., Ind., N. Dak., Kans., Colo., Calif.

SAPINDUS.

Sapindus sp., soapberry.

- Cylindrosporium griseum* Heald and Wolf, leaf-blight. Tex.
Glomerella cingulata (Ston.) Spauld. and Schrenk, bitter-rot. Fla.
Phoradendron flavescens (Pursh) Nutt., mistletoe. Tex.
Uncinula circinata Cke. and Pk., powdery mildew. Tex.

SASSAFRAS.

S. variifolium, sassafras.

- Actinothyrium gloeosporioides* L. R. Tehon, leaf-spot. Ill.
Armillaria mellea (Vahl) Quel., root-rot. Pa.
Fomes ribis (Schum.) Fr., root-rot. Mo.
Gloeosporium affine Ell. and Kell., leaf-spot. N. J., S. C., Ohio, Ind., Ill.
G. officinale Ell. and Ev., leaf-spot. Del.
Phoradendron flavescens (Pursh) Nutt., mistletoe. Tex.
Phyllosticta sassafras Cke., leaf-spot. Widespread in Eastern States.

SCHINUS.

S. molle, California peppertree.

- Armillaria mellea* (Vahl) Quel., root-rot. Calif.
Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.
Trametes schini J. G. Brown, timber-rot. Ariz.

SECALE.

S. cereale, rye.

- Bacterium translucens secalis* C. S. Reddy, James Godkin, and A. G. Johnson, bacterial blight. Ind., Ill., Wis.
B. translucens undulosum EFS., L. R. Jones, and C. S. Reddy, black-chaff. Wis.
Claviceps purpurea (Fr.) Tul., ergot. General.
Colletotrichum graminicolum (Ces.) Wils., anthracnose. General east of Great Plains.
Erysiphe graminis DC., powdery mildew. General.
Fusarium culmorum leleuxii Sherb., seedling blight. Oreg.
Gibberella saubinetii (Mont.) Sacc., scab. Occurs scatteringly east of Great Plains.
Helminthosporium sativum Pam., King, and Bak., leaf-spot, foot-rot. Pa., N. Y., Minn.; probably other States.
Marasmius tritici P. A. Young. Ill.
Ophiobolus graminis Sacc., take-all. N. Y.
Puccinia dispersa Eriks., leaf-rust. General, especially east of Rocky Mts.
P. glumarum (Schm.) Eriks. and Henn., stripe-rust. Wash., Oreg.
P. graminis Pers., stem-rust. General.

SECALE—Continued.*S. cereale*, rye—Continued.

Rhynchosporium secalis (Oud.) Davis, scald. Upper Miss. Valley; Wash., Calif.

Septoria secalis Prill. and Del., leaf-blotch. East of Great Plains.

Tilletia tritici (Bjerk.) Wint., covered smut. N. Y., Wash.

Tylenchus tritici (Stein.) Bast., nematode. Va., S. C., Kans.

Urocystis occulta (Wallr.) Rabh., stem-smut. General east of Great Plains, occasional in West.

Ustilago tritici (Pers.) Rostr., loose-smut. Infrequent in Eastern States.

Mosaic, undet. Va., Ill.

SEQUOIA.*S. gigantea*, big tree.

Cercospora sequoiae Ell. and Ev., leaf-spot. Pa.

S. sempervirens, redwood.

Botrytis cinerea Auct., gray-mold.

Hexagonia carbonaria Berk. and Curt., brown-rot. Calif.

Polyporus amorphus Fr., rot. Calif.

Trameles isabellina Fr., white-rot. Calif.

T. odorata (Wulf.) Fr., brown-rot. Calif.

SHEPHERDIA.*S. argentea*, silver buffaloberry.

Phyllactinia corylea (Pers.) Karst., powdery mildew. Oreg.

Puccinia coronata Cda., crown-rust. N. Dak., Nebr., Mont., Wyo., Colo.

Sphaerotheca humuli (DC.) Burr., powdery mildew. Colo., Mont.

S. canadensis, russet buffaloberry.

Puccinia caricis-shepherdiae Davis, rust. N. Y., Mich., Wis., S. Dak.

P. coronata Cda., crown-rust. Occurs scatteringly in Rocky Mt. States and Pacific Coast States.

SOJA.*S. max*, soybean.

Bacillus lathyri Manns and Taub., streak. Del., Nebr.

Bacterium glycineum Coerper, bacterial blight. Pa., Ind., Ill., Wis., Miss., La.

B. phaseoli sojense Hedges, bacterial pustule. Va., N. C., S. C., La., Tex., Ark., Kans.

B. sojae Wolf, bacterial blight. Conn., N. C., S. C., La., Mich., Wis., Nebr.

B. solanacearum EFS. (*Phytophthora solanacearum* (EFS.) Com. S. A. B.), bacterial wilt. N. C.

Botrytis cinerea Auct., gray-mold. Ohio.

Cercospora cruenta Sacc., leaf-spot. Ala.

Corticium vagum Berk. and Curt., stem-rot. Tenn., Ind.

Diaporthe sojae S. G. Lehman, pod-blight. N. C., Ind.

Erysiphe polygoni DC., powdery mildew. N. C.

Fusarium vasinfectum tracheiphilum EFS., wilt. Southern States; Ind.

Glomerella cingulata (Ston.) Spauld. and Schrenk, anthracnose. N. C.

Heterodera radiculicola (Greef) Muell., root-knot. S. C., Ark., Calif.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Peronospora sojae Wolf, downy mildew. Del., Ky., N. C.

P. trifoliorum D By., downy mildew. N. Y.

Sclerotinia sclerotiorum (Lib.) Mass., stem-rot. Va.

Sclerotium rolfsii Sacc., stem-rot. N. C., S. C., Ala., Miss., La.

Septoria glycines Takewo Hemmi, leaf-spot. Del.

Mosaic, undet. Widespread where crop is grown.

SOLANUM.*S. melongena*, eggplant.

Alternaria solani (Ell. and Mart.) L. R. Jones and A. J. Grout, leaf-spot. Conn., N. Y., Pa., Del., Md., W. Va., Fla.

Ascochyta lycopersici Brun. (*Didymella lycopersici* Kleb., *Diplodina destructiva* (Plow.) Petr.), leaf-spot. N. Y., Del., Ind.

SOLANUM—Continued.*S. melongena*, eggplant—Continued.

Bacterium solanacearum EFS. (*Phytomonas solanaceara* (EFS.) Com. S. A. B.), bacterial wilt. Southern States.

Botrytis cinerea Auct., gray-mold rot. Mass., Conn., Wash., Calif.

Colletotrichum atramentarium (Berk. and Br.) Taub., fruit-rot, anthracnose.

General east of Great Plains; Utah.

Corticium vagum Berk. and Curt., stem-rot. Pa., N. J., Fla., Ala., La., Tex., Wash.

Diplodia sp., fruit-spot. Fla.

Fusarium spp., wilt. East of Miss. River; La., N. Mex.

Gloeosporium melongenae Ell. and Hals., anthracnose. General east of Great Plains.

Heterodera radicola (Greef) Muell., root-knot. Southern States; Nev.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Phoma sp., storage rot. Me.

Phomopsis vexans (Sacc. and Syd.) Harter, leaf-spot, fruit-rot. East of Great Plains; Calif.

Phyllosticta solani Ell. and Mart., leaf-spot. La.

Phytophthora sp., rot. Ind.

P. infestans (Mont.) D By., late-blight. N. Y.

Pythium debaryanum Hesse, damping-off. Conn.

Sclerotinia sclerotiorum (Lib.) Mass., stem-rot. Va., La.

Sclerotium rolfsii Sacc., stem-rot. Ga., La.

Verticillium alboatrum Reinke and Berth., wilt. Mass., N. Y., N. J., Pa., Va., S. C., Ga.

Mosaic, undet. Conn., Fla., Iowa.

S. tuberosum, potato.

Actinomyces scabies (Thax.) Gues., scab. General.

Alternaria solani (Ell. and Mart.) L. R. Jones and A. J. Grout, early-blight. General, especially east of Rocky Mts.

Armillaria mellea (Vahl) Quel., root-rot. Mich., Wis., Wyo., Pacific coast.

Bacillus spp., soft-rot. General, especially on harvested crop in transit and market.

B. phytophthorus O. Appel (*Erwinia atroseptica* (Van Hall) Com. S. A. B.), black-leg. General; reported from practically every State.

Bacterium solanacearum EFS. (*Phytomonas solanaceara* (EFS.) Com. S. A. B.), bacterial wilt. Southern States.

Cercospora concors (Casp.) Sacc., leaf-blotch. Vt., N. Y., W. Va., Ga., Ind., Mich.

C. solanicola Atk., leaf-spot. Ala.

Colletotrichum atramentarium (Berk. and Br.) Taub., anthracnose. N. Y., W. Va., Ohio, Fla., Utah, Oreg.

Corticium vagum Berk. and Curt., scurf, stem-rot. General; in every State.

Fusarium spp., dry rots. Common everywhere on harvested tubers.

F. eumartii C. W. Carpenter, stem-end rot. N. Y., Pa., Ohio, Nebr.

F. oxysporum Schl., wilt. Widespread; less destructive in more northern late-potato sections.

F. radicola Woll., jelly end-rot. N. Y., Va., D. C., Miss.; not uncommon in Western States.

F. sulphureum Schl., tuber rot. N. Dak.

F. trichothecioides Woll., powdery dry-rot. Ohio, Wis., Minn., Iowa, S. Dak., Nebr., Ariz., Utah, Idaho, Wash., Oreg.

Heterodera radicola (Greef) Muell., root-knot. Southern States, Kans., Nev., Pacific coast.

Oidium sp., powdery mildew. Pa.

Phoma sp. (doubtfully parasitic), stem-blight. New England, N. Y., Pa., Mich.

P. tuberosa I. E. Melhus, J. Rosenbaum, and E. S. Schultz, storage rot. Me., Pa.

Phytophthora infestans (Mont.) D By., late-blight. East of Great Plains; Pacific coast. Most frequent in Northeast.

Pythium debaryanum Hesse, leak. Conn., N. Y., N. J., N. C., Miss., La., Minn., Nebr., Mont., Wyo., Colo., Idaho, Wash., Calif.

SOLANUM—Continued.*S. tuberosum*, potato—Continued.*Rhizoctonia crocorum* (Pers.) DC., violet root-rot. N. Y., Nebr., Wash., Oreg.*Rhizopus nigricans* Ehr., leak. Ohio, Idaho, Wash., Calif.*Sclerotinia sclerotiorum* (Lib.) Mass., stem-rot. N. Y., Fla.*Sclerotium rolfsii* Sacc., stem-rot. South Atlantic and Gulf Coast States.*Spondylocidium atrovirens* C. O. Harz, silver scurf. General.*Spongospora subterranea* (Wallr.) T. Johnson, powdery scab. Me., Conn., N. Y., Pa., S. C., Fla., Minn., Wyo., Wash., Oreg.*Synchytrium endobioticum* (Schilb.) Perc., wart. Pa., Md., W. Va.*Verticillium alboatrum* Reinke and Berth., wilt. New England, N. Y., Pa., Ohio, Wis., N. Dak., Colo., Utah, Idaho, Pacific coast.

Black-heart, nonpar. Frequent, especially as a result of overheating in transit.

Curly-dwarf, undet. Frequent.

Hollow-heart, nonpar. Frequently reported, especially in overgrown tubers.

Hopperburn, undet. East of Great Plains.

Internal brown-spot, nonpar. Frequent.

Leaf-roll, undet. General.

Leak, various causes. General (see also *Pythium debaryanum* and *Rhizopus nigricans*).

Malnutrition, nonpar. New England, Middle Atlantic States, Minn., Wash.

Mosaic, undet. General.

Net-necrosis, undet. Frequent.

Russet dwarf, undet. Idaho.

Spindle-tuber, undet. Vt., N. Y., N. J., S. Dak., Nebr., Ky.

Streak, undet. Atlantic Coast States, Mich., Minn., Idaho, Wash., Calif.

Tipburn, nonpar. General.

Witches'-broom, undet. Me., Mich., Minn.; Pacific Northwest.

Yellow dwarf, undet. Vt., N. Y., N. J., Pa., Fla., S. Dak.

SOPHORA.*S. japonica*, *pagoda tree.*Microsphaera alni* (Wallr.) Wint., powdery mildew. Conn.*Ozonium omivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.**SORBUS.***Sorbus* sp., mountain-ash.*Bacillus amylovorus* (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. Mich., Minn.*Cytospora chrysosperma* (Pers.) Fr., canker. Mont.*Gymnosporangium blasdaleanum* (Diet. and Holw.) Kern, rust. Oreg.*G. cornutum* (Pers.) Arth., rust. New England, N. Y., Mich., Wis., Mont., Wyo., Colo.*G. globosum* Farl., rust. New England, N. Y.*G. juniperinum* (L.) C. Martius, rust. Mont., Wyo., Colo., Wash.*G. nootkatensis* (Trel.) Arth., rust. Wash., Alaska.*Nummularia discreta* (Schw.) Tul., blister canker. Mass., Pa.*Phyllosticta sorbi* West., leaf-spot. Mo.*Physalospora malorum* (Pk.) Shear (*P. cydoniae* Am. Auct.), black-rot.*Polystictus hirsutus* Fr., white sap-rot.*Valsa leucostoma* (Pers.) Fr., die-back. Mont.*Venturia inaequalis* (Cke.) Aderh., scab. Minn.**SPINACIA.***S. oleracea*, spinach.*Cercospora beticola* Sacc., leaf-spot. Mass., Tex.*C. dubia* (H. Riess) Wint., leaf-spot. Mo.*C. flagelliformis* Ell. and Hals., leaf-spot. N. J., Del., Ga., Ind., Minn.*Cladosporium macrocarpum* Preuss, scab, black-mold. Pa., Del., Tex.*Colletotrichum spinaciae* Ell. and Hals., anthracnose. Conn., N. Y., N. J., Va., La., Tex.*Corticium vagum* Berk. and Curt., stem-rot. Pa., Ill., Wash.*Entyloma ellisii* Hals., smut. N. J.

SPINACIA—Continued.*S. oleracea*, spinach—Continued.*Fusarium* sp., root-rot. Tex., Oreg.*F. spinaciae* Sherb., wilt. Idaho.*Heterodera radicola* (Greef) Muell., root-knot. Ind.*Heterosporium variabile* Cke., leaf-spot. Conn., Mass., Va., Mo., Tex., Calif.*Macrosporium cheiranthi* (Lib.) Fr., leaf-spot. Ohio.*Ozonium omnivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.*Peronospora effusa* (Grev.) Ces., downy mildew. General.*Phyllosticta chenopodii* Sacc., leaf-spot. N. J., Del., Va.*Phytophthora* sp., crown-rot. N. J., Ind.*Puccinia sarcobati* (Pk.) Bethel, rust. Ariz., Wash., Oreg.*Pythium debaryanum* Hesse, damping-off. Conn.

Malnutrition, nonpar. Va.

Mosaic, undet. Widespread.

SPIRAEA.*Spiraea* sp., spirea.*Bacillus amylovorus* (Burr.) Trev. (*Erwinia amylovora* (Burr.) Com. S. A. B.), blight. Md.*Cercospora rubigo* Cke. and Hark., leaf-spot. Kans.*Cylindrosporium filipendulae* Thuem., leaf-spot. Wis., Wash., Oreg.*Ozonium omnivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.*Podosphaera oxyacanthae* DC., powdery mildew. New England, Ohio, Mo., Wyo., Wash.*Rhopalidium cercosporelloides* Dearn., leaf-spot. Idaho.*Septoria salicifolia* (Trel.) Ell. and Ev., leaf-spot. Wis., Kans.*Sphaerotheca humuli* (DC.) Burr., powdery mildew. Conn., N. Y.**STAPHYLEA**.*S. trifolia*, American bladdernut.*Ramularia isarioides* Sacc., leaf-spot. Pa., Wis., Mo.*Septoria cirrhosa* Wint., leaf-spot. Mo.**STIZOLOBIUM**.*S. deeringianum*, velvetbean.*Aplanobacter stizobii* Wolf, bacterial leaf-spot. N. C.*Bacterium vignae* M. W. Gardner and J. B. Kendrick, bacterial spot. Ind.*Cercospora mucunae* Syd., leaf-spot. Ala.*C. stizobii* Syd., leaf-spot. N. C., S. C., Fla.*Corticium vagum* Berk. and Curt., stem-rot. Ga., La.*Heterodera radicola* (Greef) Muell., root-knot. Calif.*Phyllosticta mucunae* Ell. and Ev., leaf-spot. Ala.*Sclerotium rolfsii* Sacc., stem-rot. Ga., Ala.**SYMPHORICARPOS**.*Symphoricarpos* sp., snowberry, coralberry.*Cercospora symphoricarpi* Ell. and Ev., leaf-spot. Nebr., Kans., Mont.*Fomes ribis* (Schum.) Fr., root-rot. N. Dak., Kans.*Microsphaera diffusa* Cke. and Pk., powdery mildew. General.*Nectria coryli* Fckl., canker. N. Dak.*Puccinia abundans* (Pk.) Jack., rust. Great Plains and westward.*P. symphoricarpi* Hark., rust. Western States.*Septoria oedospora* F. E. Clements, leaf-spot. Colo.*S. symphoricarpi* Ell. and Ev., leaf-spot. N. Dak., Nebr., Mont., Idaho, Wash.**SYRINGA**.*Syringa* sp., lilac.*Hymenochaete agglutinans* Ell., stem girdling. Conn.*S. persica*, Persian lilac.*Cercospora lilacis* (Desm.) Sacc., leaf-spot. Miss.*Microsphaera alni* (Wallr.) Wint., powdery mildew. Iowa.

SYRINGA—Continued.*S. vulgaris*, lilac.*Botrytis cinerea* Auct., blossom blight. Wash.*Cercospora lilacis* (Desm.) Sacc., leaf-spot. S. C., Miss., Minn.*C. macromaculans* Heald and Wolf, leaf-spot. Tex., Okla.*Corticium stevensii* Burt (*Hypochnus ochroleucus* F. Noack), leaf-blight. N. C., Miss.*Microsphaera alni* (Wallr.) Wint., powdery mildew. General.*Phyllosticta halstedii* Ell. and Ev., leaf-spot. N. J.*P. syringae* West., leaf-spot. Wis.*Polystictus versicolor* (L.) Fr., yellowish wood-rot. Mo.*Sphaeropsis* sp. Pa.*Uncinula clintonii* Pk., powdery mildew. Ind.**TAGETES.***Tagetes* sp., marigold.*Botrytis cinerea* Auct., rot. Conn.**TAXODIUM.***T. distichum*, baldcypress.*Fomes geotropus* Cke., pecky heartwood rot. Southern States.*Pestalozzia funerea* Desm., leaf-blight. Tex.**THUJA.***T. occidentalis*, arborvitae.*Keithia thujina* E. J. Durand, leaf-blight. Va., Wis.*Lophodermium pinastri* (Schröd.) Chev., leaf-cast. Me.*Pestalozzia funerea* Desm., leaf-blight. Me., Mass., Va., N. C.*Phomopsis juniperovora* G. Hahn, nursery blight. Pa.*T. orientalis*, oriental arborvitae.*Diplodia* sp., die-back. Fla.*Phomopsis juniperovora* G. Hahn, nursery blight. Pa., Ind., Ill., Iowa, Nebr., Kans.*T. plicata*, giant arborvitae.*Armillaria mellea* (Vahl) Quel., root-rot.*Coryneum thujinum* Dearn., leaf-blight. Oreg.*Herpotrichia nigra* Hartig, matting. Idaho.*Keithia thujina* E. J. Durand, leaf-blight. Pacific Northwest.*Mycosphaerella thujae* Petr., leaf-spot. Idaho.*Polyporus schweinitzii* Fr., red-brown butt-rot. Idaho.*Poria weirii* Murr., brown laminated rot. Throughout range of host.*Trametes pini* (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot.**TILIA.***Tilia* sp., linden, basswood.*Actinonema tiliae* Allesch., leaf-spot. Ind.*Cercospora microsora* Sacc., leaf-blight. Widespread east of Great Plains.*C. tiliae* Pk., leaf-spot. N. Y.*Gloeosporium tiliae* Oud., anthracnose. N. Y., Mass., Va., Minn.*Hydnum septentrionale* Fr., white sapwood rot. Ala.*Melanconium tiliae* Pk., twig-blight. Ohio.*Phoradendron flavescens* (Pursh) Nutt., mistletoe.*Phyllosticta tiliae* Sacc. and Speg., leaf-spot. N. Y., Pa., W. Va.*Uncinula clintonii* Pk., powdery mildew. Widespread in Eastern States.**TRAGOPOGON.***T. porrifolius*, vegetable-oyster, salsify.*Albugo tragopogonis* (DC.) S. F. Gray, white-rust. General in northern half of United States.*Bacillus carotovorus* L. R. Jones (*Erwinia carotovora* (Jones) Com. S. A. B.), soft-rot. Conn.*Corticium vagum* Berk. and Curt., stem-rot. Wash.*Erysiphe cichoracearum* DC., powdery mildew. Nebr., Wash.*Heterodera radicola* (Greef) Muell., root-knot. Va., Ala., Wash.*Ozonium omnivorum* Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

TRAGOPOGON—Continued.

T. porrifolius, vegetable-oyster, salsify—Continued.

Sclerotinia intermedia G. B. Ramsey, root-rot. Ill.

Verticillium sp., wilt. N. Y.

TRIFOLIUM.

T. hybridum, alsike clover.

Bacillus lathyri Manns and Taub., streak. Del.

Bacterium trifoliorum L. R. Jones et al., bacterial leaf-spot. General in Eastern States.

Cercospora zebrina Pass., leaf-spot. Va., Ind.

Colletotrichum trifolii S. M. Bain, anthracnose. East of Great Plains; Colo., Wash.

Corticium vagum Berk. and Curt., root-rot. Ohio, Minn.

Erysiphe polygoni DC., powdery mildew. W. Va., Minn., Idaho.

Phyllachora trifolii (Pers.) Fekl., sooty spot. Minn.

Sclerotinia trifoliorum Eriks., root-rot. Mich.

Sphaerulina trifolii Rostr., leaf-spot. N. Y., Ill., Mo.

Thielavia basicola (Berk. and Br.) Zopf, black root-rot.

Uromyces trifolii (Hedw. f.) Lév., rust. General.

Mosaic, undet. Pa.

T. incarnatum, crimson clover.

Bacterium trifoliorum L. R. Jones et al., leaf-spot. Md., Va., N. C., Ind., Wis., Iowa.

Botrytis sp., gray-mold. Del.

Cercospora medicaginis Ell. and Ev., leaf-spot. Del.

Colletotrichum trifolii S. M. Bain., anthracnose. Ky.

Corticium vagum Berk. and Curt., root-rot. N. C.

Gloeosporium caulivorum L. Kirchner, anthracnose. N. C.

Heterodera radicola (Greef) Muell., root-knot.

Macrosporium sp., leaf-spot. Va.

Phyllachora trifolii (Pers.) Fekl., sooty spot. Pa., Md., Ga., Ala., N. Dak.

Sclerotinia trifoliorum Eriks., root-rot. N. J., Md., N. C., S. C., Ky., Tenn., Oreg.

Sclerotium rolfsii Sacc., stem-rot. La.

Sphaerulina trifolii Rostr., leaf-spot. N. Y., Ill., Mo.

Thielavia basicola (Berk. and Br.) Zopf, black root-rot.

Uromyces fallens (Desm.) Kern, rust. Del., Ind., S. Dak.

U. trifolii (Hedw. f.) Lév., rust. Pa., Ala., Wis., Minn.

Mosaic, undet. Pa.

**T. medium*, zigzag clover.

Bacterium trifoliorum L. R. Jones et al., leaf-spot. Md., Va., N. C., Ind., Wis., Iowa.

Uromyces trifolii (Hedw. f.) Lév., rust.

T. pratense, red clover (includes **T. medium perenne*, mammoth clover).

Bacillus lathyri Manns and Taub., streak. Del.

Bacterium trifoliorum L. R. Jones et al., leaf-spot. D. C., Md., Va., N. C., Ind., Wis., Iowa.

Botrytis sp., gray-mold. Pa.

Cercospora medicaginis Ell. and Ev., leaf-spot. N. J.

C. zebrina Pass., leaf-spot. N. Y., Mich., Ind., Iowa.

Colletotrichum destructivum P. J. O'Gara, anthracnose. Utah.

C. trifolii S. M. Bain, anthracnose. N. Y., Pa., N. J., Del., Va., Ohio, Ind., Mich., Wis., Ky., Tenn.

Cuscuta spp., dodder. N. Y., Del., Md., La., Ohio, Nebr.

Erysiphe polygoni DC., powdery mildew. Widespread, especially abundant east of Great Plains.

Gloeosporium caulivorum L. Kirchner, anthracnose. N. Y. to Va., westward to Mo.

Heterodera radicola (Greef) Muell., root-knot. W. Va., La., Utah.

Macrosporium sarcinaeforme Cav., leaf-spot. East of Great Plains.

Phyllachora trifolii (Pers.) Fekl., sooty spot. East of Great Plains; Wash., Oreg.

Pseudopeziza trifolii (Biv.) Fekl., leaf-spot. Widespread in Northern States.

TRIFOLIUM—Continued.

T. pratense, red clover, etc.—Continued.

Sclerotinia trifoliorum Eriks., root-rot. Pa., N. C., Ky., Tenn., La., Ind., Mich., Oreg.

Septoria trifolii Ell., leaf-spot. N. J.

Sphaerulina trifolii Rostr., leaf-spot. N. Y., Ill., Mo.

Thielavia basicola (Berk. and Br.) Zopf, black root-rot.

Tylenchus dipsaci (Kuehn) Bast., stem nematode. Utah, Pacific Northwest.

Uromyces fallens (Desm.) Kern, rust. General.

Mosaic, undet. Conn., Pa., Ark., Ind., Minn.

T. repens, white clover.

Bacterium trifoliorum L. R. Jones et al., leaf-spot. D. C., Md., Va., N. C., Ind., Wis., Iowa.

Brachysporium trifolii C. H. Kauffman, blight. D. C.

Cercospora helvola Sacc., leaf-spot. N. J.

Corticium vagum Berk. and Curt., stem-rot. N. Y.

Erysiphe polygoni DC., powdery mildew. W. Va., Idaho.

Heterodera radicola (Greef) Muell., root-knot.

Peronospora trifoliorum D By., downy mildew. Nebr.

Phyllachora trifolii (Pers.) Fekl., sooty spot. East of Great Plains; Utah.

Pseudopeziza trifolii (Biv.) Fekl., leaf-spot. Miss.

Sclerotinia trifoliorum Eriks., root-rot. Miss., Oreg.

Sphaerulina trifolii Rostr., leaf-spot. N. Y., N. J., Ill., Mo.

Thielavia basicola (Berk. and Br.) Zopf, black root-rot. Conn.

Uromyces nerviphila (Grognot) Lagh., rust. N. Y., Wis., Colo.

U. trifolii (Hedw. f.) Lév., rust. General.

TRITICUM.

T. aestivum, wheat.

Bacterium atrofaciens McC. (*Phytomonas atrofaciens* (McC.) Com. S. A. B.), basal glume-rot. General in States between Miss. River and Rocky Mts.; Mich., N. Y., Pa.

B. translucens undulosum EFS., L. R. Jones, and C. S. Reddy, black-chaff. Miss. Valley; Great Plains area; Mont., Idaho.

Claviceps purpurea (Fr.) Tul., ergot. Occurs scatteringly, especially on durum wheat.

Colletotrichum graminicolum (Ces.) Wils., anthracnose. East of Great Plains.

Erysiphe graminis DC., powdery mildew. General.

Fusarium culmorum letius Sherb., seedling blight. Oreg.

Gibberella saubinetii (Mont.) Sacc., scab. General east of Great Plains; most prevalent in Corn Belt.

Helminthosporium sativum Pam., King, and Bak., foot-rot. Widely distributed, especially in upper Miss. Valley States.

Leptosphaeria tritici (S. Garovaglio) Pass., leaf-spot. N. Y., Iowa, Nebr.

Marasmius tritici P. A. Young, root-rot. Ill.

Mycosphaerella tulasnei Jacz. (*Cladosporium herbarum* (Pers.) Lk.; *Hormodendrum cladosporioides* Sacc.), sooty mold. N. Y., N. J., W. Va., Ohio, Mich., Pacific coast.

Ophiobolus graminis Sacc., take-all. N. Y., Md., Va., N. C., Tenn., Ark., Ind., Kans., Wash., Oreg., Calif.

Podosporiella verticillata P. J. O'Gara, seedling blight. Utah.

Puccinia glumarum (Schm.) Eriks. and Henn., stripe-rust. West of Great Plains.

P. graminis Pers., stem-rust. General; most prevalent in the spring-wheat area.

P. tritici Eriks., leaf-rust. General, especially prevalent in South.

Sclerospora macrospora Sacc., downy mildew. Del., Ky., Tenn., Calif.

Sclerotium rhizodes Auer., *Sclerotium* blight. Idaho, Mont., Wash.

Septoria nodorum Berk., glume-blotch. General; most prevalent in South and Ohio Valley States.

S. tritici Desm., speckled leaf-blotch. General.

Tilletia laevis Kuehn, bunt. General.

T. tritici (Bjerk.) Wint., bunt. Mostly west of Miss. River. Serious in Pacific Northwest.

TRITICUM—Continued.

T. aestivum, wheat—Continued.

Tylenchus tritici (Stein.) Bast., nematode. N. Y., Va., W. Va., N. C., S. C., Ga., Calif.

Urocystis tritici Koern., flag-smut. Ill., Mo., Kans.

Ustilago tritici (Pers.) Rostr., loose-smut. Widespread.

Wojnowicia graminis (McAlp.) Sacc. and D. Sacc., foot-rot. N. Y., Kans., Ark., Oreg.

Black-point, associated with *Helminthosporium* spp. N. Dak.

Mosaic, undet. Nebr., Ind., Ill.

Rosette, undet. Ill., Ind.

Sterility, undet. Minn., N. Dak., Kans., Mo., Tex.

Stripe, undet. N. Y., Pa., Ill.

**T. dicoccum*, emmer.

Ophiobolus graminis Sacc., take-all. N. Y.

Puccinia glumarum (Schm.) Eriks. and Henn., stripe-rust. Idaho.

P. graminis Pers., stem-rust. Tex., Okla., Mo., Wash.

P. triticea Eriks., leaf-rust. Ga., Tex.

Septoria nodorum Berk., glume-blotch. Md.

Tilletia tritici (Bjerk.) Wint., bunt. N. Dak.

Tylenchus tritici (Stein.) Bast., nematode. Va.

**T. spelta*, spelt.

Colletotrichum graminicolum (Ces.) Wils., anthracnose. Ill.

Gibberella saubinetii (Mont.) Sacc., scab. Md., Ill., Minn.

Ophiobolus graminis Sacc., take-all. N. Y.

Puccinia glumarum (Schm.) Eriks. and Henn., stripe-rust. Idaho.

P. graminis Pers., stem-rust. Mo., Minn., N. Dak., Kans., Wyo., Wash.

P. triticea Eriks., leaf-rust. Wis., Minn., Tex.

Tilletia tritici (Bjerk.) Wint., bunt. N. Dak.

Tylenchus tritici (Stein.) Bast., nematode. Va.

Ustilago tritici (Pers.) Rostr., loose-smut. N. Y.

TROPAEOLUM.

T. majus, nasturtium.

Alternaria sp. (associated with *Pleospora*), black-mold. N. J.

Bacterium aptatum N. Brown and Jamieson (*Phytophthora aptatum* (Brown and Jamieson) Com. S. A. B.), leaf-spot. N. J., Miss.

B. solanacearum EFS. (*Phytophthora solanacearum* (EFS.) Com. S. A. B.), bacterial wilt. Md., Va., Fla.

Cercospora tropaeoli Atk., leaf-spot. Ala.

Heterodera radiculicola (Greef) Muell., root-knot. N. J.

Pleospora tropaeoli Hals., leaf-spot. Ohio.

Puccinia sarcobati (Pk.) Bethel, rust. Utah (a single report).

TSUGA.

T. canadensis, Canada hemlock.

Armillaria mellea (Vahl) Quel., root-rot. Pa.

Fomes pinicola Fr., brown crumbly rot. General in Northeastern States.

F. roseus (Alb. and Schw.) Fr., brown pocket rot. N. Y.

Hymenochaete agglutinans Ell. Pa.

Hypoderma brachysporum (Rostr.) Tub., leaf-cast. Vt.

Keithia tsugae (Farl.) E. J. Durand, leaf-blight. N. H., Wis.

Melampsora abietis-canadensis (Farl.) C. A. Ludwig, rust. New England; N. Y., and westward to northern Wis.

M. farlowii (Arth.) Davis, rust. New England.

Polyporus borealis Fr., cuboidal rot. N. Y.

Pucciniastrum hydrangeae (Berk. and Curt.) Arth., rust. Va.

P. myrtilli (Schum.) Arth., rust. New England, N. Y., Pa., Va., W. Va., N. C., Ind.

Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot.

T. heterophylla, western hemlock.

Botrytis sp., seedling twig-blight. Idaho.

Caecoma dubium C. A. Ludwig, rust. Wash., Oreg.

Dimerosporium tsugae Dearn., sooty mold. Wash., Oreg.

TSUGA—Continued.

T. heterophylla, western hemlock—Continued.

Echinodontium tinctorium Ell. and Ev., brown stringy rot. Mont., Idaho, Oreg.

Fomes annosus Fr., white spongy rot. Wash.

F. officinalis Fr., reddish brown-rot.

F. pinicola Fr., brown crumbly rot. General with host.

Polyporus dryadeus (Pers.) Fr., white root-rot. Pacific Northwest.

P. schweinitzii Fr., red-brown butt-rot. Oreg.

P. sulphureus (Bull.) Fr., brown-checked wood-rot.

Razoumofskya tsugensis C. O. Rosendahl, hemlock mistletoe. Pacific Northwest; Alaska.

Rhizina inflata (Schaeff.) Sacc., seedling blight. Idaho.

Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Wash.

Uredo holwayi Arth., needle rust. Mont., Idaho.

T. mertensiana, mountain hemlock.

Echinodontium tinctorium Ell. and Ev., brown stringy rot. Wash.

Fomes officinalis Fr., brown heartwood rot. Idaho.

F. pinicola Fr., brown crumbly rot. General with host.

Herpotrichia nigra Hartig, matting. Oreg.

Polyporus schweinitzii Fr., red-brown butt-rot.

Pythium debaryanum Hesse, damping-off.

Razoumofskya tsugensis C. O. Rosendahl, mistletoe. Idaho, Wash.

Trametes pini (Brot.) Fr. (*Fomes pini* (Thore) Karst.), honeycomb rot. Oreg.

TULIPA.

Tulipa sp., tulip.

Botrytis tulipae (Lib.) E. F. Hopkins, botrytis blight. General.

Rhizoctonia tuliparum (Kleb.) Whet. and John M. Arthur, gray bulb-rot. N. Y.

Sclerotinia sp., white-rot. Ohio.

ULMUS.

Ulmus spp., elms.

Cercospora sphaeriaeformis Cke., leaf-spot. La.

Coniothyrium ulmi Tharp. Tex.

Cylindrosporium tenuisporum Heald and Wolf, leaf-spot. Tex.

Fomes applanatus (Pers.) Wallr. (*Ganoderma applanatum* (Pers.) Pat.), white-streaked rot. Northeastern States.

F. igniarius (L.) Fr., white heart-rot. Northeastern States.

Gloeosporium inconspicuum Cav., leaf-spot. Mass., Ohio, Mich., Minn., Okla.

Gnomonia ulmea (Sacc.) Thuem., black-spot. General east of Great Plains.

Heterodera radicola (Greef) Muell., root-knot. Calif.

Microsphaera ulni (Wallr.) Wint., powdery mildew. Ill., Iowa.

Mycosphaerella ulmi Kleb., leaf-spot. East of Great Plains.

Nectria cinnabarina (Tode) Fr., canker. Ohio.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex., Ariz.

Phoradendron flavescens (Pursh) Nutt., mistletoe. Ind., Southwest.

Phyllactinia corylea (Pers.) Karst., powdery mildew. S. C., Ala., Miss., Tex., Ill.

Phyllosticta confertissima Ell. and Ev., leaf-spot. Pa., Kans.

P. erratica Ell. and Ev., leaf-spot. Ala.

P. melaleuca Ell. and Ev., leaf-spot. Ala., Miss., Okla., Mich., Iowa, Mo., Nebr.

Pleurotus ulmarius Bull., white sap-rot. Widespread east of Great Plains.

Poronidulus conchifer (Schw.) Murr., twig-blight. Ohio.

Sphaeropsis ulmicola Ell. and Ev., canker. Pa., Md., Ohio, Wis.

Taphrina ulmi (Fekl.) Johans., leaf blister. Mass.

Uncinula macrospora Pk., powdery mildew. General east of Great Plains.

VACCINIUM.

Vaccinium spp., blueberry.

Bacterium tumefaciens EFS. and Town. (*Phytoplasma tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. Miss.

Calyptospora columnaris (Alb. and Schw.) Kuehn, stem-rust. Frequent in the Northern States and Pacific coast.

Exobasidium vaccinii (Fekl.) Wor., red leaf-spot. General.

Microsphaera alni vaccinii (Schw.) Salm., powdery mildew. North Atlantic States; Ohio, Ill., Wash.

Pucciniastrum myrtilli (Schum.) Arth., leaf-rust. General, especially in Northern States.

Ramularia effusa Pk., leaf-spot. Wis.

Rhytisma vaccinii (Schw.) Fr., tar-spot. Widespread.

Sclerotinia vaccinii-corymbosi J. M. Reade, fruit-rot, twig-blight. North Atlantic States; Ind., Mich.

Septoria albopunctata Cke., leaf-spot. S. C.

S. difformis Cke. and Pk., leaf-spot. N. Y.

Venturia compacta Pk. Me., Mass., La., Wis.

V. macrocarpon, cranberry.

Acanthorhynchus vaccinii Shear, blotch-rot. Me., Mass., N. J., W. Va., N. C., Mich., Wis., Wash.

Botrytis cinerea Auct., gray-mold rot. Wash.

Ceuthospora lunata Shear, black-rot. Mass., N. J., Wash.

Exobasidium oxycocci Rostr., rose-bloom. Mass., Wash., Oreg.

E. vaccinii (Fekl.) Wor., red leaf-spot. Atlantic Coast States; W. Va., Wis., Minn., Wash.

Fusicoccum putrefaciens Shear, end-rot. Me., Mass., N. J., N. C., Mich., Wis., Wash., Oreg.

Glomerella cingulata vaccinii Shear, bitter-rot. Mass., N. J., N. C., Wis., Wash., Oreg.

Guignardia vaccinii Shear, early-rot. Mass., Conn., N. Y., N. J., Va., N. C., Wis., Wash.

Leptothyrium oxycocci Shear. Mass.

Microsphaera alni vaccinii (Schw.) Salm., powdery mildew. N. J., Ohio.

Phomopsis sp., fruit-rot. Wash.

Phyllosticta putrefaciens Shear, rot. N. J.

Pucciniastrum myrtilli (Schum.) Arth., rust. Wash.

Sclerotinia oxycocci Wor., hard-rot and tip-blight. Wis., Wash.

Synchytrium vaccinii F. Thomas, red-gall. N. J.

Venturia compacta Pk. Me., Mass., N. Y., N. J., Wis., Wash.

False-blossom, undet. Mass., N. J., Wis., Wash., Oreg.

VIBURNUM.

Viburnum spp., viburnum.

Botrytis sp., blight. Wash.

Cercospora varia Pk., leaf-spot. Mass., N. Y., Ind., Ill., Iowa, Colo.

Coleosporium viburni Arth., rust. Md., Wis., Iowa.

Corticium stevensii Burt (*Hypochnus ochroleucus* F. Noack), leaf-blight. N. C.

Helminthosporium beaumontii Sacc., leaf-spot. Ala.

Hendersonia foliorum Fekl., leaf-spot. Tex.

Hymenochaete agglutinans Ell. Pa.

Microsphaera alni (Wallr.) Wint., powdery mildew. Widespread.

Phyllosticta lantanoides Pk., leaf-spot. N. Y.

P. punctata Ell. and Dearn., leaf-spot. Wis.

P. tineola Sacc., leaf-spot. Wis.

Plasmopara viburni Pk., downy mildew. N. Y., W. Va., Ga., Ala., Minn.

Puccinia linkii J. F. Klotzsch, rust. Mich., Idaho, Mont., Wash.

VICIA.

V. faba, broadbean.

Ascochyta pisi Lib. (*Mycosphaerella pinodes* (Berk. and Blox.) R. E. Stone), leaf-spot. Calif.

Corticium vagum Berk. and Curt., stem-rot. Mich., Wash.

Fusarium culmorum W. G. Smith, root-rot. Utah.

F. trichothecioides Woll., root-rot. Utah.

Heterodera radicolica (Greef) Muell., root-knot. Calif.

Uromyces fabae (Pers.) D By., rust. Ala., La.

VICIA—Continued.

V. sativa, common vetch.

Ascochyta pisi Lib. (*Mycosphaerella pinodes* (Berk. and Blox.) R. E. Stone), leaf-spot. Miss.

Cercospora viciae Ell. and Holw., leaf-spot. Iowa, Kans.

Peronospora viciae D By., downy mildew. Va., Miss.

V. villosa, hairy vetch, winter vetch.

Ascochyta pisi Lib. (*Mycosphaerella pinodes* (Berk. and Blox.) R. E. Stone), leaf-spot. Occurs scattering in widely separated States.

Heterodera radiculicola (Greef) Muell., root-knot. Calif.

Pseudopeziza medicaginis (Lib.) Sacc., leaf-spot. Miss.

Sclerotinia sp. Calif.

VIGNA.

V. catjang, catjang-pea.

Cercospora vignae Ell. and Ev., leaf-spot. Del., Fla., Tenn., La., Ark.

Fusarium vasinfectum tracheiphilum EFS., wilt. Widespread in Southern States.

Heterodera radiculicola (Greef) Muell., root-knot.

V. sinensis, cowpea.

Amerosporium oeconomicum Ell. and Tr., leaf-spot. Southern States; Del., Conn.

Bacillus lathyri Manns and Taub., streak. Del.

Bacterium phaseoli EFS. (*Phytomonas phaseoli* (EFS.) Com. S. A. B.), bacterial blight. Okla.

B. vignae M. W. Gardner and J. B. Kendrick (*Phytomonas vignae* (Gardner, and Kendrick) Com. S. A. B.), bacterial spot. Ind., Kans.

Cercospora cruenta Sacc., leaf-spot. General east of Great Plains, especially in South.

C. dolichii Ell. and Ev., leaf-spot. Conn., N. J., Ala., Miss.

*C. vignae*³ M. Raciborski, leaf-spot. Tex.

Cladosporium vignae M. W. Gardner, leaf-spot. Ind.

Colletotrichum lindemuthianum (Sacc. and Magn.) Briosi and Cav. (*Glomerella lindemuthiana* Shear), anthracnose. Del., Fla.

Corticium vagum Berk. and Curt., stem-rot. Southern States.

Erysiphe polygoni DC., powdery mildew. Southern States.

Fusarium martii phaseoli Burk., root-rot.

F. vasinfectum tracheiphilum EFS., wilt. Southern States; Ind., Kans., Calif.

Heterodera radiculicola (Greef) Muell., root-knot. Southern States.

Microsphaera alni (Wallr.) Wint., powdery mildew. Ga.

M. euphorbiae Berk. and Curt., powdery mildew. S. C., Fla.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Okla., Tex.

Phyllosticta phaseolina Sacc., leaf-spot. Del., Va., N. C., S. C., Ind., Ill., Mich.

Sclerotium rolfsii Sacc., stem-rot. N. C.

Thielavia basicola (Berk. and Br.) Zopf, black root-rot. Md., D. C.

Uromyces vignae A. Barclay, rust. Md., Va., Fla., Ala., Tex., Ind., Iowa, Mo., Calif.

Mosaic, undet. Ark., Okla., Ind.

VIOLA.

V. odorata, sweet violet.

Alternaria violae Gall. and P. H. Dorsett, leaf-spot. Conn., Tex.

Cercospora violae Sacc., leaf-spot. Conn., Pa., Ala., Tex.

Colletotrichum violae-tricoloris R. E. Smith, leaf-spot. Conn.

Corticium vagum Berk. and Curt., stem-rot. N. Y.

Gloeosporium violae Berk. and Br., leaf-spot. Miss.

Heterodera radiculicola (Greef) Muell., root-knot.

Phyllosticta violae Desm., leaf-spot. Conn., Mass.

Sclerotium rolfsii Sacc., stem-rot. Md.

Thielavia basicola (Berk. and Br.) Zopf, black root-rot. Mass., Conn., N. Y., Md., D. C.

³ *Cercospora vignae* Rac. was described from Java in 1898; *Cercospora vignae* Ell. and Ev. from Louisiana in 1886. Heald and Wolf reported *C. vignae* Rac. from Texas in 1912.

VIOLA—Continued.

V. tricolor and *V. cornuta*, pansy.

Alternaria violae Gall. and P. H. Dorsett, leaf-spot. N. Y., Pa.

Cercospora violae Sacc., leaf-spot. N. Y.

Colletotrichum sp., anthracnose. Fla.

C. violae-tricoloris R. E. Smith, leaf-spot. Mass., N. Y., N. J.

Corticium vagum Berk. and Curt., root-rot. Del., Minn.

Fusarium sp., wilt. Ohio, Mich.

F. violae Wolf, stem-rot, root-rot. Nebr.

Heterodera radiculicola (Greef) Muell., root-knot. N. Y.

Peronospora violae D By., downy mildew. Miss., Ill., Nebr.

Puccinia ellisiana Thuem., rust. Kans., Nebr.

P. violae (Schum.) DC., rust. S. C., N. Dak.

Pythium debaryanum Hesse, damping-off. Conn.

Ramularia sp., leaf-spot. Wash.

Sphaerotheca humuli fuliginea (Schl.) Salm., powdery mildew. Wash.

Thielavia basicola (Berk. and Br.) Zopf, black root-rot. Conn.

VITIS.

Vitis sp., grape.

Armillaria mellea (Vahl) Quel., root-rot. Wash.

Bacterium tumefaciens EFS. and Town. (*Phytomonas tumefaciens* (EFS. and Town.) Com. S. A. B.), crown-gall. General.

Botrytis cinerea Auct., gray-mold rot. General on fruit.

Clitocybe monadelpha (Morgan) Sacc., root-rot. Mo., Ark., Okla., Tex.

Cryptosporella viticola (Reddick) Shear, dead-arm. General east of Great Plains.

Glomerella cingulata (Ston.) Spauld. and Schrenk, ripe-rot. Widespread east of Miss. River.

Guignardia bidwellii (Ell.) Viala and Ravaz, black-rot. General east of Great Plains.

Heterodera radiculicola (Greef) Muell., root-knot. Calif.

Isariopsis clavispora (Berk. and Curt.) Sacc. (*Cercospora viticola* (Ces.) Sacc.), leaf-spot. Southern States; Mich.

Melanconium fuligineum (Scrib. and Viala) Cav., bitter-rot. Widespread east of Great Plains, especially South.

Ozonium omnivorum Shear (*Phymatotrichum omnivorum* (Shear) Dug.), root-rot. Tex.

Pestalozzia unicola Speg., leaf-blight. Fla., Miss., Tenn., Ohio, Ind., Nebr.

Phakopsora vitis (Thuem.) Syd., rust. S. C., Fla.

Plasmopara viticola (Berk. and Curt.) Berl. and DeToni, downy mildew. General east of Great Plains; Calif.

Rhytisma vitis Schw., tar-spot. N. Y., Pa.

Roesleria hypogaea Thuem. and Pass., root-rot. N. Y., Pa., Ohio, Ill., Mo., N. Mex.

Septoria ampelina Berk. and Curt., leaf-spot. N. Y., Va., Tex., Kans.

Sphaceloma ampelinum D By. (*Gloeosporium ampelophagum* (Pass.) Sacc.), anthracnose. General but erratic east of Great Plains.

Uncinula necator (Schw.) Burr., powdery mildew. General.

California vine disease, undet. Calif.

Chlorosis. Eastern States.

Little-leaf, undet. Calif.

Spanish measles, undet. Calif.

WISTERIA.

Wisteria sp., wisteria.

Phyllosticta wistariae Sacc., leaf-spot. N. J., Tex.

Septoria wistariae Tharp, leaf-spot. Tex.

YUCCA.

Y. filamentosa, common yucca.

Cercospora concentrica Cke. and Ell., leaf-spot. N. J.

Coniothyrium concentricum (Desm.) Sacc., leaf-spot. Ala., N. Y., Idaho.

Gloeosporium yuccaegenum Ell. and Ev., leaf-spot. Mo.

Leptosphaeria filamentosa Ell. and Ev., leaf-spot. N. J.

ZEA.

Z. mays, maize, Indian corn.

- Aplanobacter stewartii* (EFS.) McC. (*Phytomonas stewartii* (EFS.) Com. S. A. B.), bacterial wilt. General east of Great Plains; Southwest; especially on certain varieties of sweet corn.
- Aspergillus* sp., ear-mold. La., Tex., Calif.
- Bacterium dissolvens* H. R. Rosen, bacterial root-rot. Miss., La., Ark., Ill.
- Basisporium gallarum* M. Molliard (*Coniosporium gecevi* Bub.), cob-rot. Conn., Ohio, Iowa, Kans.
- Cephalosporium acremonium* Cda., black bundle. East of Great Plains, Calif.
- Cercospora sorghi* Ell. and Ev., leaf-spot. S. C., Ala., La.
- Colletotrichum graminicolum* (Ces.) Wils., anthracnose. Conn., N. J., S. C., W. Va.
- Diplodia zeae* Lév., dry-rot. General where corn is grown.
- Fusarium culmorum leteius* Sherb., seedling blight. Oreg.
- F. moniliforme* Sheld., ear-rot and seed-rot. General.
- Gibberella saubinetii* (Mont.) Sacc., root-rot. General east of Great Plains.
- Helminthosporium* sp., stalk canker. Ga.
- H. turcicum* Pass., leaf-blight. General east of Great Plains.
- Heterodera radiculicola* (Greef) Muell., root-knot. S. C., Ala.
- Ophiobolus heterostrophus* Drechs., leaf-spot. Fla.
- Phyllosticta* sp., leaf-spot. Fla.
- Phyoderma zeae-maydis* Shaw, brown-spot. N. J., Md. to Nebr. and south. Most important in South.
- Phytophthora cactorum* (Leb. and Cohn) Schroet., root-rot. Conn.
- Puccinia sorghi* Schw., rust. General, especially east of Rocky Mts.
- Pythium* sp., root-rot. Fla.
- Rhizoctonia* sp., root-rot. Ohio, Minn., Wash.
- Rhizopus nigricans* Ehr., seed-blight.
- Sclerospora graminicola* (Sacc.) Schroet., downy mildew. Iowa.
- Sclerotium rolfsii* Sacc., stem-rot. Fla., Ala., Tex.
- Sorosporium reilianum* (Kuehn) McAlp., head-smut. N. Y., Ohio, Kans., Idaho, Wash., Calif.
- Ustilago zeae* (Beckm.) Ung., smut. General.
- Mosaic, undet. Ga., Ark., La.
- Sheath-spot, *Fusaria* and other fungi. General.

ZINNIA.

Z. elegans, zinnia.

- Cercospora atricincta* Heald and Wolf, leaf-spot. Fla., Tex.
- C. zinniae* Ell. and Mart., leaf-spot. Fla., Ala.
- Choanephora* sp., blossom blight. Fla.
- Erysiphe cichoracearum* DC., powdery mildew. Conn., W. Va., Ill., Tex.
- Fusarium* sp., wilt. Wash.
- Paratylenchus nanus* Cobb, nematode. Md.

INDEX TO COMMON NAMES OF HOSTS

	Page		Page
Acacia.....	13	Cabbage.....	27-28
Alaska cedar.....	34	Calendula.....	30
Alder.....	18	California poppy.....	45
Alfalfa.....	64-65	Callistephus.....	30
Almond.....	20	Camphor-tree.....	35
Amaryllis. See Belladonna-lily.		Cane.....	23
American beautyberry.....	30	Canna.....	31
Ampelopsis.....	19	Cantaloupe.....	40
Apple.....	62-64	Cape-jasmine.....	48
Apricot.....	84	Carnation.....	44
Arborvitae.....	100	Carolina-jessamine.....	48
Arrowwood. See Viburnum.		Carpet bentgrass.....	17
Artichoke, globe.....	42	Carrot.....	43
Jerusalem.....	50	Cassava.....	64
Ash.....	47-48	Castor-bean.....	91
Asparagus.....	24	Catalpa.....	32-33
Aspen.....	80-81	Catjang-pea.....	106
Aster. See Callistephus.		Cauliflower.....	28
Avocado.....	70	Cedar, Alaska.....	34
Azalea.....	24	incense.....	59
		red.....	56-57
Baldcypress.....	100	white.....	54
Balsam. See Fir.		Celeriac.....	22
Bamboo.....	23, 24-73	Celery.....	22
Japanese timber.....	73	Chard, Swiss.....	26
Banana.....	66	Chayote.....	34
Barberry.....	25	Cherimoya.....	21
Barley.....	52-53	Cherry.....	81-82
Basswood.....	100	Chess.....	30
Bayberry.....	66	Chestnut.....	32
Bay, red.....	70	Chicory.....	35
sweet.....	62	China-aster.....	63
Beachgrass.....	19	Chinaberry.....	50
Bean.....	70-71	Chinese cabbage.....	29
broad.....	105	Chinquapin.....	32
Scarlet Runner.....	71	Chives.....	18
soy.....	96	Chokeberry.....	23
velvet.....	99	Chokecherry.....	83-84
Bearberry.....	23	Christmasberry.....	72
Beautyberry, American.....	30	Christmas-rose.....	50
Beech.....	46	Chrysanthemum.....	34
blue.....	32	Citrus.....	35
Beet.....	25-26	Clematis.....	37
Begonia.....	24-25	Clover.....	101-102
Belladonna-lily.....	19	Japan.....	58
Bentgrass.....	17	southern bur.....	64
Bermuda grass.....	31	white sweet.....	65
Big tree.....	99	Coconut.....	38
Birch.....	26-27	Coffee.....	38
Bittersweet.....	33	Coffeetree, Kentucky.....	49
Blackberry.....	92-93	Collards.....	28
Logan.....	93	Columbine.....	22
Black locust.....	91	Coneflower.....	94
Bladdernut.....	99	Coralberry.....	99
Blueberry.....	105	Coralree.....	45
Blueflag iris.....	55	Corn, Indian.....	108
Bluegrass.....	80	Cosmos.....	39
Boston ivy.....	20	Cotton.....	49
Box.....	30	Cottonwood.....	80-81
Boxelder.....	14	Couchgrass.....	16
Broadbean.....	105	Cowpea.....	106
Bromegrass, Hungarian.....	29	Crab apple.....	62
Broomcorn.....	52	Cranberry.....	105
Broomcorn millet.....	69	Crapemyrtle.....	57
Brussels sprouts.....	29	Crossvine.....	27
Buckeye.....	16	Croton.....	39
Buckthorn.....	89	Cucumber.....	40
Buckwheat.....	46	Cucumbertree.....	62
Buffaloberry.....	96	Currant.....	90-91
Buffalo grass.....	30	Cyees.....	41
Bugloss.....	21	Cyclamen.....	41
Bumelia.....	30	Cypress.....	41
Bur clover, southern.....	64	bald.....	100
Burningbush.....	45	Nootka.....	34
Bush-honeysuckle.....	44		
Butterflybush.....	30	Dahlia.....	42-43
Butterflyweed.....	23	Dasheen.....	38
Butternut.....	55	Date palm.....	72
Buttonbush.....	33	Daylily.....	50

	Page		Page
Deutzia.....	43	Jetbead.....	90
Devils-walkingstick.....	23	Johnson grass.....	52
Dewberry.....	93	Jonquil.....	67
Dill.....	21	Judas-tree.....	33
Dogwood.....	38	Juneberry.....	19
Douglas fir.....	86	Juniper.....	56
Easter lily.....	59	Kale.....	28
Eggplant.....	96-97	Kentucky coffeetree.....	49
Elderberry.....	95	Kerria.....	57
Elm.....	104	Kohlrabi.....	29
Emmer.....	103	Larch.....	58
Endive.....	35	Larkspur.....	43
Eucalyptus.....	45	Laurel, mountain.....	57
Fescue.....	46	Lemon.....	36
Fig.....	46	Lettuce.....	57
Filbert.....	39	Lilac.....	99-100
Fir.....	11-13	Lily.....	69
Douglas.....	86	Lima bean.....	71
Firethorn, scarlet.....	86	Lime.....	35
Flax.....	59	Linden.....	100
Flowering dogwood.....	38	Loblolly-bay.....	49
Forsythia.....	47	Locust, black.....	91
Four-o'clock.....	66	Logan blackberry.....	93
Foxglove.....	44	Loquat.....	44
Fringetree.....	34	Lupine.....	60
Gaillardia.....	48	Madrone.....	23
Garlic.....	18	Magnolia.....	62
Geranium.....	70	Maize.....	108
Ginseng.....	69	Maleberry.....	61
Gladiolus.....	48	Mangel-wurzel.....	26
Globe artichoke.....	42	Mango.....	64
Goldenbell.....	47	Manzanita.....	23
Goldenglow.....	94	Maple.....	18-15
Goldenseal.....	53	Marigold.....	100
Gooseberry.....	91	Meadowsweet.....	47
Gourd.....	40	Mesquite.....	81
Grape.....	107	Mignonette.....	89
Oregon.....	62	Millet.....	34
Grapefruit.....	36	broomcorn.....	69
Grass, Bermuda.....	31	proso.....	69
buffalo.....	30	Mockorange.....	72
Johnson.....	52	Monkshood.....	16
orchard.....	42	Moonseed.....	66
Sudan.....	52	Moosewood.....	14
Guava.....	86	Morning-glory.....	54
Gumbo.....	50-51	Mountain-ash.....	98
Hackberry.....	33	Mountain-laurel.....	57
Hawthorn.....	39	Mulberry.....	66
Hazelnut.....	39	Muskmelon.....	40
Hemlock.....	103-104	Narcissus.....	67
Hemp.....	31	Nasturtium.....	103
Hickory.....	51	Nectarine.....	21
Holly.....	53-54	Ninebark.....	73
Hollygrape.....	62	Nootka cypress.....	34
Hollyhock.....	18-19	Oak.....	87-88
Honeylocust.....	49	Oats.....	24
Honeysuckle.....	60	Oceanspray.....	52
bush.....	44	Okra.....	50-51
Hop.....	53	Oleander.....	67
Hophornbeam.....	68	Olive.....	67-68
Hoptree.....	86	Russian.....	44
Hornbeam.....	31-32	Onion.....	17
hop.....	68	Orange.....	35-37
Horsechestnut.....	16	king.....	36
Horseradish.....	88	Osage.....	61
Huckleberry.....	48	Satsuma.....	36-37
Hungarian bromegrass.....	29	Seville.....	35-36
Hyacinth.....	53	Orchard grass.....	42
Hydrangea.....	53	Oregon grape.....	62
Incense-cedar.....	59	Osage-orange.....	61
India rubber tree.....	47	Osmanthus.....	68
Indigobush.....	19	Pagoda tree.....	98
Iris.....	55	Pansy.....	107
Ivy, Boston.....	20	Papaw.....	23
English.....	50	Papaya.....	31
Japan clover.....	58	Parkinsonia.....	69
Japanese barberry.....	25	Parsley.....	70
Japanese creeper.....	70	Parsnip.....	69
Japanese timber bamboo.....	73	Pea.....	79
Jasmine, Cape.....	48	sweet.....	58
Jerusalem-artichoke.....	50	Peach.....	20-21
Jessamine.....	48	Peanut.....	22-23

	Page		Page
Pear.....	86-87	Soapberry.....	95
Pecan.....	51	Sorgo.....	52
Peony.....	68	Sorghum.....	52
Pepper.....	31	Sour gum.....	67
Pepperidge.....	67	Sourwood.....	68
Peppermint.....	66	Southern bur clover.....	64
Peppertree.....	95	Soybean.....	96
Peppervine.....	19	Spearmint.....	66
Persimmon.....	44	Spelt.....	103
Petsai cabbage.....	29	Spicebush.....	25
Petunia.....	70	Spinach.....	98-99
Phlox.....	72	Spirea.....	99
Pigeon-pea.....	30	Spruce.....	73-74
Pine.....	74-79	Squash.....	40
Pineapple.....	21	Strawberry.....	47
Planetree.....	80	Sudan grass.....	52
Pleurisy root.....	23	Sugar-apple.....	21
Plum.....	85-86	Sugar beet.....	25-26
Poinciana.....	80	Sugarberry.....	33
Pomegranate.....	86	Sugarcane.....	94
Poplar.....	80-81	Sugar sorghum.....	52
Poppy, California.....	45	Sumac.....	90
Potato.....	97-98	Summer squash.....	40
sweet.....	54	Sunflower.....	50
Primrose.....	81	Sweetbay.....	62
Princess tree.....	70	Sweetclover, white.....	65
Privet.....	59	Sweetfern.....	38
Proso millet.....	69	Sweetgum.....	59
Prune.....	84	Sweet pea.....	58
Pumpkin.....	41	Sweetpotato.....	54
Quackgrass.....	16	Sweet-william.....	43
Quince.....	41-42	Swiss chard.....	25
Radish.....	89	Sycamore.....	80
Raspberry.....	93-94	Syringa.....	72
Redbay.....	70	Tamarack.....	58
Redbud.....	33	Tangerine.....	36
Redcedar.....	56-57	Teosinte.....	45
Redpepper.....	31	Timothy.....	72
Redtop.....	17	Tobacco.....	67
Redwood.....	96	Tomato.....	60-61
Rhododendron.....	89-90	Toyon. See Christmasberry.....	
Rhodora.....	90	Treemallow.....	58
Rhubarb.....	89	Tree-of-heaven.....	17
Rice.....	63	Trumpet creeper.....	27
Rose.....	91-92	Tulip.....	104
Rose-of-Sharon.....	51	Tuliptree.....	59-60
Rubbertree, India.....	47	Tupelo.....	67
Russian-olive.....	44	Turnip.....	29
Rutabaga.....	27	Umbrella-tree, Texas.....	65
Rye.....	95-96	Vegetable-oyster.....	100-101
Ryegrass.....	60	Velvetbean.....	99
Sago cycas.....	41	Vetch.....	106
Salal.....	48	Viburnum.....	105
Salsify.....	100-101	Violet.....	106
Sapodilla.....	15	Virginia creeper.....	20
Sapote.....	15	Virgins-bower.....	37
Saskatoon.....	19	Wahoo.....	45
Sassafras.....	95	Walnut.....	55-56
Satsuma orange.....	36-37	Watermelon.....	35
Scarlet Runner bean.....	71	Waxmyrtle.....	66
Serviceberry.....	19	Wheat.....	102-103
Seville orange.....	35-36	Whitecedar.....	34
Shadblow.....	19	Willow.....	94
Shadbush.....	19	Winterberry.....	53-54
Shallot.....	17	Wisteria.....	107
Silktassel-bush.....	48	Witch-hazel.....	49-50
Silverberry.....	44	Wormseed.....	34
Smoketree.....	90	Yellow poplar. See Tuliptree.....	
Snakeroot, Virginia.....	23	Yucca.....	107
Snapdragon.....	22	Yucca.....	107
Snowball. See Viburnum.....		Zinnia.....	108
Snowberry.....	99		
Snow-on-the-mountain.....	45		

ORGANIZATION OF THE UNITED STATES DEPARTMENT OF AGRICULTURE

April 1, 1926

<i>Secretary of Agriculture</i>	W. M. JARDINE.
<i>Assistant Secretary</i>	R. W. DUNLAP.
<i>Director of Scientific Work</i>	_____
<i>Director of Regulatory Work</i>	WALTER G. CAMPBELL.
<i>Director of Extension Work</i>	C. W. WARBURTON.
<i>Director of Information</i>	NELSON ANTRIM CRAWFORD.
<i>Director of Personnel and Business Administration</i>	W. W. STOCKBERGER.
<i>Solicitor</i>	R. W. WILLIAMS.
<i>Weather Bureau</i>	CHARLES F. MARVIN, <i>Chief</i> .
<i>Bureau of Agricultural Economics</i>	THOMAS P. COOPER, <i>Chief</i> .
<i>Bureau of Animal Industry</i>	JOHN R. MOHLER, <i>Chief</i> .
<i>Bureau of Plant Industry</i>	WILLIAM A. TAYLOR, <i>Chief</i> .
<i>Forest Service</i>	W. B. GREELEY, <i>Chief</i> .
<i>Bureau of Chemistry</i>	C. A. BROWNE, <i>Chief</i> .
<i>Bureau of Soils</i>	MILTON WHITNEY, <i>Chief</i> .
<i>Bureau of Entomology</i>	L. O. HOWARD, <i>Chief</i> .
<i>Bureau of Biological Survey</i>	E. W. NELSON, <i>Chief</i> .
<i>Bureau of Public Roads</i>	THOMAS H. MACDONALD, <i>Chief</i> .
<i>Bureau of Home Economics</i>	LOUISE STANLEY, <i>Chief</i> .
<i>Bureau of Dairying</i>	C. W. LARSON, <i>Chief</i> .
<i>Fixed Nitrogen Research Laboratory</i>	F. G. COTTRELL, <i>Director</i> .
<i>Office of Experiment Stations</i>	E. W. ALLEN, <i>Chief</i> .
<i>Office of Cooperative Extension Work</i>	C. B. SMITH, <i>Chief</i> .
<i>Library</i>	CLARIBEL R. BARNETT, <i>Librarian</i> .
<i>Federal Horticultural Board</i>	C. L. MARLATT, <i>Chairman</i> .
<i>Insecticide and Fungicide Board</i>	J. K. HAYWOOD, <i>Chairman</i> .
<i>Packers and Stockyards Administration</i>	JOHN T. CAINE, <i>in Charge</i> .
<i>Grain Futures Administration</i>	J. W. T. DUVEL, <i>in Charge</i> .

This bulletin is a contribution from

<i>Bureau of Plant Industry</i>	WILLIAM A. TAYLOR, <i>Chief</i> .
<i>Office of Plant-Disease Survey and Pathological Collections.</i>	C. L. SHEAR, <i>Senior Pathologist in Charge</i> .

112

ADDITIONAL COPIES

OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.

AT

15 CENTS PER COPY



alpinus

Crowell & Lavalée (xvii, 70) p. 48

